

Structure-Based Identification and Neutralization Mechanism of Tyrosine Sulfate Mimetics That Inhibit HIV-1 Entry

Priyamvada Acharya,[†] Cajetan Dogo-Isonagie,^{‡,▽} Judith M. LaLonde,^{§,▽} Son N. Lam,^{‡,▽} George J. Leslie,^{¶,▽} Mark K. Louder,^{‡,▽} Leah L. Frye,^{||} Asim K. Debnath,[#] Jeremy R. Greenwood,^{||} Timothy S. Luongo,[†] Loïc Martin,[⊥] K. Shawn Watts,^{||} James A. Hoxie,[¶] John R. Mascola,[†] Carole A. Bewley,^{*,‡} and Peter D. Kwong^{*,†}

[†]Vaccine Research Center, National Institute of Allergy and Infectious Diseases and [‡]Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892, United States

[§]Department of Chemistry, Bryn Mawr College, Bryn Mawr, Pennsylvania 19010, United States

[¶]Department of Medicine, Hematology-Oncology Division, University of Pennsylvania, Philadelphia, Pennsylvania 19104, United States


^{||}Schrödinger, LLC, Portland, Oregon 97204, United States

[#]Laboratory of Molecular Modeling and Drug Design, Lindsley F. Kimball Research Institute, New York Blood Center, 310 E 67th Street, New York, New York 10021, United States

[⊥]CEA, iBiTecS, Service d'Ingénierie Moléculaire des Protéines, Gif sur Yvette F-91191, France

S Supporting Information

ABSTRACT: Tyrosine sulfate-mediated interactions play an important role in HIV-1 entry. After engaging the CD4 receptor at the cell surface, the HIV-1 gp120 glycoprotein binds to the CCR5 co-receptor *via* an interaction that requires two tyrosine sulfates, at positions 10 and 14 in the CCR5-N terminus. Building on previous structure determinations of this interaction, here we report the targeting of these tyrosine sulfate binding sites for drug design through *in silico* screening of small molecule libraries, identification of lead compounds, and characterization of biological activity. A class of tyrosine sulfate-mimicking small molecules containing a “phenyl sulfonate-linker-aromatic” motif was identified that specifically inhibited binding of gp120 to the CCR5-N terminus as well as to sulfated antibodies that recognize the co-receptor binding region on gp120. The most potent of these compounds bound gp120 with low micromolar affinity and its CD4-induced conformation with K_D 's as tight as ~ 50 nM. Neutralization experiments suggested the targeted site to be conformationally inaccessible prior to CD4 engagement. Primary HIV-1 isolates were weakly neutralized, preincubation with soluble CD4 enhanced neutralization, and engineered isolates with increased dependence on the N terminus of CCR5 or with reduced conformational barriers were neutralized with IC₅₀ values as low as ~ 1 μ M. These results reveal the potential of targeting the tyrosine sulfate interactions of HIV-1 and provide insight into how mechanistic barriers, evolved by HIV-1 to evade antibody recognition, also restrict small-molecule-mediated neutralization.



Human immunodeficiency virus type 1 (HIV-1) continues to be a major global health problem resulting in millions of deaths each year. Although a variety of anti-HIV-1 therapeutics have been developed, the emergence of HIV-1 strains resistant to existing therapies and the side effects of drug regimens make identification of new targets a continuing priority.¹ Entry inhibitors are an emerging class of therapeutics that interfere with attachment, fusion, or entry of HIV-1 into cells [reviewed in ref 2]. The trimeric HIV-1 envelope protein is composed of two glycoproteins: the attachment-mediating gp120 and the fusion-inducing gp41. The engagement of the cell surface CD4 receptor by gp120 induces structural rearrangements in the viral spike that lead to formation of a highly conserved co-receptor binding site

on gp120.^{3–5} Chemokine receptors, CCR5 or CXCR4, function as HIV-1 co-receptors [reviewed in ref 6]; upon binding to the CD4-induced conformation of gp120, they trigger additional conformational changes that lead to displacement of the fusogenic N terminus of gp41 into the target cell membrane and subsequent fusion of viral and cell membranes. Although each step in the entry pathway is a potential target for intervention, a number of hurdles, including difficulties with blocking protein–protein interactions, have complicated the development of entry

Received: February 24, 2011

Accepted: July 27, 2011

Published: July 27, 2011




Figure 1. Target site. A critical step of HIV entry involves interaction with sulfated tyrosine residues in the CCR5 co-receptor, which the antibody 412d mimics to neutralize HIV-1. (a) gp120-CCR5-N terminus interaction. HIV-1 gp120 (gray) interacts with its cell surface receptor CD4 (yellow) and co-receptor CCR5 (magenta). CCR5-N terminus binds in an α -helical conformation at the base of the V3 loop (orange) of gp120. Two sulfated tyrosine residues (stick representation) are critical for binding of the CCR5-N terminus to gp120. (b) Mimicry of gp120-CCR5-N terminus interaction by the CDR H3 loop of 412d antibody. The CDR H3 loop (red) of antibody 412d (green) is a functional mimic of the CCR5-N terminus with two sulfated tyrosine residues (sticks) dominating its interaction with gp120.

inhibitors. Nonetheless, the promise of entry inhibitors as antiretrovirals is demonstrated by two licensed drugs: fuzeon,⁷ a fusion inhibitor, and maraviroc,⁸ a CCR5 antagonist.

This study focuses on the interaction of the HIV-1 gp120 and the CCR5 co-receptor. The critical nature of the interaction with CCR5 is demonstrated by the resistance to HIV-1 infection of individuals homozygous for a naturally occurring variant of CCR5, in which 32 residues of the CCR5-N terminus are missing.⁹ CCR5 is an integral membrane protein with an extracellular N terminus, seven membrane-spanning helices, and three extracellular loops. Interactions between gp120 and CCR5 involve at least the N terminus and the second extracellular loop (ECL-2).^{10,11} We previously determined the structure of the CCR5-N terminus bound to gp120 by a combination of NMR, X-ray crystallography, and docking techniques.¹² The CCR5-N terminus adopts an α -helical conformation and binds to a highly conserved region at the base of the third variable loop (V3) on HIV-1 gp120 (Figure 1, panel a; Supplementary Figure 1). This interaction is dominated by two sulfated tyrosines at positions 10 and 14 of CCR5. Disruption of tyrosine sulfation results in loss of binding and viral entry. The importance of this pair of tyrosine sulfates is further demonstrated by the monoclonal antibody 412d; this antibody also has two tyrosine sulfates in its heavy chain third complementarity-determining region (CDR H3).^{13,14} The 412d CDR H3 region can substitute for the N terminus of CCR5 to make a functional, entry-competent co-receptor.¹⁵ Interestingly, the structure of antibody 412d in complex with HIV-1 gp120 reveals that the CDR H3 of 412d forms an extended conformation¹² (Figure 1, panel b). Despite different modes of binding, the involvement and importance of tyrosine sulfates is a common feature of both interactions.

Tyrosine sulfates at positions 10 and 14 in CCR5 and in the CDR H3 loop of antibody 412d occupy similar binding pockets on gp120.¹² Tyrosine sulfation is a post-translational modification that has been implicated in mediating high-affinity protein–protein interactions in a number of systems [reviewed in ref 16]; similar to phosphorylation,^{17–19} these post-translational modifications provide a handle by which molecular mimicry can be




Figure 2. Inhibitor identification flowchart. Starting from atomic level coordinates of gp120-CD4-CCR5-N terminus and gp120-CD4-412d Fab, two methods of *in silico* screening were used in combination with an ELISA-based screen to identify binding chemotypes. Selected compounds were characterized further for their gp120 binding and HIV-1 neutralization properties.

used to inhibit biology. Here we present a structure-guided search for small, drug-like molecules that inhibit viral entry by mimicking critical tyrosine sulfate mediated interactions made by the CCR5-N terminus. Structural information provided the basis for *in silico* screening of compound libraries. Results from the computational analyses were experimentally tested for their ability to bind gp120 as well as to compete with the binding of the CCR5-N terminus and antibodies that recognize a CD4-induced epitope on gp120. Molecules able to inhibit binding of the CCR5-N terminus to gp120 were further evaluated in viral entry inhibition assays to determine potency and mechanism of neutralization.

RESULTS AND DISCUSSION

***In Silico* Screening and CCR5-N Terminus Competition Assay.** To identify small molecule inhibitors of CCR5-N terminus binding to gp120, we coupled structure-based *in silico* screening to an ELISA-based assay (Figure 2 and Supplementary Figure 2). Two *in silico* screening protocols were used to screen two different small molecule libraries. In the first approach, the ZINC database V7²⁰ of drug-like compounds was searched using a consensus method involving the ROCS shape- and chemical




Figure 3. Tyrosine sulfate mimicry in inhibitor identification. The critical role of two N terminal tyrosine sulfate residues in CCR5 binding and their unique chemistry suggested the utility of the tyrosine sulfate binding sites as drug targets. (a) Tyrosine sulfate binding sites on gp120. Sequences of CCR5-N terminus and 412d CDR H3 loop are shown above with the tyrosine residues that are sulfated and critical for binding colored red. The CCR5-N terminus (magenta) and the CDR H3 loop (red) of 412d antibody interact with the base of the V3 loop (orange) of gp120 (gray). The region surrounding the tyrosine sulfate residues was targeted for small molecule screening. (b) Ligand-based probes for *in silico* screening. Structure-based probes (shown as sticks) centered around the tyrosine sulfate moieties were used for shape-based screening of the ZINC small molecule library. (c) Tyrosine sulfate analogues. The rank order of binding affinity of CCR5 peptides containing modified tyrosines at positions 10 and 14 are shown. Compounds containing phenyl sulfonate and phenyl sulfonamide groups were screened in this study.

similarity-matching routine²¹ coupled to docking analysis using the Gold algorithm²² in an effort to identify CCR5-N terminus mimetics. In parallel, the Glide molecular docking algorithm²³ was used to search a Schrödinger database of commercially available screening compounds for small molecules that bind to the co-receptor binding site on gp120.

Structure-guided queries for the shape-based search were generated by centering on the critical tyrosine sulfate residues of the ligand (Figure 3, panels a and b; and Supplementary Figure 3). Previous biochemical²⁴ and NMR²⁵ data showed that the CCR5-N terminus binding site is selective for tyrosine sulfate over tyrosine phosphate (Figure 3, panel c). Therefore, compounds that contained phosphates were removed from further analysis. Compounds were filtered for the presence of a

sulfur–oxygen bond, and using this criterion sulfates, sulfonates, and sulfonamides were shortlisted. Search of the ZINC database revealed that it contained very few phenyl sulfates, although there were a large number of phenyl sulfonates (Figure 3, panel c). Although missing the oxygen atom linking the SO_3^- and phenyl groups in tyrosine sulfate, phenyl sulfonate retains the aromatic and anionic properties of the tyrosine sulfate side chain.

Compounds were screened for their ability to inhibit the interaction of gp120 with a CCR5-N terminus analogue in an ELISA-based inhibition assay²⁵ (Supplementary Figure 2). Compound **1.1** (Supplementary Figures 3–5), a high-scoring hit appearing in both CCR5 Nt:Y14–Y15 and 412D:Y107–A108–P109 queries, showed the best inhibition and was chosen for the next round of ROCS screening. A secondary ROCS search of the ZINC database was conducted using **1.1** (Supplementary Figure 6) followed by ELISA screening to obtain additional hits (Supplementary Figure 7). These results allowed identification of a chemotype, comprising a phenyl sulfonic acid moiety (Figure 3, panel c) connected to an aryl group by a variable linker, capable of inhibiting gp120–CCR5-N terminus interaction. Chemical structure comparisons on commercial compound libraries were carried out (Supplementary Figure 6) to select compounds on the basis of their similarities with the identified pharmacophore. These were screened in the CCR5-N terminus inhibition ELISA assay to identify additional hits (Supplementary Figure 8).

In an alternative approach, Schrödinger's in-house database of commercially available screening compounds were docked²³ to the CCR5-N terminus binding site on gp120 (Figure 2). Having already identified a binding chemotype, we used this information to filter the hits obtained in the Glide search. A panel of aryl sulfonate compounds were docked to the CCR5-N terminus/412d CDR H3 loop binding site on gp120, and the top-scoring compounds were visually inspected and ranked according to Glide scores. ELISA screening of 60 of these compounds resulted in additional hits (Supplementary Figure 9).

Chemotype Identification. Earlier we showed that a CCR5-N terminus analog, in which the tyrosine sulfate (TYS) residues were replaced by tyrosine sulfonate (Figure 3, panel c), retained significant binding to gp120.²⁵ In this study, the requirement of a phenyl sulfonate headgroup in the drug-like compounds was established in the first few cycles of screening. This moiety shares the aromatic and anionic properties of tyrosine sulfate and the criteria used for *in silico* screening, as well as the direct competition with a CCR5-N terminus analogue, ensured that the determinants of this critical interaction were retained in the selected molecules. Although the phenyl sulfonate group was critical for binding, the contacts made by this moiety alone were not sufficient to inhibit CCR5-N terminus binding to gp120. A panel of aryl sulfonates that showed less than 20% inhibition in the ELISA assay are shown in Supplementary Figure 10. This is consistent with previous biochemical analyses carried out on CCR5-N terminus binding to gp120²⁴ that showed that not only the tyrosine sulfates but also the residues that flanked them were important for binding. The compounds that showed greater than 70% inhibition of CCR5-N terminus peptide binding in the ELISA assay all contained a phenyl sulfonate moiety linked at the *para* position via a diazo or a hydrazino linker to an aromatic group (Supplementary Figures 7–9). A complete list of compounds screened in the ELISA assay with percent inhibition of gp120–CCR5 N terminus interaction is presented in Supplementary Figure 11.

Assessment of Compound Purity and Structure. During characterization, we observed that some compounds were




Figure 4. Small molecule inhibitors of CCR5-N terminus binding to gp120. Small molecules were selected on the basis of their ability to inhibit binding of gp120 to a CCR5-N terminus peptide analogue. All inhibitors contained a phenyl sulfonate moiety, which retains the anionic and aromatic properties of tyrosine sulfate. The two classes of compounds shown were selected for further analyses.

light-sensitive and/or isomerized. Two subgroups whose structures contained the desired pharmacophore and were not prone to isomerization were identified (Figure 4). Compounds 1–3 and 7 (Figure 4 and Table 1) were selected for further analysis. These were subjected to purity evaluation and structural verification by LC–MS and ¹H and ¹³C NMR (Supplementary Figures 12–27). Compounds that were less than 95% pure by LC–MS were purified by RP-HPLC to greater than 95% purity, and their structures confirmed again by NMR and high resolution mass spectroscopy (HR-MS).

Binding Affinity of Compounds to gp120 by Surface Plasmon Resonance (SPR). Binding of compounds to immobilized gp120 was measured in the presence and absence of M48-U1²⁶ (Figure 5, Table 1; Supplementary Figures 28–31). M48-U1 is a member of a class of CD4-peptide mimetics that we previously showed induces the CD4-bound conformation of gp120.^{27–29} The binding kinetics of all of the compounds were too fast for association and dissociation rates to be quantified by SPR. Binding affinities determined by steady-state analyses ranged from 1.5 to 6 μM (Table 1), and binding was enhanced 15- to 40-fold in the presence of the CD4-mimetic peptide M48-U1. Compound 1 showed the tightest binding to gp120 in the CD4-bound conformation with a *K_D* of 49 ± 1 nM.

Inhibition of CD4-Induced Antibody Binding. On the basis of their ability to inhibit CCR5-N terminus binding to gp120, the compounds were expected to occupy one or both tyrosine sulfate binding pockets on gp120 (Figure 1 and Figure 3, panel a). Specificity of the compounds to this site was established by their ability to compete with an antibody that targets the CD4-induced (CD4i) epitope while not affecting the binding of an antibody that binds to an unrelated epitope. The CD4i antibody, 412d was used for the competition assays since its CDR H3 region recognizes the target site¹² (Figure 1 and Figure 2, panel a). The compounds showed dose-dependent inhibition of 412d binding





to gp120 (Figure 5, panel b). No inhibition was observed for binding of antibody C11, which binds outside the CD4i epitope.

Inhibition of HIV-1 Entry. Compounds 1, 2, 3, and 7 were tested for their ability to inhibit HIV entry, and the CD4i antibody 17b was used as positive control. Compounds were tested against a panel of 27 HIV-1 primary isolates along with HIV-2 strains 7312A and 7312A.V434M, a simian strain SIV-mac251 and murine leukemia virus (MuLV) (Figure 6, panel a and Supplementary Table 1). Compound concentrations from 0 to 250 μM were tested. While compounds 3 and 7 did not neutralize any of the viruses tested, compounds 1 and 2 inhibited 25 and 13 viruses, respectively, with 1 showing greater breadth and potency than 2. The activity of the compounds on most strains was weak with IC₅₀ values greater than 100 μM; except for HIV-1 Clade B viruses SS1196.01 and 6535.3 which were inhibited by 1 with IC₅₀ values of 50.3 and 94 μM, respectively. HIV-2 7312A was inhibited by 1, an effect further enhanced in HIV-2 7312A.V434M, a strain that contains a valine to methionine mutation at position 434 leading to greater sensitivity to HIV-1 CD4i antibodies.³⁰ A similar enhancement of neutralization of HIV-2 7312A.V434M compared to its parental virus was seen with 2 and 17b IgG.

Effect of CD4 Induction on Neutralization Potency. The compounds target a region on gp120 that forms only after CD4 engagement. To test if induction of the CD4-bound conformation of the viral spike would enhance the efficacy of the tyrosine sulfate mimetic compounds to inhibit entry, we tested the activity of the compounds on viruses that were first triggered with either soluble CD4 (sCD4) or NBD-556, a CD4 binding site-specific small molecule that preferentially binds to the CD4-bound conformation.³¹ Three viruses, ZA012 (HIV-1, Clade C), JR-FL (HIV-1, Clade B) and 7312A.V434M (HIV-2) were used for the triggering experiments (Figure 6, panel b; Supplementary Tables 2 and 3). 17b IgG was used as positive control.^{30,32} Compounds 1 and 2 showed enhanced neutralization of all three viruses tested when pretriggered with sCD4 with the most significant dose-dependent response seen with ZA012. The effect of triggering with NBD-556 was less significant both for the tyrosine sulfate mimetic compounds and 17b IgG. This observation is consistent with the fact that sCD4 can bind to the viral spike and induce the CD4-bound conformation. NBD-556, however, binds preferentially to the CD4-bound conformation, hence is a probe of the CD4-bound conformation but does not induce a similar conformational change.

Neutralization of Engineered HIV-1 Strains. The compounds were tested in two additional virological assays, each designed to maximize sensitivity for detection of activity of inhibitors that target the co-receptor interactive region of gp120. The first assay utilized a virus with a serine to alanine change (S199A) at the base of the V2 loop of gp120 that disrupts a glycosylation sequon thereby preventing N-linked glycosylation at residue 197.³³ This mutation conferred ability to enter CD4⁻ cells and increased sensitivity to neutralization by sCD4 and various gp120-directed antibodies.^{33,34} HIV-1 ADA Env-pseudovirus with the S199A mutation showed enhanced neutralization by the CD4i antibody 17b (Supplementary Figure 35) compared to the wild-type ADA virus. The efficiency of neutralization by 17b increased when the assay was performed using CD4⁻ target cells, further enhancing sensitivity of the assay for characterization of lead stage inhibitors. We tested 1–3 and 7 (Figure 4) for their ability to inhibit entry of ADA S199A. Whereas 3 and 7 did not show neutralization at the concentration

Table 1. Binding and Entry Inhibition

Compound	K_D (unliganded gp120) (μM)	K_D (gp120 in CD4-bound conformation) (μM)	Geometric IC_{50} mean of primary HIV-1 neutralization (μM)
	1.2 ± 0.1	0.049 ± 0.001	157.6*
	1.6 ± 0.1	0.094 ± 0.002	123.7*
	3.9 ± 0.2	0.112 ± 0.003	> 250
	5.8 ± 0.4	0.181 ± 0.004	> 250

* Geometric means were calculated for viruses that neutralized with IC_{50} value <250 μM ; compound 1 neutralized 23 HIV-1 isolates, and compound 2 neutralized 12 HIV-1 isolates.

range tested, 1 and 2 were able to inhibit viral entry with IC_{50} values of 1.5 ± 0.5 and $44 \pm 7 \mu\text{M}$, respectively (Figure 6, panel c). Toxicity assays showed that 1 and 2 were not toxic to the CF2-Luc cells up to a concentration of 250 μM (Supplementary Figure 36).

A second assay used an engineered virus, CEMAX, adapted for entry using the CCR5-(or the CXCR4-) N terminus (fused to a membrane spanning anchoring peptide) in the complete absence of co-receptor extracellular loops.³⁵ CEMAX remained fully competent for mediating infection on either wildtype CCR5 or CXCR4. In light of CEMAX's exclusive dependence on the co-receptor N terminus for entry, we expected that this virus would be more susceptible to inhibition by compounds that target gp120 interaction with CCR5-N terminus. For the assays with CEMAX we tested compounds 2 and 7, one from each class shown in Figure 4. Both 2 and 7 inhibited CEMAX entry into CCR5⁺ CD4⁺ CF2-Luc reporter cells, with an IC_{50} of $\sim 40 \mu\text{M}$ and showed greater than 80% inhibition at 100 μM (Figure 6, panel c). In the concentration range tested (0–100 μM), no effect was seen on entry of MuLV (Supplementary Figure 32). Both 2 and 7 also inhibited entry of the dual-tropic HIV-1 isolate, HIV-1/R3A and the R5-tropic viruses YU2 and SF162, albeit to a lesser extent (Supplementary Figure 33). Toxicity assays showed that 2 and 7 were not toxic to CF2-Luc cells up to a concentration of 100 μM (Supplementary Figure 34). Interestingly, the

compounds inhibited not only CCR5-mediated entry but also CEMAX entry into cells expressing CXCR4 suggesting targeting of sites common to both R5- and X4-tropic viruses.

All compounds tested (Table 1) bound to monomeric gp120 with binding affinities within 5-fold of each other, yet 1 and 2 neutralized primary isolates and the engineered virus ADA S199A to varying degrees, whereas 3 and 7 did not show neutralization of these viruses in the concentration range tested. These observations suggest isolate-specific barriers in the trimeric HIV-1 spike that further modulate activity and lead to differential activity of these compounds. When tested on CEMAX, an isolate engineered for increased dependence on the co-receptor N termini, 2 and 7 inhibit entry with similar IC_{50} values. This suggests alleviation of barriers that cause differential reactivity of the compounds against native HIV isolates.

Summary and Implications for Development of Entry Inhibitors That Target the HIV Envelope—Co-receptor Interaction. The co-receptor binding site, especially the region spanning the bridging sheet and the base of the V3 loop (Supplementary Figure 1),⁵ is highly conserved. Antibodies that bind to this region in monomeric gp120 have broad binding specificity across different viral isolates. While the high degree of conservation is a desirable property of this potential target, this region forms only upon CD4 engagement of the HIV-1 spike. After CD4 binds, the proximity of the viral and cell membranes




Figure 5. gp120 binding. (a) Direct binding to gp120. Binding of compounds 1, 2, 3, and 7 to gp120 in the unliganded (top panel) and in the CD4-bound conformation (bottom panel) was measured by SPR and affinity calculated by steady state analysis. (b) Inhibition of binding of CD4i antibody 412d to gp120 in the CD4-bound conformation by the compounds.

renders the co-receptor binding site sterically inaccessible to molecules as large as antibodies,³⁶ though antibody fragments and single chain antibody variants retain access.³⁷ Small molecules, therefore, may present a viable strategy for intervention at the site of co-receptor binding on HIV-1.

We have used structural information for *in silico* screening and coupled this to experimental screening to identify small molecule inhibitors of HIV-1 binding to its co-receptor. This led to the identification of a novel phenylsulfonate-linker-aromatic motif with HIV entry inhibition properties. Since information from ligands (CCR5-N terminus and 412d antibody) that specifically target CCR5-using viruses was used to screen for small molecule mimetics, we expected these mimetics to be active only against CCR5-mediated entry but found that they inhibited entry independent of whether it occurred with CCR5 or CXCR4 (Figure 6, panel c). While somewhat unexpected, one potential explanation is the high degree of conservation of the tyrosine sulfate binding regions on gp120 across all HIV isolates, irrespective of tropism.

HIV uses a number of mechanisms to evade antibody-mediated neutralization. These include variable loop occlusion,³⁸ glycan shielding,³⁹ and conformational masking.⁴⁰ Comparison of the properties of the tyrosine sulfate mimetic small molecules

determined in this study with other known HIV-1 neutralizing molecules (Table 2 and Supplementary Table 4) show that their properties strongly correlate with those of molecules that target the HIV co-receptor binding site. First, 15- to 40-fold tighter binding of the tyrosine sulfate mimetic small molecules was observed to gp120 fixed in the CD4-bound conformation relative to gp120 that was unrestrained. Second, they competed with antibodies that bind to the CD4-induced epitope of gp120. Third, neutralization of primary isolates was enhanced by pre-triggering with sCD4. Fourth, viruses with impaired conformational masking (e.g., ADA S199A) were neutralized with greater potency than native HIV-1 isolates. These results and the strong correlation of properties with molecules targeting the CD4i epitope of gp120 (Table 2) reveal that the mechanism of action of these tyrosine sulfate mimetics involves binding to the conformationally malleable co-receptor binding region of gp120, specifically, the CD4-induced site. Conformational masking appears to hinder the ability of these small molecules to neutralize primary HIV-1 isolates. Our studies indicate that these tyrosine sulfate mimetics may work best in synergy with molecules that induce the CD4-bound state of gp120 such as sCD4 or CD4-mimetic peptides.^{27–29}

In summary, we report the structure-guided identification and characterization of a new class of tyrosine sulfate-mimicking small




Figure 6. Inhibition of HIV entry. (a) Neutralization profiles of compounds 1, 2, and monoclonal antibody 17b depicted with 29-isolate neutralization dendrograms. SIV and MuLV were used as controls. IC₅₀ values are listed in Supplementary Table 1. Insets show neutralization of two HIV-2 isolates tested. (b) Inhibition of entry in presence of two-domain CD4 (sCD4) or the small molecule NBD-556.⁴³ The *p*-values indicate statistical significance for effect of triggering with sCD4 or NBD-556 compared to untreated virus calculated using ANOVA. (c) Left, inhibition of entry of HIV-1 ADA S199A, and right, inhibition of CCR5-mediated entry (circles) and of CXCR4-mediated entry (squares) of CEMAX.

Table 2. Significance of Agreement between Phenotypes of Molecules Targeting Different Neutralization Sites on HIV-1^a

	HIV-1 neutralizing molecules					
	CD4-induced site targeting	CD4-binding site targeting	glycan targeting	gp41 targeting	V3 targeting	quaternary site targeting
TYS mimetic small molecules	0.003	<i>1</i>	0.467	<i>1</i>	0.048	0.524
CD4-induced site targeting		0.464	0.464	<i>1</i>	0.048	<i>1</i>
CD4-binding site targeting			0.067	0.196	0.52	0.464
Glycan targeting				0.464	0.167	<i>1</i>

^aNumbers in table are *p*-values from Fisher's exact test. Supplementary Table 4 lists phenotypes of neutralizing molecules; **bold** = *p*-value < 0.05, significant; *italic* = *p*-value > 0.05, not significant.

molecules that bind to the HIV-1 gp120 envelope glycoprotein and inhibit entry. Recognition of and inhibition by these tyrosine

sulfate mimetics provide an example for how the *O*-sulfation post-translational modification can be exploited therapeutically.

Analysis of the neutralization mechanism of these compounds indicates that the conformational barriers evolved by HIV-1 to protect the highly conserved co-receptor binding region from antibody-mediated neutralization also restricts neutralization by small molecules that target this region. Despite this evasion tactic, the ability of antibody fragments to potently neutralize HIV-1³⁷ provides proof-of-principle that, with sufficient affinity, effective neutralization by this class of tyrosine sulfate mimetics can be achieved. It will be interesting to see if improved mimicry increases neutralization potency, for example, with small molecules that contain tyrosine sulfate as opposed to tyrosine sulfonate. Moreover, compounds **1** and **2** neutralized primary HIV-1 isolates far better than compounds **3** and **7**, despite less than 5-fold difference in K_D , suggesting that small changes in affinity may have more dramatic effects on neutralization potency.

METHODS

Additional details for each section are available in Supporting Information.

Virtual Screening. gp120-bound coordinates of 412d and CCR5-N terminus were used for virtual screening. ROCS²¹ searches were done on the ZINC database, and hits were rescored and docked with Gold.²² In a complementary approach the druglike subset of a database of commercially available compounds was docked to the protein models using Glide v4.5 (Schrödinger, LLC).²³

ELISA-Based Screening Assay. The ELISA-based screening was performed as described earlier²⁵ using 500 μ M compound concentration (unless otherwise indicated).

Surface Plasmon Resonance. Different compound concentrations were injected onto a gp120-coated surface in absence or presence of M48-U1. For competition assays, 412d and C11 IgG were immobilized on different flow cells. A gp120-M48-U1 complex with or without compound was injected and initial slopes were calculated for the on-phase.

HIV Entry Inhibition Assays. The ability of the compounds to inhibit entry of CEMAX³⁵ was evaluated on canine thymocyte cells (CF2-luc) stably expressing a luciferase reporter under control of an HIV-1 LTR (kindly provided by D. Gabuzda, Harvard).

All other viral entry inhibition assays were performed as previously described,⁴¹ except assays with ADA S199A, where target cells were canine thymocytes (Cf2Th) stably expressing human CCR5 or both CD4 and CCR5.⁴² The triggering assays were performed as described earlier.³²

Target cell viability was determined in a format similar to the neutralization assays using a luminescence-based assay measuring cellular expression of adenosine triphosphate (CellTiter-Glo Luminescent Cell Viability Assay, Promega Corp.).

ASSOCIATED CONTENT

Supporting Information. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: caroleb@mail.nih.gov; pdkwong@nih.gov.

Author Contributions

[∇]These authors contributed equally to this work.

ACKNOWLEDGMENT

We thank J. Lloyd for HR-MS data, members of the Structural Biology Section and Structural Bioinformatics Core at the NIH

Vaccine Research Center for comments on the manuscript, J. Stuckey for assistance with figures, and I. Georgiev for discussions on statistical methods. sCD4 and Cf2Th cells stably expressing human CD4 and CCR5 were obtained through the AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH. Support for this work was provided by the Intramural AIDS Targeted Antiviral Program (C.A.B. and P.D.K.), by the Intramural Research Program of the NIH (NIDDK and NIAID), by grants from the European Community's Sixth Framework programme (EMPRO) under grant agreement no. LSHP-2003-503558, by a grant from the French National Agency of Research (ANR-07-EMPB-019-01) (L.M.) and by a grant from the NIH (GM 56550) (J.M.L.). C.D.-I. was supported by an Intramural AIDS Research Fellowship.

REFERENCES

- (1) Grant, P. M., and Zolopa, A. R. (2010) Optimal antiretroviral therapy: HIV-1 treatment strategies to avoid and overcome drug resistance. *Curr. Opin. Invest. Drugs* 11, 901–910.
- (2) Qian, K., Morris-Natschke, S. L., and Lee, K. H. (2009) HIV entry inhibitors and their potential in HIV therapy. *Med. Res. Rev.* 29, 369–393.
- (3) Chen, B., Vogan, E. M., Gong, H. Y., Skehel, J. J., Wiley, D. C., and Harrison, S. C. (2005) Structure of an unliganded simian immunodeficiency virus gp120 core. *Nature* 433, 834–841.
- (4) Kwong, P. D., Wyatt, R., Robinson, J., Sweet, R. W., Sodroski, J., and Hendrickson, W. A. (1998) Structure of an HIV gp120 envelope glycoprotein in complex with the CD4 receptor and a neutralizing human antibody. *Nature* 393, 648–659.
- (5) Rizzuto, C. D., Wyatt, R., Hernandez-Ramos, N., Sun, Y., Kwong, P. D., Hendrickson, W. A., and Sodroski, J. (1998) A conserved HIV gp120 glycoprotein structure involved in chemokine receptor binding. *Science* 280, 1949–1953.
- (6) Berger, E. A., Murphy, P. M., and Farber, J. M. (1999) Chemokine receptors as HIV-1 co-receptors: Roles in viral entry, tropism, and disease. *Annu. Rev. Immunol.* 17, 657–700.
- (7) Matthews, T., Salgo, M., Greenberg, M., Chung, J., DeMasi, R., and Bolognesi, D. (2004) Enfuvirtide: The first therapy to inhibit the entry of HIV-1 into host CD4 lymphocytes. *Nat. Rev. Drug Discovery* 3, 215–225.
- (8) Dorr, P., Westby, M., Dobbs, S., Griffin, P., Irvine, B., Macartney, M., Mori, J., Rickett, G., Smith-Burchnell, C., Napier, C., Webster, R., Armour, D., Price, D., Stammen, B., Wood, A., and Perros, M. (2005) Maraviroc (UK-427,857), a potent, orally bioavailable, and selective small-molecule inhibitor of chemokine receptor CCR5 with broad-spectrum anti-human immunodeficiency virus type 1 activity. *Antimicrob. Agents Chemother.* 49, 4721–4732.
- (9) Huang, Y. X., Paxton, W. A., Wolinsky, S. M., Neumann, A. U., Zhang, L. Q., He, T., Kang, S., Ceradini, D., Jin, Z. Q., Yazdanbakhsh, K., Kunstman, K., Erickson, D., Dragon, E., Landau, N. R., Phair, J., Ho, D. D., and Koup, R. A. (1996) The role of a mutant CCR5 allele in HIV-1 transmission and disease progression. *Nat. Med.* 2, 1240–1243.
- (10) Cormier, E. G., Persuh, M., Thompson, D. A. D., Lin, S. W., Sakmar, T. P., Olson, W. C., and Dragic, T. (2000) Specific interaction of CCR5 amino-terminal domain peptides containing sulfotyrosines with HIV-1 envelope glycoprotein gp120. *Proc. Natl. Acad. Sci. U.S.A.* 97, 5762–5767.
- (11) Cormier, E. G., and Dragic, T. (2002) The crown and stem of the V3 loop play distinct roles in human immunodeficiency virus type 1 envelope glycoprotein interactions with the CCR5 co-receptor. *J. Virol.* 76, 8953–8957.
- (12) Huang, C. C., Lam, S. N., Acharya, P., Tang, M., Xiang, S. H., Hussan, S. S., Stanfield, R. L., Robinson, J., Sodroski, J., Wilson, I. A., Wyatt, R., Bewley, C. A., and Kwong, P. D. (2007) Structures of the CCR5 N terminus and of a tyrosine sulfated antibody with HIV-1 gp120 and CD4. *Science* 317, 1930–1934.
- (13) Huang, C. C., Venturi, M., Majeed, S., Moore, M. J., Phogat, S., Zhang, M. Y., Dimitrov, D. S., Hendrickson, W. A., Robinson, J., Sodroski, J., Wyatt, R., Choe, H., Farzan, M., and Kwong, P. D.

- (2004) Structural basis of tyrosine sulfation and VH-gene usage in antibodies that recognize the HIV type 1 co-receptor-binding site on gp120. *Proc. Natl. Acad. Sci. U.S.A.* 101, 2706–2711.
- (14) Choe, H., Li, W., Wright, P. L., Vasilieva, N., Venturi, M., Huang, C. C., Grundner, C., Dorfman, T., Zwick, M. B., Wang, L., Rosenberg, E. S., Kwong, P. D., Burton, D. R., Robinson, J. E., Sodroski, J. G., and Farzan, M. (2003) Tyrosine sulfation of human antibodies contributes to recognition of the CCR5 binding region of HIV-1 gp120. *Cell* 114, 161–170.
- (15) Xiang, S. H., Farzan, M., Si, Z., Madani, N., Wang, L., Rosenberg, E., Robinson, J., and Sodroski, J. (2005) Functional mimicry of a human immunodeficiency virus type 1 co-receptor by a neutralizing monoclonal antibody. *J. Virol.* 79, 6068–6077.
- (16) Kehoe, J. W., and Bertozzi, C. R. (2000) Tyrosine sulfation: a modulator of extracellular protein-protein interactions. *Chem. Biol.* 7, R57–R61.
- (17) Gunning, P. T., Glenn, M. P., Siddiquee, K. A. Z., Katt, W. P., Masson, E., Sebt, S. M., Turkson, J., and Hamilton, A. D. (2008) Targeting protein-protein interactions: Suppression of Stat3 dimerization with rationally designed small-molecule, nonpeptidic SH2 domain binders. *ChemBioChem* 9, 2800–2803.
- (18) Panigrahi, K., Eggen, M., Maeng, J. H., Shen, Q. R., and Berkowitz, D. B. (2009) The alpha, alpha-difluorinated phosphonate L-pSer-analogue: An accessible chemical tool for studying kinase-dependent signal transduction. *Chem. Biol.* 16, 928–936.
- (19) Sun, J. P., Wu, L., Fedorov, A. A., Almo, S. C., and Zhang, Z. Y. (2003) Crystal structure of the Yersinia protein-tyrosine phosphatase YopH complexed with a specific small molecule inhibitor. *J. Biol. Chem.* 278, 33392–33399.
- (20) Irwin, J. J., and Shoichet, B. K. (2005) ZINC - A free database of commercially available compounds for virtual screening. *J. Chem. Inf. Model.* 45, 177–182.
- (21) Rush, T. S., 3rd, Grant, J. A., Mosyak, L., and Nicholls, A. (2005) A shape-based 3-D scaffold hopping method and its application to a bacterial protein-protein interaction. *J. Med. Chem.* 48, 1489–1495.
- (22) Jones, G., Willett, P., Glen, R. C., Leach, A. R., and Taylor, R. (1997) Development and validation of a genetic algorithm for flexible docking. *J. Mol. Biol.* 267, 727–748.
- (23) Halgren, T. A., Murphy, R. B., Friesner, R. A., Beard, H. S., Frye, L. L., Pollard, W. T., and Banks, J. L. (2004) Glide: A new approach for rapid, accurate docking and scoring. 2. Enrichment factors in database screening. *J. Med. Chem.* 47, 1750–1759.
- (24) Cormier, E. G., Persuh, M., Thompson, D. A., Lin, S. W., Sakmar, T. P., Olson, W. C., and Dragic, T. (2000) Specific interaction of CCR5 amino-terminal domain peptides containing sulfotyrosines with HIV-1 envelope glycoprotein gp120. *Proc. Natl. Acad. Sci. U.S.A.* 97, 5762–5767.
- (25) Lam, S. N., Acharya, P., Wyatt, R., Kwong, P. D., and Bewley, C. A. (2008) Tyrosine sulfate isosteres of CCR5 N-terminus as tools for studying HIV-1 entry. *Bioorg. Med. Chem.* 16, 10113–10120.
- (26) Van Herrewege, Y., Morellato, L., Descours, A., Aerts, L., Michiels, J., Heyndrickx, L., Martin, L., and Vanham, G. (2008) CD4 mimetic miniproteins: potent anti-HIV compounds with promising activity as microbicides. *J. Antimicrob. Chemother.* 61, 818–826.
- (27) Huang, C. C., Stricher, F., Martin, L., Decker, J. M., Majeed, S., Barthe, P., Hendrickson, W. A., Robinson, J., Roumestand, C., Sodroski, J., Wyatt, R., Shaw, G. M., Vita, C., and Kwong, P. D. (2005) Scorpion-toxin mimics of CD4 in complex with human immunodeficiency virus gp120: Crystal structures, molecular mimicry, and neutralization breadth. *Structure* 13, 755–768.
- (28) Martin, L., Stricher, F., Misse, D., Sironi, F., Pugniere, M., Barthe, P., Prado-Gotor, R., Freulon, I., Magne, X., Roumestand, C., Menez, A., Lusso, P., Veas, F., and Vita, C. (2003) Rational design of a CD4 mimic that inhibits HIV-1 entry and exposes cryptic neutralization epitopes. *Nat. Biotechnol.* 21, 71–76.
- (29) Stricher, F., Huang, C. C., Descours, A., Duquesnoy, S., Combes, O., Decker, J. M., Do Kwon, Y., Lusso, P., Shaw, G. M., Vita, C., Kwong, P. D., and Martin, L. (2008) Combinatorial optimization of a CD4-mimetic miniprotein and cocrystal structures with HIV-1 gp120 envelope glycoprotein. *J. Mol. Biol.* 382, 510–524.
- (30) Decker, J. M., Bibollet-Ruche, F., Wei, X., Wang, S., Levy, D. N., Wang, W., Delaporte, E., Peeters, M., Derdeyn, C. A., Allen, S., Hunter, E., Saag, M. S., Hoxie, J. A., Hahn, B. H., Kwong, P. D., Robinson, J. E., and Shaw, G. M. (2005) Antigenic conservation and immunogenicity of the HIV co-receptor binding site. *J. Exp. Med.* 201, 1407–1419.
- (31) Madani, N., Schon, A., Princiotta, A. M., Lalonde, J. M., Courter, J. R., Soeta, T., Ng, D., Wang, L., Brower, E. T., Xiang, S. H., Kwon, Y. D., Huang, C. C., Wyatt, R., Kwong, P. D., Freire, E., Smith, A. B., 3rd, and Sodroski, J. (2008) Small-molecule CD4 mimics interact with a highly conserved pocket on HIV-1 gp120. *Structure* 16, 1689–1701.
- (32) Wu, X., Sambor, A., Nason, M. C., Yang, Z. Y., Wu, L., Zolla-Pazner, S., Nabel, G. J., and Mascola, J. R. (2008) Soluble CD4 broadens neutralization of V3-directed monoclonal antibodies and guinea pig vaccine sera against HIV-1 subtype B and C reference viruses. *Virology* 380, 285–295.
- (33) Kolchinsky, P., Kiprilov, E., Bartley, P., Rubinstein, R., and Sodroski, J. (2001) Loss of a single N-linked glycan allows CD4-independent human immunodeficiency virus type 1 infection by altering the position of the gp120 V1/V2 variable loops. *J. Virol.* 75, 3435–3443.
- (34) Li, Y., Cleveland, B., Klots, I., Travis, B., Richardson, B. A., Anderson, D., Montefiori, D., Polacino, P., and Hu, S. L. (2008) Removal of a single N-linked glycan in human immunodeficiency virus type 1 gp120 results in an enhanced ability to induce neutralizing antibody responses. *J. Virol.* 82, 638–651.
- (35) Leslie, G. J., Acharya, P., Haggarty, B., Jordan, A., Romano, J., Kwong, P. D., Hoxie, J. (2011) Derivation of a dual-tropic HIV-1 isolate that can utilize the co-receptor amino terminus in the absence of extracellular loops. *Keystone HIV Pathogenesis Meeting*, Whistler, Canada.
- (36) Labrijn, A. F., Poignard, P., Raja, A., Zwick, M. B., Delgado, K., Franti, M., Binley, J., Vivona, V., Grundner, C., Huang, C. C., Venturi, M., Petropoulos, C. J., Wrin, T., Dimitrov, D. S., Robinson, J., Kwong, P. D., Wyatt, R. T., Sodroski, J., and Burton, D. R. (2003) Access of antibody molecules to the conserved co-receptor binding site on glycoprotein gp120 is sterically restricted on primary human immunodeficiency virus type 1. *J. Virol.* 77, 10557–10565.
- (37) Chen, W. Z., Zhu, Z. Y., Feng, Y., and Dimitrov, D. S. (2008) Human domain antibodies to conserved sterically restricted regions on gp120 as exceptionally potent cross-reactive HIV-1 neutralizers. *Proc. Natl. Acad. Sci. U.S.A.* 105, 17121–17126.
- (38) Wyatt, R., Moore, J., Accola, M., Desjardin, E., Robinson, J., and Sodroski, J. (1995) Involvement of the V1/V2 Variable Loop Structure in the exposure of Human-Immunodeficiency-Virus Type-1 Gp120 epitopes induced by receptor-binding. *J. Virol.* 69, 5723–5733.
- (39) Wei, X. P., Decker, J. M., Wang, S. Y., Hui, H. X., Kappes, J. C., Wu, X. Y., Salazar-Gonzalez, J. F., Salazar, M. G., Kilby, J. M., Saag, M. S., Komarova, N. L., Nowak, M. A., Hahn, B. H., Kwong, P. D., and Shaw, G. M. (2003) Antibody neutralization and escape by HIV-1. *Nature* 422, 307–312; Erratum: *Nature* 423, 197.
- (40) Kwong, P. D., Doyle, M. L., Casper, D. J., Cicala, C., Leavitt, S. A., Majeed, S., Steenbeke, T. D., Venturi, M., Chaiken, I., Fung, M., Katinger, H., Parren, P. W. L. H., Robinson, J., Van Ryk, D., Wang, L. P., Burton, D. R., Freire, E., Wyatt, R., Sodroski, J., Hendrickson, W. A., and Arthos, J. (2002) HIV-1 evades antibody-mediated neutralization through conformational masking of receptor-binding sites. *Nature* 420, 678–682.
- (41) Li, Y. X., Migueles, S. A., Welcher, B., Svehla, K., Phogat, A., Louder, M. K., Wu, X. L., Shaw, G. M., Connors, M., Wyatt, R. T., and Mascola, J. R. (2007) Broad HIV-1 neutralization mediated by CD4-binding site antibodies. *Nat. Med.* 13, 1032–1034.
- (42) Mirzabekov, T., Bannert, N., Farzan, M., Hofmann, W., Kolchinsky, P., Wu, L. J., Wyatt, R., and Sodroski, J. (1999) Enhanced expression, native purification, and characterization of CCR5, a principal HIV-1 co-receptor. *J. Biol. Chem.* 274, 28745–28750.
- (43) Zhao, Q., Ma, L., Jiang, S., Lu, H., Liu, S., He, Y., Strick, N., Neamati, N., and Debnath, A. K. (2005) Identification of N-phenyl-N'-(2,2,6,6-tetramethyl-piperidin-4-yl)-oxalamides as a new class of HIV-1 entry inhibitors that prevent gp120 binding to CD4. *Virology* 339, 213–225.