


Antibody-Recruiting Molecules: An Emerging Paradigm for Engaging Immune Function in Treating Human Disease

Patrick J. McEnaney,^{†,§} Christopher G. Parker,^{†,§} Andrew X. Zhang,^{†,§} and David A. Spiegel^{*,†,‡}

[†]Department of Chemistry, Yale University, New Haven, Connecticut 06520, United States

[‡]Department of Pharmacology, Yale School of Medicine, New Haven, Connecticut 06510, United States

ABSTRACT: Synthetic immunology, the development of synthetic systems capable of modulating and/or manipulating immunological functions, represents an emerging field of research with manifold possibilities. One focus of this area has been to create low molecular weight synthetic species, called antibody-recruiting molecules (ARMs), which are capable of enhancing antibody binding to disease-relevant cells or viruses, thus leading to their immune-mediated clearance. This article provides a thorough discussion of contributions in this area, beginning with the history of small-molecule-based technologies for modulating antibody recognition, followed by a systematic review of the various applications of ARM-based strategies. Thus, we describe ARMs capable of targeting cancer, bacteria, and viral pathogens, along with some of the scientific discoveries that have resulted from their development. Research in this area underscores the many exciting possibilities at the interface of organic chemistry and immunobiology and is positioned to advance both basic and clinical science in the years to come.



The introduction of cowpox (*vaccinia*) virus immunization by Edward Jenner in 1796 was a landmark moment in the history of medicine.¹ Not only did Jenner's vaccination strategy ultimately lead to the eradication of smallpox, it made clear the extraordinary power of a person's own immune system for warding off deadly illness. Subsequent advances, including the implementation of passive antibody therapy or "serum therapy" in the late 1800s,² and Milstein and Kohler's report of the first engineered monoclonal antibody in the 1970s,³ have paved the way for a new revolution in immunotherapeutics focused on monoclonal-antibody-based drugs. The first member of this class, Muromonab-CD3 (anti-CD3 or OKT3),³ was cleared by the FDA in 1986 for treating transplant rejection, and since that time, the number of antibody drugs has increased dramatically; 31 agents are currently approved for clinical use, and more than 300 are undergoing clinical trials.^{4–7} Furthermore, in 2011 alone, antibody-based therapies grossed \$44.6 billion worldwide, and this number has been predicted to increase in the upcoming years.^{6,8,9}

The surge in popularity of monoclonal antibodies can be easily understood in light of their many extraordinary properties. Antibody molecules are readily generated against a variety of disease-relevant targets, some of which have been conventionally considered "undruggable".¹⁰ Additionally, because antibodies often interact with their targets with excellent affinity and specificity, undesirable side effects related to off-target binding are thought to be low relative to traditional small-molecule-based therapeutics. Finally, antibodies may elicit therapeutic responses by a variety of mechanisms including inhibition of protein function,^{11,12} targeting the delivery of cytotoxic drugs,^{13–15} and triggering immune effector responses.^{16–18}

There are two mechanisms by which antibodies can trigger immune-mediated cytotoxicity: engagement of plasma complement proteins and/or direct activation of immune effector cells. The former process, termed complement-dependent cytotoxicity (CDC), begins upon cell-surface immobilization of certain members of the complement protein family (*e.g.*, C1q) by opsonizing antibodies. This event initiates a downstream proteolytic cascade culminating in direct cell lysis or recruitment of complement-receptor-expressing effector cells, ultimately leading to target cell clearance.¹⁹ Alternatively, binding of the antibody's crystallizable fragment (Fc) to Fc-receptors expressed on the surface of various immune cells can lead to receptor cross-linking, followed by target cell phagocytosis or the release of potent oxidizing agents and protein toxins (*e.g.*, granzyme and perforin).²⁰ These processes are termed antibody-dependent cellular phagocytosis (ADCP) and antibody-dependent cellular cytotoxicity (ADCC), respectively.¹⁹ Importantly, because Fc receptor-mediated cytotoxicity can enhance the processing and presentation of disease-relevant antigens, these cellular mechanisms have the capacity to give rise to long-lasting adaptive immunity.^{21–24}

These advantages notwithstanding, antibody-based therapeutics suffer from certain limitations that arise primarily from their high molecular weights and peptide structures. For example, antibody administration can result in systemic inflammatory response syndrome, IgE-mediated acute anaphylactic reactions, serum sickness, and cytokine release syndrome, all of which

Received: March 15, 2012

Accepted: June 1, 2012

Published: July 3, 2012




Figure 1. Antibody recruiting small molecules (ARMs). ARM s are bifunctional small molecules that function by forming ternary complexes with disease-relevant targets and endogenous antibodies. Ternary complex assembly leads to the activation of immune effector functions, followed by immune-mediated cytotoxicity and/or clearance of disease-causing species.

have the potential to be life-threatening.²⁵ In addition, because antibody therapeutics contain “non-self” protein sequences, they can elicit host immune reactions, thus counteracting their efficacy.²⁵ Although the widespread “humanization” of antibody protein sequences has drastically reduced the potential for undesired immune responses, such reactions remain problematic clinically.^{26,27} Another limiting factor is that individuals can express different allelic variants of Fc receptors, which can directly impact the efficacy of “naked” monoclonal antibodies intended to act through interaction with these receptors.²⁸ Antibody–toxin conjugates, developed to address such difficulties, are associated with other drawbacks. These include premature drug release arising from antibody–drug linker instability, which can lead to toxic effects in normal tissues.²⁹ In addition, identifying optimal levels of drug conjugation can be difficult; low loadings may render treatment ineffective, whereas high loadings may lead to excess toxicity or a loss of antigen specificity.³⁰ Other limitations of antibody-based therapeutics include a lack of oral bioavailability, difficulties in standardization and characterization, and high costs.^{25,30–32}

Researchers have begun to investigate the development of low molecular weight species (“small molecules”) that possess the complex functional properties of antibodies and other biologics. Because small molecules are generally inexpensive to produce, optimizable for oral bioavailability,²⁹ and unlikely to cause unwanted allergic or immunogenic responses,³¹ these agents may be able to address many of the limitations of biologics without compromising their advantages. These studies offer insight into how man-made systems for controlling human immune function might broadly enable the development of a synthetic immunology for use in both basic and clinical science.^{33,34}

We focus here on antibody-recruiting small molecules (ARMs), which we define as synthetic, bifunctional molecules capable of inducing antibodies to bind disease-relevant proteins, cells, or organisms (Figure 1). Simultaneous association of ARM s with antibodies and surface-exposed receptors results in the formation of ternary complexes, which can elicit antibody-dependent immune effector responses. By convention, we define the two regions of ARM s as the target-binding terminus (TBT), which recognizes the disease-associated protein target (either present on cell/viral surfaces or free in solution), and the antibody-binding terminus (ABT), which associates with anti-hapten antibodies. Notably, we have chosen to limit the scope of this Review to technologies in which synthetic (*i.e.*, non-recombinant), organic ligands are employed to control the immunological functions of antibody proteins; therefore, a variety of interesting and important systems, including antibody–drug conjugates,³⁵ antibody-based recombinant constructs (*e.g.*, bispecific antibodies, diabodies, and others),^{36–38} synthetic vaccines,³⁹ and ligand-templated supramolecular assemblies,^{40–44} are not covered in detail herein.

As early as the 1970s, artificially engineered systems were shown to redirect antibody responses to the surfaces of cells that are ordinarily non-immunogenic. For example, liver, spleen, and red blood cells covalently modified with small molecule haptens, such as trinitrophenyl (TNP) groups, were shown to serve as targets for antibody-dependent cell-mediated immune responses.^{45,46} Furthermore, anti-DNP antibodies of IgG and IgM isotypes were shown to associate specifically with liposomes labeled with dinitrophenyl (DNP) groups, leading to the induction of a complement-mediated lytic response.⁴⁷ In all of these cases, antibody targeting and cytotoxicity were shown to be hapten-dependent, that is, antibodies were directed as a result of covalent DNP or TNP labeling. Although hapten labeling in these cases was nonspecific, these early studies were critical in demonstrating that antibody-mediated immune responses could be templated by “non-native”, synthetic materials.

Subsequently, strategies utilizing rationally designed bifunctional systems to redirect the immune response to disease-relevant targets began to emerge. For example, chimeric proteins consisting of IgG or IgM Fc domains fused to human CD4, the cell-surface receptor target of HIV gp120, were shown to bind both to the complement protein C1q and to Fc-receptors.^{48–50} These “immunoadhesins” were further shown to enhance immune effector responses selectively against HIV infected cells in the presence of peripheral blood mononuclear cells (PBMCs), while inflicting minimal background cytotoxicity. These pioneering studies represent the first evidence that rationally designed, bifunctional molecules could specifically target immune-mediated functions to pathogenic proteins.

Subsequent research efforts expanded on these findings by demonstrating that proteins derivatized with small-molecule haptens could also possess immunomodulatory properties. For example, soluble CD4 covalently modified with the dinitrophenyl (DNP) motif was shown to mediate the formation of a quaternary complex between gp120, anti-DNP antibodies, and soluble complement protein C1q.⁵¹ Similarly, a dimeric Fab fragment (F(ab')₂) directed against the T-cell marker antithymocyte globulin (ATG), covalently labeled with fluorescein, was shown to be selectively cytotoxic to T-cells in the presence anti-fluorescein antibodies and complement proteins.⁵² Additionally, treatment of fluorescein-immunized mice with this molecule resulted in the clearance of peripheral ATG-expressing T-cells. Together these studies were critical in setting the stage for developing bifunctional ARM s.

Anti-hapten Antibodies Used in ARM Strategies. To date, ARM s have incorporated two types of functionality at the ABT: (1) small molecule ligands for “endogenous” antibodies or (2) rationally designed functional handles, which require delivery of preformed antibody-small molecule conjugates or preimmunization for induction of selective antibody responses. Perhaps the most common targets in the first category include the galactosyl-(1–3)-galactose (α -Gal) carbohydrate epitope




Figure 2. Chemical structures of cancer-targeting ARMs. TBTs are highlighted in red boxes, and ABTs are in blue boxes.

and the 2,4-dinitrophenyl (DNP) motif. Intriguingly, 2–8% of circulating antibodies in the bloodstream are believed to recognize the α -Gal trisaccharide,^{80,81} and these are believed to arise following exposure to this carbohydrate on the surfaces of cells derived from prokaryotes and non-primate eukaryotes.⁸² Unlike α -Gal, DNP and other nitroarenes are not likely produced biosynthetically.⁸³ Despite this, approximately 1% of circulating antibodies in the human bloodstream have been shown to recognize nitroarene epitopes.^{83,84} Although the origin of anti-DNP antibodies is not known, one potential route of human inoculation involves exposure to DNP-containing dyes, preservatives, and/or pesticides,⁸⁵ which have been detected as environmental contaminants throughout the United States.^{86,87} An alternative hypothesis is that dietary ingestion of proteins and/or peptides containing nitroaromatic amino acids, formed in foods during the cooking process,⁸⁸ leads to adaptive

immune responses against hapten-containing neo-epitopes,^{89–91} perhaps in a TLR-independent manner.⁹²

The second class of ABT that has been used in ARM strategies includes non-native or synthetic antigens. Although such motifs by definition are not expected to bind pre-existing antibody proteins, humoral immune responses against these epitopes are readily induced by immunization with the hapten of interest conjugated to a carrier protein. One advantage of this approach is that haptens with useful chemical and/or physical properties can be chosen. For example, anti-fluorescein antibodies can easily be induced through immunization with protein conjugates of fluorescein isothiocyanate (FITC),^{62,93} and the haptens' photophysical properties provide a convenient handle for binding and imaging studies. Another useful immunization-dependent strategy exploits the unique properties of catalytic aldolase antibodies, which can form covalent adducts with 1,3-diketones (Figure 2, 1 \rightarrow 3; 2 \rightarrow 4) or β -

Table 1. Summary of the Applications of Antibody-Recruiting Small Molecules to Disease Targets

disease/pathogen	target	antibody-binding moiety	molecule type	<i>in vitro</i> response	<i>in vivo</i> model	Ab source
<i>E. coli</i> ^{53,54}	mannose receptor	avidin	small molecule-protein conjugate	CDC, phagocytosis		commercial
<i>E. coli</i> ⁵⁵	mannose receptor	α -Gal	peptide	inhibition of agglutination		endogenous
Gram-positive bacteria ^{56,57}	D-Ala-D-Ala	fluorescein	polymer	phagocytosis		commercial
HIV ⁵⁸	gp41	α -Gal	peptide	viral inhibition		endogenous
HIV ⁵⁹	gp120	α -Gal	peptide	CDC, ADCC		endogenous
HIV ⁶⁰	gp120	DNP	small molecule	CDC, viral inhibition		commercial
HIV ⁶¹	CCRS	β -lactam	mAb-small molecule conjugate	viral inhibition		monoclonal antibody
lung cancer ^{62–64}	folate receptor	fluorescein, DNP	small molecule	CDC, ADCC, phagocytosis	syngeneic female Balb/c	immunization
Karposi's sarcoma, colon cancer, melanoma ^{65–67}	integrin receptors ($\alpha_v\beta_3$ and $\alpha_v\beta_5$)	1,3-diketone	mAb-small molecule conjugate	CDC, ADCC	female nude and SCID xenografts	monoclonal antibody
prostate cancer ⁶⁸	ET _A	1,3-diketone	mAb-small molecule conjugate	opsonization	nude xenograft	monoclonal antibody
breast cancer, melanoma, osteosarcoma, Karposi's sarcoma ^{69,70}	$\alpha_v\beta_5$	α -Gal	small molecule	CDC		endogenous
B cell lymphoma ^{71,72}	CD22	nitrophenol	small molecule, polymer	opsonization		commercial
colon cancer, melanoma ⁷³	integrin receptors ($\alpha_v\beta_3$ and $\alpha_v\beta_5$)	1,3-diketone	small molecule	ADCC	syngeneic female BALB/c, and C57BL6	reactive immunization
melanoma, ovarian adenocarcinoma ⁷⁴	integrin receptors ($\alpha_v\beta_3$ and $\alpha_v\beta_5$) and LHRH receptor	β -lactam	mAb-small molecule conjugate	opsonization		monoclonal antibody
prostate cancer ⁷⁵	PSMA	DNP	small molecule	ADCC		commercial
umbilical cord endothelial cells (HUVEC) ⁷⁶	VEGF	β -lactam	RNA aptamer	inhibition of cell migration	female athymic nude mice (pharmacokinetic)	monoclonal antibody
colon adenocarcinoma, breast carcinoma ⁷⁷	VEGF and Ang2	β -lactam (azetidinone)	mAb-small molecule conjugate	inhibition of receptor binding	female athymic nude mice	monoclonal antibody
prostate cancer ⁷⁸	PSMA	DNP	small molecule	ADCC	hu-PBL-NOD/SCID	immunization
colon adenocarcinoma, glioblastoma ⁷⁹	uPAR	DNP	small molecule-protein conjugate	ADCC, phagocytosis		commercial

lactam (Figure 2, 5 \rightarrow 6) functional groups.^{65,94–97} Antibodies specific for the 1,3-diketone functionality can either be administered passively or generated through reactive immunization with a 1,3-diketone-KLH conjugate.⁹⁸

Applications of Antibody Recruiting Small Molecules in Disease Targeting. Advances in synthetic and biophysical chemistry have enabled researchers to construct bifunctional small molecules against a broad variety of structurally unrelated disease-relevant targets. Due to their modular nature, ARMs are able to form immuno-modulatory ternary complexes with various disease-relevant species, simply as a function of the structure and recognition properties of the TBT. Thus, applications of the ARM strategy have included both cancer and infectious disease (Table 1) and suggest a number of additional possibilities for future therapeutic development.

Cancer. Cancer is one of the leading causes of death worldwide and is believed to be responsible for one in every four deaths in the United States.⁹⁹ Traditional treatment options for patients suffering with malignancies include surgical resection, direct irradiation, and cytotoxic chemotherapy, all of which are frequently associated with severe side effects.^{100–103} Recently, biologic agents and cellular immunotherapies have emerged as popular alternatives in cancer treatment, and due to their high specificity, these modalities have the potential to address many of the problems associated with traditional chemotherapies (e.g., off-target effects, etc.).¹⁰⁴ Indeed, with the inclusion of monoclonal antibodies into the repertoire, cancer therapeutics grossed \$18.5 billion in sales in 2009 alone.

Despite these successes, available treatments remain inadequate for most patients suffering with malignant neoplasms, and the demand for novel, targeted therapies is growing.¹⁰⁵ ARM technologies may represent promising alternatives for these patients with the potential both to complement and improve upon available anti-cancer modalities.

Barbas and colleagues were among the first investigators to demonstrate the benefits of combining small molecules with antibody proteins for cancer-relevant applications. These researchers have primarily exploited a “catalytic monoclonal antibody”, mAb 38C2, capable of reacting with the 1,3-diketone motif to form an enaminone (Figure 2, 1 \rightarrow 3; 2 \rightarrow 4). mAb 38C2 has been conjugated to β -diketone-containing TBTs to generate “chemically programmed antibodies” capable of recognizing various cell-surface targets. For example, the conjugate cp38C2 (Figure 2, 3)^{65–67,73} targets the $\alpha_v\beta_3$ and $\alpha_v\beta_5$ integrins,^{106,107} cell-surface proteins that are highly overexpressed in a wide variety of cancers, including ovarian, cervical, breast, and melanoma.^{108,109} This construct has been shown to mediate CDC and ADCC against human M21 melanoma cells in culture and has demonstrated remarkable efficacy in M21-based murine xenografts (81% average reduction in tumor growth after 42 days).⁶⁷ Compound cp38C2 has also been shown to inhibit metastasis of M21 tumor cells in female SCID mice, more than doubling their median survival *versus* untreated controls.

Subsequent studies from the Barbas group eliminated the requirement to use exogenous antibody proteins in their

chemically programmed antibody strategy. Using syngeneic murine colon cancer (CT26) and melanoma (B16) models, it was demonstrated that wild-type BALB/c mice, preimmunized with a diketone hapten (JW-KLH), produced aldolase antibodies capable of forming covalent adducts with synthetic diketone groups. Once “programmed,” these animals were treated with either **1** or cRGD-dk, a cyclic-peptide-based integrin-targeting conjugate, and tumor-specific ADCC responses were observed, resulting in approximately 75% tumor reduction compared with vehicle control.⁷³

Another useful function of small molecule conjugates with aldolase antibodies, termed “CovX-bodies”, has been to improve the pharmacokinetic properties of therapeutic compounds. For example, mAb 38C2 conjugate **4** has been employed as a delivery vehicle for a metabolically labile small-molecule inhibitor of endothelin A (ET_A),⁶⁸ a receptor involved in neovascularization and implicated in cancer, renal failure, heart failure, and hypertension.^{110–112} This conjugate exhibited efficacy in a murine xenograft model of human prostate cancer (PC-3) and demonstrated up to a 45% inhibition of tumor growth *versus* controls. This antibody-attachment strategy has also been employed to minimize toxic side effects of the HIV entry inhibitor Aplavirac⁶¹ and to stabilize peptide-based targeting agents such as VEGF⁷⁷ and angiogenesis inhibitors of thrombospondin-1.¹¹³ Indeed, a CovX-body containing a peptide-based angiopoietin-2 inhibitor (CVX-060) is currently being evaluated in a phase II clinical trial in patients with advanced renal cell carcinoma.^{2,114}

Recent work from the Barbas lab has taken the ARM concept into a variety of novel directions. For example, these authors have demonstrated that the catalytic antibody mAb 38C2 can be conjugated with bifunctional small molecules, enabling them to target two different surface macromolecules: the integrin receptors ($\alpha_v\beta_3$ and $\alpha_v\beta_5$) and the luteinizing hormone releasing hormone receptors (LHRH-R).⁷⁴ More recent investigations have shown that conjugating mAb 38C2 to ARC245 (**5**), an RNA aptamer capable of binding VEGF, can increase the ARC245 serum half-life from minutes to 21 h.⁷⁶ Given the flexibility of available screening methods for identifying selective, high-affinity aptamers,^{115,116} this strategy has the potential to greatly accelerate the process of TBT discovery against numerous disease-relevant targets.

Reports from the Paulson and Bundle groups have disclosed a class of ARMs for targeting B-cell lymphomas through multivalent interactions with CD22,^{71,72} a cell-surface regulator of B-cell signaling overexpressed in malignant cells.¹¹⁷ To this end, a bifunctional molecule (**7**) was constructed containing an *o*-nitro phenol element at the ABT to bind endogenous anti-nitrophenol (anti-NP) antibodies and a glycan sequence at the TBT for binding CD22. By complexing this construct with decavalent anti-NP IgM, these authors demonstrated an increase in binding avidity for CD22-expressing cells of approximately 2 orders of magnitude *versus* control conditions lacking anti-NP IgM. Follow-up studies demonstrated that further increases in TBT valency, through conjugation to a polymer support, led to 100-fold higher levels of anti-NP IgM recruitment to target B-cells *versus* unconjugated compounds.⁷²

The Low group has developed ARMs targeting the folate receptor (FR), a cell-surface protein overexpressed in many cancers, by utilizing folic acid at the TBT and either fluorescein (**8**) or DNP at the ABT.^{62–64} For example, when co-administered with IL-2, construct **8** enhanced median survival by 250% in wild-type female BALB/c mice grafted with FR

positive M109 lung tumors and preimmunized with BSA-FITC or KLH-FITC.⁶² Co-administration of this fluorescein-folate construct with both IL-2 and IFN- γ enhanced survival by 300%. Follow-up studies suggested that antitumor effects resulted from ADCP and ADCC mechanisms,⁶³ and substitution of the fluorescein group with DNP did not diminish efficacy in these models.⁶⁴ Notably, FR-targeted ARMs were found to induce long-lasting immunity against FR-expressing tumors; rechallenge of mice previously treated with M109 tumor cells led to rejection of tumor without introduction of any additional ARM.¹¹⁸ Although details relevant to the mechanism of this memory effect were not reported, depletion of both CD4(+) and CD8(+) T cells prevented protective immunity upon tumor re-challenge, suggesting that ARMs might be useful as therapeutic vaccines,¹¹⁹ as has also been suggested for monoclonal antibody therapies.^{120–123}

Seeking to exploit the widespread prevalence of endogenous anti- α -Gal antibodies in the human bloodstream,⁶⁹ Kiessling and co-workers synthesized bifunctional constructs containing known integrin-binding functionality at the TBT and the α -Gal trisaccharide motif at the ABT (**11**). These ARMs proved effective at inhibiting integrin-mediated cell adhesion, recruiting anti- α -Gal antibodies, and mediating complement-dependent cytotoxicity against various cancer cells using normal human serum as the sole source of anti- α -Gal antibodies and complement proteins.⁷⁰ Cytotoxicity studies comparing this ARM with a toxin-conjugate in which the α -Gal trisaccharide was replaced by doxorubicin revealed the antibody-recruiting agent to be more selective; construct **11** exhibited activity only against cells expressing high levels of integrin, whereas the doxorubicin conjugate proved cytotoxic to cells expressing both high and low levels of integrin. On the basis of these findings, the authors concluded that the high selectivities of ARM-based agents resulted from non-linear increases in antibody binding avidities due to multivalent interactions.

Anti-cancer efforts in the Spiegel laboratory have focused on the development of antibody-recruiting small molecules directed against prostate cancer cells, called ARM-Ps.^{75,124} These bifunctional molecules contain a glutamate urea moiety at the TBT for targeting the prostate specific membrane antigen (PSMA). This surface-bound protein is overexpressed in most subtypes of prostate cancer cells,¹²⁵ as well as in the neovasculature of many solid tumors (e.g., glioblastoma multiforme,¹²⁶ bladder cancer,¹²⁷ gastric and colorectal cancer¹²⁸). An ARM-P derivative containing 8 oxyethylene units in the linker, called ARM-P8 (**9**), was found to possess an optimal compromise between affinity to PSMA and ability to form ternary complex. Follow-up studies demonstrated that ARM-P8 could mediate ADCC against PSMA-expressing prostate cancer cells in the presence of anti-DNP antibodies and peripheral blood mononuclear cells (PBMCs). Interestingly, high concentrations of ARM-P8 were found to serve as auto-inhibitors of ternary complex formation, in both biophysical and cell viability assays. Analogous observations have been made previously in systems involving bifunctional molecules that template ternary linkages, and these observations support ternary complex formation as a necessary precondition for ARM-P8-mediated ADCC.^{63,71} Furthermore, the amount of ternary complex was shown to be directly dependent on the concentration of antibody, indicating that levels of anti-DNP antibody in serum can directly affect the efficacy of ARM-P8.⁷⁵




Figure 3. Bacteria-targeting ARMs, with TBTs in red boxes and ABTs in blue boxes.

DUPA, an ARM-P8 homologue, was recently evaluated in a humanized NOD/SCID mouse model. Thus, NOD/SCID animals transfused with human PBMCs and xenografted with either LNCaP (PSMA-positive) or DU145 (PSMA-negative) tumors were immunized with DNP-KLH.⁷⁸ DUPA administration was found both to inhibit tumor growth and prolong survival in an antibody- and PSMA-dependent fashion. Animals that were not preimmunized against DNP, or that were xenografted with PSMA-negative DU145 tumors, were unaffected by DUPA treatment. Taken together, these studies provide further support for the potential utility of ARMs in clinical applications.

During the course of characterizing ARM-P8, researchers in the Spiegel laboratory observed that ARM-P derivatives with short linker regions displayed surprisingly strong PSMA-binding affinities. Follow-up biochemical, crystallographic, and computational studies led to the serendipitous discovery of an arene-binding site on PSMA that can accommodate electron poor aromatic rings, such as DNP.¹²⁴ Although the interaction of the DNP moiety with the arene-binding site appears to involve only two amino acids, it affords an increase in PSMA binding potency of up to 2 orders of magnitude. Next-generation ARM-Ps that take advantage of this arene-binding site interaction are expected to show significant enhancements in PSMA binding affinity *versus* available derivatives.

More recently, researchers in the Spiegel laboratory synthesized an antibody-recruiting molecule called “ARM-U” (10), which targets the urokinase-type plasminogen activator receptor (uPAR).⁷⁹ uPAR is expressed on the surfaces of breast, colon, stomach, and bladder cancers^{129,130} and has been used as a diagnostic marker for malignancy.^{131–135} ARM-U has been shown to target uPAR at the high affinity uPA binding site, recruit anti-DNP antibodies to uPAR-expressing A172 human glioblastoma cells, and ultimately mediate ADCP and ADCC in an antibody- and uPAR-specific manner. These studies underscore the generality of the ARM strategy for cancer treatment.

Infectious Disease (Bacteria and Viruses). The World Health Organization (WHO) has estimated that infectious

agents (viruses, bacteria, and parasites) are responsible for approximately 25% (15 million) of global deaths each year and are the predominant cause of mortality in developing nations.^{136,137} Vaccines are considered to be among the most successful strategies for fighting infectious disease,¹³⁸ and have been proven effective in combating typhoid, cholera, rabies, measles, mumps, hepatitis B, rubella, tetanus, and polio. Despite these achievements, current vaccination strategies are limited by difficulties in production,¹³⁹ variable levels of immunostimulation, and high costs.^{140–142}

Although traditional small-molecule-based antibacterial and antiviral therapeutics have also shown high efficacy, their utility has been hampered by surges of rapid resistance.^{143,144} Monoclonal antibody-based therapies have demonstrated preclinical success in treating various infectious diseases;^{145–148} however, despite their therapeutic promise, only a small percent of antibodies currently in development are indicated for infectious disease treatment.¹⁴⁸ To date, only one antibody-based antiviral agent (against respiratory syncytial virus, Palivizumab) has obtained FDA approval, and there are no clinically approved monoclonal antibodies that target bacterial pathogens.^{149,150} Thus, there is a critical need to develop new therapeutic strategies for treating infectious disease.

Bacteria. The therapeutic arsenal against bacterial infection has largely consisted of natural products and synthetic small molecules. Conventional antibiotics act by targeting vital bacterial functions such as cell wall synthesis, protein synthesis, RNA transcription, and DNA replication. Because many bacterial species share common essential targets, these agents are often harmful to native flora as well as pathogenic microbes and can increase host susceptibility to certain infections.^{151,152} Furthermore, the emergence of organisms that are resistant to many, if not all, available agents has proven increasingly problematic. Indeed, there has been significant recent interest in the development of monoclonal antibodies for treating drug-resistant bacteria, in part because these agents could exploit mechanisms distinct from conventional antibiotics, making them less likely to induce cross-resistance.¹⁴⁷ Despite this, out

of 13 mAbs currently in clinical development for treating bacterial infection, none have demonstrated significant efficacy.¹⁴⁷ Antibacterial therapeutics with novel mechanisms of action would therefore be highly desirable.^{153–155}

A variety of ARM-based antibacterial strategies have been evaluated. The first example of such an approach was reported by Bednarski and colleagues and employed a rationally designed, bifunctional molecule capable of directing anti-avidin antibodies to *E. coli* (Figure 3, 12).^{53,54} Biotin was conjugated to the C-glycoside of mannose, a known ligand for bacterial mannose receptors, and this construct was shown to recruit anti-avidin antibodies to the surface of *E. coli* in a manner dependent on the presence of conjugate, avidin, and antibodies. These researchers further demonstrated that complexes between avidin, antibody, and ARM could mediate complement- and macrophage-dependent cytotoxicity in a manner competent by α -mannopyranoside. Interestingly, the inherent multivalency of avidin significantly enhanced the millimolar binding affinity of the C-glycoside ligand to the mannose receptor.

Wang and co-workers expanded this concept by developing bifunctional polymers capable of redirecting endogenous anti- α -Gal antibodies to *E. coli*.⁵⁵ Using chemo-enzymatic synthesis, these researchers constructed a polymeric ARM derivative containing poly-mannose as the TBT and poly- α -Gal as the ABT (14). These bifunctional polymers were shown to bind both *E. coli* mannose receptors and endogenous anti- α -Gal antibodies from human serum using competition ELISA experiments.

More recently, Whitesides and co-workers^{56,57} developed ARMs that target pathogenic bacteria by utilizing the potent antibiotic vancomycin. Polyvalent polymers containing fluorescein at the ABT and vancomycin at the TBT were synthesized (13) and shown to redirect anti-fluorescein antibodies to the surface of various Gram-positive bacteria (*S. epidermidis*, *S. pneumoniae*, and *S. aureus*). Using fluorescence microscopy and flow cytometry, the authors demonstrated that the antibody-recruiting polymer could mediate phagocytosis of opsonized bacteria in the presence of anti-fluorescein antibodies.

Viruses. Although vaccine-based strategies have been extremely successful against certain viral diseases, a significant number of viral pathogens have proven refractory to such approaches (e.g., HIV, herpes simplex viruses, etc.).^{156,157} For the most part, available antiviral agents function by inhibiting enzymes such as reverse transcriptase, polymerase, protease, integrase, primase, and neuraminidase,¹⁵⁸ and their utility is limited by resistance development,^{159,160} low efficacy,¹⁶¹ and the high rate of spontaneous mutation inherent to the viral lifecycle.¹⁶² Monoclonal antibody therapies targeting viruses have experienced only modest success, and only a single such agent, which targets respiratory syncytial virus (RSV), has been approved for clinical use.¹⁵⁰ Novel technologies with the potential to harness the endogenous immune response in killing viral pathogens could be profoundly useful in the fight against viral diseases.

An early example of an ARM-based antiviral strategy was described by Wang and colleagues (Figure 4).⁵⁸ Using chemo-enzymatic synthesis, these researchers prepared a bifunctional molecule designed to redirect endogenous anti- α -Gal antibodies to HIV. This agent incorporated the α -Gal trisaccharide epitope at the ABT and was linked to the 36-amino-acid gp41 fusion inhibitory peptide, T-20, at the TBT (15). The authors




Figure 4. Virus-targeting ARMs, with TBTs in red boxes and ABTs in blue boxes.

subsequently demonstrated that functionalization of T-20 had minimal effects on its ability to inhibit virus fusion, and that the bifunctional glycopeptide could bind anti- α -Gal IgG and IgM antibodies from human serum.

More recently, Valhne *et al.* developed a series of bifunctional glycopeptides capable of mediating immune responses against HIV-infected cells.⁵⁹ These constructs were derived by chemically linking the α -Gal disaccharide to a series of 15-mer oligopeptides derived from the gp120-binding region of CD4 (16). Using ELISA and immunofluorescence microscopy, the authors then showed that these bifunctional glycopeptides could redirect endogenous anti- α -Gal antibodies from human serum to both immobilized and cell-surface-expressed gp120. Additional assays then demonstrated that the presence of human antibodies enhanced the fusion inhibitory activity of the peptide by 10% *versus* control conditions. Interestingly, an additional 5–15% enhancement in inhibition was observed when complement-preserved human serum was used in HIV infectivity assays, which was attributed to the direct cytolytic action of complement proteins on HIV-infected cells. Glycopeptide-derived ARMs were also shown to mediate immune responses against chronically HIV IIIB/LAV-infected ACH2 cells in the presence of human serum and isolated natural killer (NK) cells *via* an ADCC mechanism, although some analogues proved cytotoxic even in the absence of NK cells.

The Spiegel laboratory has recently developed a non-peptidic HIV-targeted ARM, called ARM-H.⁶⁰ This bifunctional small molecule incorporates a derivative of the known small-molecule fusion inhibitor BMS-378806 at the TBT,¹⁶³ along with the DNP motif at the ABT (17). ARM-H-mediated formation of ternary complex with anti-DNP antibodies and HIV-1 Env-expressing cells was shown to induce complement-dependent destruction of these cells. Furthermore, ARM-H can bind gp120 competitively with CD4 and also inhibit the entry of HIV-1 virus into human T-cells. Thus, ARM-H has the potential to interfere with the survival of HIV through multiple complementary mechanisms.

In general, by converting virulence factors (e.g., lectins, gp120) into recognition elements for immune-mediated destruction, ARMs have the potential to target various infectious pathogens. Although still in their infancy, such




Figure 5. The development of ARMs emerged from the confluence of many disparate fields of study, including synthetic organic chemistry and immunology. Although still in early stages of development, ARMs have the potential to contribute significantly to basic and clinical sciences.

ARMs could serve as promising alternatives or adjuncts to available immunotherapies, antibiotics, and antiviral agents.

Outlook. The strategies detailed herein underscore the promise of ARM technology in a range of therapeutically relevant contexts. Despite significant progress in this arena, certain obstacles remain before this strategy can be advanced into the clinic. For example, the different ABT types mentioned above are each likely to be associated with unique advantages and disadvantages. Although approaches that exploit endogenous antibodies are anticipated to be the most straightforward to implement in practice, their utility will likely vary among patients as a function of antibody concentrations, affinities, isotype and subisotype distributions, and other factors. Comprehensive, population-wide investigations into the prevalence and properties of known endogenous antibodies, as well as the identification of such species with entirely new binding specificities, will be critical for clinical applications. Preimmunization strategies could afford greater control over these parameters but would involve additional operational complexity, which may also carry increased risks of side effects. Finally, preconjugated antibody–small molecule species can be constructed using a single antibody isotype (e.g., IgG1, a potent inducer of ADCC); however, such agents would likely carry many of the same limitations of available immunotherapies (e.g., dosing *via* injection, immunogenicity, *etc.*). Overall, the optimal ABT will likely depend on the specific patient and/or disease process being targeted.

While most reported TBT motifs interact with relatively well-characterized ligand–receptor systems, discovery strategies to enable unbiased targeting of disease-associated surface proteins could greatly extend the applicability of the ARM approach. To this end, modern techniques in rational ligand design¹⁶⁴ and high-throughput chemistry¹⁶⁵ and biology¹⁶⁶ are likely to prove enabling, and recent strategies for the discovery of novel antibody-binding carbohydrate motifs using array technologies,^{167–169} the systematic identification of antibody biomarkers for both healthy and disease states,¹⁷⁰ and the identification of compounds capable of modulating protein–protein interactions^{171–174} provide cause for optimism along these lines. Finally, novel chemical scaffolds (e.g., for targeting multiple receptors at once)^{71,72,74,175} and assembly strategies (e.g., *in vivo* bioorthogonal chemistry)^{176,177} have the potential to facilitate ARM optimization by improving receptor-binding profiles, decreasing molecular weight, enhancing oral bioavailability, and perhaps through other mechanisms as well.

By exploiting an emerging chemical understanding of complex biological systems, future efforts to rationally modulate human immunological functions have the potential to augment our ability to prevent, diagnose, and treat human disease.³³ ARM-based strategies represent an important step in this direction, bridging mechanistic features of biologic agents with a detailed understanding of small molecule structure and

function (Figure 5). Next-generation immunomodulators have the potential to move beyond the ARMs, enabling precise control over immune responses and contributing to an understanding of the molecular events underlying human disease at the resolution of atoms and molecules.

AUTHOR INFORMATION

Corresponding Author

*E-mail: david.spiegel@yale.edu.

Author Contributions

[§]These authors contributed equally to this work.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors apologize to many wonderful colleagues whose work we were unable to cite due to space limitations. Work in our laboratory on ARMs is supported by the National Institute of Health through the NIH New Innovator Award number 1DP2OD002913-01.

KEYWORDS

antibody-recruiting molecules (ARMs): synthetic bifunctional molecules capable of inducing antibodies to bind disease-relevant proteins, cells, or organisms

synthetic immunology: the creation of synthetic systems that perform complex immunological functions

antibody: protein that is produced by B-cells which identifies and neutralizes disease associated objects

immunomodulators: rationally designed molecules that can affect components of the immune system

immunotherapy: disease treatment that utilizes, induces, suppresses, or enhances components of the immune system

bifunctional molecule: a molecule that possesses two distinct binding targets

ternary complex: an assembly containing three distinct species, held together either through covalent or noncovalent bonds

REFERENCES

- (1) Riedel, S. (2005) Edward Jenner and the history of smallpox and vaccination. *Proc. (Baylor Univ. Med. Cent.)* 18, 21–25.
- (2) Casadevall, A., Dadachova, E., and Pirofski, L. A. (2004) Passive antibody therapy for infectious diseases. *Nat. Rev. Microbiol.* 2, 695–703.
- (3) Elbakri, A., Nelson, P. N., and Abu Odeh, R. O. (2010) The state of antibody therapy. *Hum. Immunol.* 71, 1243–1250.
- (4) Ledford, H. (2010) 'Biosimilar' drugs poised to penetrate market. *Nature* 468, 18–19.
- (5) Aggarwal, S. (2009) What's fueling the biotech engine-2008. *Nat. Biotechnol.* 27, 987–993.

- (6) Reichert, J. M. (2011) Antibody-based therapeutics to watch in 2011. *MAbs* 3, 76–99.
- (7) (2012) FDA Approved Drug Products, In <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm>, U.S. Food and Drug Administration.
- (8) Weiner, L. M., Surana, R., and Wang, S. (2010) Monoclonal antibodies: versatile platforms for cancer immunotherapy. *Nat. Rev. Immunol.* 10, 317–327.
- (9) Forecasting, B. R. M. (2011) Antibody Drugs: Technologies and Global Markets, BIO016H ed., BCC Research Market Forecasting.
- (10) Visintin, M., Melchionna, T., Cannistraci, I., and Cattaneo, A. (2008) In vivo selection of intrabodies specifically targeting protein-protein interactions: A general platform for an “undruggable” class of disease targets. *J. Biotechnol.* 135, 1–15.
- (11) Sunada, H., Magun, B. E., Mendelsohn, J., and MacLeod, C. L. (1986) Monoclonal antibody against epidermal growth factor receptor is internalized without stimulating receptor phosphorylation. *Proc. Natl. Acad. Sci. U.S.A.* 83, 3825–3829.
- (12) Li, S., Schmitz, K. R., Jeffrey, P. D., Wiltzius, J. J. W., Kussie, P., and Ferguson, K. M. (2005) Structural basis for inhibition of the epidermal growth factor receptor by cetuximab. *Cancer Cell* 7, 301–311.
- (13) Capala, J., and Bouchelouche, K. (2010) Molecular imaging of HER2-positive breast cancer: a step toward an individualized ‘image and treat’ strategy. *Curr. Opin. Oncol.* 22, 559–566.
- (14) Francisco, J. A., Cervený, C. G., Meyer, D. L., Mixan, B. J., Klussman, K., Chace, D. F., Rejniak, S. X., Gordon, K. A., DeBlanc, R., Toki, B. E., Law, C.-L., Doronina, S. O., Siegall, C. B., Senter, P. D., and Wahl, A. F. (2003) cAC10-vcMMAE, an anti-CD30-monomethyl auristatin E conjugate with potent and selective antitumor activity. *Blood* 102, 1458–1465.
- (15) Alley, S. C., Okeley, N. M., and Senter, P. D. (2010) Antibody-drug conjugates: Targeted drug delivery for cancer. *Curr. Opin. Chem. Biol.* 14, 529–537.
- (16) Clynes, R., Takechi, Y., Moroi, Y., Houghton, A., and Ravetch, J. V. (1998) Fc receptors are required in passive and active immunity to melanoma. *Proc. Natl. Acad. Sci. U.S.A.* 95, 652–656.
- (17) Clynes, R. A., Towers, T. L., Presta, L. G., and Ravetch, J. V. (2000) Inhibitory Fc receptors modulate in vivo cytotoxicity against tumor targets. *Nat. Med.* 6, 443–446.
- (18) Sampson, J. H., Crotty, L. E., Lee, S., Archer, G. E., Ashley, D. M., Wikstrand, C. J., Hale, L. P., Small, C., Dranoff, G., Friedman, A. H., Friedman, H. S., and Bigner, D. D. (2000) Unarmed, tumor-specific monoclonal antibody effectively treats brain tumors. *Proc. Natl. Acad. Sci. U.S.A.* 97, 7503–7508.
- (19) Murphy, K. P., Travers, P., Walport, M., and Janeway, C. (2008) *Janeway’s Immunobiology*, Garland Science, New York
- (20) Nimmerjahn, F., and Ravetch, J. V. (2008) Fcγ receptors as regulators of immune responses. *Nat Rev Immunol* 8, 34–47.
- (21) Dhodapkar, K. M., Krasovský, J., Williamson, B., and Dhodapkar, M. V. (2002) Antitumor monoclonal antibodies enhance cross-presentation of cellular antigens and the generation of myeloma-specific killer T cells by dendritic cells. *J. Exp. Med.* 195, 125–133.
- (22) Akiyama, K., Ebihara, S., Yada, A., Matsumura, K., Aiba, S., Nukiwa, T., and Takai, T. (2003) Targeting apoptotic tumor cells to FcγR provides efficient and versatile vaccination against tumors by dendritic cells. *J. Immunol.* 170, 1641–1648.
- (23) Cioca, D. P., Deak, E., Cioca, F., and Paunescu, V. (2005) Monoclonal antibodies targeted against melanoma and ovarian tumors enhance dendritic cell-mediated cross-presentation of tumor-associated antigens and efficiently cross-prime CD8+ T cells. *J. Immunother.* 29, 41–52.
- (24) Murillo, O., Dubrot, J., Palazon, A., Arina, A., Azpilikueta, A., Alfaro, C., Solano, S., Ochoa, M. C., Berasain, C., Gabari, I., Perez-Gracia, J. L., Berraondo, P., Hervás-Stubbs, S., and Melero, I. (2009) In vivo depletion of DC impairs the anti-tumor effect of agonistic anti-CD137 mAb. *Eur. J. Immunol.* 39, 2424–2436.
- (25) Hansel, T. T., Kropshofer, H., Singer, T., Mitchell, J. A., and George, A. J. T. (2010) The safety and side effects of monoclonal antibodies. *Nat. Rev. Drug Discovery* 9, 325–338.
- (26) Getts, D. R., Getts, M. T., McCarthy, D. P., Chastain, E. M., and Miller, S. D. (2010) Have we overestimated the benefit of human(ized) antibodies? *mAbs* 2, 682–694.
- (27) Harding, F. A., Stickler, M. M., Razo, J., and DuBridge, R. (2010) The immunogenicity of humanized and fully human antibodies: Residual immunogenicity resides in the CDR regions. *mAbs* 2, 256–265.
- (28) Siberil, S., Dutertre, C.-A., Fridman, W.-H., and Teillaud, J.-L. (2007) FcγR: The key to optimize therapeutic antibodies? *Crit. Rev. Oncol. Hematol.* 62, 26–33.
- (29) Wu, A., and Senter, P. (2005) Arming antibodies: prospects and challenges for immunoconjugates. *Nat. Biotechnol.* 23, 1137–1146.
- (30) Allen, T. M. (2002) Ligand-targeted therapeutics in anticancer therapy. *Nat. Rev. Cancer* 2, 750–763.
- (31) Imai, K., and Takaoka, A. (2006) Comparing antibody and small-molecule therapies for cancer. *Nat. Rev. Cancer* 6, 714–727.
- (32) Pashine, A., Valiante, N. M., and Ulmer, J. B. (2005) Targeting the innate immune response with improved vaccine adjuvants. *Nat. Med.* 11, S63–S68.
- (33) Spiegel, D. A. (2010) Grand Challenge Commentary: Synthetic immunology to engineer human immunity. *Nat. Chem. Biol.* 6, 871–872.
- (34) Kodadek, T. (2010) Development of antibody surrogates for the treatment of cancers and autoimmune disease. *Curr. Opin. Chem. Biol.* 14, 721–727.
- (35) Ducry, L., and Stump, B. (2010) Antibody–drug conjugates: Linking cytotoxic payloads to monoclonal antibodies. *Bioconjugate Chem.* 21, 5–13.
- (36) Müller, D., and Kontermann, R. E. (2010) Bispecific antibodies for cancer immunotherapy: Current perspectives. *BioDrugs* 24, 89–98.
- (37) Baeuerle, P. A., and Reinhardt, C. (2009) Bispecific T-cell engaging antibodies for cancer therapy. *Cancer Res.* 69, 4941–4944.
- (38) Xiao, J., Chen, R., Pawlicki, M. A., and Tolbert, T. J. (2009) Targeting a homogeneously glycosylated antibody Fc to bind cancer cells using a synthetic receptor ligand. *J. Am. Chem. Soc.* 131, 13616–13618.
- (39) Morelli, L., Poletti, L., and Lay, L. (2011) Carbohydrates and immunology: Synthetic oligosaccharide antigens for vaccine formulation. *Eur. J. Org. Chem.* 2011, 5723–5777.
- (40) Li, Q., So, C. R., Fegan, A., Cody, V., Sarikaya, M., Vallera, D. A., and Wagner, C. R. (2010) Chemically self-assembled antibody nanorings (CSANs): Design and characterization of an anti-CD3 IgM biomimetic. *J. Am. Chem. Soc.* 132, 17247–17257.
- (41) Li, Q., Hapka, D., Chen, H., Vallera, D. A., and Wagner, C. R. (2008) Self-Assembly of Antibodies by Chemical Induction. *Angew. Chem., Int. Ed.* 47, 10179–10182.
- (42) Kitov, P. I., Mulvey, G. L., Griener, T. P., Lipinski, T., Solomon, D., Paszkiewicz, E., Jacobson, J. M., Sadowska, J. M., Suzuki, M., Yamamura, K.-I., Armstrong, G. D., and Bundle, D. R. (2008) In vivo supramolecular templating enhances the activity of multivalent ligands: a potential therapeutic against the Escherichia coli O157 AB5 toxins. *Proc. Natl. Acad. Sci. U.S.A.* 105, 16837–16842.
- (43) Kitov, P. I., Lipinski, T., Paszkiewicz, E., Solomon, D., Sadowska, J. M., Grant, G. A., Mulvey, G. L., Kitova, E. N., Klassen, J. S., Armstrong, G. D., and Bundle, D. R. (2008) An entropically efficient supramolecular inhibition strategy for shiga toxins. *Angew. Chem., Int. Ed.* 47, 672–676.
- (44) Carlson, J. C. T., Kanter, A., Thuduppathy, G. R., Cody, V., Pineda, P. E., Mccivor, R. S., and Wagner, C. R. (2003) Designing protein dimers: the importance of ligand conformational equilibria. *J. Am. Chem. Soc.* 125, 1501–1507.
- (45) Shearer, G. M. (1974) Cell mediated cytotoxicity to trinitrophenyl-modified syngeneic lymphocytes. *Eur. J. Immunol.* 4, 527–533.
- (46) Nelson, D. L., Poplack, D. G., Holiman, B. J., and Henkart, P. A. (1979) ADCC against human erythrocyte target cells: role of the anti-

target cell antibodies in determining lymphocyte killer activity. *Clin. Exp. Immunol.* 35, 447–453.

(47) Six, H. R., Uemura, K., and Kinsky, S. C. (1973) Effect of immunoglobulin class and affinity on the initiation of complement-dependent damage to liposomal model membranes sensitized with dinitrophenylated phospholipids. *Biochemistry* 12, 4003–4011.

(48) Capon, D. J., Chamow, S. M., Mordenti, J., Marsters, S. A., Gregory, T., Mitsuya, H., Byrn, R. A., Lucas, C., Wurm, F. M., Groopman, J. E., et al. (1989) Designing CD4 immunoadhesins for AIDS therapy. *Nature* 337, 525–531.

(49) Trauneker, A., Schneider, J., Kiefer, H., and Karjalainen, K. (1989) Highly efficient neutralization of HIV with recombinant CD4-immunoglobulin molecules. *Nature* 339, 68–70.

(50) Byrn, R. A., Mordenti, J., Lucas, C., Smith, D., Marsters, S. A., Johnson, J. S., Cossup, P., Chamow, S. M., Wurm, F. M., et al. (1990) Biological properties of a CD4 immunoadhesin. *Nature* 344, 667–670.

(51) Shokat, K. M., and Schultz, P. G. (1991) Redirecting the immune response: ligand-mediated immunogenicity. *J. Am. Chem. Soc.* 113, 1861–1862.

(52) Lussow, A. R., Buelow, R., Fanget, L., Peretto, S., Gao, L., and Pouletty, P. (1996) Redirecting circulating antibodies via ligand-hapten conjugates eliminates target cells in vivo. *J. Immunother. Emphasis Tumor Immunol.* 19, 257–265.

(53) Bertozzi, C. R., and Bednarski, M. D. (1992) Antibody targeting to bacterial cells using receptor-specific ligands. *J. Am. Chem. Soc.* 114, 2242–2245.

(54) Bertozzi, C. R., and Bednarski, M. D. (1992) A receptor-mediated immune response using synthetic glycoconjugates. *J. Am. Chem. Soc.* 114, 5543–5546.

(55) Li, J., Zacharek, S., Chen, X., Wang, J., Zhang, W., Janczuk, A., and Wang, P. G. (1999) Bacteria targeted by human natural antibodies using alpha-gal conjugated receptor-specific glycopolymers. *Bioorg. Med. Chem.* 7, 1549–1558.

(56) Metallo, S. J., Kane, R. S., Holmlin, R. E., and Whitesides, G. M. (2003) Using bifunctional polymers presenting vancomycin and fluorescein groups to direct anti-fluorescein antibodies to self-assembled monolayers presenting D-alanine-D-alanine groups. *J. Am. Chem. Soc.* 125, 4534–4540.

(57) Krishnamurthy, V. M., Quinton, L. J., Estroff, L. A., Metallo, S. J., Isaacs, J. M., Mizgerd, J. P., and Whitesides, G. M. (2006) Promotion of opsonization by antibodies and phagocytosis of Gram-positive bacteria by a bifunctional polyacrylamide. *Biomaterials* 27, 3663–3674.

(58) Naicker, K. P., Li, H., Heredia, A., Song, H., and Wang, L.-X. (2004) Design and synthesis of alpha-Gal-conjugated peptide T20 as novel antiviral agent for HIV-immunotargeting. *Org. Biomol. Chem.* 2, 660–664.

(59) Perdomo, M. F., Levi, M., Saellberg, M., and Vahne, A. (2008) Neutralization of HIV-1 by redirection of natural antibodies. *Proc. Natl. Acad. Sci. U.S.A.* 105, 12515–12520.

(60) Parker, C. G., Domaal, R. A., Anderson, K. S., and Spiegel, D. A. (2009) An antibody-recruiting small molecule that targets HIV gp120. *J. Am. Chem. Soc.* 131, 16392–16394.

(61) Gavriluk, J., Uehara, H., Otsubo, N., Hessel, A., Burton, D. R., and Barbas, C. F., 3rd (2010) Potent inhibition of HIV-1 entry with a chemically programmed antibody aided by an efficient organocatalytic synthesis. *ChemBioChem* 11, 2113–2118.

(62) Lu, Y., and Low, P. S. (2002) Folate targeting of haptens to cancer cell surfaces mediates immunotherapy of syngeneic murine tumors. *Cancer Immunol. Immunother.* 51, 153–162.

(63) Lu, Y., Segal, E., and Low, Philip, S. (2005) Folate receptor-targeted immunotherapy: induction of humoral and cellular immunity against hapten-decorated cancer cells. *Int. J. Cancer.* 116, 710–719.

(64) Lu, Y., You, F., Vlahov, I., Westrick, E., Fan, M., Low, P. S., and Leamon, C. P. (2007) Folate-targeted dinitrophenyl hapten immunotherapy: Effect of linker chemistry on antitumor activity and allergic potential. *Mol. Pharmaceutics* 4, 695–706.

(65) Rader, C., Sinha, S. C., Popkov, M., Lerner, R. A., and Barbas, C. F., III (2003) Chemically programmed monoclonal antibodies for

cancer therapy: Adaptor immunotherapy based on a covalent antibody catalyst. *Proc. Natl. Acad. Sci. U.S.A.* 100, 5396–5400.

(66) Li. (2004) Chemical adaptor immunotherapy: Design, synthesis, and evaluation of novel integrin-targeting devices. *J. Med. Chem.* 47, 5630–5640.

(67) Popkov, M., Rader, C., Gonzalez, B., Sinha, S. C., and Barbas, C. F., III (2006) Small molecule drug activity in melanoma models may be dramatically enhanced with an antibody effector. *Int. J. Cancer* 119, 1194–1207.

(68) Doppalapudi, V., Tryder, N., Li, L., Aja, T., Griffith, D., Liao, F.-F., Roxas, G., Ramprasad, M., Bradshaw, C., and Barbas, C. (2007) Chemically programmed antibodies: endothelin receptor targeting CovX-Bodies. *Bioorg. Med. Chem. Lett.* 17, 501–506.

(69) Owen, R. M., Carlson, C. B., Xu, J., Mowery, P., Fasella, E., and Kiessling, L. L. (2007) Bifunctional ligands that target cells displaying the alpha v beta 3 integrin. *ChemBioChem* 8, 68–82.

(70) Carlson, C. B., Mowery, P., Owen, R. M., Dykhuizen, E. C., and Kiessling, L. L. (2007) Selective tumor cell targeting using low-affinity, multivalent interactions. *ACS Chem. Biol.* 2, 119–127.

(71) O'Reilly, M. K., Collins, B. E., Han, S., Liao, L., Rillahan, C., Kitov, P. I., Bundle, D. R., and Paulson, J. C. (2008) Bifunctional CD22 ligands use multimeric immunoglobulins as protein scaffolds in assembly of immune complexes on B cells. *J. Am. Chem. Soc.* 130, 7736–7745.

(72) Cui, L., Kitov, P. I., Completo, G. C., Paulson, J. C., and Bundle, D. R. (2011) Supramolecular complexing of membrane siglec CD22 mediated by a polyvalent heterobifunctional ligand that templates on IgM. *Bioconjugate Chem.* 22, 546–550.

(73) Popkov, M., Gonzalez, B., Sinha, S. V., and Barbas, C. F., III (2009) Instant immunity through chemically programmable vaccination and covalent self-assembly. *Proc. Natl. Acad. Sci. U.S.A.* 106, 4378–4383.

(74) Gavriluk, J., Wuellner, U., Salahuddin, S., Goswami, R., Sinha, S., and Barbas, C. (2009) An efficient chemical approach to bispecific antibodies and antibodies of high valency. *Bioorg. Med. Chem. Lett.* 19, 3716–3720.

(75) Murelli, R. P., Zhang, A. X., Michel, J., Jorgensen, W. L., and Spiegel, D. A. (2009) Chemical control over immune recognition: A class of antibody-recruiting small molecules that target prostate cancer. *J. Am. Chem. Soc.* 131, 17090–17092.

(76) Wuellner, U., Gavriluk, J., and Barbas, C. (2010) Expanding the concept of chemically programmable antibodies to RNA aptamers: chemically programmed biotherapeutics. *Angew. Chem., Int. Ed.* 49, 5934–5937.

(77) Doppalapudi, V. R., Huang, J., Liu, D., Jin, P., Liu, B., Li, L., Desharnais, J., Hagen, C., Levin, N. J., Shields, M. J., Parish, M., Murphy, R. E., Del Rosario, J., Oates, B. D., Lai, J.-Y., Matin, M. J., Ainekulu, Z., Bhat, A., Bradshaw, C. W., Woodnutt, G., Lerner, R. A., and Lappe, R. W. (2010) Chemical generation of bispecific antibodies. *Proc. Natl. Acad. Sci. U.S.A.* 107, 22611–22616.

(78) Dubrovskaya, A., Kim, C., Elliott, J., Shen, W., Kuo, T.-H., Koo, D.-I., Li, C., Tuntland, T., Chang, J., Groessl, T., Wu, X., Gorney, V., Ramirez Montagut, T., Spiegel, D., Cho, C., and Schultz, P. (2011) A chemically induced vaccine strategy for prostate cancer. *ACS Chem. Biol.* 6, 1223–1231.

(79) Jakobsche, C. E., McEnaney, P. J., Zhang, A. X., and Spiegel, D. A. (2011) Reprogramming urokinase into an antibody-recruiting anticancer agent. *ACS Chem. Biol.* 7, 316–321.

(80) Galili, U., Rachmilewitz, E. A., Peleg, A., and Flechner, I. (1984) A unique natural human IgG antibody with anti-alpha-galactosyl specificity. *J. Exp. Med.* 160, 1519–1531.

(81) Parker, W., Bruno, D., Holzknicht, Z. E., and Platt, J. L. (1994) Characterization and affinity isolation of xenoreactive human natural antibodies. *J. Immunol.* 153, 3791–3803.

(82) Galili, U., Clark, M. R., Shohet, S. B., Buehler, J., and Macher, B. A. (1987) Evolutionary relationship between the natural anti-Gal antibody and the Galalpha 1 -> 3Gal epitope in primates. *Proc. Natl. Acad. Sci. U.S.A.* 84, 1369–1373.

- (83) Ortega, E., Kostovetzky, M., and Larralde, C. (1984) Natural DNP-binding immunoglobulins and antibody multispecificity. *Mol. Immunol.* 21, 883–888.
- (84) Farah, F. S. (1973) Natural antibodies specific to the 2,4-dinitrophenyl group. *Immunology* 25, 217–226.
- (85) Ju, K.-S., and Parales, R. (2010) Nitroaromatic compounds, from synthesis to biodegradation. *Microbiol. Mol. Biol. Rev.* 74, 250.
- (86) 2,4-Dinitrophenol (CASRN 51-28-5), In <http://www.epa.gov/ncea/iris/subst/0152.htm>, United States Environmental Protection Agency.
- (87) Harris, M. O., Cocoran, J. J. (1995) Toxicological Profile for Dinitrophenols, Agency for Toxic Substances and Disease Registry. <http://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=729&tid=132>.
- (88) Lauer, K. (1990) Environmental nitrophenols and autoimmunity. *Mol. Immunol.* 27, 697–698.
- (89) Gauba, V., Grünewald, J., Gorney, V., Deaton, L. M., Kang, M., Bursulaya, B., Ou, W., Lerner, R. A., Schmedt, C., Geierstanger, B. H., Schultz, P. G., and Ramirez-Montagut, T. (2011) Loss of CD4 T-cell-dependent tolerance to proteins with modified amino acids. *Proc. Natl. Acad. Sci. U.S.A.* 108, 12821–12826.
- (90) Grünewald, J., Hunt, G. S., Dong, L., Niessen, F., Wen, B. G., Tsao, M.-L., Perera, R., Kang, M., Laffitte, B. A., Azarian, S., Ruf, W., Nasoff, M., Lerner, R. A., Schultz, P. G., and Smider, V. V. (2009) Mechanistic studies of the immunochemical termination of self-tolerance with unnatural amino acids. *Proc. Natl. Acad. Sci. U.S.A.* 106, 4337–4342.
- (91) Grünewald, J., Tsao, M.-L., Perera, R., Dong, L., Niessen, F., Wen, B. G., Kubitz, D. M., Smider, V. V., Ruf, W., Nasoff, M., Lerner, R. A., and Schultz, P. G. (2008) Immunochemical termination of self-tolerance. *Proc. Natl. Acad. Sci. U.S.A.* 105, 11276–11280.
- (92) Palm, N. W., and Medzhitov, R. (2009) Immunostimulatory activity of haptened proteins. *Proc. Natl. Acad. Sci. U.S.A.* 106, 4782–4787.
- (93) Cherukuri, A., Durack, G., and Voss, E. W., Jr. (1997) Evidence for hapten recognition in receptor-mediated intracellular uptake of a hapten-protein conjugate by murine macrophage. *Mol. Immunol.* 34, 21–32.
- (94) Wagner, J., Lerner, R. A., and Barbas, C. F., III (1995) Efficient aldolase catalytic antibodies that use the enamine mechanism of natural enzymes. *Science* 270, 1797–1800.
- (95) Barbas, C. F., Heine, A., Zhong, G., Hoffmann, T., Gramatikova, S., Bjrnstedt, R., List, B., Anderson, J., Stura, E. A., Wilson, I. A., and Lerner, R. A. (1997) Immune versus natural selection: antibody aldolases with enzymic rates but broader scope. *Science* 278, 2085–2092.
- (96) Guo, F., Das, S., Mueller, B., Barbas, C., Lerner, R., and Sinha, S. (2006) Breaking the one antibody-one target axiom. *Proc. Natl. Acad. Sci. U.S.A.* 103, 11009–11014.
- (97) Gavriluyk, J., Wuellner, U., and Barbas, C. (2009) Beta-lactam-based approach for the chemical programming of aldolase antibody 38C2. *Bioorg. Med. Chem. Lett.* 19, 1421–1424.
- (98) Tanaka, F., and Barbas, C. F. (2002) Reactive immunization: a unique approach to catalytic antibodies. *J. Immunol. Methods* 269, 67–79.
- (99) Jemal, A., Siegel, R., Xu, J., and Ward, E. (2010) Cancer statistics, 2010. *CA Cancer J. Clin.* 60, 277–300.
- (100) Chabner, B. A., and Roberts, T. G. (2005) Chemotherapy and the war on cancer. *Nat. Rev. Cancer* 5, 65–72.
- (101) Helleday, T., Petermann, E., Lundin, C., Hodgson, B., and Sharma, R. (2008) DNA repair pathways as targets for cancer therapy. *Nat. Rev. Cancer* 8, 193–204.
- (102) Tennant, D., Duran, R., and Gottlieb, E. (2010) Targeting metabolic transformation for cancer therapy. *Nat. Rev. Cancer* 10, 267–277.
- (103) Zhang, J., Yang, P., and Gray, N. (2009) Targeting cancer with small molecule kinase inhibitors. *Nat. Rev. Cancer* 9, 28–39.
- (104) Drake, C. G. (2010) Prostate cancer as a model for tumour immunotherapy. *Nat. Rev. Immunol.* 10, 580–593.
- (105) Aggarwal, S. (2010) Targeted cancer therapies. *Nat. Rev. Drug Discovery* 9, 427–428.
- (106) Nemeth, J. A., Nakada, M. T., Trikha, M., Lang, Z., Gordon, M. S., Jayson, G. C., Corringham, R., Prabhakar, U., Davis, H. M., and Beckman, R. A. (2007) Alpha-v integrins as therapeutic targets in oncology. *Cancer Invest.* 25, 632–646.
- (107) Felding-Habermann, B. (2003) Integrin adhesion receptors in tumor metastasis. *Clin. Exp. Metastasis* 20, 203–213.
- (108) Chattopadhyay, N., and Chatterjee, A. (2001) Studies on the expression of alpha(v)beta3 integrin receptors in non-malignant and malignant human cervical tumor tissues. *J. Exp. Clin. Cancer Res.* 20, 269–275.
- (109) Wong, N. C., Mueller, B. M., Barbas, C. F., Ruminski, P., Quaranta, V., Lin, E. C., and Smith, J. W. (1998) Alphav integrins mediate adhesion and migration of breast carcinoma cell lines. *Clin. Exp. Metastasis* 16, 50–61.
- (110) Nelson, J., Bagnato, A., Battistini, B., and Nisen, P. (2003) The endothelin axis: emerging role in cancer. *Nat. Rev. Cancer* 3, 110–116.
- (111) Remuzzi, G., Perico, N., and Benigni, A. (2002) New therapeutics that antagonize endothelin: promises and frustrations. *Nat. Rev. Drug Discovery* 1, 986–1001.
- (112) Attin, T., Camidge, R., Newby, D. E., and Webb, D. J. (2005) Endothelin antagonism in pulmonary hypertension, heart failure, and beyond. *Heart* 91, 825–831.
- (113) Li, L. N., Leedom, T. A., Do, J., Huang, H. H., Lai, J. Y., Johnson, K., Osothprarop, T. F., Rizzo, J. D., Doppalapudi, V. R., Bradshaw, C. W., Lappe, R. W., Woodnutt, G., Levin, N. J., and Pirie-Shepherd, S. R. (2011) Antitumor efficacy of a thrombospondin 1 mimetic CovX-Body. *Transl. Oncol.* 4, 239–247.
- (114) Huang, H., Lai, J.-Y., Do, J., Liu, D., Li, L., Del Rosario, J., Doppalapudi, V. R., Pirie-Shepherd, S., Levin, N., Bradshaw, C., Woodnutt, G., Lappe, R., and Bhat, A. (2011) Specifically targeting angiopoietin-2 inhibits angiogenesis, Tie2-expressing monocyte infiltration, and tumor growth. *Clin. Cancer Res.* 17, 1001–1011.
- (115) Ellington, A. D., and Szostak, J. W. (1990) In vitro selection of RNA molecules that bind specific ligands. *Nature* 346, 818–822.
- (116) Tuerk, C., and Gold, L. (1990) Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science* 249, 505–510.
- (117) Crocker, P. R., Paulson, J. C., and Varki, A. (2007) Siglecs and their roles in the immune system. *Nat. Rev. Immunol.* 7, 255–266.
- (118) Lu, Y., Segal, E., Leamon, C., and Low, P. (2004) Folate receptor-targeted immunotherapy of cancer: mechanism and therapeutic potential. *Adv. Drug Delivery Rev.* 56, 1161–1176.
- (119) Chames, P., Van Regenmortel, M., Weiss, E., and Baty, D. (2009) Therapeutic antibodies: successes, limitations and hopes for the future. *Br. J. Pharmacol.* 157, 220–233.
- (120) Ellenhorn, J. D., Schreiber, H., and Bluestone, J. A. (1990) Mechanism of tumor rejection in anti-CD3 monoclonal antibody-treated mice. *J. Immunol.* 144, 2840–2846.
- (121) Runyon, K., Lee, K., Zuberek, K., Collins, M., Leonard, J. P., and Dunussi Joannopoulos, K. (2001) The combination of chemotherapy and systemic immunotherapy with soluble B7-immunoglobulin G leads to cure of murine leukemia and lymphoma and demonstration of tumor-specific memory responses. *Blood* 97, 2420–2426.
- (122) Ruf, P., and Lindhofer, H. (2001) Induction of a long-lasting antitumor immunity by a trifunctional bispecific antibody. *Blood* 98, 2526–2534.
- (123) Morecki, S., Lindhofer, H., Yacovlev, E., Gelfand, Y., Ruf, P., and Slavin, S. (2008) Induction of long-lasting antitumor immunity by concomitant cell therapy with allogeneic lymphocytes and trifunctional bispecific antibody. *Exp. Hematol.* 36, 997–1003.
- (124) Zhang, A., Murelli, R., Barinka, C., Michel, J., Cocleaza, A., Jorgensen, W., Lubkowski, J., and Spiegel, D. (2010) A remote arene-binding site on prostate specific membrane antigen revealed by antibody-recruiting small molecules. *J. Am. Chem. Soc.* 132, 12711–12716.

- (125) Holmes, E. H., Greene, T. G., Tino, W. T., Boynton, A. L., Aldape, H. C., Misrock, S. L., and Murphy, G. P. (1996) Analysis of glycosylation of prostate-specific membrane antigen derived from LNCaP cells, prostatic carcinoma tumors, and serum from prostate cancer patients. *Prostate Suppl.* 7, 25–29.
- (126) Wernicke, A. G., Edgar, M., Lavi, E., Liu, H., Salerno, P., Bander, N., and Gutin, P. (2011) Prostate-specific membrane antigen as a potential novel vascular target for treatment of glioblastoma multiforme. *Archi. Pathol. Lab. Med.* 135, 1486–1489.
- (127) Samplaski, M., Heston, W., Elson, P., Magi Galluzzi, C., and Hansel, D. (2011) Folate hydrolase (prostate-specific antigen) 1 expression in bladder cancer subtypes and associated tumor neovasculature. *Mod. Pathol.* 24, 1521–1529.
- (128) Haffner, M., Kronberger, I., Ross, J., Sheehan, C., Zitt, M., Mhlmann, G., Ofner, D., Zelger, B., Ensinger, C., Yang, X., Geley, S., Margreiter, R., and Bander, N. (2009) Prostate-specific membrane antigen expression in the neovasculature of gastric and colorectal cancers. *Hum. Pathol.* 40, 1754–1761.
- (129) Blasi, F., and Carmeliet, P. (2002) uPAR: a versatile signalling orchestrator. *Nat. Rev. Mol. Cell Biol.* 3, 932–943.
- (130) Romer, J., Nielsen, B., and Ploug, M. (2004) The urokinase receptor as a potential target in cancer therapy. *Curr. Pharm. Des.* 10, 2359–2376.
- (131) Andreassen, P. A., Kjller, L., Christensen, L., and Duffy, M. J. (1997) The urokinase-type plasminogen activator system in cancer metastasis: a review. *Int. J. Cancer* 72, 1–22.
- (132) Duffy, M. J., O'Grady, P., Devaney, D., O'Siorain, L., Fennelly, J. J., and Lijnen, H. J. (1988) Urokinase-plasminogen activator, a marker for aggressive breast carcinomas. Preliminary report. *Cancer* 62, 531–533.
- (133) Sier, C. F., Stephens, R., Bizik, J., Mariani, A., Bassan, M., Pedersen, N., Frigerio, L., Ferrari, A., Dan, K., Brnner, N., and Blasi, F. (1998) The level of urokinase-type plasminogen activator receptor is increased in serum of ovarian cancer patients. *Cancer Res.* 58, 1843–1849.
- (134) Harbeck, N., Kates, R., Gauger, K., Willems, A., Kiechle, M., Magdolen, V., and Schmitt, M. (2004) Urokinase-type plasminogen activator (uPA) and its inhibitor PAI-I: novel tumor-derived factors with a high prognostic and predictive impact in breast cancer. *Thrombosis Haemostasis* 91, 450–456.
- (135) Konecny, G., Untch, M., Pihan, A., Kimmig, R., Gropp, M., Stieber, P., Hepp, H., Slamon, D., and Pegram, M. (2001) Association of urokinase-type plasminogen activator and its inhibitor with disease progression and prognosis in ovarian cancer. *Clin. Cancer Res.* 7, 1743–1749.
- (136) Mathers, C., Boerma, T., Ma Fat, D. (2008) *The Global Burden of Disease*, 2004 update, ed., p 146, World Health Organization, Geneva, Switzerland.
- (137) Rappuoli, R., Mandl, C. W., Black, S., and De Gregorio, E. (2011) Vaccines for the twenty-first century society. *Nat. Rev. Immunol.* 11, 865–872.
- (138) Hilleman, M. R. (2000) Vaccines in historic evolution and perspective: a narrative of vaccine discoveries. *Vaccine* 18, 1436–1447.
- (139) Ulmer, J. B., Valley, U., and Rappuoli, R. (2006) Vaccine manufacturing: challenges and solutions. *Nat. Biotechnol.* 24, 1377–1383.
- (140) Plotkin, S. A. (2009) Vaccines: the fourth century. *Clin. Vaccine Immunol.* 16, 1709–1719.
- (141) Guy, B. (2007) The perfect mix: recent progress in adjuvant research. *Nat. Rev. Microbiol.* 5, 505–517.
- (142) Sheridan, C. (2005) The business of making vaccines. *Nat. Biotechnol.* 23, 1359–1366.
- (143) Palumbi, S. R. (2001) Humans as the world's greatest evolutionary force. *Science* 293, 1786–1790.
- (144) Razonable, R. R. (2011) Antiviral drugs for viruses other than human immunodeficiency virus. *Mayo Clin. Proc.* 86, 1009–1026.
- (145) Saylor, C., Dadachova, E., and Casadevall, A. (2009) Monoclonal antibody-based therapies for microbial diseases. *Vaccine* 27 (Suppl 6), G38–46.
- (146) Nagy, E., Giefing, C., and von Gabain, A. (2008) Anti-infective antibodies: a novel tool to prevent and treat nosocomial diseases. *Expert Rev. Anti-Infect. Ther.* 6, 21–30.
- (147) Bebbington, C., and Yarranton, G. (2008) Antibodies for the treatment of bacterial infections: current experience and future prospects. *Curr. Opin. Biotechnol.* 19, 613–619.
- (148) ter Meulen, J. (2007) Monoclonal antibodies for prophylaxis and therapy of infectious diseases. *Expert Opin. Emerging Drugs* 12, 525–540.
- (149) Xiao, X., and Dimitrov, D. S. (2007) Monoclonal antibodies against viruses and bacteria: a survey of patents. *Recent Pat. Antiinfect. Drug Discovery* 2, 171–177.
- (150) Jan, t. M. (2011) Monoclonal antibodies in infectious diseases: Clinical pipeline in 2011. *Infect. Dis. Clin. North Am.* 25, 789–802.
- (151) Sekirov, I., Tam, N. M., Jogova, M., Robertson, M. L., Li, Y., Lupp, C., and Finlay, B. B. (2008) Antibiotic-induced perturbations of the intestinal microbiota alter host susceptibility to enteric infection. *Infect. Immun.* 76, 4726–4736.
- (152) Willing, B. P., Russell, S. L., and Finlay, B. B. (2011) Shifting the balance: antibiotic effects on host-microbiota mutualism. *Nat. Rev. Microbiol.* 9, 233–243.
- (153) Walsh, C. (2000) Molecular mechanisms that confer antibacterial drug resistance. *Nature* 406, 775–781.
- (154) Nikaido, H. (2009) Multidrug resistance in bacteria. *Annu. Rev. Biochem.* 78, 119–146.
- (155) Cooper, M. A., and Shlaes, D. (2011) Fix the antibiotics pipeline. *Nature* 472, 32.
- (156) Huang, D. B., Wu, J. J., and Tyring, S. K. (2004) A review of licensed viral vaccines, some of their safety concerns, and the advances in the development of investigational viral vaccines. *J. Infect.* 49, 179–209.
- (157) Berzofsky, J. A., Ahlers, J. D., Janik, J., Morris, J., Oh, S., Terabe, M., and Belyakov, I. M. (2004) Progress on new vaccine strategies against chronic viral infections. *J. Clin. Invest.* 114, 450–462.
- (158) Carreiras, F., Denoux, Y., Staedel, C., Lehmann, M., Sichel, F., and Gauduchon, P. (1996) Expression and localization of alpha v integrins and their ligand vitronectin in normal ovarian epithelium and in ovarian carcinoma. *Gynecol. Oncol.* 62, 260–267.
- (159) Margeridon-Thermet, S., and Shafer, R. W. (2010) Comparison of the mechanisms of drug resistance among HIV, Hepatitis B, and Hepatitis C. *Viruses* 2, 2696–2739.
- (160) Griffiths, P. D. (2009) A perspective on antiviral resistance. *J. Clin. Virol.* 46, 3–8.
- (161) De Clercq, E. (2002) Strategies in the design of antiviral drugs. *Nat. Rev. Drug Discovery* 1, 13–25.
- (162) Dessain, S. K., Adekar, S. P., and Berry, J. D. (2008) Exploring the native human antibody repertoire to create antiviral therapeutics. *Curr. Top. Microbiol. Immunol.* 317, 155–183.
- (163) Wang, T., Zhang, Z., Wallace, O. B., Deshpande, M., Fang, H., Yang, Z., Zadjura, L. M., Tweedie, D. L., Huang, S., Zhao, F., Ranadive, S., Robinson, B. S., Gong, Y. F., Ricarrdi, K., Spicer, T. P., Deminie, C., Rose, R., Wang, H. G., Blair, W. S., Shi, P. Y., Lin, P. F., Colonna, R. J., and Meanwell, N. A. (2003) Discovery of 4-benzoyl-1-[(4-methoxy-1H-pyrrolo[2,3-b]pyridin-3-yl)oxoacetyl]-2-(R)-methylpiperazine (BMS-378806): a novel HIV-1 attachment inhibitor that interferes with CD4-gp120 interactions. *J. Med. Chem.* 46, 4236–4239.
- (164) Reddy, A. S., Pati, S. P., Kumar, P. P., Pradeep, H. N., and Sastry, G. N. (2007) Virtual screening in drug discovery - A computational perspective. *Curr. Protein Pept. Sci.* 8, 329–351.
- (165) Lange, T., and Lindell, S. (2005) Recent advances in microwave-assisted combinatorial synthesis and library generation. *Comb. Chem. High Throughput Screening* 8, 595–606.
- (166) Osmond, R. I. W., Das, S., and Crouch, M. F. (2010) Development of cell-based assays for cytokine receptor signaling, using an AlphaScreen SureFire assay format. *Anal. Biochem.* 403, 94–101.
- (167) Oyelaran, O., McShane, L. M., Dodd, L., and Gildersleeve, J. C. (2009) Profiling human serum antibodies with a carbohydrate antigen microarray. *J. Proteome Res.* 8, 4301–4310.

- (168) Huflejt, M. E., Vuskovic, M., Vasiliu, D., Xu, H., Obukhova, P., Shilova, N., Tuzikov, A., Galanina, O., Arun, B., Lu, K., and Bovin, N. (2009) Anti-carbohydrate antibodies of normal sera: Findings, surprises and challenges. *Mol. Immunol.* 46, 3037–3049.
- (169) Chen, W., Gu, L., Zhang, W., Motari, E., Cai, L., Styslinger, T. J., and Wang, P. G. (2011) L-Rhamnose antigen: a promising alternative to alpha -Gal for cancer immunotherapies. *ACS Chem. Biol.* 6, 185–191.
- (170) Reddy, M. M., Wilson, R., Wilson, J., Connell, S., Gocke, A., Hynan, L., German, D., and Kodadek, T. (2011) Identification of candidate IgG biomarkers for Alzheimer'S Disease via combinatorial library screening. *Cell* 144, 132–142.
- (171) Sergeeva, A., Kolonin, M. G., Molldrem, J. J., Pasqualini, R., and Arap, W. (2006) Display technologies: Application for the discovery of drug and gene delivery agents. *Adv. Drug Delivery Rev.* 58, 1622–1654.
- (172) Udugamasooriya, D. G., Dineen, S. P., Brekken, R. A., and Kodadek, T. (2008) A peptoid “antibody surrogate” that antagonizes VEGF receptor 2 activity. *J. Am. Chem. Soc.* 130, 5744–5752.
- (173) Buller, F., Steiner, M., Frey, K., Mircsof, D., Scheuermann, J., Kalisch, M., Buhlmann, P., Supuran, C. T., and Neri, D. (2011) Selection of carbonic anhydrase IX inhibitors from one million DNA-Encoded compounds. *ACS Chem. Biol.* 6, 336–344.
- (174) Meireles, L. M. C., and Mustata, G. (2011) Discovery of modulators of protein-protein interactions: current approaches and limitations. *Curr. Top. Med. Chem.* 11, 248–257.
- (175) Mammen, M., Choi, S. K., and Whitesides, G. M. (1998) Polyvalent interactions in biological systems: Implications for design and use of multivalent ligands and inhibitors. *Angew. Chem., Int. Ed.* 37, 2755–2794.
- (176) Lim, R. K. V., and Lin, Q. (2010) Bioorthogonal chemistry: recent progress and future directions. *Chem. Commun.* 46, 1589–1600.
- (177) Sletten, E. M., and Bertozzi, C. R. (2009) Bioorthogonal chemistry: Fishing for selectivity in a sea of functionality. *Angew. Chem., Int. Ed.* 48, 6974–6998.