

# Analysis of the Tau-Associated Proteome Reveals That Exchange of Hsp70 for Hsp90 Is Involved in Tau Degradation


Andrea D. Thompson,<sup>†,||</sup> K. Matthew Scaglione,<sup>‡</sup> John Prensner,<sup>†</sup> Anne T. Gillies,<sup>†,||</sup> Arul Chinnaiyan,<sup>†</sup> Henry L. Paulson,<sup>‡</sup> Umesh K. Jinwal,<sup>⊥,\*</sup> Chad A. Dickey,<sup>\*⊥</sup> and Jason E. Gestwicki<sup>\*,†,§,||</sup>

Departments of <sup>†</sup>Pathology, <sup>‡</sup>Neurology, and <sup>§</sup>Biological Chemistry and the <sup>||</sup>Life Sciences Institute, University of Michigan, Ann Arbor, Michigan 48103, United States

Departments of <sup>⊥</sup>Pharmaceutical Sciences and <sup>⊥</sup>Molecular Medicine, University of South Florida, Tampa, Florida 33613, United States

## S Supporting Information

**ABSTRACT:** The microtubule associated protein tau (MAPT/tau) aberrantly accumulates in 15 neurodegenerative diseases, termed tauopathies. One way to treat tauopathies may be to accelerate tau clearance, but the molecular mechanisms governing tau stability are not yet clear. We recently identified chemical probes that markedly accelerate the clearance of tau in cellular and animal models. In the current study, we used one of these probes in combination with immunoprecipitation and mass spectrometry to identify 48 proteins whose association with tau changes during the first 10 min after treatment. These proteins included known modifiers of tau proteotoxicity, such as ILF-2 (NFAT), ILF-3, and ataxin-2. A striking observation from the data set was that tau binding to heat shock protein 70 (Hsp70) decreased, whereas binding to Hsp90 significantly increased. Both chaperones have been linked to tau homeostasis, but their mechanisms have not been established. Using peptide arrays and binding assays, we found that Hsp70 and Hsp90 appeared to compete for binding to shared sites on tau. Further, the Hsp90-bound complex proved to be important in initiating tau clearance in cells. These results suggest that the relative levels of Hsp70 and Hsp90 may help determine whether tau is retained or degraded. Consistent with this model, analysis of reported microarray expression data from Alzheimer's disease patients and age-matched controls showed that the levels of Hsp90 are reduced in the diseased hippocampus. These studies suggest that Hsp70 and Hsp90 work together to coordinate tau homeostasis.



Tau is primarily expressed in neurons, where it plays a central role in stabilizing microtubules within axons.<sup>1–3</sup> Tau homeostasis is regulated by its expression, phosphorylation, and turnover.<sup>4</sup> In a series of tauopathies, including Alzheimer's disease (AD), frontotemporal dementia (FTLD), progressive supranuclear palsy (PSP), and corticobasal degeneration (CBD), tau homeostasis is disrupted, leading to hyperphosphorylation and accumulation of intracellular aggregates.<sup>5–7</sup> Genetic depletion of tau restores cognitive defects in several mouse models,<sup>8–12</sup> suggesting that reducing tau levels may be an effective way to restore its normal homeostasis. Indeed, recent studies using compounds that reduce tau levels by increasing proteosomal and/or autophagic clearance<sup>13</sup> have shown that this strategy is able to partially recover cognitive defects in cellular and mouse models.<sup>14,15</sup> Together, these findings have focused attention on understanding which cellular pathways protect tau and which proteins are important for its degradation.

Important regulators of tau homeostasis include the molecular chaperones heat shock protein 70 (Hsp70) and Hsp90. Hsp70 and its constitutively expressed isoform, Hsc70, bind directly to tau in a region near the microtubule-binding domains upon release from the microtubule.<sup>16</sup> Through these interactions, Hsp70s facilitate the rebinding of tau to



microtubules and have also been implicated in blocking tau aggregation and promoting its degradation.<sup>17–19</sup> Similarly, Hsp90 and a number of other co-chaperones have been implicated in regulating tau phosphorylation, aggregation, and degradation.<sup>16,17,20–22</sup> Thus, Hsp70 and Hsp90 appear to play roles during multiple processes involved in establishing tau homeostasis. However, these studies have not yet revealed the molecular mechanisms involved, and it is unclear how these chaperones ultimately control tau stability.

Here, we have used a small molecule to acutely disrupt tau equilibrium and thereby promote degradation. A key feature of this approach is that this small molecule reduces tau levels rapidly (within ~15 min in HeLa cells), allowing identification of proteins that change in their association with tau during the first few minutes of triage. Using mass spectrometry and quantitative spectral analysis, we found that only 48 tau-associated proteins are released or enriched during the switch to a degradation fate. Interestingly, Hsp70 is released during the early stages of tau degradation and is replaced by Hsp90. Knockdown analysis revealed that Hsp90 plays a role in


Received: May 25, 2012

Accepted: July 8, 2012

Published: July 8, 2012

**(A) Strategy to define changes in the tau-associated proteome during acute degradation****(C) Confirmation of Hsp70 to Hsp90 switch by western blot****(B) 48 proteins differentially associate with tau**

Increased tau binding			Decreased tau binding		
Bayes Factor	Fold Change	Gene Name	Bayes Factor	Fold Change	Gene Name
8.37E+04	2.27	EPPK1	1.65E+06	3.29	KHSRP
5.22E+03	2.85	HNRNPUL1	7.42E+05	15.94	ATXN2
1.91E+03	2.38	SEC16A	2.00E+04	6.03	SNRPD1
1.90E+03	2.83	HIST1H1D	1.14E+03	7.99	ATXN2L
1.43E+03	2.82	CAD	281.18	4.49	ILF2
777.16	2.38	NCL	99.47	3.27	NHP2L1
363.17	8.86	VIM	93.31	2.98	LSM12
304.15	5.73	AHCY	<b>68.63</b>	<b>2.22</b>	<b>HSPA1B</b>
261.25	5.42	KRT17	57.03	2.59	FAM98B
<b>244.74</b>	<b>2.42</b>	<b>HSP90AB1</b>	50.93	3.95	SMARCE1
100.75	9.15	RPS4X	30.57	2.12	SMARCA4
91.64	5.12	RPL7	24.08	2.41	ILF3
86.70	4.51	RPS24	22.42	2.89	TNRC6B
68.23	3.23	NUDT21	18.07	3.92	HNRNPAB
49.20	4.06	S100A8	15.78	3.38	GSTM3
47.63	3.56	RPL34	15.08	2.62	HLA-DRA
29.67	2.22	DST	13.02	4.17	CD59
29.03	3.00	RPS8	12.07	3.35	GRPEL1
26.95	3.80	KRT79	12.07	3.35	COTL1
25.14	2.82	G3BP1	10.32	2.81	SLC3A2
22.09	3.19	MRPS28			
19.77	2.55	HRNR			
19.48	3.54	FASN			
18.09	2.44	RPL32			
16.82	2.95	RPL7A			
14.83	5.23	RPL10A			
14.19	3.50	S100A9			
10.23	3.52	TIA1			



**Figure 1.** Proteomic analysis of the proteins associated with tau during the acute switch to a degradation fate in response to methylene blue (MB) treatment. (A) Schematic of the experimental strategy for identifying tau-associated proteins that alter tau stability. (B) Using spectral counting, 456 of co-immunoprecipitated proteins did not change in their association with tau in response to MB treatment. However, 20 proteins decreased their association with tau, and 28 preferentially bound. (C) Western blots on freshly immunoprecipitated samples confirmed that MB causes Hsp70 to be released from tau, while Hsp90 binding increases.

targeting tau for degradation. Further, we find that Hsp70 competes with Hsp90 for binding to tau, suggesting that the levels of Hsp70 and Hsp90, in some cases, may dictate tau stability, a result supported by analysis of expression data from AD patients and age matched controls. These studies suggest possible new strategies for the development of therapeutics that target tau for clearance.

## RESULTS AND DISCUSSION

**Identification of Proteins Involved in Tau Homeostasis.** Tau is an intrinsically disordered protein<sup>23</sup> that is thought to engage in protein–protein interactions that govern its localization, activity, and stability. Thus, we reasoned that there are likely tau-associated proteins that stabilize tau within the cell, while other complexes may be critical for its clearance.

We also reasoned that a better understanding of these tau-binding factors might reveal potential new drug targets and provide insights into the mechanisms of chaperone-mediated tau triage. Toward these goals, an important advance is the recent discovery of molecules that acutely disrupt tau equilibrium and favor a rapid change in tau stability. One such molecule, methylene blue (MB), has been shown to reduce tau levels in a variety of cellular and animal models of tauopathies<sup>13–15</sup> and reduce polyQ levels in polyglutamine-expansion disorders.<sup>24</sup> To test whether this compound could be used to explore changes in the tau-associated proteome, we first confirmed that MB reduces total tau levels by ~80% in HeLa (C3) cells stably expressing V5-4RON tau (Supplemental Figure 1A). Loss of tau in this model is rapid and a new equilibrium is reached within ~10–20 min (Supplemental Figure 1B). The speed of this switch is important because it allows insights into the acute changes that occur in the tau-associated protein complexes, while avoiding complications originating from global cellular responses to MB.

To identify proteins that change in their association with tau during the compound-initiated switch to a degradation fate, we pretreated with the proteasome inhibitor bortezomib for 4 h and then applied MB (50  $\mu$ M) or a vehicle control. Bortezomib was used to trap complexes destined for the proteasome, facilitating their subsequent identification by mass spectrometry (Supplemental Figure 1C). Immunoprecipitations with a V5 antibody were performed 10 min after MB or vehicle treatment, and the precipitated material was subject to analysis by LC-MS/MS to identify the tau-associated proteome (Figure 1A). These experiments were performed in two independent, biological replicates, and each replicate was analyzed by mass spectrometry in triplicate. Similar levels of tau (MAPT) were immunoprecipitated in each sample (Supplemental Figure 2), and a total of 504 proteins were identified to co-immunoprecipitate, including tubulin, TDP-43, Hsp70, and Hsp90 (Supplemental Table 1).


Next, quantitative spectral counting was utilized to detect proteins that change in their association with tau in response to MB. To be considered differentially associated with tau, proteins were selected if they had greater than a 2-fold change in abundance and a Bayes factor greater than 10. While the vast majority of interacting proteins, including tubulin, did not change their association with tau upon MB treatment, this criteria identified 48 differentially associated proteins, 20 of which decreased their binding to tau and 28 that increased their binding (Figure 1C, Supplemental Table 1). Interestingly, this list includes a number of factors previously identified as modifiers of tau toxicity, such as ataxin-2, IL-2, and IL-3,<sup>25,26</sup> suggesting that some of these factors might, in part, alter tau proteotoxicity by influencing its turnover. This list also included a number of proteins involved in gene regulation, such as the SWI-SNF components SMARCE1 (BAF57) and SMARCA4 (BRG1), and ribosome-associated proteins, such as RPS4X.

#### Hsp90 Is an Important Factor in Tau Degradation.

One of the most striking observations from the tau interactome analysis was that the association with stress-inducible Hsp70 (HSPA1B) in response to MB was significantly reduced, whereas binding to Hsp90 (HSP90AB1) was increased (Figure 1C). To confirm this finding, the V5-immunoprecipitations were repeated on freshly treated HeLa(C3) cells, and Western blots for Hsp70 and Hsp90 were performed. Consistent with the mass spectrometry findings, Hsp70 decreased while Hsp90 increased its association with tau after MB treatment (Figure

1D). This switch also occurred in the absence of a proteasome inhibitor (Figure 1D). These results suggest that interplay between Hsp70 and Hsp90 may be involved in targeting tau for clearance. Consistent with this idea, it had been previously shown that overexpression of Hsp70 enhances MB-mediated clearance of tau.<sup>13</sup>

To better understand the specific role of Hsp90 in this system, we examined how changes in Hsp90 levels influence MB-initiated tau degradation. Hsp90 is best known for its ability to stabilize many substrates, such as nuclear hormone receptors and kinases, protecting them from degradation.<sup>27,28</sup> However, Hsp90 also promotes the degradation of other substrates, such as von Hippel Lindau factor and high-density lipoprotein.<sup>29,30</sup> To explore the role of Hsp90 within the tau system, we performed siRNA knock-down of Hsp90. Consistent with previous reports,<sup>17,20,31,32</sup> partial knockdown of Hsp90 (by ~80%) did not significantly change the levels of total tau. However, it did suppress the ability of MB to clear tau (Figure 2). This result supports a model in which Hsp90 is




**Figure 2.** Methylene blue (MB)-initiated tau degradation is dependent on Hsp90. Hsp90 siRNA does not cause a significant change in total tau levels in untreated cells but attenuates tau clearance in response to MB. Quantifications of band intensities from two independent experiments were performed using Image J. (\* Student's two-tailed unpaired *t* test, *p*-value <0.05).


involved in targeting tau for degradation, at least in response to MB. This result is in agreement with a previous study using Hsp90 inhibitors, which suggested an active role for Hsp90 in targeting tau for degradation.<sup>20</sup>

**Hsp90 Competes with Hsp70 for Binding to Tau.** On the basis of these results, we hypothesized that exchange of Hsp70 for Hsp90 on tau might occur *via* competition for shared binding sites. At least two binding sites for Hsp70 have been identified on tau,<sup>33</sup> but the binding sites for Hsp90 have not yet been described. To study this question, we developed peptide microarrays composed of 15-mer peptides covering the longest isoform of tau, which is expressed in the peripheral nervous system tau. These tau peptides were assembled with 4 amino acid overlaps, for a total of 194 spots that were each present in triplicate on glass slides. Binding of Hsp70-His and Hsp90-His to these peptide arrays was measured using fluorescent anti-His antibodies, and the negative controls were antibody alone and a nucleotide-binding domain (NBD) of Hsp70, which does not bind substrates. As expected, the negative controls bound nonspecifically to only a few peptides, which were excluded from subsequent analyses (Figure 3A, Supplemental Table 2). In contrast, Hsp70 and Hsp90 were bound to a number of tau-derived peptides (Figure 3A). Restricting our analysis of potential binding sites to those present in the common tau isoform of the central nervous system (4R0N), we identified four Hsp70 binding sites and two for Hsp90 (Figure 3B). Strikingly, both of the Hsp90 binding sites were shared by Hsp70. Previous work has determined that

## (A) Hsp70 and Hsp90 bind discrete and partially overlapping sites in tau



## (B) Schematic of Hsp70 and Hsp90 binding sites




**Figure 3.** Hsp70 and Hsp90 bind discrete and partially overlapping sites on tau, as measured by peptide microarrays. (A) Representative array results using Hsp70 nucleotide binding domain (NBD), Hsp90, and Hsp70 (all 10  $\mu\text{M}$ ) are shown. Further, the fluorescence intensities at a wavelength of 532 nm, after background subtraction, for each array spot tested in triplicate are shown. Peptides with intensities 3 standard errors of mean (SEM) above the mean for the total data set are colored in red. Peptides that bound either Hsp70 NBD or antibody alone are shown in gray. One false-positive spot (array ID: 161) was removed for clarity. (B) Schematic of the Hsp70 and Hsp90 binding sites mapped onto 4RON tau, highlighting the four microtubule-binding repeats (R1–4).

deletion of residues Ile219, Ile220 and Ile250, Val251 reduces Hsp70 binding to tau.<sup>33</sup> Consistent with this result, these residues were present in the third and fourth Hsp70 binding sites, as measured by peptide microarray. These Hsp70 binding sites were further validated using a software program<sup>34</sup> developed to predict binding sites for the prokaryotic Hsp70, DnaK. Using this approach, sites 2, 3, and 4 were positively identified.


The peptide microarray findings indicate that Hsp90 binding sites are shared by Hsp70 and suggest that these chaperones might compete for binding to tau. To test this model, we first synthesized peptides ( $\sim 7$ –15 amino acids) corresponding to the binding sites predicted by the microarray. However, these peptides showed weak ( $>100 \mu\text{M}$ ) binding, which was difficult to accurately measure. Therefore, instead we aimed to validate this model by studying the binding of Hsp70 and Hsp90 in the context of 4RON tau in an ELISA-like platform. Immobilized

Hsp70 bound tau with an affinity of  $2.9 \pm 0.2 \mu\text{M}$  and Hsp90 bound with a slightly weaker affinity of  $7.0 \pm 1.0 \mu\text{M}$  (Figure 4A). As predicted by the array results, Hsp70 competed for


**(A) Hsp70 and Hsp90 bind to tau by ELISA**



**(B) Hsp70 competes with Hsp90**



**(C) Hsp90 only weakly competes with Hsp70**



**Figure 4.** Hsp70 competes with Hsp90 for binding to tau. (A) Immobilized Hsp70 and Hsp90 bind to 4RON tau. (B) Soluble Hsp70 can inhibit binding of tau to immobilized Hsp90 with an  $\text{IC}_{50}$  value of  $4 \mu\text{M}$ . As a control, free Hsp90 also competes with this interaction. (C) Soluble Hsp90 only weakly competes with immobilized Hsp70 for binding to tau with an  $\text{IC}_{50}$  value greater than  $50 \mu\text{M}$ . As a control, free Hsp70 inhibits binding with an  $\text{IC}_{50}$  of  $8 \mu\text{M}$ . Similar values were seen using different Hsp70 and Hsp90 isoforms (see Table 2). All of the experiments were performed in triplicate. Results are shown as the average and standard error of the mean.

binding with Hsp90 ( $\text{IC}_{50} \sim 4 \mu\text{M}$ ) (Figure 4B). Conversely, Hsp90 was less effective at competing with Hsp70 ( $\text{IC}_{50} > 50 \mu\text{M}$ ), perhaps because of the two unique Hsp70 binding sites (Figure 4C). These same relationships were observed with the major stress-inducible (Hsp70, Hsp90 $\alpha$ ) and the constitutively expressed (Hsc70, Hsp90 $\beta$ ) isoforms (Table 1). On the basis of these results, we concluded that the MB-initiated increase in Hsp90 binding observed by mass spectrometry may be the

**Table 1. Competition between Hsp70 and Hsp90 for Binding to Tau**

protein	Hsp70-tau binding		Hsp90 $\alpha$ -tau binding	
	$\text{IC}_{50}^a$ ( $\mu\text{M}$ )	$\text{SEM}^b$	$\text{IC}_{50}^a$ ( $\mu\text{M}$ )	$\text{SEM}^b$
Hsp70 (HSPA1B)	8	4	4	2
Hsc70 (HSPA8)	17	9	1.9	0.5
Hsp90 $\alpha$ (HSP90AA1)	>100	NA	4	2
Hsp90 $\beta$ (HSP90AB1)	59	42	3	1
buffer	>100	NA	>100	NA
Hsp70 NBD	>100	NA	NA	NA


<sup>a</sup> $\text{IC}_{50}$  denotes the concentration which reaches 50% inhibition. <sup>b</sup> $\text{SEM}$  denotes the standard error of the mean.

result of exchange of Hsp70 for Hsp90 on shared binding sites within tau.

**Hsp90 Is Decreased in Human Brains with Alzheimer's Disease.** These results suggest that tau degradation may be sensitive to the relative levels of Hsp70 and Hsp90. More specifically, reduced Hsp90 levels might favor net retention of tau, perhaps unbalancing tau homeostasis. To explore this idea, we examined the GDS810 data set, which collected microarray expression data from the hippocampal tissue of patients with incipient, moderate, and severe AD and age-matched controls.<sup>35</sup> Similarly, we queried the GSE5281 data set that includes expression results from dissected, histopathologically normal hippocampal neurons of AD patients and age-matched controls.<sup>36,37</sup> Previous analyses of these data sets had already noted AD-related changes in protein folding genes, such as chaperones. However, we wanted to specifically ask how the relative levels of Hsp70 and Hsp90 might change as a function of disease. Accordingly, we analyzed both data sets (Supplemental Figure 3) and observed statistically significant changes and trends that indicated an AD-associated decrease in Hsp90 (Figure 5). Although the samples sizes are relatively low, we did not observe consistent changes in stress-inducible Hsp70 levels (HSPA1B, HSPA2, HSPA6, HSPA14), but Hsc70 (HSPA8) was decreased in both data sets (Table 2). Importantly for the current study, the ratio of stress-inducible Hsp70s and Hsp90s was significantly changed in AD, suggesting that imbalance in this chaperone ratio might contribute to the accumulation of tau. Interestingly, these findings are consistent with an observation made in an AD mouse model in which Hsp90 protein levels were found to be inversely correlated with tau levels.<sup>17</sup> Taken together, a model emerges in which Hsp90 levels and the Hsp70:Hsp90 ratio help determine tau stability.

**Discussion.** Tau homeostasis is important in neurodegenerative tauopathies and enhancing tau degradation may be a promising therapeutic strategy.<sup>10,13–16</sup> Thus, it is important to understand the pathways that stabilize tau and those that favor its turnover. Using a chemical biology approach, we specifically studied the early changes in the tau-associated proteome that occur during the acute switch to a degradation fate.

From the list of tau-associated proteins, a striking observation was the loss of Hsp70 (HSPA1B) binding, coupled with an increase in Hsp90 (HSP90AB1) binding. What role is Hsp70 playing in this process? Previous findings suggest that Hsp70 is important in stabilizing tau. For example, Hsc70 is required to recycle tau on microtubules, perhaps restricting its availability to the degradation pathways.<sup>19</sup> However, it is also clear that Hsp70s are not exclusively devoted to stabilizing tau. For



**Figure 5.** Hsp90 levels are decreased in hippocampal samples from Alzheimer's disease (AD) patients. Analysis of the (A) GDS810 and (B) GSE5281 data sets reveals a decrease in the mRNA levels of both isoforms of Hsp90 (HSPAA1 and HSP90AB1) in AD samples. There were no statistically significant changes in the levels of most of the stress-inducible Hsp70s, such as HSPAB1 (C and D), HSPA1B, HSPA2, HSPA4, or HSPA14 (Table 2 and Supplemental Figure 4). However, the levels of Hsc70 (HSPA8) were decreased in AD patient samples (C and D).

example, if this were the case, then Hsp70 knockdown should reduce tau levels, phenocopying MB treatment. Instead, only mild changes in tau levels are typically observed upon knockdown or overexpression of Hsp70.<sup>17</sup> One possible way to account for these observations is that Hsp70 could play an active role in "hand-off" of tau to the degradation pathway. This process might be initiated by MB in these studies, because although MB is certainly not a selective Hsp70 inhibitor,<sup>38,39</sup> overexpression of Hsp70 enhances compound-induced degradation of tau.<sup>13</sup> Thus, MB might cause a structural transition in Hsp70, linking the chaperone to the degradation pathway. Such a mechanism could involve co-chaperones, such as Hop, which bridge the Hsp70 and Hsp90 systems, or alternatively, acute release of tau from Hsp70 may generate a type of tau structure

that binds particularly well to Hsp90. In this model, the activity of Hsp70 that is favored by chemical inhibitors might not produce the same phenotype as knockdown or overexpression, which removes the entire protein. Rather, the transition might require a chemical trigger, such as MB or other Hsp70 inhibitors, which alter Hsp70s nucleotide state, substrate affinity, and/or conformation.<sup>16,40</sup>

What role is Hsp90 playing in this process? Our observations point to a role of Hsp90 in the degradation of tau (see Figure 2). Hsp90 is best known for its ability to stabilize ~200 "client" proteins, such as the glucocorticoid receptor (GR).<sup>28</sup> In those systems, Hsp90 is normally found in the final, high affinity complex that protects the active protein fold.<sup>27,41–43</sup> Accordingly, inhibitors of Hsp90 relieve the protective effect and favor degradation of the clients, usually through a process involving Hsp70. Thus, Hsp90 is typically considered a protective chaperone, while Hsp70 is considered a triage chaperone. However, there are clear examples of clients in the literature that do not follow this paradigm. For example, Hsp90 promotes the degradation of the von Hippel-Lindau tumor-suppressor protein and high-density lipoprotein,<sup>29,30</sup> and in those systems Hsp70 appears to play an upstream role in folding. Our results suggest that tau might fall into this latter category, for which Hsp90 assumes the task of promoting turnover. However, tau seems to differ in some important ways. For example, Hsp90 inhibitors block the degradation of Von Hippel-Lindau tumor-suppressor protein and high-density lipoprotein, whereas these same compounds reduce mutant and phosphorylated tau in a pathway dependent on CHIP, but not HSF-1 or Hsp70.<sup>20</sup> These findings point to a model in which Hsp90 stabilizes some forms of tau, protecting them from clearance,<sup>20,31,44</sup> while also participating in the clearance of tau when Hsp90 or Hsp70 is inhibited. Clearly, the roles played by Hsp90 in tau homeostasis are complex.

One compelling way to rationalize these observations is to invoke different "pools" of Hsp90. Hsp90 engages in protein–protein interactions with many different co-chaperones, and recent work has elegantly shown that distinct Hsp90 complexes exist in the cytosol.<sup>45</sup> We can speculate that some of these Hsp90 complexes may be involved in stabilizing tau while others, such as the ones apparently favored by MB treatment, might target it for degradation. Indeed, some Hsp90 co-chaperones are known to promote tau stabilization, such as cdc37 and FKBP51, while others accelerate degradation, such as FKBP52 and CHIP.<sup>16</sup> Further, the acetylation state of Hsp90 may also influence whether it targets tau for degradation.<sup>46,47</sup> Unfortunately, there was no obvious change in co-chaperone levels in the tau mass spectrometry studies (Supplemental Table 1), but these factors may not have been abundant enough or might bind weakly. Further work, perhaps using small molecules that target individual Hsp90 complexes, will likely be required to better understand this mechanism.

Some of the biggest unresolved issues in understanding tau degradation are what happens during the handoff of tau from Hsp70 to Hsp90 and what happens after formation of the Hsp90-tau complex? One key observation could be that Hsp70 efficiently competes with Hsp90 for binding tau, but not the inverse (see Figure 4 and Table 1). Thus, Hsp90-mediated degradation might depend on a prior signal to release Hsp70, and moreover, this signal might induce a conformation of Hsp70 that actively recruits Hsp90. Despite these speculations and open questions, these findings suggest that exchange of Hsp70 for Hsp90 is one aspect of the mechanism. This model

Table 2. Gene expression level changes for Hsp70 and Hsp90 in Alzheimer's disease (AD)

Gene	Array ID*	GSE5281		GSD810					
		direction of change	p-value	direction of change	p-value	direction of change	p-value	direction of change	p-value
HSP90AA1	210211_s_at, 211968_s_at, 214328_s_at	down	0.088	down	0.031	down	0.236	down	0.088
HSP90AB1	200064_at, 214359_s_at for both and 1557910_at for GSE5281	down	0.017	down	0.114	down	0.046	down	0.008
HSPA1B	200799_at, 200800_s_at, 202581_at	-	0.438	down	0.252	-	0.743	-	0.272
HSPA8	208687_x_at, 210338_s_at, 221891_x_at for both and 224187_x_at for GSE5281	down	0.028	down	0.211	down	0.236	down	0.066
HSPA2	211538_s_at	-	0.227	-	0.470	-	0.370	up	0.224
HSPA4	208815_x_at, 211015_s_at, 211016_x_at	-	0.515	-	1.000	-	0.321	-	0.864
HSPA6	117_at, 213418_at	up	0.032	down	0.252	-	0.321	-	0.181
HSPA14	219212_at for both and 226887_at, 227650_at for GSE5281	-	0.687	-	0.711	down	0.167	down	0.114
NUP37	218622_at	-	0.403	-	0.837	-	0.321	-	0.776

\*Unless otherwise noted the Array IDs given were present in both data sets. Abbreviations are used to indicate the direction of change in gene expression; d denotes a decrease in gene expression in AD patients, and u denotes an increase. Gray denotes gene expression changes with a Mann–Whitney U test *p*-value of <0.05.

is supported by patient expression data and results from mouse models,<sup>17</sup> in which decreased Hsp90 transcript level is linked to tau accumulation and AD. Additional insights into this process will likely accelerate drug discovery in tauopathies, perhaps by suggesting ways of using Hsp70 and Hsp90 inhibitors to maximize the restoration of tau homeostasis. We suggest that rapidly altering the fate of tau using chemical probes will be a particularly valuable strategy toward that goal.

Finally, this work identified a list of tau-associated proteins sensitive to MB treatment (see Figure 1B) and it included a number of factors, including ILF-2 (NFAT), ILF-3, and ataxin 2, that were previously linked to tau proteotoxicity. While our work focused on the contributions of Hsp70 and Hsp90 in regulating tau protein homeostasis, this list of tau-associated factors may be enriched in other proteins that contribute to the regulation of tau homeostasis. For example, although not the focus of this work, the results with ataxin-2 (ATXN2) and ataxin-2-like protein (ATXN2L) were particularly striking, with ~16- and 8-fold reductions in binding to tau during degradation, respectively. These observations are interesting because ataxin-2 is linked to spinocerebellar ataxia type 2 (SCA2), a polyglutamine expansion neurodegenerative disorder,<sup>48</sup> so direct contacts between these disease-associated proteins might be speculated to be important in their pathophysiology.

## METHODS

**Reagents, Cell Lines, and General Methods.** Tetramethylthionine (methylene blue) was purchased from Sigma. siRNA (Qiagen) was transfected at 20 nM.<sup>20,21</sup> Antibodies utilized are as follows: V5 (Sigma, V8137),  $\beta$ -actin (Anaspec, 54591), tau-WB/ELISA (Santa Cruz Biotech, sc5587), pTau (pS396/S404, provided by Dr. Peter Davies Albert Einstein College of Medicine), Hsp70 (Assay Designs), Hsp90 (Santa Cruz Biotech, sc7947), goat anti-rabbit HRP (Anaspec, 28177), and goat anti-mouse HRP (Anaspec 28173). All cells were maintained according to ATCC guidelines. Stably transfected HeLa (C3) cells overexpressing wild-type 4RON human tau were previously generated by clonal selection with G418 (Invitrogen).<sup>13</sup> Absorbance measurements were performed using a SpectraMax M5 multimode plate reader (Molecular Devices).

**Western Blot Analysis.** Samples were separated under reducing and denaturing conditions using 10–20% Tris-Tricine gels (Invitrogen). After transfer to nitrocellulose membrane (Whatman), the membranes were blocked with 5% w/v milk in TBS-T (25 mM Tris-HCl, pH 7.4, 140 mM NaCl, 0.1% v/v Tween 20) for at least 3 h.

Protein levels were detected by incubating the membrane overnight at 4 °C with primary antibodies diluted to 1:1,000, unless otherwise noted, in TBS-T containing 2% w/v bovine serum albumin (BSA, Sigma). Finally, the membrane was incubated for 1 h with the appropriate HRP-conjugated secondary antibody diluted 1:10,000 in TBS-T with 5% w/v BSA. Membranes were washed three times for 5 min with TBS-T at each step. Membranes were developed using Supersignal West Pico chemiluminescence kit (Thermo Scientific). For the Western blot analysis of immunoprecipitated samples titers of 1:250 Hsp70 antibody and 1:500 Hsp90 antibody were utilized.

**Immunoprecipitation of V5-Tau.** HeLa cells stably transfected with V5-4RON tau<sup>13</sup> were grown to 90% confluence and subsequently treated with 5  $\mu$ M Bortezomib for 4 h, followed by a 10 min treatment with either 50  $\mu$ M MB or vehicle (DMSO) control (1% v/v). Cells were lysed with M-Per lysis buffer with protease and phosphatase inhibitor cocktails,<sup>19</sup> and 5 mg of lysate was immunoprecipitated by incubating with 75  $\mu$ L of goat anti-V5 conjugated to agarose beads (Bethyl Laboratories, S190-119) at 4 °C overnight in the dark. Beads were subsequently washed with 100  $\mu$ L of PBS (10 mM Na<sub>2</sub>HPO<sub>4</sub>, 2 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4, 137 mM NaCl, and 2.7 mM KCl) + 0.1% v/v Tween 20, followed by washes with 100  $\mu$ L and then 200  $\mu$ L of PBS. Finally proteins associated were eluted using 45  $\mu$ L of 0.1 M glycine (pH 2.7) and neutralized by adding 5  $\mu$ L 1 M Tris-HCl (pH 8.0). Twenty-five microliters of these samples was separated on 10–20% Tris-Tricine gel (Invitrogen) for Western blot or mass spectrometry analysis. Previous work utilizing this HeLa model has demonstrated that the V5-4RON tau is present in several phosphorylated forms including pS212/p231, pS262/S456, and pS396/S404.<sup>13,21</sup> Further, MB reduces both total tau and tau phosphorylated at S396/S404.<sup>13</sup> Thus, the V5 immunoprecipitated samples likely include a number of tau and phospho-tau variants, providing an overview of the changes in the tau-associated proteome.

**Mass Spectrometry.** Immunoprecipitations were performed on two independent biological samples and separated, as described, using 10–20% Tris-Tricine gels (Invitrogen). Each sample was subjected to in-gel digestion with trypsin and proteins were identified by liquid chromatography-high resolution tandem MS (LTQ-Orbitrap XL, ThermoFisher) at the University of Michigan's Proteomics Resource facility, using standard protocol as previously described.<sup>49</sup> Mass spectrometry analysis was performed in technical triplicates on each sample, providing a total of six data sets per condition. ThermoFisher RAW files were converted to mzXML format using msconvert<sup>50</sup> and were searched against the human UniProt database (release version: 2010\_09). This database fasta file contained 20286 protein entries concatenated with reverse (decoy) sequences and common contaminants. Searches were performed using X!Tandem with the *k*-score plug-in.<sup>51,52</sup> Search results were postprocessed using PeptideProphet

and ProteinProphet from the Trans-Proteomic Pipeline (TPP) software suite (Supplemental Table 1).

**Quantitative Spectral Analysis.** Total spectral counts were obtained from each of the six data sets using Abacus.<sup>55</sup> The abacus parameters used were maxIniProbTH = 0.97, minCombinedFilePw = 0.5, all other parameters were kept at their default values. Using these options, a protein-level false-discovery rate (FDR) of 0.05 was achieved. At this threshold 504 proteins were identified to co-immunoprecipitate with V5-tau across all six data sets. Differential protein expression was assessed by label-free quantification using QSpec.<sup>56</sup> From the QSpec results, a protein was considered to be differentially associated with tau if it had a Bayes Factor greater than 10 and exhibited a fold change greater than 2. Using this definition, 48 of the 504 co-immunoprecipitating proteins were found to be differentially associated upon MB treatment (Figure 1C, Supplemental Table 1).

**Protein Purification.** Human Hsp70 (HSPA1A), Hsc70 (HSPA8), and Hsp70 NBD (1–383) were purified as previously described for DnaK using a His column, including cleavage of the His tag *via* TEV protease as indicated, and a final purification on an ATP-agarose column.<sup>57,58</sup> Human Hsp90-beta (HSP90AB1) and Hsp90-alpha (HSP90AA1) were purified as previously described.<sup>59</sup> N-Terminal His-tagged Human 4RON tau was purified according to the previously developed protocol.<sup>60</sup>

**Tau Peptide Microarray.** The tau peptide microarray was designed using 15-mer peptides with 4 amino acid overhangs, spanning the full sequence of PNS-tau (P10636-9). In addition full-length proteins expected to bind Hsp70 and Hsp90 were printed as positive controls, including human and mouse IgG and tau. Empty spots were used as negative controls. The arrays were printed on single microscope slides in triplicate (JPT peptide technologies). Binding was tested per manufacturer's protocol using 10  $\mu$ M Hsp70-His, Hsp90-His, or Hsp70 nucleotide binding domain (NBD) in binding buffer (25 mM HEPES pH 7.2, 150 mM NaCl, 20 mM KCl, 5 mM MgCl<sub>2</sub>, and 0.5 mM DTT). Binding was detected using 1:1,000 titer of Hilyte555 anti-His antibody (Anaspec) in TBS-T with 1% w/v BSA (Sigma) and scanning the arrays at a fluorescence emission of 532 nm using a GenePix 4100A Microarray Scanner (Molecular Devices) and standard local background subtraction. Binding was defined as peptides with fluorescence signals greater than 3 standard errors above the mean of the total data set. Peptides that appear as positive in the Hsp70 NBD or antibody alone tests were excluded as false positives.

**ELISA-Based Tau Binding.** The procedure for chaperone binding to tau was adapted from a previous report.<sup>61</sup> Briefly, 0.1 mg mL<sup>-1</sup> of either human Hsp70 or Hsp90 (30  $\mu$ L) in immobilization buffer (50 mM MES, pH 5.5, and 0.5 mM DTT) was added to each well of 96-well plates (Thermo Fisher, clear, nonsterile, flat bottom), and these plates were incubated for 6 h at 37 °C. The wells were then washed with 100  $\mu$ L of TBS-T (3  $\times$  3 min, rocking). To these wells was added 50  $\mu$ L of 4RON tau solution (at indicated concentrations) in binding buffer (25 mM HEPES, pH 7.2, 150 mM NaCl, 20 mM KCl, 5 mM MgCl<sub>2</sub>, and 0.5 mM DTT) with 1 mM ATP, and the plates were incubated at RT with gentle rocking overnight. Competition experiments were performed in the presence of 7.5  $\mu$ M tau. Plates were developed using the rabbit anti-tau primary antibody (Santa Cruz, sc5587) (1:2,000 dilution, in TBS-T, 50  $\mu$ L/well) and the goat anti-rabbit HRP conjugated secondary antibody (Anaspec, 28177) (1:2,000 dilution in TBS-T, 50  $\mu$ L/well). The TMB substrate kit (Cell Signaling Technology) was used to detect binding. In each experiment, the signal from nonspecific binding of 4RON to empty control wells was used as a negative control, but this signal was minimal. Binding curves were fit using hyperbolic fits with a nonzero intercept in GraphPad Prism version 5.0 for Windows (GraphPad Software).

**Analysis of Hsp70 and Hsp90 Expression Levels in AD Samples.** The GDS810<sup>35</sup> and GSE5281<sup>36</sup> gene expression microarray data sets were downloaded from the NCBI Gene Expression Omnibus. Normalized data for relevant microarray probes in all samples were extracted for the genes listed in Table 2. The sample GSM21206 was excluded from the GDS810 data set due to artifact rendering which made this sample incomparable to the rest. Probe expression values

were standardized as a z-score ( $z = (\text{value} - \text{mean})/(\text{standard deviation})$ ). Individual probe z-scores belonging to a given gene were averaged within a single patient's sample. Further, the average and variance of expression for a given gene was compared across sample groups (control patients versus patients with disease). Statistical significance ( $p$ -value < 0.05) for gene expression changes between two sample groups was determined with a Mann–Whitney U test (see Supplemental Figure 3).

## ■ ASSOCIATED CONTENT

### ● Supporting Information

This material is available free of charge *via* the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [ujinwal@health.usf.edu](mailto:ujinwal@health.usf.edu); [cdickey@health.usf.edu](mailto:cdickey@health.usf.edu); [gestwick@umich.edu](mailto:gestwick@umich.edu).

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

The authors would like to thank W. Pratt, A. Lieberman, and Y. Osawa for helpful discussions. We would also like to acknowledge D. Southworth for help in purifying Hsp90. The University of Michigan's Department of Pathology Proteomics Core performed the mass spectrometry studies. A.D.T. was supported by a predoctoral fellowship from the National Institute of Health (F30AG035464). This work was additionally supported by NIH R01NS059690 and an anonymous gift from the Michigan Alzheimer's Research Center (MADC). Additional support was from NIH R01NS073899, CurePSP and the Alzheimer's Association to C. Dickey.

## ■ REFERENCES

- (1) Drechsel, D. N., Hyman, A. A., Cobb, M. H., and Kirschner, M. W. (1992) Modulation of the dynamic instability of tubulin assembly by the microtubule-associated protein tau. *Mol. Biol. Cell* 3, 1141–1154.
- (2) Drubin, D. G., Feinstein, S. C., Shooter, E. M., and Kirschner, M. W. (1985) Nerve growth factor-induced neurite outgrowth in PC12 cells involves the coordinate induction of microtubule assembly and assembly-promoting factors. *J. Cell Biol.* 101, 1799–1807.
- (3) Kempf, M., Clement, A., Faissner, A., Lee, G., and Brandt, R. (1996) Tau binds to the distal axon early in development of polarity in a microtubule- and microfilament-dependent manner. *J. Neurosci.* 16, 5583–5592.
- (4) Drewes, G., Ebnet, A., Preuss, U., Mandelkow, E. M., and Mandelkow, E. (1997) MARK, a novel family of protein kinases that phosphorylate microtubule-associated proteins and trigger microtubule disruption. *Cell* 89, 297–308.
- (5) Spillantini, M. G., and Goedert, M. (1998) Tau protein pathology in neurodegenerative diseases. *Trends Neurosci.* 21, 428–433.
- (6) Williams, D. R. (2006) Tauopathies: classification and clinical update on neurodegenerative diseases associated with microtubule-associated protein tau. *Int. Med. J.* 36, 652–660.
- (7) Kidd, M. (1963) Paired helical filaments in electron microscopy of Alzheimer's disease. *Nature* 197, 192–193.
- (8) Santacruz, K., Lewis, J., Spire, T., Paulson, J., Kotilinek, L., Ingelsson, M., Guimaraes, A., DeTure, M., Ramsden, M., McGowan, E., Forster, C., Yue, M., Orne, J., Janus, C., Mariash, A., Kuskowski, M., Hyman, B., Hutton, M., and Ashe, K. H. (2005) Tau suppression in a neurodegenerative mouse model improves memory function. *Science* 309, 476–481.
- (9) Sydow, A., Van der Jeugd, A., Zheng, F., Ahmed, T., Balschun, D., Petrova, O., Drexler, D., Zhou, L., Rune, G., Mandelkow, E., D'Hooge,



R., Alzheimer, C., and Mandelkow, E. M. (2011) Tau-induced defects in synaptic plasticity, learning, and memory are reversible in transgenic mice after switching off the toxic Tau mutant. *J. Neurosci.* 31, 2511–2525.

(10) Roberson, E. D., Halabisky, B., Yoo, J. W., Yao, J., Chin, J., Yan, F., Wu, T., Hamto, P., Devidze, N., Yu, G. Q., Palop, J. J., Noebels, J. L., and Mucke, L. (2011) Amyloid-beta/Fyn-induced synaptic, network, and cognitive impairments depend on tau levels in multiple mouse models of Alzheimer's disease. *J. Neurosci.* 31, 700–711.

(11) Roberson, E. D., Scarce-Levie, K., Palop, J. J., Yan, F., Cheng, I. H., Wu, T., Gerstein, H., Yu, G. Q., and Mucke, L. (2007) Reducing endogenous tau ameliorates amyloid beta-induced deficits in an Alzheimer's disease mouse model. *Science* 316, 750–754.

(12) Vessel, K. A., Zhang, K., Brodbeck, J., Daub, A. C., Sharma, P., Finkbeiner, S., Cui, B., and Mucke, L. (2010) Tau reduction prevents Abeta-induced defects in axonal transport. *Science* 330, 198.

(13) Jinwal, U. K., Miyata, Y., Koren, J., 3rd, Jones, J. R., Trotter, J. H., Chang, L., O'Leary, J., Morgan, D., Lee, D. C., Shults, C. L., Rousaki, A., Weeber, E. J., Zuiderweg, E. R., Gestwicki, J. E., and Dickey, C. A. (2009) Chemical manipulation of hsp70 ATPase activity regulates tau stability. *J. Neurosci.* 29, 12079–12088.

(14) Congdon, E. E., Wu, J. W., Myeku, N., Figueroa, Y. H., Herman, M., Marinec, P. S., Gestwicki, J. E., Dickey, C. A., Yu, W. H., and Duff, K. (2012) Methylthioninium chloride (methylene blue) induces autophagy and attenuates tauopathy in vitro and in vivo. *Autophagy* 8.

(15) O'Leary, J. C., 3rd, Li, Q., Marinec, P., Blair, L. J., Congdon, E. E., Johnson, A. G., Jinwal, U. K., Koren, J., 3rd, Jones, J. R., Kraft, C., Peters, M., Abisambra, J. F., Duff, K. E., Weeber, E. J., Gestwicki, J. E., and Dickey, C. A. (2010) Phenothiazine-mediated rescue of cognition in tau transgenic mice requires neuroprotection and reduced soluble tau burden. *Mol. Neurodegener.* 5, 45.

(16) Miyata, Y., Koren, J., Kiray, J., Dickey, C. A., and Gestwicki, J. E. (2011) Molecular chaperones and regulation of tau quality control: strategies for drug discovery in tauopathies. *Future Med. Chem.* 3, 1523–1537.

(17) Dou, F., Netzer, W. J., Tanemura, K., Li, F., Hartl, F. U., Takashima, A., Gouras, G. K., Greengard, P., and Xu, H. (2003) Chaperones increase association of tau protein with microtubules. *Proc. Natl. Acad. Sci. U.S.A.* 100, 721–726.

(18) Petrucelli, L., Dickson, D., Kehoe, K., Taylor, J., Snyder, H., Grover, A., De Lucia, M., McGowan, E., Lewis, J., Prihar, G., Kim, J., Dillmann, W. H., Browne, S. E., Hall, A., Voellmy, R., Tsuboi, Y., Dawson, T. M., Wolozin, B., Hardy, J., and Hutton, M. (2004) CHIP and Hsp70 regulate tau ubiquitination, degradation and aggregation. *Hum. Mol. Genet.* 13, 703–714.

(19) Jinwal, U. K., O'Leary, J. C., 3rd, Borysov, S. I., Jones, J. R., Li, Q., Koren, J., 3rd, Abisambra, J. F., Vestal, G. D., Lawson, L. Y., Johnson, A. G., Blair, L. J., Jin, Y., Miyata, Y., Gestwicki, J. E., and Dickey, C. A. (2010) Hsc70 rapidly engages tau after microtubule destabilization. *J. Biol. Chem.* 285, 16798–16805.

(20) Dickey, C. A., Kamal, A., Lundgren, K., Klosak, N., Bailey, R. M., Dunmore, J., Ash, P., Shoraka, S., Zlatkovic, J., Eckman, C. B., Patterson, C., Dickson, D. W., Nahman, N. S., Jr., Hutton, M., Burrows, F., and Petrucelli, L. (2007) The high-affinity HSP90-CHIP complex recognizes and selectively degrades phosphorylated tau client proteins. *J. Clin. Invest.* 117, 648–658.

(21) Jinwal, U. K., Koren, J., 3rd, Borysov, S. I., Schmid, A. B., Abisambra, J. F., Blair, L. J., Johnson, A. G., Jones, J. R., Shults, C. L., O'Leary, J. C., 3rd, Jin, Y., Buchner, J., Cox, M. B., and Dickey, C. A. (2010) The Hsp90 cochaperone, FKBP51, increases Tau stability and polymerizes microtubules. *J. Neurosci.* 30, 591–599.

(22) Shimura, H., Miura-Shimura, Y., and Kosik, K. S. (2004) Binding of tau to heat shock protein 27 leads to decreased concentration of hyperphosphorylated tau and enhanced cell survival. *J. Biol. Chem.* 279, 17957–17962.

(23) Uversky, V. N., Oldfield, C. J., and Dunker, A. K. (2008) Intrinsically disordered proteins in human diseases: introducing the D2 concept. *Annu. Rev. Biophys.* 37, 215–246.

(24) Wang, A. M., Morishima, Y., Clapp, K. M., Peng, H. M., Pratt, W. B., Gestwicki, J. E., Osawa, Y., and Lieberman, A. P. (2010) Inhibition of hsp70 by methylene blue affects signaling protein function and ubiquitination and modulates polyglutamine protein degradation. *J. Biol. Chem.* 285, 15714–15723.

(25) Bilen, J., and Bonini, N. M. (2007) Genome-wide screen for modifiers of ataxin-3 neurodegeneration in *Drosophila*. *PLoS Genet.* 3, 1950–1964.

(26) Shulman, J. M., and Feany, M. B. (2003) Genetic modifiers of tauopathy in *Drosophila*. *Genetics* 165, 1233–1242.

(27) Caplan, A. J., Mandal, A. K., and Theodoraki, M. A. (2007) Molecular chaperones and protein kinase quality control. *Trends Cell Biol.* 17, 87–92.

(28) Pratt, W. B., Morishima, Y., Murphy, M., and Harrell, M. (2006) Chaperoning of glucocorticoid receptors. *Handb. Exp. Pharmacol.* 111–138.

(29) McClellan, A. J., Scott, M. D., and Frydman, J. (2005) Folding and quality control of the VHL tumor suppressor proceed through distinct chaperone pathways. *Cell* 121, 739–748.

(30) Gusarova, V., Caplan, A. J., Brodsky, J. L., and Fisher, E. A. (2001) Apoptin B degradation is promoted by the molecular chaperones hsp90 and hsp70. *J. Biol. Chem.* 276, 24891–24900.

(31) Luo, W., Dou, F., Rodina, A., Chip, S., Kim, J., Zhao, Q., Moullick, K., Aguirre, J., Wu, N., Greengard, P., and Chiosis, G. (2007) Roles of heat-shock protein 90 in maintaining and facilitating the neurodegenerative phenotype in tauopathies. *Proc. Natl. Acad. Sci. U.S.A.* 104, 9511–9516.

(32) Sahara, N., Maeda, S., Yoshiike, Y., Mizoroki, T., Yamashita, S., Murayama, M., Park, J. M., Saito, Y., Murayama, S., and Takashima, A. (2007) Molecular chaperone-mediated tau protein metabolism counteracts the formation of granular tau oligomers in human brain. *J. Neurosci. Res.* 85, 3098–3108.

(33) Sarkar, M., Kuret, J., and Lee, G. (2008) Two motifs within the tau microtubule-binding domain mediate its association with the hsc70 molecular chaperone. *J. Neurosci. Res.* 86, 2763–2773.

(34) Rudiger, S., Germeroth, L., Schneider-Mergener, J., and Bukau, B. (1997) Substrate specificity of the DnaK chaperone determined by screening cellulose-bound peptide libraries. *EMBO J.* 16, 1501–1507.

(35) Blalock, E. M., Geddes, J. W., Chen, K. C., Porter, N. M., Markesbery, W. R., and Landfield, P. W. (2004) Incipient Alzheimer's disease: microarray correlation analyses reveal major transcriptional and tumor suppressor responses. *Proc. Natl. Acad. Sci. U.S.A.* 101, 2173–2178.

(36) Liang, W. S., Reiman, E. M., Valla, J., Dunckley, T., Beach, T. G., Grover, A., Niedzielko, T. L., Schneider, L. E., Mastroeni, D., Caselli, R., Kukull, W., Morris, J. C., Hulette, C. M., Schmechel, D., Rogers, J., and Stephan, D. A. (2008) Alzheimer's disease is associated with reduced expression of energy metabolism genes in posterior cingulate neurons. *Proc. Natl. Acad. Sci. U.S.A.* 105, 4441–4446.

(37) Liang, W. S., Dunckley, T., Beach, T. G., Grover, A., Mastroeni, D., Walker, D. G., Caselli, R. J., Kukull, W. A., McKeel, D., Morris, J. C., Hulette, C., Schmechel, D., Alexander, G. E., Reiman, E. M., Rogers, J., and Stephan, D. A. (2007) Gene expression profiles in anatomically and functionally distinct regions of the normal aged human brain. *Physiol. Genomics* 28, 311–322.

(38) Oz, M., Lorke, D. E., and Petroianu, G. A. (2009) Methylene blue and Alzheimer's disease. *Biochem. Pharmacol.* 78, 927–932.

(39) Schirmer, R. H., Adler, H., Pickhardt, M., and Mandelkow, E. (2011) Lest we forget you—methylene blue... *Neurobiol. Aging* 32 (2325), e2327–2316.

(40) Rousaki, A., Miyata, Y., Jinwal, U. K., Dickey, C. A., Gestwicki, J. E., and Zuiderweg, E. R. (2011) Allosteric drugs: the interaction of antitumor compound MKT-077 with human Hsp70 chaperones. *J. Mol. Biol.* 411, 614–632.

(41) Morishima, Y., Murphy, P. J., Li, D. P., Sanchez, E. R., and Pratt, W. B. (2000) Stepwise assembly of a glucocorticoid receptor.hsp90 heterocomplex resolves two sequential ATP-dependent events involving first hsp70 and then hsp90 in opening of the steroid binding pocket. *J. Biol. Chem.* 275, 18054–18060.

- (42) Billecke, S. S., Draganov, D. I., Morishima, Y., Murphy, P. J., Dunbar, A. Y., Pratt, W. B., and Osawa, Y. (2004) The role of hsp90 in heme-dependent activation of apo-neuronal nitric-oxide synthase. *J. Biol. Chem.* 279, 30252–30258.
- (43) Schumacher, R. J., Hurst, R., Sullivan, W. P., McMahon, N. J., Toft, D. O., and Matts, R. L. (1994) ATP-dependent chaperoning activity of reticulocyte lysate. *J. Biol. Chem.* 269, 9493–9499.
- (44) Dickey, C. A., Dunmore, J., Lu, B., Wang, J. W., Lee, W. C., Kamal, A., Burrows, F., Eckman, C., Hutton, M., and Petrucelli, L. (2006) HSP induction mediates selective clearance of tau phosphorylated at proline-directed Ser/Thr sites but not KXGS (MARK) sites. *FASEB J.* 20, 753–755.
- (45) Moullick, K., Ahn, J. H., Zong, H., Rodina, A., Cerchietti, L., Gomes DaGama, E. M., Caldas-Lopes, E., Beebe, K., Perna, F., Hatzi, K., Vu, L. P., Zhao, X., Zatorska, D., Taldone, T., Smith-Jones, P., Alpaugh, M., Gross, S. S., Pillarsetty, N., Ku, T., Lewis, J. S., Larson, S. M., Levine, R., Erdjument-Bromage, H., Guzman, M. L., Nimer, S. D., Melnick, A., Neckers, L., and Chiosis, G. (2011) Affinity-based proteomics reveal cancer-specific networks coordinated by Hsp90. *Nat. Chem. Biol.* 7, 818–826.
- (46) Cook, C., Gendron, T. F., Scheffel, K., Carlomagno, Y., Dunmore, J., Deture, M., and Petrucelli, L. (2012) Loss of HDAC6, a novel CHIP substrate, alleviates abnormal tau accumulation. *Hum. Mol. Genet.* 21, 2936–2945.
- (47) Kovacs, J. J., Murphy, P. J., Gaillard, S., Zhao, X., Wu, J. T., Nicchitta, C. V., Yoshida, M., Toft, D. O., Pratt, W. B., and Yao, T. P. (2005) HDAC6 regulates Hsp90 acetylation and chaperone-dependent activation of glucocorticoid receptor. *Mol. Cell* 18, 601–607.
- (48) Lastres-Becker, I., Rub, U., and Auburger, G. (2008) Spinocerebellar ataxia 2 (SCA2). *Cerebellum* 7, 115–124.
- (49) Brady, G. F., Galban, S., Liu, X., Basrur, V., Gitlin, J. D., Elenitoba-Johnson, K. S., Wilson, T. E., and Duckett, C. S. (2010) Regulation of the copper chaperone CCS by XIAP-mediated ubiquitination. *Mol. Cell. Biol.* 30, 1923–1936.
- (50) Kessner, D., Chambers, M., Burke, R., Agus, D., and Mallick, P. (2008) ProteoWizard: open source software for rapid proteomics tools development. *Bioinformatics* 24, 2534–2536.
- (51) Craig, R., and Beavis, R. C. (2004) TANDEM: matching proteins with tandem mass spectra. *Bioinformatics* 20, 1466–1467.
- (52) MacLean, B., Eng, J. K., Beavis, R. C., and McIntosh, M. (2006) General framework for developing and evaluating database scoring algorithms using the TANDEM search engine. *Bioinformatics* 22, 2830–2832.
- (53) Keller, A., Nesvizhskii, A. I., Kolker, E., and Aebersold, R. (2002) Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search. *Anal. Chem.* 74, 5383–5392.
- (54) Nesvizhskii, A. I., Keller, A., Kolker, E., and Aebersold, R. (2003) A statistical model for identifying proteins by tandem mass spectrometry. *Anal. Chem.* 75, 4646–4658.
- (55) Fermin, D., Basrur, V., Yocum, A. K., and Nesvizhskii, A. I. (2011) Abacus: a computational tool for extracting and pre-processing spectral count data for label-free quantitative proteomic analysis. *Proteomics* 11, 1340–1345.
- (56) Choi, H., Fermin, D., and Nesvizhskii, A. I. (2008) Significance analysis of spectral count data in label-free shotgun proteomics. *Mol. Cell. Proteomics* 7, 2373–2385.
- (57) Chang, L., Thompson, A. D., Ung, P., Carlson, H. A., and Gestwicki, J. E. (2010) Mutagenesis reveals the complex relationships between ATPase rate and the chaperone activities of Escherichia coli heat shock protein 70 (Hsp70/DnaK). *J. Biol. Chem.* 285, 21282–21291.
- (58) Chang, L., Bertelsen, E. B., Wisen, S., Larsen, E. M., Zuiderweg, E. R., and Gestwicki, J. E. (2008) High-throughput screen for small molecules that modulate the ATPase activity of the molecular chaperone DnaK. *Anal. Biochem.* 372, 167–176.
- (59) Southworth, D. R., and Agard, D. A. (2008) Species-dependent ensembles of conserved conformational states define the Hsp90 chaperone ATPase cycle. *Mol. Cell* 32, 631–640.
- (60) Barghorn, S., Biernat, J., and Mandelkow, E. (2005) Purification of recombinant tau protein and preparation of Alzheimer-paired helical filaments in vitro. *Methods Mol. Biol.* 299, 35–51.
- (61) Miyata, Y., Chang, L., Bainor, A., McQuade, T. J., Walczak, C. P., Zhang, Y., Larsen, M. J., Kirchhoff, P., and Gestwicki, J. E. (2010) High-throughput screen for Escherichia coli heat shock protein 70 (Hsp70/DnaK): ATPase assay in low volume by exploiting energy transfer. *J. Biomol. Screening* 15, 1211–1219.