


Exploiting preQ₁ Riboswitches To Regulate Ribosomal FrameshiftingChien-Hung Yu,[†] Jinghui Luo,[‡] Dirk Iwata-Reuyl,[§] and René C. L. Olsthoorn^{*,†}[†]Department of Molecular Genetics and [‡]Biophysical Structural Chemistry, Leiden Institute of Chemistry, Leiden University, PO Box 9502, Leiden, The Netherlands[§]Department of Chemistry, Portland State University, Portland, Oregon 97201, United States

Supporting Information

ABSTRACT: Knowing the molecular details of the interaction between riboswitch aptamers and their corresponding metabolites is important to understand gene expression. Here we report on a novel *in vitro* assay to study preQ₁ riboswitch aptamers upon binding of 7-aminomethyl-7-deazaguanine (preQ₁). The assay is based on the ability of the preQ₁ aptamer to fold, upon ligand binding, into a pseudoknotted structure that is capable of stimulating -1 ribosomal frameshifting (-1 FS). Aptamers from three different species were found to induce between 7% and 20% of -1 FS in response to increasing preQ₁ levels, whereas preQ₁ analogues were 100–1000-fold less efficient. In depth mutational analysis of the *Fusobacterium nucleatum* aptamer recapitulates most of the structural details previously identified for preQ₁ aptamers from other bacteria by crystallography and/or NMR spectroscopy. In addition to providing insight into the role of individual nucleotides of the preQ₁ riboswitch aptamer in ligand binding, the presented system provides a valuable tool to screen small molecules against bacterial riboswitches in a eukaryotic background.



Standard translation of a mRNA proceeds by linear decoding of non-overlapping triplets but for functional reasons can sometimes be overruled to permit translation of an alternative reading frame, a process referred to as ribosomal frameshifting (reviewed in refs 1–3). Most reported examples involve -1 ribosomal frameshifting (-1 FS) where translating ribosomes slip one nucleotide (nt) into the 5'-direction (-1 reading frame) on the mRNA and generate an alternative protein. It is well-known that two *cis*-acting RNA elements are the main signals to induce -1 FS: (i) a heptameric nucleotide sequence called the slip site where the ribosome changes reading frame with consensus X XXY YYZ [where X, Y are any nucleotide, Z \neq Y, and spaces denote the initial reading frame]⁴ and (ii) a stimulatory RNA structure, a hairpin or a pseudoknot, downstream of the slip site [reviewed in refs 5 and 6]. The length of the spacer between slip site and downstream structure, generally 6–9 nts, is also crucial for efficient -1 FS. The appropriate spacer length presumably serves to fine-tune the tension generated by the downstream RNA structure, thereby eliciting the appropriate fraction of frameshifting.^{7,8} Although primarily found in mammalian, plant, and bacterial viruses and a number of transposons,^{9,10} a growing body of evidence shows that -1 FS is involved in decoding bacterial, worm, and mammalian cellular genes.^{11–15} Moreover, the propagation and infectivity of some clinically relevant RNA viruses, such as *Human immunodeficiency virus* type-1 (HIV-1), are negatively affected by modulating -1 FS efficiency,¹⁶ representing -1 FS as a promising therapeutic target to interfere with viral infections. Recently, ligands have been selected that are able to bind the *Severe acute respiratory*

syndrome coronavirus (SARS-CoV) FS pseudoknot and inhibit -1 FS.¹⁷

Therefore, it is of importance to understand how the downstream stimulatory RNA structures affect -1 FS efficiency. It has been suggested by structural and single molecule studies that the mechanical strength of downstream structures correlates with -1 FS efficiency.^{18–20} Intriguingly, for simple perfect hairpin structures and antisense oligonucleotide-forming duplexes, it has been shown that their ability to promote -1 FS positively correlates with the calculated Gibbs free energy,^{21–23} while there is no clear correlation between stability and FS efficiency in frameshifter pseudoknots.²⁴ The latter may be due to the hard-to-predict loop-stem interactions within the pseudoknot frameshifters that play a critical role in enhancing mechanical strength and reducing brittleness of these structures.^{25–30}

High-resolution structural data is required to specifically define the loop-stem interactions within RNA pseudoknots. In reported frameshifter pseudoknots, however, structural details are available only for those with a short stem 1 [≤ 6 base-pair (bp)].⁶ Importantly, they all share a common structural feature, the presence of loop 3 (L3) and stem 1 (S1) (see Figure 1A for nomenclature of stems and loops of a pseudoknot in this study) tertiary interactions close to the helical junction of the stems. Furthermore, a recently defined hTPK-DU177 pseudoknot derived from human telomerase RNA was shown to rely on

Received: November 20, 2012

Accepted: January 17, 2013

Published: January 17, 2013




Figure 1. Structural comparison of preQ₁ riboswitch aptamers and a frameshifting pseudoknot. (A) Schematic representation of the secondary structure of hairpin (H)-type pseudoknots. The “S” denotes the stem region, and the “L” denotes the loop regions. (B) Secondary structures representation of preQ₁-bound aptamers from three indicated bacteria and the *Sugar cane yellow leaf virus* (ScYLV) P1-P2 frameshifting pseudoknot (ScYLV frameshift). Q stands for preQ₁.




Figure 2. PreQ₁ riboswitch aptamers form H-type pseudoknots upon preQ₁ binding and induce -1 ribosomal frameshifting (-1 FS) in a concentration-dependent manner. (A–C) preQ₁ riboswitch aptamers from *Fusobacterium nucleatum* (Fnu), *Bacillus subtilis* (Bsu), and *Thermoanaerobacter tengcongensis* (Tte), respectively. Note that a C (underlined) was inserted in loop L3 of Fnu and Bsu aptamers to prevent ribosomes from encountering a stop codon 5 nts downstream. Bracketed sequences indicate the formation of base pairs in the presence of preQ₁. Slippery sequences are underlined. SDS-PAGE analysis of ³⁵S-methionine labeled translation products in the presence or absence of preQ₁ (0–200 μM) in rabbit reticulocyte lysates (RRL). -1 ribosomal frameshifting is monitored by appearance of the 65-kD product (FS). The non-shifted zero-frame product is indicated by NFS. Quantitative analysis of frameshifting efficiency [FS (%)] is described in Methods. Reported values of FS (%) and SD are from at least three independent experiments.

loop 1 (L1)-stem 2 (S2) triples near the helical junction to promote efficient -1 FS,³⁰ once more highlighting the crucial role of loop-helix interactions.

Riboswitches are gene regulation elements generally located in the 5′-untranslated regions (5′-UTR) of bacterial mRNAs to control gene expression by forming mutually exclusive structures elicited by environmental changes such as the binding of small metabolites or temperature change.³¹ Among riboswitches with available structural information, three of them [*S*-adenosylmethionine class II (SAM-II), *S*-adenosylhomocysteine (SAH), and 7-aminomethyl-7-deazaguanine class I (preQ₁-I)] can adopt a pseudoknot conformation upon ligand binding.³² Interestingly, the SAH riboswitch aptamer can induce SAH-dependent -1 FS³³ to a maximal efficiency of 4%. The relatively inefficient -1 FS may be due to the unusual

pseudoknot topology of the SAH aptamer.³⁴ In contrast, the preQ₁-I and SAM-II riboswitch aptamers adopt the typical hairpin (H)-type pseudoknot structure, which is selected for most frameshifting signals. The preQ₁-I riboswitch controls a set of genes associated with preQ₁ biosynthesis.³⁵ PreQ₁ is a biosynthetic precursor of queuosine (Q), which is a hypermodified nucleotide found in the wobble position of GUN anticodons of tRNA^{Tyr}, tRNA^{His}, tRNA^{Asn}, and tRNA^{Asp} and is important for translational fidelity.³⁶ The preQ₁-I riboswitch is only 34 nt in size, and detailed structural information is available for two of them.^{37–39}

In the present work, we demonstrate that several wild-type (wt) preQ₁ riboswitch aptamers³⁵ can also function as true ligand-responsive frameshifter pseudoknots. By stabilizing S1 of the preQ₁ aptamer of *Fusobacterium nucleatum* (Fnu), about




Figure 3. The stabilized version of Fnu (stab-Fnu) induces highly efficient -1 FS upon preQ₁ binding. stab-Fnu was constructed by replacing the three A-U and U-A pairs by C-G and G-C pairs in S1 of Fnu as shown. SDS-PAGE analysis of ³⁵S-methionine labeled translation products in RRL using stab-Fnu mRNA in the presence or absence of preQ₁. See legend to Figure 2 for more details.




Figure 4. Observation of nucleobase discrimination by preQ₁ riboswitch aptamer using -1 FS as a reporter. (A) Schematic representation of Watson-Crick base pairing of preQ₁ with C17, the determinant nucleotide of preQ₁ riboswitch aptamers. The chemical structure of preQ₁-related compounds tested in FS assays is also presented. (B) SDS-PAGE analysis of ³⁵S-methionine labeled translation products in RRL using stab-Fnu mRNA in the absence or presence of preQ₁ and preQ₁-related nucleobases at indicated concentrations. See legend to Figure 2 for more details. (C) Graph showing the dose-response curves of -1 FS of stab-Fnu induced by various concentrations of preQ₁, guanine, or 2,6-diaminopurine.

40% of ribosomes shifted frame upon addition of 20 μM preQ₁. In depth mutational analysis of the Fnu aptamer recapitulated most of the structural details previously identified for preQ₁ aptamers from other bacteria by crystallography and/or NMR spectroscopy. The preQ₁ aptamer-based frameshifting system presented here may be used in high throughput screening of putative antibacterial drugs.

RESULTS

PreQ₁ Riboswitch Aptamers Can Stimulate -1 FS. The preQ₁ riboswitch aptamers that fold into H-type pseudoknot structures upon ligand binding are structurally similar to typical frameshifting pseudoknots.^{6,40} The three preQ₁ aptamers used in this study are shown (Figure 1B) in comparison with the *Sugar cane yellow leaf virus* (ScYLV) frameshifter pseudoknot.²⁷

Since it has been reported that upon binding of preQ₁ these aptamer pseudoknots are greatly stabilized, we asked whether these ligand-induced pseudoknots could promote FS as well. The preQ₁ aptamers were cloned into our frameshift reporter construct and subsequently assayed for -1 FS in a rabbit reticulocyte lysate expression system (see Methods). In the absence of preQ₁ the *Bacillus subtilis* (Bsu) and Fnu aptamers showed close to zero -1 FS (Figure 2, lanes 1, 9). However -1 FS increased steadily with increasing concentrations of added preQ₁, reaching 7% for the Bsu aptamer (Figure 2, lane 14) and 20% for the Fnu aptamer (Figure 2, lane 8) at the highest concentration of preQ₁ still compatible with efficient translation (200 μM). Interestingly, the *Thermoanaerobacter tengcongensis* (Tte) aptamer exhibited 3.5% of -1 FS in the absence of preQ₁ (Figure 2, lane 15) and peaked at 20 μM with 20% (Figure 2,

lane 18). This result is consistent with the finding that the Tte aptamer forms a pseudoknot structure in the absence of preQ₁.³⁹ In control assays (Supplementary Figure S1) the addition of preQ₁ did not affect FS induced by a non-preQ₁-binding 12-bp hairpin, while disruption of the S2 stem resulted in a non-frameshifting hairpin, indicating that the pseudoknot conformation induced by preQ₁ is responsible for the recorded FS.

Since the Fnu aptamer displayed both high FS efficiency and high preQ₁ sensitivity, we decided to perform additional studies with this aptamer. By substituting G-C for A-U bps in stem S1 of the Fnu aptamer (stab-Fnu), the FS efficiency could be enhanced to 39.1% allowing a better read-out in further analyses (Figure 3). The enhanced FS efficiency is in line with previous data on frameshifter pseudoknots showing that stability of S1 is a major determinant for -1 FS.⁴¹

We note that in order to preserve the reading frame after -1 FS, a C-residue was inserted into L3 of the Bsu and Fnu aptamers. Although the effect of the C in the corresponding position within the A-rich loop (corresponding to L3 in preQ₁ aptamer) of known frameshifter pseudoknots is ambiguous,^{6,42,43} we demonstrated that the additional nucleotide in this position has a negligible effect on the preQ₁ aptamer in our frameshifting-based system since (1) substitution of C by G showed nearly identical FS activity as stab-Fnu (C28G, Supplementary Figure S2A), and (2) simultaneous substitution of A10-U33 by G10-C33 in S2 and removal of the inserted C (A10G+U33C+ΔC28, Supplementary Figure S2B) did induce similar levels of frameshifting as the stab-Fnu construct. These results demonstrate that the preQ₁-induced pseudoknot is an efficient frameshifter.

Ligand Recognition by the *F. nucleatum* preQ₁ Aptamer. The aptamer domain of a riboswitch is responsible for highly selective binding of the target ligand rather than suboptimal analogues to regulate gene expression. The binding affinity for their cognate ligands generally varies by over 2 orders of magnitude to avoid mis-regulation.⁴⁴ To demonstrate that this frameshifting-based assay system can be applied to study ligand selectivity of preQ₁ aptamers, several preQ₁ analogues (Figure 4A) were incubated with the stab-Fnu aptamer to assess their ability to induce frameshifting. As shown in Figure 4B, addition of 2 μM preQ₁ induced 35.4% of frameshifting (Figure 4B, lane 2), 2.2-fold higher than the level (16.2%) induced by 200 μM guanine (Figure 4B, lane 3), which has minor differences in chemical features at the 7 position (Figure 4A). 2,6-Diaminopurine, a preQ₁ analogue capable of inducing structural changes in the preQ₁ aptamer,³⁵ induced 10.3% frameshifting (Figure 4B, lane 4) when present at 200 μM. This 1.5-fold lower efficiency of 2,6-diaminopurine compared to guanine could be due to the fact that only the 2-amino proton can interact with C17 (Figure 4A). However, xanthine and hypoxanthine, both of which can form two hydrogen bonds with C17, did not promote frameshifting (Figure 4B, lanes 6 and 7), consistent with previous findings that the 2-amino group is crucial in ligand recognition and structure modulation.³⁵ Adenine, which cannot base pair with cytidine, did not result in frameshifting at a concentration of 200 μM (Figure 4B, lane 5), as expected. However, 2,4-diaminopyrimidine, although capable of forming identical hydrogen bonds with C17 as 2,6-diaminopurine, exhibited no significant frameshifting (Figure 4B, lane 8) even at a concentration of 200 μM, indicating that the purine moiety is important in molecular recognition and/or base stacking.

On the basis of the observation that the discriminator of purine riboswitches relies in part on Watson-Crick (WC) base pair formation for their ligand selection,^{45,46} we substituted U17 for C17 to test the selectivity for adenine analogues. In a panel of test compounds (Supplementary Figure S3A), we found that 2,6-diaminopurine, which can base pair with U17 by three H-bonds, and preQ₁, which forms a non-canonical wobble pair with U17, were able to promote weak but significant frameshifting (Supplementary Figure S3B). This result agrees in part with previous in-line probing analysis³⁵ showing that 2,6-diaminopurine but not preQ₁ can induce significant structural changes in the mutant C17U preQ₁ aptamer of Bsu. The inability to detect preQ₁ binding may be due to the lower concentration (1 μM) used in that study versus 200 μM in our assays. Our data, in combination with the results of the Bsu preQ₁ aptamer in ligands recognition, recapitulate the importance of 2-amino and 7-deaza-7-aminomethyl groups as well as interactions between the discriminator base and ligand in this type of compact aptamer.

To further compare our frameshifting-based assay in ligand binding to typical in-line probing assays, we plotted the frameshifting efficiency as a function of the concentration of preQ₁ along with two selected ligands to calculate their potency (EC₅₀) (Figure 4C). Although the concentrations of guanine and 2,6-diaminopurine could not be raised high enough to obtain saturating levels of FS, it is clear from Figure 4c that the EC₅₀ of preQ₁ (180.7 nM) is at least 3 orders of magnitude lower than that of either guanine or 2,6-diaminopurine. Previous in-line probing analyses showed a 25-fold difference between preQ₁ and guanine (K_d of preQ₁ and guanine were determined as 20 and 500 nM, respectively³⁵). This suggests that our assay system more closely resembles the natural riboswitch aptamer in discriminating between analogues with over 2 orders of magnitude in binding affinity, despite the fact that our assay primarily responds to the thermodynamic properties of the aptamer and lacks the kinetic properties of riboswitches needed for regulating transcription termination.⁴⁷

Role of Stems of the *F. nucleatum* preQ₁ Aptamer in Frameshifting. It has been suggested that in frameshifting pseudoknots the lower stems (S1) play a major role in stalling elongating ribosomes while the upper stems (S2) may provide torsional restraints to resist ribosome unwinding.⁴⁸ Therefore, variations of residues in either stem that result in a change in ligand binding affinity or stability should affect frameshifting efficiency. We showed in the prior construct (stab-Fnu) that stabilizing S1 could increase FS about 2-fold. Disruption of 1 bp in S2 dramatically reduced the responsiveness to preQ₁ (U33C, Figure 5). The compensatory mutant (A10G+U33C, Figure 5) in which this base pair was restored again acted as a preQ₁-dependent frameshifter. Interestingly, this S2 stabilized mutant resulted in even higher levels of FS (47%). In the absence of preQ₁ this construct showed already some FS (0.8%), suggesting that the S2 formed in the absence of the ligand. These results indicate that the overall stability and integrity of stems affect the ability to induce FS.

We next focused on residues that are located in stems that may be directly involved in the preQ₁-binding pocket. The C18U substitution (Figure 5), which results in the formation of a G5-U18 wobble base pair, was detrimental to FS (less than 1%) even though there is only one WC H-bond missing compared to G5-C18. Moreover, flipping the G5-C18 bp (G5C+C18G, Figure 5) was also detrimental to FS. These results indicate that the highly conserved G5-C18 is crucial for




Figure 5. Effect of mutations in the stem regions on -1 FS. Mutations are indicated in the secondary structure representation. SDS-PAGE analysis of ^{35}S -methionine labeled translation products in RRL using mRNA of each mutant in the presence or absence of preQ₁. See legend to Figure 2 for more details.

formation of the preQ₁-binding pocket. The G11-C32 bp also showed greatly suppressed FS when flipped into C11-G32 (G11C+C32G, Figure 5), consistent with the analogous G11-C32 bp of Bsu aptamer being part of the preQ₁ binding pocket.^{37,38}

We noticed that the conserved YUAR sequence, which forms the 5' half of stem S2 in preQ₁ riboswitches, exhibits the so-called "U-turn" YUNR (Y = pyrimidine, R = purine) motif. This motif has been shown to provide a scaffold for rapid interaction with complementary RNA, forming a paradigm in antisense RNA/target recognition.⁴⁹ In preQ₁ riboswitches it may have evolved to rapidly interact with the downstream complementary sequence to facilitate pseudoknot formation upon ligand binding. To test this hypothesis, we made mutations in the CUAG sequence (and its complement to preserve S2 integrity) of the Fnu pseudoknot and studied their FS efficiency in response to preQ₁. The results, however, indicate that the conserved YUAR sequence in S2 does not

function as a U-turn since CAUG and CAAG sequences were equally efficient as CUUG, which does have the YUNR motif (Supplementary Figure S4).

Effects of Loop Mutants on preQ₁-Responsive Frameshifting. The two stems of the preQ₁ aptamer pseudoknot are separated by three loops (L1 to L3, Figure 1A). These loops strongly resemble the loops of frameshifter pseudoknots either in length or sequence except for the 6-nt L2, which is generally one or zero nucleotide in frameshifter pseudoknots.⁶ Since it has been shown that the loop-stem tertiary interactions are crucial in stabilizing frameshifter pseudoknots,^{25–30} we further investigated the contribution of these tertiary interactions within the preQ₁ aptamer to FS. The results are summarized in Figure 6.

In L1, the sixth nucleotide is found exclusively to be U in all class I type II preQ₁ aptamers. Moreover, in high resolution structures, U6 is involved in the quadruple interaction together with preQ₁, C17, and A31 to form part of the preQ₁ binding pocket.^{37,38} We made a U6A substitution to disrupt the quadruple interaction and found its ability to induce frameshifting is dramatically impaired, confirming that the highly conserved U6 is a major component of the preQ₁-responsive pseudoknot.

The presence of a 6-nt loop between stems S1 and S2 in preQ₁ aptamers is rarely seen in frameshifter pseudoknots (see below) as such a loop would have a large destabilizing effect. Possibly in the ligand-bound form, reorganization of L2 compensates for the destabilizing effect. The only known frameshifter pseudoknot with a large L2 is found in the ovine *Visna-Maedi lentivirus* in which 7 nts (5'CGUCCGC3') are located between two stems of 7 bp each.⁵⁰ It is reasonable to propose that the L2 plays some role in ligand recognition. To investigate the involvement of L2 in the pseudoknot structure in frameshifting, we first deleted all the nucleotides in L2 except C17, the preQ₁ binding nucleotide. This construct (Δ C12-A16), although still capable of inducing a substantial level of FS (7.8%), is not responsive to preQ₁, indicating the L2 is indeed important in trapping preQ₁ in the binding pocket. A16, which forms a base triple with G11-C32 in the Bsu aptamer³⁷ and is the only other highly conserved residue in L2, is also crucial in inducing FS as evidenced by the fact that the A16G substitution is more than 3-fold less efficient (11.5%) in FS compared to stab-Fnu at 200 μM preQ₁. Structural studies indicate that the other nucleotides in L2 are not involved in any interactions with preQ₁,^{37,38} and our mutational analysis of these residues is consistent with that; mutations C12U, C12A, and A15C all displayed a similar level of frameshifting as stab-Fnu. The relatively high tolerance toward nucleotide changes in L2 also agrees with the low phylogenetic conservation of these nucleotides in preQ₁ aptamers.³⁵

L3 of preQ₁ aptamers resembles L2 of frameshifter pseudoknots by its high number of adenosines. It has been shown in structural studies that some of the adenosines form interactions with bases in the minor groove of S1 exclusively via A-amino kissing motifs, in which the Watson-Crick edge of adenines are involved in the interaction, or via mixed A-amino kissing and A-minor motifs, in which the sugar edge of adenines is involved.^{37,38,51} To investigate whether such interactions are important for the stability of the Fnu preQ₁ aptamer, mutations were made in this region and assayed for their FS efficiencies. As shown in Figure 6, the A31U and A30U nucleotide substitutions that were designed to disrupt base quadruples at the junction of the preQ₁ aptamer pseudoknot are both inactive




Figure 6. FS efficiency of loop mutants. Mutations are indicated in the secondary structure representation. The reported -1 FS efficiency is in the presence of $200 \mu\text{M}$ preQ₁.

in frameshifting (<1%). These data indicate that the role of these adenosines of the Fnu aptamer is very similar to that at the corresponding position of the Bsu aptamer for which similar A to U changes also dramatically reduce preQ₁ binding.³⁷ Changing other nucleotides in L3 showed less dramatic changes in FS efficiency suggesting that their individual contributions to the stability of the pseudoknot are less important (see Figure 6). Similar observations have been reported for the *Simian retrovirus* type-I (SRV-1) frameshifter pseudoknot where single changes in L3 did not strongly affect FS but the combined changes did.²⁸

Conclusions. We have shown here that preQ₁ riboswitch aptamers from *B. subtilis*, *F. nucleatum*, and *T. tencongensis* can induce significant levels of -1 FS upon ligand binding. Structurally and functionally these aptamers resemble the small luteovirus frameshifter pseudoknots, and the levels of -1 FS are comparable: 20% for Fnu versus 22% for ScYLV (after correcting for the 1.5-fold more efficient UUUAAAC slippery sequence in our system; GGGAAAC is used by luteoviruses). In the ScYLV and *Beet western yellows luteovirus* (BWYV) pseudoknots²⁵ the structure is stabilized by interactions between nucleotides in L1 with S2, and L3 with S1 (preQ₁ aptamer nomenclature). However, the absence of a large L2 loop in luteovirus pseudoknots makes these pseudoknots more stable and frameshifting independent of ligand-binding.

It has been suggested that due to their relatively small size preQ₁ aptamers are less likely to be detected by automated searching methods and may comprise a substantial fraction of yet to be discovered riboswitches.³⁵ Pseudoknots like that of the Tte aptamer would be especially difficult to predict due to the presence of non-canonical base pairs in S2,³⁹ and it is conceivable that there exists an undiscovered frameshift mechanism exploiting riboswitch-like ligand-induced conformational changes to regulate gene expression. Algorithms aimed at detecting potential frameshifting elements involving this type of pseudoknot should take this into account.

In our frameshift reporter constructs, the preQ₁ aptamers are flanked by long strands of RNA but are nonetheless fully responsive to ligand addition. This may explain why we need a higher concentration of preQ₁ than expected (given a reported K_d of $\sim 280 \text{ nM}$ ⁵² versus EC_{50} of $\sim 2 \mu\text{M}$ for wt Fnu] to induce frameshifting, since alternative structures may form in this

situation. However, these constructs may be closer to the natural situation than the small synthetic RNAs used in structural studies and binding assays. Moreover, frameshifting assays for detecting ligand-aptamer interactions, although not quantitative, show not only ligand-dependent but also reasonable sensitivity (between 20 and 200 nM) and a broad dynamic range (20 to 200 μM). Since the preQ₁ riboswitch is responsible for regulating the expression of genes involved in queuosine synthesis, which is essential for survival at least in stationary growth phase,⁵³ we think it is likely that we can utilize frameshift assays for selecting compounds that can bind to the preQ₁ aptamer and inhibit the growth of pathogens. Furthermore, using a eukaryotic cell-free translation system to monitor prokaryotic RNA–ligand interaction is an advantage for antibacterial drug discovery, since we can simultaneously monitor potential adverse effects on eukaryotic translation. Thus, using frameshift assays in analyzing preQ₁ aptamers may have great potential in high throughput selection of compounds with antibacterial activity.

METHODS

Frameshift Reporter Constructs and Oligonucleotides. The -1 FS was monitored by the SF reporter construct described earlier.²⁸ Mutants were constructed by ligating pairs of complementary oligonucleotides (Eurogentec and Sigma-Aldrich) into SpeI–NcoI digested SF reporter plasmids. Constructs were verified by automated dideoxy sequencing using chain terminator dyes (LGTC, Leiden, The Netherlands). A list of oligonucleotide sequences is available upon request.

Chemicals. PreQ₁ was synthesized and purified as described.³⁵ Adenine, uracil, guanine, hypoxanthine, xanthine, 2,6-diaminopurine, and 2,4-diaminopyrimidine were purchased from Sigma-Aldrich. Guanine was dissolved in 0.15 N KOH, 2,6-diaminopurine was dissolved in RNase-free water, and the other compounds were dissolved in DMSO.

In Vitro Transcription. DNA templates were linearized by BamHI (Fermentas) digestion and purified by successive phenol/chloroform extraction. SP6 polymerase directed transcription was carried out in a 50 μL reaction containing $\sim 2 \mu\text{g}$ linearized DNA, 1 mM NTPs, 40 mM Tris-HCl (pH 7.9), 10 mM NaCl, 10 mM DTT, 6 mM MgCl₂, 2 mM spermidine, 6 units of RNase inhibitor (RNAguard, Pharmacia), and 15 units of SP6 polymerase (Promega). After an incubation period of 2 h at 37 °C, samples were taken and run on agarose gels to determine the quality and quantity of the transcripts. Appropriate

dilutions of the reaction mix in sterilized water were directly used for *in vitro* translations.

In Vitro Translation. Experiments were carried out in duplicate using ~30 ng of mRNA in the presence or absence of the indicated compounds (typically tested at a concentration between 0 and 200 μ M). Reactions contained 4 μ L of rabbit reticulocyte lysate (RRL, Promega), 0.25 μ L of 35 S-methionine (Amersham, *in vitro* translation grade), 0.25 μ L of 1 mM amino acids without methionine, and mRNA and water to a final volume of 10 μ L. After incubation for 60 min at 28 °C, samples were boiled for 3 min in 10 μ L of 2X Laemmli buffer followed and resolved on 13% SDS polyacrylamide gels. Gels were dried and exposed to phosphorimager screens (Biorad). After scanning (Molecular Imager FX, Biorad), band intensity of 0-frame and -1 frameshift products was quantified by Quantity One software (Biorad). Frameshift percentages were calculated as the amount of -1 frameshift product divided by the sum of 0 and -1 frame products, corrected for the number of methionines (10 in the 0-frame product and 28 in the fusion product), multiplied by 100.

■ ASSOCIATED CONTENT

📄 Supporting Information

This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: olsthoor@chem.leidenuniv.nl.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank A. P. Gulyaev for pointing out the possibility of a U-turn motif in preQ₁ riboswitches.

■ REFERENCES

- (1) Farabaugh, P. J. (2000) Translational frameshifting: implications for the mechanism of translational frame maintenance. *Prog. Nucleic Acid Res. Mol. Biol.* 64, 131–170.
- (2) Baranov, P. V., Gesteland, R. F., and Atkins, J. F. (2002) Recoding: translational bifurcations in gene expression. *Gene* 286, 187–201.
- (3) Namy, O., Rousset, J. P., Naphine, S., and Brierley, I. (2004) Reprogrammed genetic decoding in cellular gene expression. *Mol. Cell* 13, 157–168.
- (4) Brierley, I., Jenner, A. J., and Inglis, S. C. (1992) Mutational analysis of the “slippery-sequence” component of a coronavirus ribosomal frameshifting signal. *J. Mol. Biol.* 227, 463–479.
- (5) Brierley, I., and Pennell, S. (2001) Structure and function of the stimulatory RNAs involved in programmed eukaryotic-1 ribosomal frameshifting. *Cold Spring Harbor Symp. Quant. Biol.* 66, 233–248.
- (6) Giedroc, D. P., and Cornish, P. V. (2009) Frameshifting RNA pseudoknots: structure and mechanism. *Virus Res.* 139, 193–208.
- (7) Namy, O., Moran, S. J., Stuart, D. I., Gilbert, R. J. C., and Brierley, I. (2006) A mechanical explanation of RNA pseudoknot function in programmed ribosomal frameshifting. *Nature* 441, 244–247.
- (8) Lin, Z., Gilbert, R. J. C., and Brierley, I. (2012) Spacer-length dependence of programmed -1 or -2 ribosomal frameshifting on a U6A heptamer supports a role for messenger RNA (mRNA) tension in frameshifting. *Nucleic Acids Res.* 40, 8674–8689.
- (9) Brierley, I., Pennell, S., and Gilbert, R. J. (2007) Viral RNA pseudoknots: versatile motifs in gene expression and replication. *Nat. Rev. Microbiol.* 5, 598–610.
- (10) Baranov, P. V., Fayet, O., Hendrix, R. W., and Atkins, J. F. (2006) Recoding in bacteriophages and bacterial IS elements. *Trends Genet.* 22, 174–181.
- (11) Larsen, B., Gesteland, R. F., and Atkins, J. F. (1997) Structural probing and mutagenic analysis of the stem-loop required for

Escherichia coli dnaX ribosomal frameshifting: programmed efficiency of 50%. *J. Mol. Biol.* 271, 47–60.

- (12) Baranov, P. V., Wills, N. M., Barriscale, K. A., Firth, A. E., Jud, M. C., Letsou, A., Manning, G., and Atkins, J. F. (2011) Programmed ribosomal frameshifting in the expression of the regulator of intestinal stem cell proliferation, adenomatous polyposis coli (APC). *RNA Biol.* 8, 637–647.

- (13) Manktelow, E., Shigemoto, K., and Brierley, I. (2005) Characterization of the frameshift signal of Edr, a mammalian example of programmed-1 ribosomal frameshifting. *Nucleic Acids Res.* 33, 1553–1563.

- (14) Wills, N. M. (2006) A functional -1 ribosomal frameshift signal in the human paraneoplastic Ma3 gene. *J. Biol. Chem.* 281, 7082–7088.

- (15) Clark, M. B., Janicke, M., Gottesbuhren, U., Kleffmann, T., Legge, M., Poole, E. S., and Tate, W. P. (2007) Mammalian gene PEG10 expresses two reading frames by high efficiency -1 frameshifting in embryonic-associated tissues. *J. Biol. Chem.* 282, 37359–37369.

- (16) Shehu-Xhilaga, M., Crowe, S. M., and Mak, J. (2001) Maintenance of the Gag/Gag-Pol ratio is important for human immunodeficiency virus type-1 RNA dimerization and viral infectivity. *J. Virol.* 75, 1834–1841.

- (17) Park, S.-J., Kim, Y.-G., and Park, H.-J. (2011) Identification of RNA pseudoknot-binding ligand that inhibits the -1 ribosomal frameshifting of SARS-Coronavirus by structure-based virtual screening. *J. Am. Chem. Soc.* 133, 10094–10100.

- (18) Hansen, T. M., Reihani, S. N. S., Oddershede, L. B., and Sørensen, M. A. (2007) Correlation between mechanical strength of messenger RNA pseudoknots and ribosomal frameshifting. *Proc. Natl. Acad. Sci. U.S.A.* 104, 5830–5835.

- (19) Green, L., Kim, C.-H., Bustamante, C., and Tinoco, I., Jr. (2008) Characterization of the mechanical unfolding of RNA pseudoknots. *J. Mol. Biol.* 375, 511–528.

- (20) Chen, G., Chang, K.-Y., Chou, M.-Y., Bustamante, C., and Tinoco, I., Jr. (2009) Triplex structures in an RNA pseudoknot enhance mechanical stability and increase efficiency of -1 ribosomal frameshifting. *Proc. Natl. Acad. Sci. U.S.A.* 106, 12706–12711.

- (21) Yu, C.-H., Noteborn, M. H. M., and Olsthoorn, R. C. L. (2010) Stimulation of ribosomal frameshifting by antisense LNA. *Nucleic Acids Res.* 38, 8277–8283.

- (22) Yu, C.-H., Noteborn, M. H., Pleij, C. W. A., and Olsthoorn, R. C. L. (2011) Stem-loop structures can effectively substitute for an RNA pseudoknot in -1 ribosomal frameshifting. *Nucleic Acids Res.* 39, 8952–8959.

- (23) Bidou, L., Stahl, G., Grima, B., Liu, H., and Cassan, M. (1997) In vivo HIV-1 frameshifting efficiency is directly related to the stability of the stem-loop stimulatory signal. *RNA* 3, 1153–1158.

- (24) Cornish, P. V., Stammer, S. N., and Giedroc, D. P. (2006) The global structures of a wild-type and poorly functional plant luteoviral mRNA pseudoknot are essentially identical. *RNA* 12, 1959–1969.

- (25) Su, L., Chen, L., Egli, M., Berger, J. M., and Rich, A. (1999) Minor groove RNA triplex in the crystal structure of a ribosomal frameshifting viral pseudoknot. *Nat. Struct. Mol. Biol.* 6, 285–292.

- (26) Michiels, P. J., Versleijen, A. A., Verlaan, P. W., Pleij, C. W., Hilbers, C. W., and Heus, H. A. (2001) Solution structure of the pseudoknot of SRV-1 RNA involved in ribosomal frameshifting. *J. Mol. Biol.* 310, 1109–1123.

- (27) Cornish, P. V., Hennig, M., and Giedroc, D. P. (2005) A loop 2 cytidine-stem 1 minor groove interaction as a positive determinant for pseudoknot-stimulated -1 ribosomal frameshifting. *Proc. Natl. Acad. Sci. U.S.A.* 102, 12694–12699.

- (28) Olsthoorn, R. C. L., Reumerman, R., Hilbers, C. W., Pleij, C. W. A., and Heus, H. A. (2010) Functional analysis of the SRV-1 RNA frameshifting pseudoknot. *Nucleic Acids Res.* 38, 7665–7672.

- (29) Kim, Y. G., Maas, S., Wang, S. C., and Rich, A. (2000) Mutational study reveals that tertiary interactions are conserved in ribosomal frameshifting pseudoknots of two luteoviruses. *RNA* 6, 1157–1165.

- (30) Chou, M.-Y., and Chang, K.-Y. (2010) An intermolecular RNA triplex provides insight into structural determinants for the pseudoknot stimulator of -1 ribosomal frameshifting. *Nucleic Acids Res.* 38, 1676–1685.
- (31) Roth, A., and Breaker, R. R. (2009) The structural and functional diversity of metabolite-binding riboswitches. *Annu. Rev. Biochem.* 78, 305–334.
- (32) Deigan, K. E., and Ferré-D'Amaré, A. R. (2011) Riboswitches: discovery of drugs that target bacterial gene-regulatory RNAs. *Acc. Chem. Res.* 44, 1329–1338.
- (33) Chou, M.-Y., Lin, S.-C., and Chang, K.-Y. (2010) Stimulation of -1 programmed ribosomal frameshifting by a metabolite-responsive RNA pseudoknot. *RNA* 16, 1236–1244.
- (34) Edwards, A. L., Reyes, F. E., Héroux, A., and Batey, R. T. (2010) Structural basis for recognition of S-adenosylhomocysteine by riboswitches. *RNA* 16, 2144–2155.
- (35) Roth, A., Winkler, W. C., Regulski, E. E., Lee, B. W. K., Lim, J., Jona, I., Barrick, J. E., Ritwik, A., Kim, J. N., Welz, R., Iwata-Reuyl, D., and Breaker, R. R. (2007) A riboswitch selective for the queuosine precursor preQ1 contains an unusually small aptamer domain. *Nat. Struct. Mol. Biol.* 14, 308–317.
- (36) Iwata-Reuyl, D. (2008) An embarrassment of riches: the enzymology of RNA modification. *Curr. Opin. Chem. Biol.* 12, 126–133.
- (37) Kang, M., Peterson, R., and Feigon, J. (2009) Structural Insights into riboswitch control of the biosynthesis of queuosine, a modified nucleotide found in the anticodon of tRNA. *Mol. Cell* 33, 784–790.
- (38) Klein, D. J., Edwards, T. E., and Ferré D'Amaré, A. R. (2009) Cocystal structure of a class I preQ1 riboswitch reveals a pseudoknot recognizing an essential hypermodified nucleobase. *Nat. Struct. Mol. Biol.* 16, 343–344.
- (39) Jenkins, J. L., Krucinska, J., McCarty, R. M., Bandarian, V., and Wedekind, J. E. (2011) Comparison of a preQ1 riboswitch aptamer in metabolite-bound and free states with implications for gene regulation. *J. Biol. Chem.* 286, 24626–24637.
- (40) Rieder, U., Lang, K., Kreutz, C., Polacek, N., and Micura, R. (2009) Evidence for pseudoknot formation of class I preQ1 riboswitch aptamers. *Chembiochem* 10, 1141–1144.
- (41) ten Dam, E. B., Verlaan, P. W., and Pleij, C. W. (1995) Analysis of the role of the pseudoknot component in the SRV-1 gag-pro ribosomal frameshift signal: loop lengths and stability of the stem regions. *RNA* 1, 146–154.
- (42) Egli, M., Minasov, G., Su, L., and Rich, A. (2002) Metal ions and flexibility in a viral RNA pseudoknot at atomic resolution. *Proc. Natl. Acad. Sci. U.S.A.* 99, 4302–4307.
- (43) Pallan, P. S., Marshall, W. S., Harp, J., Jewett, F. C., 3rd, Wawrzak, Z., Brown, B. A., 2nd, Rich, A., and Egli, M. (2005) Crystal structure of a luteoviral RNA pseudoknot and model for a minimal ribosomal frameshifting motif. *Biochemistry* 44, 11315–11322.
- (44) Breaker, R. R. (2011) Prospects for riboswitch discovery and analysis. *Mol. Cell* 43, 867–879.
- (45) Mandal, M., Boese, B., Barrick, J. E., Winkler, W. C., and Breaker, R. R. (2003) Riboswitches control fundamental biochemical pathways in *Bacillus subtilis* and other bacteria. *Cell* 113, 577–586.
- (46) Gilbert, S. D., Stoddard, C. D., Wise, S. J., and Batey, R. T. (2006) Thermodynamic and kinetic characterization of ligand binding to the purine riboswitch aptamer domain. *J. Mol. Biol.* 359, 754–768.
- (47) Wickiser, J. K., Winkler, W. C., Breaker, R. R., and Crothers, D. M. (2005) The speed of RNA transcription and metabolite binding kinetics operate an FMN riboswitch. *Mol. Cell* 18, 49–60.
- (48) Plant, E. P., and Dinman, J. D. (2005) Torsional restraint: a new twist on frameshifting pseudoknots. *Nucleic Acids Res.* 33, 1825–1833.
- (49) Franch, T., Petersen, M., Wagner, E. G., Jacobsen, J. P., and Gerdes, K. (1999) Antisense RNA regulation in prokaryotes: rapid RNA/RNA interaction facilitated by a general U-turn loop structure. *J. Mol. Biol.* 294, 1115–1125.
- (50) Pennell, S., Manktelow, E., Flatt, A., Kelly, G., Smerdon, S. J., and Brierley, I. (2008) The stimulatory RNA of the Visna-Maedi retrovirus ribosomal frameshifting signal is an unusual pseudoknot with an interstem element. *RNA* 14, 1366–1377.
- (51) Nissen, P., Ippolito, J. A., Ban, N., Moore, P. B., and Steitz, T. A. (2001) RNA tertiary interactions in the large ribosomal subunit: the A-minor motif. *Proc. Natl. Acad. Sci. U.S.A.* 98, 4899–4903.
- (52) Rieder, U., Kreutz, C., and Micura, R. (2010) Folding of a transcriptionally acting preQ1 riboswitch. *Proc. Natl. Acad. Sci. U.S.A.* 107, 10804–10809.
- (53) Iwata-Reuyl, D. (2003) Biosynthesis of the 7-deazaguanosine hypermodified nucleosides of transfer RNA. *Bioorg. Chem.* 31, 24–43.