

Synthesis of 3,4-Dihydropyridin-2(1*H*)-ones and 3,4-Dihydro-2*H*-pyrans via Four-Component Reactions of Aromatic Aldehydes, Cyclic 1,3-Carbonyls, Arylamines, and Dimethyl Acetylenedicarboxylate

Jing Sun, Er-Yan Xia, Qun Wu, and Chao-Guo Yan*

College of Chemistry & Chemical Engineering, Yangzhou University, Yangzhou 225002, China

Supporting Information

ABSTRACT: A protocol has been developed for the efficient synthesis of structurally diverse 3,4-dihydropyridin-2(1*H*)-ones and 3,4-dihydro-2*H*-pyrans via four-component reactions of arylamines, acetylenedicarboxylate, aromatic aldehydes and cyclic 1,3-diketones. The selective formation of the very different pyridinone or pyran derivatives depends on the structure of cyclic 1,3-diketone. The key steps are proposed to involve Michael addition of the enamino ester formed in situ from the reaction of arylamine with dimethyl acetylenedicarboxylate to arylidene cyclic 1,3-diketones.

KEYWORDS: multicomponent reaction, 1, 3-dipolar addition, pyridinone, pyran, β -enamino ester


INTRODUCTION

Multicomponent reactions have received considerable attention because of their wide range of applications in organic and medical chemistry. Such reactions are able to create combinatorial libraries of complex and diverse structures in efficient fashion by virtue of their convergent nature.^{1,2} Several new designed multicomponent reactions have been reported recently,^{3,4} including processes that take advantage of Huisgen 1,4-dipoles formed from the addition of nitrogen heterocycles or amines to electron-deficient alkynes.^{5–10} Thus, domino reactions involving primary amines, acetylenedicarboxylates, and a third component have been provided elegant procedures for the synthesis of various N- and N, O-heterocycles.^{11–13} We have reported related four-component reactions of aromatic aldehydes, arylamines, acetylenedicarboxylate and acetonitrile derivatives, providing quick access to polysubstituted dihydropyridines.¹⁴ To obtain more information about this novel multicomponent reaction and to examine its substrate scope and limitations, we replaced nitrile-activated components with cyclic 1,3-dicarbonyl compounds. We report here efficient four-component assemblies with these building blocks, producing structurally diverse 3,4-dihydropyridin-2(1*H*)-one and 3,4-dihydro-2*H*-pyran derivatives.

RESULTS AND DISCUSSION

At first we investigated the reactivity of Meldrum acid in the four-component reaction. Meldrum's acid proved to be an active component in reactions performed in ethanol solution in the presence of triethylamine as base catalyst at elevated temperature (50–60 °C) for six hours. 3,4-Dihydropyridin-2(1*H*)-ones 1a–1h were produced in moderate yields (41–60%) as shown in Table 1. To increase the yields of 3,4-dihydropyridin-2(1*H*)-ones, the reaction conditions were also examined. But these trying did not provided much better results. The lower yields of

Table 1. Synthesis of Polysubstituted 3,4-Dihydropyridin-2(1*H*)-ones



entry	compd	Ar	Ar'	yield (%) ^a
1	1a	C ₆ H ₅	<i>p</i> -CH ₃ C ₆ H ₄	41
2	1b	<i>p</i> -(CH ₃) ₂ CHC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	46
3	1c	<i>p</i> -ClC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	58
4	1d	<i>m</i> -ClC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	53
5	1e	<i>m</i> -NO ₂ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	60
6	1f	C ₆ H ₅	C ₆ H ₅	45
7	1g	C ₆ H ₅	<i>p</i> -ClC ₆ H ₄	47
8	1h	<i>p</i> -ClC ₆ H ₄	<i>p</i> -ClC ₆ H ₄	52

^a Isolated yields. All compounds were characterized by ¹H, ¹³C NMR. Reaction conditions: Et₃N(0.5 equiv.), EtOH, 50–60 °C, 6 h.

the products might be due to a competitive ring-opening and subsequent decarboxylation of the Meldrum's acid moiety in hot basic solution. The four-component reaction proceeded smoothly with normal and electron-deficient aromatic amines and aldehydes. The structures of the two representative compounds (1c and 1h) were determined by X-ray diffraction; the structure of 1c is shown in Figure 1. The reaction mechanism (Scheme 1) is believed to involve Michael addition of the 1,3-dipole

Received: March 3, 2011

Revised: April 26, 2011

Published: May 15, 2011





Figure 1. Molecular structure of 3,4-dihydropyridin-2(1H)-one **1c** and 3,4-dihydro-2H-pyrans **2f**, **4b**.

Scheme 1. Proposed Mechanism of the Formation of 3,4-Dihydropyridin-2(1H)-ones



intermediate (B) to the arylidene Meldrum's acid condensate (A), followed by intramolecular nucleophilic substitution in intermediate (C) to form the cyclic amide with elimination of acetone and carbon dioxide. Dihydropyridinone ring system constitutes a group of natural biologically active compounds such as alkaloids which have attracted much attention owing to their wide range of biological activities. The most synthetic approaches have been concentrated on the 1,4-dihydropyridinone and its derivatives. But the synthesis of 3,4-dihydropyridin-2-one has not been explored widely and there were only limited literature precedents for their synthesis.^{15,16} Here we are successfully provided an efficient route to polysubstituted 3,4-dihydropyridin-2-ones starting from readily available starting materials.


The replacement of Meldrum's acid with dimedone produced complex mixtures of products when triethylamine was used to catalyze the four-component reaction. An exploration of several parameters (including base catalyst, solvent, temperature, the sequence of addition of substrates) provided dramatic improvements. Optimal results were obtained by carrying out the four component reaction at room temperature for about one day without using any catalyst. The resulting precipitates were collected by filtration and were found to be novel polysubstituted 3,4-dihydro-2H-pyrans such as **2a** (Table 2). 3,4-Dihydro-2H-pyran moiety is a fundamental structural unit of biologically active compounds. These types of compounds have previously been accessed only by more cumbersome multistep procedures.^{17–20} We are pleased to find a convenient method for the synthesis of polysubstituted 3,4-dihydro-2H-pyrans via a four-component reaction. A variety of electron-poor and electron-rich aromatic aldehydes and amines were found to react very smoothly to provide the analogous dihydro-2H-pyrans in satisfactory yields (64–85%) as shown in Table 2.

The structures of 3,4-dihydro-2H-pyrans **2a–2m** were fully supported by elemental analysis, ¹H and ¹³C NMR, MS, and IR spectra, and were further confirmed by single-crystal X-ray diffraction studies performed on compounds **2f** (Figure 1) and **2j**. The most surprising feature of the structure of **2a–2m** is the fact that one carbonyl group of dimedone took part in cyclization, rather than the amino group, giving the oxygen-containing six-membered ring. Two diastereomers in each product were clearly identified in ¹H NMR spectra in ratios of 5:1 to 7:1, reflecting the cis/trans relationship of the two ester groups. For example, **2a** showed a doublet at 4.20 ppm for the proton at C-4 of the major (trans) isomer and a corresponding doublet at 4.33 ppm for the minor (cis) isomer; other resonances showed similar differences. The X-ray single-crystal molecular structure of **2f** and **2j** showed the aryl group at C-4, ester group at C-3 and ester group at C-2 to be in the all trans-configuration, corresponding to the major isomer of the product.

A similar reactivity pattern was observed with 1,3-cyclohexanedione and 4-hydroxycoumarin as dicarbonyl components, giving the corresponding polysubstituted 3,4-dihydro-2H-pyrans (**3a–3e**) in 66–78% yields (Table 3) and tricyclic compounds (**4a–4e**) in 62–75% yields (Table 3). Benzylamine was also incorporated in place of anilines in reactions with dimedone, giving 3,4-dihydro-2H-pyrans **5a–5j** in 63–81% yields. Taken together, these reactions provided 20 new 3,4-dihydro-2H-pyran derivatives, of which three were characterized by X-ray crystallography (**3b**, **4b** (Figure 1) and **5b**). ¹H NMR spectra indicated a strong preference for only one isomer, presumably that observed by crystallography having trans relationships between the two ester groups and between the aryl group at C-4 and ester at C-3.

Scheme 2 shows a plausible reaction mechanism for these multicomponent transformations, starting with the same type of Knoevenagel and Michael condensation intermediates (A and B) employed in the Meldrum's acid reactions above.¹⁷ Michael addition of 1,3-dipolar intermediate B to A should give C, which is isomerized to D by ketone-enol tautomerization. Intramolecular addition of the enolate to the electron-deficient alkene unit (or its imine/iminium tautomer, not shown) provides intermediate E, which in turn is converted to the final 3,4-dihydro-2H-pyran 2 by protonation. The stereochemistry of the 3,4-dihydro-2H-pyran is controlled by the cis/trans configuration of the in situ formed intermediate (B) and the three sequential reaction steps in the reaction mechanism. The addition of arylamines to but-2-enedioates has been reported to give both

Table 2. Synthesis of Polysubstituted 3,4-Dihydro-2H-pyrans



entry	compd	Ar	Ar'	yield (% trans/cis) ^a
1	2a	C ₆ H ₅	<i>p</i> -CH ₃ C ₆ H ₄	71 (6:1)
2	2b	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	67 (6:1)
3	2c	<i>p</i> -(CH ₃) ₂ CHC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	72 (5:1)
4	2d	<i>p</i> -CH ₃ OC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	64 (6:1)
5	2e	<i>p</i> -FC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	69 (7:1)
6	2f	<i>p</i> -ClC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	78 (7:1)
7	2g	<i>p</i> -BrC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	75 (7:1)
8	2h	<i>m</i> -NO ₂ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	85 (7:1)
9	2i	<i>p</i> -CH ₃ C ₆ H ₄	C ₆ H ₅	62 (7:1)
10	2j	<i>p</i> -CH ₃ C ₆ H ₄	<i>m</i> -ClC ₆ H ₄	74 (6:1)
11	2k	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -ClC ₆ H ₄	75 (6:1)
12	2l	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -CH ₃ OC ₆ H ₄	76 (6:1)
13	2m	<i>p</i> -(CH ₃) ₂ NC ₆ H ₄	<i>p</i> -CH ₃ OC ₆ H ₄	71 (6:1)

^a Isolated yields. All compounds were characterized by ¹H, ¹³C NMR. The ratios of trans/cis were estimated by ¹H NMR. Reaction conditions: EtOH, r.t., 24 h.

cis- and trans-isomers, depending on the nature of the solvent, substituent, and concentration of the reactants.²¹ Recently Jiang reported 2-arylamino fumarates (trans-isomers) to be favored in protic solvents such as methanol and ethanol and the corresponding maleates (cis-isomers) to be favored in nonprotic media such as DMF.²² In our case, the two Michael addition steps and the protonation step in Scheme 2 are all reversible and in thermodynamic equilibrium, giving rise to the more stable 2,3-trans/3,4-trans isomer as the major product. A direct hetero-Diels–Alder addition of the 1,3-dipolar intermediate (B) to arylidene dimedone (A),^{23,24} could be ruled out as an alternative mechanism, since the stereochemical outcome should be different.


CONCLUSION

In summary, we have developed an interesting four-component reaction of aromatic aldehydes, cyclic 1,3-diketones, aryl amines, and dimethyl acetylenedicarboxylate providing two different classes of products depending on the nature of the dicarbonyl component. Mechanisms for the formation of the resulting 3,4-dihydropyridin-2(1H)-ones and 3,4-dihydro-2H-pyrans derivatives are proposed in which stereochemistry is controlled by thermodynamic factors. The potential uses of the reaction in synthetic and medicinal chemistry may be significant, since the products share structural and functional group properties with a variety of biologically active molecules. Further expansion of the reaction scope and synthetic applications of this methodology are in progress in our laboratory.

EXPERIMENTAL SECTION

1. Typical Procedure for the Four-Component Reaction of Arylamines, Dimethyl Acetylenedicarboxylate, Aromatic Aldehydes and Meldrum Acid. A mixture of aromatic aldehyde

Table 3. Synthesis of 3,4-Dihydro-2H-pyran Derivatives




entry	compd	Ar	Ar'	yield ^a (%)
1	3a	<i>p</i> -ClC ₆ H ₄	C ₆ H ₅	66
2	3b	<i>p</i> -ClC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	71
3	3c	<i>m</i> -NO ₂ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	78
4	3d	<i>p</i> -ClC ₆ H ₄	<i>p</i> -ClC ₆ H ₄	75
5	3e	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -CH ₃ OC ₆ H ₄	69
6	4a	<i>p</i> -ClC ₆ H ₄	<i>p</i> -CH ₃ OC ₆ H ₄	66
7	4b	<i>m</i> -ClC ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	62
8	4c	<i>p</i> -CH ₃ C ₆ H ₄	<i>p</i> -ClC ₆ H ₄	70
9	4d	<i>m</i> -NO ₂ C ₆ H ₄	<i>p</i> -CH ₃ C ₆ H ₄	72
10	4e	<i>m</i> -NO ₂ C ₆ H ₄	<i>p</i> -ClC ₆ H ₄	75
11	5a	C ₆ H ₅	CH ₂ C ₆ H ₅	65
12	5b	<i>p</i> -CH ₃ C ₆ H ₄	CH ₂ C ₆ H ₅	68
13	5c	<i>p</i> -(CH ₃) ₂ CHC ₆ H ₄	CH ₂ C ₆ H ₅	68
14	5d	<i>m</i> -(CH ₃) ₃ CC ₆ H ₄	CH ₂ C ₆ H ₅	73
15	5e	<i>p</i> -CH ₃ OC ₆ H ₄	CH ₂ C ₆ H ₅	68
16	5f	<i>p</i> -FC ₆ H ₄	CH ₂ C ₆ H ₅	70
17	5g	<i>m</i> -ClC ₆ H ₄	CH ₂ C ₆ H ₅	63
18	5h	<i>p</i> -ClC ₆ H ₄	CH ₂ C ₆ H ₅	73
19	5i	<i>p</i> -BrC ₆ H ₄	CH ₂ C ₆ H ₅	70
20	5j	<i>m</i> -NO ₂ C ₆ H ₄	CH ₂ C ₆ H ₅	81

^a Isolated yields. All compounds were characterized by ¹H, ¹³C NMR. Reaction conditions: EtOH, r.t. 24 h.

(2.0 mmol), meldrum acid (2.0 mmol, 0.288 g), and triethylamine (1.0 mmol, 0.101 g) in 5.0 mL ethanol was stirred at room for ten minutes. Then a solution of arylamine (2.0 mmol) and dimethyl acetylenedicarboxylate (2.0 mmol, 0.284 g) in 5.0 mL ethanol was added to it. The whole solution was stirred the elevated temperature (50–60 °C) for six hours. In most cases the resulting precipitates were collected by filtration and washed with cold alcohol to give the pure product for analysis. In fewer cases the products were further purified with SiO₂ thin-layer chromatograph with ethyl acetate and light petroleum (V/V = 1/3) as elute to give the product **1a**: yellow solid, 41%, mp 147–149 °C; ¹H NMR (600 MHz, CDCl₃) δ 7.38–7.34 (m, 4H, ArH), 7.27 (t, *J* = 6.6 Hz, 1H, ArH), 7.19 (d, *J* = 7.8 Hz, 2H, ArH), 7.05 (d, *J* = 6.6 Hz, 2H, ArH), 4.31 (d, *J* = 7.2 Hz, 1H, CH), 3.68 (s, 3H, OCH₃), 3.50 (s, 3H, OCH₃), 3.21 (dd, *J*₁ = 7.8 Hz, *J*₂ = 16.2 Hz, 1H, CH₂), 2.96 (d, *J* = 16.2 Hz, 1H, CH₂), 2.35 (s, 3H, CH₃); ¹³C NMR (150 MHz, CDCl₃) δ 168.5, 165.6, 163.3, 144.7, 139.9, 139.2, 133.3, 129.8, 129.1, 128.8, 127.5, 126.7, 109.4, 52.6, 52.3, 38.6, 36.7, 21.3; IR(KBr) *v* 3021, 2953, 1739, 1707, 1633, 1506, 1433, 1347, 1300, 1257, 1204, 1146, 1112, 1069, 987, 849, 803, 754 cm⁻¹; MS (*m/z*) 378.26 ([M - 1]⁺) 100%. Anal. Calcd for C₂₂H₂₁NO₅: C 69.64, H 5.58, N 3.69; Found C 69.78, H 5.81, N 3.54.

2. Typical Procedure for the Four-Component Reaction of Arylamines, Dimethyl Acetylenedicarboxylate, Aromatic Aldehydes, and Dimedone. A mixture of arylamine (2.0 mmol) and dimethyl acetylenedicarboxylate (2.0 mmol, 0.284 g) in 5.0 mL ethanol was stirred at room for ten minutes. Then aromatic aldehyde (2.0 mmol), dimedone (2.0 mmol, 0.284 g) was added. The whole

Scheme 2. Formation Mechanism of Polysubstituted 3,4-Dihydro-2H-pyran



solution was stirred at room temperature for additional about 24 h. The resulting precipitates were collected by filtration and recrystallized in a mixture of ethanol and dimethyl formamide ($V/V = 10/1$) to give the pure product for analysis. **2a**: white solid, 71%, mp 171–172 °C; ^1H NMR (600 MHz, CDCl_3) δ 7.26–7.24 (m, 3H, ArH), 7.13 (d, $J = 7.8$ Hz, 2H, ArH), 6.98 (d, $J = 8.4$ Hz, 2H, ArH), 6.74 (d, $J = 8.4$ Hz, 2H, ArH), 5.44 (brs, 1H, NH), 4.20 (d, $J = 10.2$ Hz, 1H, CH), 3.56 (s, 3H, OCH_3), 3.51 (s, 3H, OCH_3), 3.18 (d, $J = 10.2$ Hz, 1H, CH), 2.60 (d, $J = 17.4$ Hz, 1H, CH_2), 2.34 (d, $J = 17.4$ Hz, 1H, CH_2), 2.24 (s, 3H, CH_3), 2.17 (s, 2H, CH_2), 1.15 (s, 3H, CH_3), 1.01 (s, 3H, CH_3); ^{13}C NMR (150 MHz, CDCl_3) δ 195.9, 171.5, 168.7, 166.5, 141.2, 140.8, 129.7, 128.6, 128.2, 127.2, 126.9, 126.8, 125.9, 115.6, 113.3, 87.8, 54.1, 53.1, 52.6, 50.7, 42.0, 38.8, 31.7, 31.4, 29.2, 27.6, 20.5; IR (KBr) ν 3391, 3022, 2960, 2876, 1759, 1729, 1643, 1592, 1519, 1434, 1370, 1348, 1317, 1265, 1232, 1168, 1047, 1006, 940, 820, 756 cm^{-1} ; MS (m/z) 478.55 ($[\text{M} + 1]^+$) 100%. Anal. Calcd for $\text{C}_{28}\text{H}_{31}\text{NO}_6$: C 70.42, H 6.54, N 2.93; Found C 70.35, H 6.82, N 2.71. The same reaction procedure was carried out by using other cyclic 1,3-dicarbonyl compounds and other amines in the reactions to give 3,4-dihydro-2H-pyran derivatives **2b–2m**, **3a–3e**, **4a–4e**, and **5a–5j**.

ASSOCIATED CONTENT

S Supporting Information. The experimental details and the spectroscopic data, including crystallographic data (CIF) of all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: cgyan@yzu.edu.cn.

Funding Sources

This work was financially supported by the National Natural Science Foundation of China (Grant No. 20972132).

REFERENCES

(1) (a) Armstrong, R. W.; Combs, A. P.; Tempest, P. A.; Brown, S. D.; Keating, T. A. Multiple-Component Condensation Strategies for Combinatorial Library Synthesis. *Acc. Chem. Res.* **1996**, *29*, 123–131. (b) Dömling, A.

Recent Developments in Isocyanide Based Multicomponent Reactions in Applied Chemistry. *Chem. Rev.* **2006**, *106*, 17–89.

(2) (a) Zhu, J.; Bienayme, H. *Multicomponent Reactions*; Wiley: Weinheim, 2005; (b) Tietze, L. F.; Kinzel, T.; Brazel, C. C. The Domino Multicomponent Allylation Reaction for the Stereoselective Synthesis of Homoallylic Alcohols. *Acc. Chem. Res.* **2009**, *42*, 367–378.

(3) (a) Touré, B. B.; Hall, D. G. Natural Product Synthesis Using Multicomponent Reaction Strategies. *Chem. Rev.* **2009**, *109*, 4439–4486. (b) Candeias, N. R.; Montalbano, F.; Cal, P. M.; Gois, P. M. P. Boronic Acids and Esters in the Petasis-Borono Mannich Multicomponent Reaction. *Chem. Rev.* **2010**, *110*, 6169–6193.

(4) (a) Sunderhaus, J. D.; Dockendorff, C.; Martin, S. F. Synthesis of Diverse Heterocyclic Scaffolds via Tandem Additions to Imine Derivatives and Ring-Forming Reactions. *Tetrahedron* **2009**, *65*, 6454–6469. (b) Wang, K.; Nguyen, K.; Huang, Y. J.; Dömling, A. Cyanoacetamide Multicomponent Reaction (I): Parallel Synthesis of Cyanoacetamides. *J. Comb. Chem.* **2009**, *11*, 920–927. (c) Bonne, D.; Coquerel, Y.; Constantieux, T.; Rodriguez, J. 1,3-Dicarbonyl Compounds in Stereoselective Domino and Multicomponent Reactions. *Tetrahedron: Asymmetry* **2010**, *21*, 1085–1109.

(5) (a) Huisgen, R.; Morikawa, M.; Herbig, K.; Brunn, E. 1,4-Dipolar Acycloadditionen, Dreikomponenten-Reaktionen Des Isochinolins mit Acetylenedicarbonsäureester und verschiedenen Dopolarophilen. *Chem. Ber* **1967**, *100*, 1094–1106. (b) Huisgen, R. *Topics in Heterocyclic Chemistry*; Castle, R., Ed.; John Wiley and Sons: New York, NY, 1969; Chapter 8, p 223.

(6) (a) Nair, V.; Rajesh, C.; Vinod, A. U.; Bindu, S.; Sreekanth, A. R.; Mathen, J. S.; Balagopal, L. Strategies for Heterocyclic Construction via Novel Multicomponent Reactions Based on Isocyanides and Nucleophilic Carbenes. *Acc. Chem. Res.* **2003**, *36*, 899–907. (b) Nair, V.; Menon, R. S.; Sreekanth, A.; Abhilash, N.; Biju, A. T. Engaging Zwitterions in Carbon–Carbon and Carbon–Nitrogen Bond-Forming Reactions: A Promising Synthetic Strategy. *Acc. Chem. Res.* **2006**, *39*, 520–530. (c) Nair, V.; Devipriya, S.; Suresh, E. Construction of Heterocycles via 1,4-Dipolar Cycloaddition of Quinoline–DMAD Zwitterion with Various Dipolarophiles. *Tetrahedron* **2008**, *64*, 3567–3577. (d) Shibata, Y.; Noguchi, K.; Hirano, M.; Tanaka, K. Rhodium-Catalyzed Highly Enantio- and Diastereoselective Cotrimerization of Alkenes and Dialkyl Acetylenedicarboxylates Leading to Furylcyclopropanes. *Org. Lett.* **2008**, *10*, 2825–2828. (e) Harju, K.; Vesterinen, J.; Yli-Kauhala, J. Solid-Phase Synthesis of Amino Acid Derived N-Unsubstituted Pyrazoles via Sydnonones. *Org. Lett.* **2009**, *11*, 2219–2221.

(7) (a) Liu, W. B.; Jiang, H. F.; Huang, L. B. One-Pot Silver-Catalyzed and PIDA-Mediated Sequential Reactions: Synthesis of Polysubstituted Pyrroles Directly from Alkynoates and Amines. *Org. Lett.* **2010**, *12*, 312–315. (b) Shen, Y. X.; Jiang, H. F.; Chen, Z. W. $\text{PdCl}_2(\text{HNMe}_2)_2$ -Catalyzed Highly Selective Cross $[2 + 2 + 2]$ Cyclization of Alkynoates and Alkenes under Molecular Oxygen. *J. Org. Chem.* **2010**, *75*, 1321–1324. (c) Liu, W. B.; Jiang, H. F.; Zhang, M.; Qi, C. R. Synthetic Approach to Polysubstituted Furans: An Efficient Addition/Oxidative Cyclization of Alkynoates and 1,3-Dicarbonyl Compounds. *J. Org. Chem.* **2010**, *75*, 966–968. (d) Bayat, M.; Imanieh, H.; Hassanzadeh, F. Triphenylphosphine-catalyzed one-pot synthesis of γ -butyrolactone derivatives and highly substituted enones via reaction of dimethyl acetylenedicarboxylate and aryl aldehydes. *Tetrahedron Lett.* **2010**, *51*, 1873–1875. (e) Terzidis, M. A.; Stephanidou-Stephanatou, J.; Tsoleridis, C. A.; Terzis, A.; Raptoulou, C. P.; Psycharis, V. Expedient One-Pot Synthesis of Highly Substituted Thiazolo[3,2-*a*]pyridines Involving Chromones. *Tetrahedron* **2010**, *66*, 947–954.

(8) (a) Yavari, I.; Piltan, M.; Moradi, L. Synthesis of Pyrrolo[2,1-*a*]-isoquinolines from Activated Acetylenes, Benzoylnitromethanes, and Isoquinoline. *Tetrahedron* **2009**, *65*, 2067–2071. (b) Kumaraswamy, G.; Rambabu, D.; Jayaprakash, N.; Rao, G. V.; Sridhar, B. Cu- and Mo-Catalyzed Expedient Synthesis of Alkynyl-Substituted Derivatives of 1,2-Dihydropyridines, -quinolines and -isoquinolines. *Eur. J. Org. Chem.* **2009**, 4158–4164. (c) Yavari, I.; Mirzaei, A.; Moradi, L.; Khalili, G. Diastereoselective Synthesis of Spiro-Functionalized Tetraalkyl Benzoisoquinopyrrolonaphthyridine-Tetracarboxylates from Isoquinoline, Dialkyl

Acetylenedicarboxylates, and Indane-1,3-dione. *Tetrahedron Lett.* **2010**, *51*, 396–398. (d) Yavari, I.; Seyfi, S.; Hossaini, Z. Formation of Trialkyl Quinoline-2,3,4-tricarboxylates by Reaction of Isatin, Dialkyl Acetylenedicarboxylates, and Sodium *O*-alkyl Carbonodithioates. *Tetrahedron Lett.* **2010**, *51*, 2193–2194.

(9) (a) Alizadeh, A.; Rostamnia, S.; Hu, M. L. A Novel Four-Component Reaction for the Synthesis of 2,5-Diaminofuran Derivatives. *Synlett* **2006**, 1592–1594. (b) Alizadeh, A.; Rostamnia, S.; Zoreh, N.; Oskueyan, Q. A Novel Synthesis of Aminofurans Using a Four-Component Reaction. *Synlett* **2007**, 1610–1612. (c) Alizadeh, A.; Rostamnia, S.; Zhu, L. G. A Novel Pseudo-Seven-Component Diastereoselective Synthesis of λ 5-Phosphanylidene Bis(2,5-dioxotetrahydro-1*H*-pyrrole-3-carboxylates) via Binucleophilic Systems. *Tetrahedron Lett.* **2010**, *51*, 4750–4754. (d) Xia, E. Y.; Sun, J.; Yao, R.; Yan, C. G. Synthesis of Zwitterionic Salts via Three Component Reactions of Nitrogen-Containing Heterocycles, Acetylenedicarboxylate and Cyclic 1,3-Dicarbonyl Compounds. *Tetrahedron* **2010**, *66*, 3569–3574.

(10) (a) Çavdar, H.; Saraçoğlu, N. Synthesis of New 2-Vinylolation Products of Indole via a Michael-Type Addition Reaction with Dimethyl Acetylenedicarboxylate and Their Diels–Alder Reactivity as Precursors of New Carbazoles. *J. Org. Chem.* **2006**, *71*, 7793–7799. (b) Ding, H. F.; Zhang, Y. P.; Bian, M.; Yao, W. J.; Ma, C. Concise Assembly of Highly Substituted Furan-Fused 1,4-Thiazepines and Their Diels–Alder Reactions with Benzynes. *J. Org. Chem.* **2008**, *73*, 578–584. (c) Yadav, J. S.; Reddy, B. V. S.; Yadav, N. N.; Gupta, M. K.; Sridhar, B. Gold(III) Chloride-Catalyzed Three-Component Reaction: A Facile Synthesis of Alkynyl Derivatives of 1,2-Dihydroquinolines and Isoquinolines. *J. Org. Chem.* **2008**, *73*, 6857–6859.

(11) (a) Fan, M. Q.; Yan, Z. Y.; Liu, W. M.; Liang, Y. M. DABCO-Catalyzed Reaction of α -Halo Carbonyl Compounds with Dimethyl Acetylenedicarboxylate: A Novel Method for the Preparation of Poly-substituted Furans and Highly Functionalized 2*H*-Pyrans. *J. Org. Chem.* **2005**, *70*, 8204–8207. (b) Naidu, B. N.; Sorenson, M. E. Facile One-Pot Synthesis of 2,3,5-Substituted 1,2,4-Oxadiazolines from Nitriles in Aqueous Solution. *Org. Lett.* **2005**, *7*, 1391–1393. (c) Nguyen, T. B.; Martel, A.; Dhal, R.; Dujardin, G. *N*-Benzyl Aspartate Nitrones: Unprecedented Single-Step Synthesis and [3 + 2] Cycloaddition Reactions with Alkenes. *Org. Lett.* **2008**, *10*, 4493–4496.

(12) (a) Lu, L.; Wei, J. M.; Chen, J.; Zhang, J. P.; Deng, H. M.; Shao, M.; Zhang, H.; Cao, W. G. A Simple and Convenient Synthesis of 2-(Perfluoroalkyl)-4*H*-chromenes from Salicyl *N*-Tosylimines or Salicylaldehydes and Methyl 2-Perfluoroalkynoates. *Tetrahedron* **2009**, *65*, 9152–9156. (b) Alizadeh, A.; Rostamnia, S.; Hosseinpour, N. A Simple and Effective Approach to the Synthesis of Rhodanine Derivatives via Three-Component Reactions in Water. *Tetrahedron Lett.* **2009**, *50*, 1533–1535. (c) Bezenšek, J.; Koleša, T.; Grošelj, U.; Wagger, J.; Stare, K.; Meden, A.; Svete, J.; Stanovnik, B. [2 + 2] Cycloaddition of Electron-Poor Acetylenes to (*E*)-3-Dimethylamino-1-heteroaryl-prop-2-en-1-ones: Synthesis of Highly Functionalized 1-Heteroaryl-1,3-butadienes. *Tetrahedron Lett.* **2010**, *51*, 3392–3397.

(13) (a) Zewge, D.; Chen, C. Y.; Deer, C.; Dormer, P. G.; Hughes, D. L. A Mild and Efficient Synthesis of 4-Quinolones and Quinolone Heterocycles. *J. Org. Chem.* **2007**, *72*, 4276–4279. (b) Teimouri, M. B.; Abbasi, T.; Mivehchi, H. Novel Multicomponent Reactions of Primary Amines and Alkyl Propiolates with Alloxan Derivatives in Water. *Tetrahedron* **2008**, *64*, 10425–10430. (c) Singh, P.; Sharma, P.; Bisetty, K.; Mahajan, M. Cycloaddition Reactions of Cross-Conjugated Enamines. *Tetrahedron* **2009**, *65*, 8478–8485. (d) Teimouri, M. B.; Abbasi, T. Facile Synthesis of Oxaspirobicyclic Butenolides via a Domino Michael Addition/Aldol Reaction/ γ -Lactonization Sequence. *Tetrahedron* **2010**, *66*, 3795–3800.

(14) Sun, J.; Xia, E. Y.; Wu, Q.; Yan, C. G. Synthesis of Polysubstituted Dihydropyridines by Four-Component Reactions of Aromatic Aldehydes, Malononitrile, Arylamines, and Acetylenedicarboxylate. *Org. Lett.* **2010**, *12*, 3678–3681.

(15) (a) Rodríguez, H.; Suárez, M.; Pérez, R.; Petit, A.; Loupy, A. Solvent-Free Synthesis of 4-Aryl Substituted 5-Alkoxy-carbonyl-6-methyl-3,4-dihydropyridones under Microwave Irradiation. *Tetrahedron Lett.*

2003, *44*, 3709–3712. (b) Rodríguez, H.; Martín, O.; Ochoa, E.; Suárez, M.; Reyes, O.; Garay, H.; Albericio, F.; Martín, N. Solid-Phase Synthesis and Structural Study of Substituted 1,4,5,6-Tetrahydro-6-oxopyridine-3-carboxylic Acids. *QSAR Comb. Sci.* **2006**, *25*, 921–927. (c) Huang, X. J.; Broadbent, S.; Dvorak, C.; Zhao, S. H. Pilot-Plant Preparation of 3,4-Dihydropyridin-2-one Derivatives, the Core Structures of P2 \times 7 Receptor Antagonists. *Org. Proc. Res. Devel* **2010**, *14*, 612–616.

(16) (a) Galanski, P.; Ko, Y. K.; Lee, K. L. Facile Carbamoylation of 3,4-Dihydropyridin-2-one with *N*-Chlorosulfonyl Isocyanate. *Mendeleev Commun.* **2006**, *16*, 46–48. (b) Yang, L.; Zheng, Q. Y.; Wang, D. X.; Huang, Z. T.; Wang, M. X. Reversal of Nucleophilicity of Enamides in Water: Control of Cyclization Pathways by Reaction Media for the Orthogonal Synthesis of Dihydropyridinone and Pyrrolidinone Clausena Alkaloids. *Org. Lett.* **2008**, *10*, 2461–2464. (c) Scheffelaar, R.; Paravidino, M.; Znabet, A.; Schmitz, R. F.; de Kanter, F. J.; Lutz, M.; Spek, A. L.; Guerra, C. F.; Bickelhaupt, F. M.; Groen, M. B.; Ruijter, E.; Orru, R. V. A. Scope and Limitations of an Efficient Four-Component Reaction for Dihydropyridin-2-ones. *J. Org. Chem.* **2010**, *75*, 1723–1732.

(17) (a) Yadav, J. S.; Reddy, B. V. S.; Rao, K. V. R.; Narendar, R. InBr₃-Catalyzed Annulations of Cyclic 1,3-Diketones with Aryl Propargyl Alcohols: A Novel Synthesis of 2,4-Diaryldihydropyrans. *Tetrahedron Lett.* **2009**, *50*, 3963–3965. (b) Nandi, G. C.; Samai, S.; Kumar, R.; Singh, M. S. An Efficient One-Pot Synthesis of Tetrahydrobenzo-[*a*]xanthene-11-one and Diazabenz[*a*]anthracene-9,11-dione Derivatives under Solvent Free Condition. *Tetrahedron* **2009**, *65*, 7129–7134. (c) Reddy, B. V. S.; Reddy, M. R.; Narasimhulu, G.; Yadav, J. S. InCl₃-Catalyzed Three-Component Reaction: A Novel Synthesis of Dihydropyrano[3,2-*b*]chromenediones under Solvent-Free Conditions. *Tetrahedron Lett.* **2010**, *51*, 5677–5679. (d) Kumar, R.; Raghuvanshi, K.; Verma, K. R.; Singh, M. S. Application of Cyclic-1,3-diketones in Domino and Multicomponent Reactions: Facile Route to Highly Functionalized Chromeno[2,3-*d*]pyrimidinones and Diazabenzob[*b*]fluorenones under Solvent-Free Conditions. *Tetrahedron Lett.* **2010**, *51*, 5933–5936.

(18) (a) Ziegert, R. E.; Toräng, J.; Knepper, K.; Bräse, S. The Recent Impact of Solid-Phase Synthesis on Medicinally Relevant Benzoannulated Oxygen Heterocycles. *J. Comb. Chem.* **2005**, *7*, 147–169. (b) Núñez, M. G.; García, P.; Moro, R. F.; Diez, D. Asymmetric Organocatalytic Synthesis of Six-Membered Oxygenated Heterocycles. *Tetrahedron* **2010**, *66*, 2089–2109. (c) Liu, W.; Zhou, J.; Zheng, C. W.; Chen, X. K.; Xiao, H.; Yang, Y. Q.; Guo, Y. L.; Zhao, G. Tandem Cross-Rauhut–Currier/Cyclization Reactions of Activated Alkenes to Give Densely Functionalized 3,4-Dihydropyrans. *Tetrahedron* **2011**, *67*, 1768–1773.

(19) (a) Wang, L. M.; Shao, J. H.; Tian, H.; Wang, Y. H.; Liu, B. Rare Earth Perfluorooctanoate [RE(PFO)₃] catalyzed one-pot synthesis of benzopyran derivatives. *J. Fluorine Chem.* **2006**, *127*, 97–100. (b) Mori, K.; Kawasaki, T.; Sueoka, S.; Akiyama, T. Expedient Synthesis of Benzopyrans via Lewis Acid-Catalyzed C–H Functionalization: Remarkable Enhancement of Reactivity by an Ortho Substituent. *Org. Lett.* **2010**, *12*, 1732–1735. (c) Spivey, A. C.; Laraia, L.; Bayly, A. R.; Rzepa, H. S.; White, A. J. P. Stereoselective Synthesis of *cis*- and *trans*-2,3-Disubstituted Tetrahydrofurans via Oxonium–Prins Cyclization: Access to the Cordigol Ring System. *Org. Lett.* **2010**, *12*, 900–903.

(20) (a) Boivin, T. L. B. Synthetic Routes to Tetrahydrofuran, Tetrahydropyran, and Spiroketal Units of Polyether Antibiotics and a Survey of Spiroketal of Other Natural Products. *Tetrahedron* **1987**, *43*, 3309–3362. (b) Elliott, M. C. Saturated Oxygen Heterocycles. *J. Chem. Soc., Perkin Trans. 1* **2002**, 2301–2323. (c) Clarke, P. A.; Santos, S. Strategies for the Formation of Tetrahydropyran Rings in the Synthesis of Natural Products. *Eur. J. Org. Chem.* **2006**, 2045–2053. (d) Tang, Y.; Oppenheimer, J.; Song, Z.; You, L.; Zhang, X.; Hsung, R. P. Strategies and Approaches for Constructing 1-Oxadecalins. *Tetrahedron* **2006**, *62*, 10785–10813. (e) Larrosa, I.; Romea, P.; Urpi, F. Synthesis of Six-Membered Oxygenated Heterocycles through Carbon–Oxygen Bond-Forming Reactions. *Tetrahedron* **2008**, *64*, 2683–2723.

(21) (a) Schmidt, R. R.; Kast, J.; Speer, H. Erzeugung von Vinylithium–Verbindungen aus Acetylenedicarbonsäure-ester-Addukten. *Synthesis* **1983**, 725–727. (b) Kandeel, A. K.; Vernon, J. M. M. Reactions of

Tertiary Allylamines with Dimethyl Acetylenedicarboxylate. *J. Chem. Soc., Perkin Trans. 1* **1987**, 2023–2026. (c) Nuvole, A.; Paglietti, G. Reaction of (*E*)- β -Enamino Amides with Dimethyl Acetylenedicarboxylate (DMAD): Formation of Benzene Derivatives, Enamino Esters, and 2-Pyridones. *J. Chem. Soc., Perkin Trans. 1* **1989**, 1007–1011. (d) Cho, C. S. An efficient dealkylative addition of trialkylamines to dialkyl acetylenedicarboxylates in the presence of a metallic chloride. *Tetrahedron Lett.* **2005**, *46*, 1415–1417. (e) Ziyaei-Halimehjani, A.; Saidi, M. R. Synthesis of Aza-Henry Products and Enamines in Water by Michael Addition of Amines or Thiols to Activated Unsaturated Compounds. *Tetrahedron Lett.* **2008**, *49*, 1244–1248.

(22) Zhu, Q. H.; Jiang, H. F.; Li, J. H.; Zhang, M.; Wang, X. J.; Qi, C. R. Practical Synthesis and Mechanistic Study of Polysubstituted Tetrahydropyrimidines with Use of Domino Multicomponent Reactions. *Tetrahedron* **2009**, *65*, 4604–4613.

(23) (a) Jung, M. J.; Buszek, K. R. Alkenyltrialkylammonium Salts As Dienophiles in Diels–Alder Reactions: Preparation, Cycloadditions, and Further Reactions β -(Dimethylamino)acrylonitrile Equivalent in Cycloadditions. *J. Org. Chem.* **1985**, *50*, 5440–5441. (b) Tietze, L. F. Domino Reactions in Organic Synthesis. *Chem. Rev.* **1996**, *96*, 115–136. (c) Behforouz, M.; Ahmadian, M. Diels–Alder Reactions of 1-Azadienes. *Tetrahedron* **2000**, *56*, 5259–5288. (d) List, B.; Castello, A. Novel Proline-Catalyzed Three-Component Reaction of Ketones, Aldehydes, and Meldrum's Acid. *Synlett* **2001**, 1687–1689.

(24) (a) Ramachary, D. B.; Chowdari, N. S.; Barbas, C. F., III The First Organocatalytic Hetero-Domino Knoevenagel-Diels-Alder-Epimerization Reactions: Diastereoselective Synthesis of Highly Substituted Spiro[cyclohexane-1,2'-indan]-1',3',4-triones. *Synlett* **2003**, 1910–1914. (b) Ramachary, D. B.; Anebuselvy, K.; Chowdari, N. S.; Barbas, C. F., III Direct Organocatalytic Asymmetric Heterodomino Reactions: The Knoevenagel/Diels–Alder/Epimerization Sequence for the Highly Diastereoselective Synthesis of Symmetrical and Nonsymmetrical Synthesis of Benzoannulated Centropolyquinanes. *J. Org. Chem.* **2004**, *69*, 5838–5849. (c) Jiménez-Alonso, S.; Estévez-Braun, A.; Ravelo, Á. G.; Zárate, R.; López, M. Double Domino Knoevenagel hetero Diels–Alder Strategy Towards Bis-pyrano-1,4-benzoquinones. *Tetrahedron* **2007**, *63*, 3066–3074.