


Efficient Solution-Phase Synthesis of 4,5,7-Trisubstituted Pyrrolo[3,2-*d*]pyrimidines

Weihe Zhang, Jing Liu, Michael A. Stashko, and Xiaodong Wang\*

Center for Integrative Chemical Biology and Drug Discovery, Division of Chemical Biology and Medicinal Chemistry, UNC Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, North Carolina 27599, United States

## Supporting Information

**ABSTRACT:** We have developed an efficient and robust route to synthesize 4,5,7-trisubstituted pyrrolo[3,2-*d*]pyrimidines as potent kinase inhibitors. This solution-phase synthesis features a  $S_NAr$  substitution reaction, cross-coupling reaction, one-pot reduction/reductive amination and *N*-alkylation reaction. These reactions occur rapidly with high yields and have broad substrate scopes. A variety of groups can be selectively introduced into the N5 and C7 positions of 4,5,7-trisubstituted pyrrolopyrimidines at a late stage of the synthesis, thereby providing a highly efficient approach to explore the structure–activity relationships of pyrrolopyrimidine derivatives. Four synthetic analogs have been profiled against a panel of 48 kinases and a new and selective FLT3 inhibitor **9** is identified.




**KEYWORDS:** pyrrolopyrimidine,  $S_NAr$  displacement, coupling reaction, reductive amination, *N*-alkylation


## INTRODUCTION

Pyrrolopyrimidines have been broadly used as synthetic pharmacophores in drug discovery due to their structural similarity to purines with numerous associated biological activities. Accordingly, pyrrolopyrimidines have been exploited as inhibitors of kinases (VEGFR,<sup>1</sup> FGFR,<sup>2</sup> HGF,<sup>3</sup> Akt,<sup>4</sup> MMP,<sup>5</sup> JAK,<sup>6</sup> Lck,<sup>7</sup> etc.), methylthioadenosine phosphorylase (MTAP),<sup>8</sup> purine nucleoside phosphorylase (PNP),<sup>9</sup> and HIV replication.<sup>10</sup> In one of our drug discovery programs, we are interested in 4,5,7-trisubstituted pyrrolo[3,2-*d*]pyrimidines as potential kinase inhibitors. Consequently, it is important to develop facile and efficient methods to synthesize this family of molecules. Retrosynthetic analysis suggests that 4,5,7-trisubstituted pyrrolo[3,2-*d*]pyrimidines can be synthesized through two general approaches (Scheme 1).

## Scheme 1. Approaches for Synthesis of Pyrrolopyrimidine Derivatives



FG = NO<sub>2</sub>, NH<sub>2</sub>, etc.  
X = Cl, Me



X = I, NO<sub>2</sub>, NH<sub>2</sub>, etc

Approach A is focused on the construction of the pyrrole ring using substituted pyrimidines as the starting materials.<sup>11</sup>

Although these methods are useful, the yields are poor for some cases.<sup>11b</sup> In addition, the substituents R<sup>1</sup> and R<sup>2</sup> on the pyrrolopyrimidine core have to be introduced prior to the ring formation, which renders further modification of these substituents difficult. In contrast, approach B starts with a simple pyrrolopyrimidine core and the substituents are introduced late in the synthesis, enabling exploration of diverse analogues possible at lower cost and in a timely fashion. However, to date there has been no reported method for this approach. In our efforts to employ pyrrolopyrimidines as selective kinase inhibitors, an efficient and robust method has been developed to synthesize 4,5,7-trisubstituted pyrrolo[3,2-*d*]pyrimidine analogues through approach B.

## RESULTS AND DISCUSSION

Compound **1** (Scheme 2) was chosen as the key building block for our synthesis for the following reasons. First, the chloride on the C4 position could be potentially displaced by various nucleophiles to introduce diversified R<sup>1</sup> substituents. Second, the NO<sub>2</sub>/NH<sub>2</sub> group at the C7 position could be used as a handle for introduction of N-containing R<sup>2</sup> substituents. Third, the R<sup>3</sup> group could be attached through *N*-alkylation at the N5 position.


The key building block **1** was synthesized by chlorination of the known compound 7-nitro-3*H*-pyrrolo[3,2-*d*]pyrimidin-4(5*H*)-one with neat phosphorus(V) oxychloride under reflux (Scheme 2). For 2-halopyrimidines, the halogen atom could be replaced smoothly by various nucleophiles.<sup>12</sup> However, the additional pyrrole ring in pyrrolopyrimidine makes the replacement reaction more difficult. With 2-(trimethylsilyl)ethoxymethyl (SEM) protecting group at nitrogen of the pyrrole ring,  $S_NAr$

Received: September 10, 2012

Revised: November 19, 2012

Published: November 26, 2012

Scheme 2. Synthesis of the Key Building block 1 and Its Transformations



displacement of chloride at the C4 position of **2** with 4-phenylphenol went smoothly at room temperature in high yield. The direct displacement of chloride in the key building block **1** with phenols, however, was challenging and only occurred after heating at 175 °C for 50 min under microwave irradiation. Because direct displacement of chloride in **1** would circumvent the need for protection and deprotection steps, this reaction was further explored (Table 1).

In the presence of potassium carbonate, phenol (entry 1), *para*-substituted phenols (entries 2 and 5), and *meta*-substituted phenol (entry 3) underwent the S<sub>N</sub>Ar substitution reactions smoothly with **1** in good yields, while *ortho*-substituted phenol (entry 4) gave poor yield possibly because of the steric hindrance. The ester functional group did not tolerate the reaction conditions (entry 6) and an aliphatic alcohol was not reactive under these conditions. However, conversion to an alkoxide (entry 7) completed the transformation at lower temperature (120 vs 175 °C) and in a shorter reaction time (20 min vs 50 min). Amines including aniline (entries 8–10) were better nucleophiles and reacted under much milder reaction conditions (conventional heating at 80 °C for overnight) with higher yields. More importantly, a few cross-coupling reactions also worked using **1** as a reactant which further diversified the available R<sup>1</sup> substituents. Accordingly, the R<sup>1</sup> substituents of **4** could be introduced either from a boronic acid/ester via Suzuki–Miyaura coupling conditions<sup>13</sup> (entries 11–15) or from a zinc reagent via the Negishi coupling reaction<sup>14</sup> (entries 16 and 17). In general, electron donating groups increase while electron withdrawing groups decrease reactivity of boronic acid in a Suzuki–Miyaura coupling reaction. For example, (2-methoxyphenyl)boronic acid (entry 12) provided product **4k** with higher yield than (3-fluorophenyl)boronic acid (*para*-methoxyl) (entry 13). In addition, furan could be introduced to the R<sup>1</sup> position (entry 14) and boronic ester (entry 15) also worked under the same reaction conditions. While Suzuki–Miyaura coupling reaction is suitable to introduce aromatic groups at the R<sup>1</sup> position, Negishi coupling reaction works better when R<sup>1</sup> is an alkyl group. As shown in Table 1, decent yields were obtained when two different zinc reagents were used (entries 16 and 17).

Next, we explored the introduction of R<sup>2</sup> substituents at the C7 position of pyrrolo[3,2-*d*]pyrimidines. Since coupling reactions had been successfully applied to introduce the C-substituents, such as aryl, heteroaryl, vinyl, alkynyl and benzyl, at this position in a similar system,<sup>15</sup> our focus was mainly on the introduction of the diversified N-substituents. The most efficient way to achieve this goal was through a one-pot reduction of the nitro group and in situ reductive amination with aldehydes under hydrogen atmosphere in the presence of palladium on activated charcoal. Using this one-pot protocol, **3** was converted to **5** in high yield (Scheme 3). The NHEt group at the C7 position in **5** was protected as a *tert*-butyl carbamate (Boc) (**6**) to minimize the potential interference with the next *N*-alkylation on the N5 position. Subsequently, the SEM protecting group in **6** was selectively removed by treatment with TBAF to provide **7**.

We were also able to introduce tertiary amines and amide groups at the C7 position of pyrrolopyrimidines by stepwise approaches as illustrated in Scheme 4. First, the nitro group in **4g** was reduced to the corresponding amine in **8**. Next, alkylation or acylation of **8** provided **9** and **10**, respectively. Although there are three NH groups in **8**, the newly formed primary amine is more reactive likely due to the low electron density and/or steric hindrance on the other two basic nitrogen atoms. The yield for the alkylated **9** was 81% while those for acylated **10** with various acid chlorides were lower because of competing *N*-diacylation at the same position.

Finally, R<sup>3</sup> substituents at the N5 position of pyrrolopyrimidines were introduced by *N*-alkylation (Table 2). Primary alkyl chlorides reacted with **7** at 150 °C for 15 min under microwave irradiation to provide the desired *N*-alkylated products (entries 1–3). For secondary chlorides (entry 4), a larger excess of halide (3 equivalents) and longer reaction time (30 min) were needed. Primary bromides (entries 5–8) also gave the desired *N*-alkylation products under the same reaction conditions as for primary chlorides. However, cyclohexyl bromide or iodide (entry 9) did not provide any desired alkylation product, presumably due to the elimination reaction of the bromide/iodide. The final products **11** were obtained by removal of the Boc group under acidic conditions.

A selectivity issue arises for compounds with a NHR group at the C4 position such as **4g** and **4i** (Table 1) in the final *N*-alkylation reaction because the nitrogen at the C4 position of the pyrimidine demonstrated reactivity similar to the pyrrole nitrogen. Accordingly, the nitrogen at the C4 position had to be protected early in the synthesis. For example, compound **4h** (Table 1) could serve as an intermediate for this purpose as the nitrogen at the C4 position was protected by a benzyl (Bn) group. *N*-alkylation reaction of **4h** with *trans-tert*-butyl (4-(bromomethyl)cyclohexyl)carbamate proceeded smoothly in good yield (Scheme 5). After the reduction of the nitro group in **12** and *N*-acylation of **13**, compound **14** was obtained. The Bn protecting group in **12** survived under the reductive hydrogenation conditions but could subsequently be removed under acidic conditions with higher temperature. The BOC protecting group was removed by trifluoroacetic acid (TFA) after the removal of the Bn group in **14** to provide **15**. Through this alternative route, the pyrrole nitrogen could be selectively alkylated.

We have also investigated the potential applications of the synthetic analogs as kinase inhibitors. Four structurally different compounds **9**, **10b**, **11a**, **15** were selected for the initial study. Each of the four compounds was tested for its capacity to inhibit a panel of 48 kinases using the Profiler Pro Kit (Caliper Life Sciences) at a concentration of 10 μM. All the experiments were carried out in duplicate. To our delight, the fms-like receptor tyrosine kinase-3 (FLT3) was the only kinase that was inhibited by more than 50% and only by **9** (details in Supporting Information). FLT3 is a validated target for drug discovery for a variety of diseases.<sup>16</sup> We thus further tested the inhibitory activity of **9** in a concentration-dependent manner. The inhibition of FLT3











kinase activity was measured at the ATP  $K_m$  using a microfluidic capillary electrophoresis (MCE) assay,<sup>17</sup> in which phosphorylated and unphosphorylated substrate peptides were separated and analyzed

through a LabChip EZ Reader. The  $IC_{50}$  of **9** in this assay was  $1.92 \mu M$ . Further SAR exploration to improve the potency of **9** is currently ongoing and will be reported in due course.

Table 1.  $S_NAr$  Substitution and Coupling Reactions of **1**


Entry	Reagent	Product	<b>4</b>	Yield (%)
1			<b>4a</b>	75 <sup>a</sup>
2			<b>4b</b>	76 <sup>a</sup>
3			<b>4c</b>	74 <sup>a</sup>
4			<b>4d</b>	32 <sup>a</sup>
5			<b>4e</b>	81 <sup>a</sup>
6				No desired product
7	n-PrOH		<b>4f</b>	72 <sup>b</sup>
8	n-BuNH <sub>2</sub>		<b>4g</b>	92 <sup>c</sup>
9			<b>4h</b>	90 <sup>c</sup>
10			<b>4i</b>	85 <sup>c</sup>
11			<b>4j</b>	87 <sup>d</sup>
12			<b>4k</b>	90 <sup>d</sup>

Table 1. continued


Entry	Reagent	Product	4	Yield (%)
13			4l	62 <sup>d</sup>
14			4m	90 <sup>d</sup>
15			4n	87 <sup>d</sup>
16			4o	64 <sup>e</sup>
17			4p	45 <sup>e</sup>

<sup>a</sup>1.5 equiv phenol, 3.0 equiv K<sub>2</sub>CO<sub>3</sub>, DMF,  $\mu$ W, 175 °C, 50 min. <sup>b</sup>1.2 equiv NaH, 2.0 equiv propanol, THF,  $\mu$ W, 120 °C, 20 min. <sup>c</sup>1.2 equiv aniline/amine, *i*-PrOH, 80 °C, overnight. <sup>d</sup>1.5 equiv boronic acid, 5% Pd(PPh<sub>3</sub>)<sub>4</sub>, DMF/H<sub>2</sub>O,  $\mu$ W, 150 °C, 15 min. <sup>e</sup>3 equiv benzylzinc bromide, toluene, 10% Pd(PPh<sub>3</sub>)<sub>4</sub>, 100 °C, overnight.

### Scheme 3. Introduction of Secondary Amine Groups at the C7 Position of Pyrrolopyrimidines



### Scheme 4. Introduction of Tertiary Amines and Amide Groups at the C7 Position of 8



## CONCLUSION

In summary, we have developed an efficient and robust route to prepare 4,5,7-trisubstituted pyrrolopyrimidines from the common intermediate **1**. The synthesis features a S<sub>N</sub>Ar substitution

reaction, cross-coupling reaction, one-pot reduction/reductive amination and *N*-alkylation reaction. These reactions occur rapidly with high yields and have broad substrate scopes. A variety of R<sup>2</sup> and R<sup>3</sup> groups can be selectively introduced into

Table 2. N-Alkylation at the N5 Position of 7

Entry	R <sup>3</sup> X	Product	11	Yield (%)
1			<b>11a</b>	93 <sup>a</sup>
2			<b>11b</b>	74 <sup>a</sup>
3			<b>11c</b>	72 <sup>a, b</sup>
4			<b>11d</b>	92 <sup>c</sup>
5			<b>11e</b>	74 <sup>a</sup>
6			<b>11f</b>	75 <sup>a</sup>
7			<b>11g</b>	76 <sup>a</sup>
8			<b>11h</b>	65 <sup>a</sup>
9				No desired product <sup>c</sup>


<sup>a</sup>1.5 equiv R<sup>3</sup>X, DMF,  $\mu$ W, 150 °C, 15 min. <sup>b</sup>In product **11c**, the Boc group on the aniline nitrogen has been removed as well under the treatment of TFA. <sup>c</sup>3 equiv halide, DMF,  $\mu$ W, 150 °C, 30 min.

4,5,7-trisubstituted pyrrolopyrimidines at a late stage of the synthesis. This synthetic strategy is highly desirable when large numbers of 4,5,7-trisubstituted pyrrolopyrimidine analogues are needed in high yields and purity for SAR development via biological evaluation. In addition, a new and selective FLT3 inhibitor **9** has been identified from the synthesized pyrrolopyrimidine analogues.

## EXPERIMENTAL PROCEDURES

**General Experimental Details.** All reagents were commercially available and used without any further purification. Microwave reaction was carried out using a Discover-S reactor with a vertically focused IR external temperature sensor and an Explorer

Scheme 5. Selective N-Alkylation at N5 Position in 4h



72 autosampler. The dynamic mode was used to set up the desired temperature and hold time with the following fixed parameters: PreStirring, 1 min; Pressure, 200 psi; Power, 200 W; PowerMax, off; Stirring, high. Flash chromatography was carried out on Teledyne ISCO Combi Flash Rf using prepacked silica gel disposable columns. A gradient from 0% to 100% ethylacetate (100–0% hexane) for nonpolar compounds or to 20% methanol (100–80% CH<sub>2</sub>Cl<sub>2</sub>) for polar compounds was used as elutes. Analytical thin-layer chromatography (TLC) was performed with silica gel 60 F<sub>254</sub> 0.25 mm precoated TLC plates. TLC plates were visualized using UV<sub>254</sub> or phosphomolybdic acid with charring. All <sup>1</sup>H NMR spectra were obtained with a 400 MHz spectrometer and <sup>13</sup>C NMR spectra were obtained with a 101 MHz spectrometer. Preparative HPLC was performed with the UV detection at 220 or 254 nm. LC-MS was performed with the UV detection at 220 nm, 254 nm, and 280 nm, and a single quadrupole mass spectrometer using electrospray ionization (ESI) source. High-resolution (positive ion) mass spectra (HRMS) were acquired using a LCMS-TOF mass spectrometer.

**4-Chloro-7-nitro-5H-pyrrolo[3,2-d]pyrimidine (1).** The solution of 7-nitro-3H-pyrrolo[3,2-d]pyrimidin-4(5H)-one (5.0 g, 27.8 mmol) in POCl<sub>3</sub> (12.6 g, 83.3 mmol) was heated under reflux for 5.0 h. Then the mixture was cooled to room temperature and poured onto the chipped ice with vigorous stirring, then, basified with K<sub>2</sub>CO<sub>3</sub> to pH 8–9. The resulting mixture was filtered and washed with water (2×) and EtOAc (3×) to provide the title compound (4.5 g, 82%) as yellow powder. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sup>6</sup>) δ 9.11 (s, 1H), 8.90 (s, 1H); <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sup>6</sup>) δ 153.1, 144.7, 142.8, 137.1, 128.1, 124.6; HRMS (TOF, ESI+) *m/z* [M + H]<sup>+</sup> calculated for C<sub>6</sub>H<sub>4</sub>ClN<sub>4</sub>O<sub>2</sub>, 199.0023; found 198.9869

**4-([1,1'-Biphenyl]-4-yloxy)-7-nitro-5-((2-(trimethylsilyl)ethoxy)methyl)-5H-pyrrolo[3,2-d]pyrimidine (3).** To a DMF (10 mL) solution of 1 (1.0 g, 5.0 mmol) and SEMCl (1.1 mL, 6.0 mmol) was added NaH (0.4 g, 60 wt % in mineral oil, 10 mmol) at 0 °C. The resulting mixture was warmed to room temperature and stirred overnight. The reaction was quenched by water. The aqueous phase was extracted by ether (2×) and EtOAc (2×). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was passed a short silica pad to provide 4-chloro-7-nitro-5-((2-(trimethylsilyl)ethoxy)methyl)-5H-pyrrolo[3,2-d]-

pyrimidine (2) (1.6 g), which was used directly for the next step. A mixture of 2 (1.6 g, 4.88 mmol), [1,1'-biphenyl]-4-ol (1.66 g, 9.76 mmol), and K<sub>2</sub>CO<sub>3</sub> (2.02 g, 14.64 mmol) in 25 mL of DMF was stirred for 5.0 h at room temperature. The reaction was quenched by water and extracted with ether (3×). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was purified by column chromatography with ISCO system to provide the title compound 3 (2.1 g, 91% over 2 steps) as a light yellow solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.78 (s, 1H), 8.45 (s, 1H), 7.73–7.67 (m, 2H), 7.63–7.58 (m, 2H), 7.49–7.43 (m, 2H), 7.40–7.35 (m, 1H), 7.34–7.29 (m, 2H), 5.87 (s, 2H), 3.77–3.65 (m, 2H), 1.03–0.93 (m, 2H), –0.02 (s, 9H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 156.3, 154.0, 150.8, 144.1, 140.2, 139.8, 133.8, 129.1, 129.0, 128.7, 127.7, 127.3, 122.1, 115.2, 79.4, 67.8, 18.0, –1.3; LC-MS (ESI+) *t*<sub>R</sub> = 6.285 min, *m/z* 463.2 [M + 1]<sup>+</sup>.

**7-Nitro-4-phenoxy-5H-pyrrolo[3,2-d]pyrimidine (4a).** *General Procedure A.* A 10 mL microwave tube was charged with 1 (92 mg, 0.46 mmol), K<sub>2</sub>CO<sub>3</sub> (193 mg, 1.4 mmol), anhydrous DMF (2.5 mL), and phenol (66 mg, 0.70 mmol). The resulting mixture was heated at 175 °C for 50 min under microwave irradiation. After it was cooled to room temperature, the reaction mixture was diluted with EtOAc and water. The aqueous phase was extracted with EtOAc (3×). The combined organic extracts were washed with brine and dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated under reduced pressure. The residue was purified by column chromatography with ISCO system to provide the title compound 4a (88 mg, 75%) as a white solid. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sup>6</sup>) δ 13.83 (s, 1H), 8.94 (s, 1H), 8.56 (s, 1H), 7.52–7.45 (m, 2H), 7.35–7.28 (m, 3H); <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sup>6</sup>) δ 156.2, 153.0, 152.1, 143.3, 134.3, 130.3, 128.3, 126.4, 122.5, 115.1; HRMS (TOF, ESI+) *m/z* [M + H]<sup>+</sup> calculated for C<sub>12</sub>H<sub>9</sub>N<sub>4</sub>O<sub>3</sub>, 257.0675; found 257.0659.

**7-Nitro-4-propoxy-5H-pyrrolo[3,2-d]pyrimidine (4f).** To a suspension of NaH (23.1 mg, 60% in mineral oil, 0.57 mmol) in anhydrous THF (2.0 mL) was added propanol (57.6 mg, 0.96 mmol) at 0 °C in a 10 mL microwave tube. After it was stirred for 30 min at 0 °C, 1 (95 mg, 0.48 mmol) was added, and the resulting mixture was heated at 120 °C for 20 min under microwave irradiation. After it was cooled to room temperature, the reaction was diluted with EtOAc and water. The aqueous phase was extracted with EtOAc (3×). The combined organic

extracts were dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The residue was purified by column chromatography with ISCO system to provide the title compound **4f** (77 mg, 72%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  13.49 (s, 1H), 8.76 (s, 1H), 8.61 (s, 1H), 4.49 (t,  $J = 6.6$  Hz, 2H), 1.86–1.76 (m, 2H), 1.01 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{DMSO}-d_6$ )  $\delta$  156.6, 153.3, 142.2, 133.2, 128.2, 115.0, 68.4, 22.2, 10.7; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_9\text{H}_{11}\text{N}_4\text{O}_3$ , 223.0831; found 223.0810.

**N-Butyl-7-nitro-5H-pyrrolo[3,2-d]pyrimidin-4-amine (4g).** *General Procedure B.* To a solution of **1** (200 mg, 1.01 mmol) in anhydrous *i*-PrOH (15 mL) was added butylamine (89 mg, 1.21 mmol). The resulting mixture was heated at 80 °C for overnight under nitrogen atmosphere. After cooling to room temperature, the reaction was diluted with EtOAc and water. The aqueous phase was extracted with EtOAc (3 $\times$ ). The combined organic extracts were washed with brine and dried ( $\text{Na}_2\text{SO}_4$ ), concentrated under reduced pressure. The residue was purified by column chromatography with ISCO system to provide the title compound **4g** (215 mg, 92%) as a yellow solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3 + \text{CD}_3\text{OD}$ )  $\delta$  8.21 (s, 1H), 8.05 (s, 1H), 3.55 (t,  $J = 7.1$  Hz, 2H), 1.64–1.56 (m, 2H), 1.42–1.33 (m, 2H), 0.89 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3 + \text{CD}_3\text{OD}$ )  $\delta$  151.8, 147.4, 129.9, 127.3, 125.2, 112.8, 41.1, 30.6, 19.6, 12.6; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{10}\text{H}_{14}\text{N}_5\text{O}_2$ , 236.1147; found 236.1128.

**7-Nitro-4-phenyl-5H-pyrrolo[3,2-d]pyrimidine (4j).** *General procedure C.* A 10 mL microwave tube was charged with **1** (150 mg, 0.76 mmol),  $\text{K}_2\text{CO}_3$  (262 mg, 1.90 mmol), phenylboronic acid (139 mg, 1.14 mmol),  $\text{Pd}(\text{PPh}_3)_4$  (44 mg, 0.038 mmol), DMF (2.0 mL) and  $\text{H}_2\text{O}$  (1.0 mL). The resulting mixture was stirred at room temperature for 3.0 min and then heated at 150 °C for 15 min. After it was cooled to room temperature, the mixture was partitioned in  $\text{H}_2\text{O}$  and  $\text{Et}_2\text{O}$ . The aqueous phase was extracted with  $\text{Et}_2\text{O}$  (3 $\times$ ). The combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The residue was purified by column chromatography with ISCO system to provide the title compound **4j** (159 mg, 87%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.14 (s, 1H), 9.00 (s, 1H), 8.05–8.01 (m, 2H), 7.64–7.60 (m, 3H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{DMSO}-d_6$ )  $\delta$  153.7, 151.0, 142.8, 136.8, 135.0, 131.4, 129.5, 129.4, 129.2, 127.7; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{12}\text{H}_9\text{N}_4\text{O}_2$ , 241.0726; found 241.0719.

**4-Benzyl-7-nitro-5H-pyrrolo[3,2-d]pyrimidine (4o).** *General Procedure D.* To a solution of **1** (96 mg, 0.48 mmol) and  $\text{Pd}(\text{PPh}_3)_4$  (112 mg, 0.097 mmol) in anhydrous toluene (12 mL) was added a 0.5 M solution of benzylzinc(II) bromide in THF (2.0 mL) at room temperature. The resulting mixture was heated at 110 °C for 12 h and quenched by water (4.0 mL). The solvent was removed under the reduced pressure. The residue was dissolved in water and extracted with EtOAc (3 $\times$ ). The combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated under reduced pressure. The residue was purified by column chromatography with ISCO system to give the title compound **4o** (78 mg, 64%) as a yellow solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD} + \text{CDCl}_3$ )  $\delta$  9.00 (s, 1H), 8.43 (s, 1H), 7.25–7.11 (m, 5H), 4.35 (s, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD} + \text{CDCl}_3$ )  $\delta$  154.5, 153.4, 141.3, 136.1, 133.8, 128.7, 128.7, 127.1, 125.7, 39.2; LC-MS (ESI+)  $t_R = 4.558$  min,  $m/z$  255.10  $[\text{M} + 1]^+$ ; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{13}\text{H}_{11}\text{N}_4\text{O}_2$ , 255.0882; found 255.0879.

**tert-Butyl (4-([1,1'-biphenyl]-4-yloxy)-5H-pyrrolo[3,2-d]pyrimidin-7-yl)(ethyl)carbamate (7).** A mixture of **3** (46 mg,

0.10 mmol) and Pd/C (4.6 mg, 10 wt % Pd/C) in MeOH (5.0 mL) was added a solution of acetaldehyde (4.4 mg, 0.10 mmol) in MeOH (0.5 mL) drop-wisely under hydrogen atmosphere. After 4.0 h, the resulting mixture was filtered over a Celite pad and washed with MeOH. After removal of MeOH solvent, the crude product was dissolved in 2.0 mL  $\text{CH}_2\text{Cl}_2$ . To this solution was added  $\text{Et}_3\text{N}$  (20 mg, 0.20 mmol) and  $\text{Boc}_2\text{O}$  (65 mg, 0.3 mmol). The reaction was stirred 6.0 h at room temperature and quenched with a saturated aqueous  $\text{NH}_4\text{Cl}$  solution. After extraction with  $\text{CH}_2\text{Cl}_2$  (3 $\times$ ), the combined organic layers were concentrated. The resulting residue was purified by a short silica column to provide **6** (50 mg), which was used directly for the next step. To a solution of **6** (50 mg, 0.088 mmol) in THF (0.5 mL) was added a solution of TBAF (0.22 mL, 1.0 M in THF, 0.22 mmol) and ethylenediamine (6.6 mg, 0.11 mmol) at room temperature. After refluxing for 1.0 h, the reaction mixture was diluted with EtOAc and washed with water. The combined organic layers are dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The residue was purified by column chromatography with ISCO system to provide the title compound **7** (25 mg, 58% over 3 steps) as a light yellow solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.50 (bs, 1H), 7.86 (bs, 1H), 7.77–7.70 (m, 2H), 7.68–7.62 (m, 2H), 7.49–7.43 (m, 2H), 7.42–7.33 (m, 3H), 3.74 (q,  $J = 7.1$  Hz, 2H), 1.46 (bd, 9H), 1.20 (bs, 3H); LC-MS (ESI+)  $t_R = 5.927$  min,  $m/z$  431.2  $[\text{M} + 1]^+$ .

***N*<sup>4</sup>-Butyl-5H-pyrrolo[3,2-d]pyrimidine-4,7-diamine (8).** The solution of **4g** (85 mg, 0.36 mmol) and 10% palladium on activated carbon (53.9 mg, 0.30 mmol) in MeOH (15.0 mL) was stirred under hydrogen atmosphere at room temperature for 3.0 h. Then the reaction mixture was filtered through a pad of Celite to afford the title compound **8** (68 mg, 92%) as a brown oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.20 (s, 1H), 7.03 (s, 1H), 3.58 (t,  $J = 7.1$  Hz, 2H), 1.75–1.60 (m, 3H), 1.56–1.43 (m, 2H), 0.99 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  150.0, 147.5, 135.2, 121.6, 114.9, 112.6, 40.1, 31.2, 19.7, 12.7; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{10}\text{H}_{16}\text{N}_5$ , 206.1406; found 206.1396.

***N*-Butyl-7-(piperazin-1-yl)-5H-pyrrolo[3,2-d]pyrimidin-4-amine (9).** Bis(2-chloroethyl)amine (95.6 mg, 0.54 mmol) was added into the solution of **8** (110.1 mg, 0.54 mmol) in *i*-PrOH (10.0 mL) at room temperature. Then  $\text{Na}_2\text{CO}_3$  (114.5 mg, 1.1 mmol) was added, the resulting reaction mixture was stirred overnight at reflux. Then water (20 mL) was added and extracted with dichloromethane (3 $\times$ ). The combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and condensed. The residue was purified by column chromatography with ISCO system to provide the title compound **9** (118.9 mg, 81%) as red oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.13 (s, 1H), 6.98 (s, 1H), 3.52 (t,  $J = 7.1$  Hz, 2H), 3.17–3.08 (m, 4H), 3.09–2.99 (m, 4H), 1.72–1.62 (m, 2H), 1.51–1.42 (m, 2H), 0.98 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  150.0, 148.4, 137.9, 130.8, 113.8, 113.5, 51.7, 44.7, 40.0, 31.2, 19.8, 12.7; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{14}\text{H}_{23}\text{N}_6$ , 275.1984; found 275.1978.

***N*-(4-(Butylamino)-5H-pyrrolo[3,2-d]pyrimidin-7-yl)-benzamide (10a).** *General Procedure E.* To the solution of **8** (0.30 mmol, 1.0 equiv), DIEA (0.45 mmol, 1.5 equiv) and dichloromethane (3.0 mL) was added benzoyl chloride (13 mg, 0.090 mmol) at room temperature. After it was stirred for 8.0 h at room temperature, the reaction was quenched with a sat.  $\text{NaHCO}_3$  solution (5.0 mL) and extracted with dichloromethane (3 $\times$ ). The combined organic phases were dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated under the reduced pressure. The residue was purified through prep-HPLC to provide the title compound **10a** (10 mg, 39%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$

12.78 (s, 1H), 10.45 (s, 1H), 8.85 (d,  $J = 5.5$  Hz, 1H), 8.23 (s, 1H), 8.02–7.92 (m, 2H), 7.81 (d,  $J = 2.9$  Hz, 1H), 7.49 (t,  $J = 7.4$  Hz, 1H), 7.38 (t,  $J = 7.7$  Hz, 2H), 3.64 (dd,  $J = 13.0, 7.2$  Hz, 2H), 1.69–1.62 (m, 2H), 1.40 (dq,  $J = 14.6, 7.3$  Hz, 2H), 0.93 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  166.5, 151.6, 142.9, 132.4, 132.3, 128.6, 127.7, 124.3, 120.9, 112.7, 112.5, 41.5, 30.9, 19.9, 13.6; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{17}\text{H}_{20}\text{N}_5\text{O}$ , 310.1668; found 310.1664.

**4-([1,1'-Biphenyl]-4-yloxy)-*N*-ethyl-5-(3-morpholinopropyl)-5H-pyrrolo[3,2-*d*]pyrimidin-7-amine (11a).** General Procedure F. A mixture of **7** (65 mg, 0.15 mmol),  $\text{K}_2\text{CO}_3$  (62 mg, 0.45 mmol), 4-(3-chloropropyl)morpholine (40 mg, 0.23 mmol), and DMF (2 mL) in a microwave tube was heated under microwave irradiation at 150 °C for 10 min. After it was cooled to room temperature, the mixture was washed with brine and extracted with EtOAc (3 $\times$ ). The combined organic layers was dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The residue was purified by column chromatography with ISCO system to provide the desired intermediate. This intermediate was dissolved in a mixture of 2.0 mL of  $\text{CH}_2\text{Cl}_2$  and 0.5 mL of TFA. After it was stirred at room temperature for 2.0 h, the solution was concentrated and purified through prep-HPLC to provide the title compound **11a** (64 mg, 93%) as a yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.43 (s, 1H), 8.08 (s, 1H), 7.77–7.70 (m, 2H), 7.67–7.60 (m, 2H), 7.48–7.32 (m, 5H), 4.67 (t,  $J = 7.0$  Hz, 2H), 3.97 (bs, 2H), 3.76 (bs, 2H), 3.65 (q,  $J = 7.3$  Hz, 2H), 3.48 (bs, 2H), 3.30–3.24 (m, 2H), 3.11 (bs, 2H), 2.54–2.39 (m, 2H), 1.40 (t,  $J = 7.3$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  157.0, 152.6, 151.2, 145.23, 141.4, 140.6, 130.0, 129.4, 128.6, 128.0, 123.5, 116.6, 116.0, 113.6, 64.9, 55.4, 53.1, 47.9, 47.1, 27.1, 11.7; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{27}\text{H}_{32}\text{N}_5\text{O}_2$ , 458.2556; found 458.2536.

***tert*-Butyl ((1*r*,4*r*)-4-((4-(Benzyl(ethyl)amino)-7-nitro-5H-pyrrolo[3,2-*d*]pyrimidin-5-yl)methyl)cyclohexyl)carbamate (12).** A 10 mL microwave tube was charged with **4h** (297 mg, 1.0 mmol),  $\text{K}_2\text{CO}_3$  (345 mg, 2.5 mmol), DMF (2.0 mL), and *trans-tert*-butyl 4-(bromomethyl)cyclohexylcarbamate (436 mg, 1.5 mmol). The resulting mixture was heated at 150 °C for 45 min under microwave irradiation. After it was cooled to room temperature, the reaction was diluted with EtOAc and washed with brine. The aqueous phase was extracted with EtOAc (3 $\times$ ). The combined organic extracts were dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure. The residue was purified by column chromatography with ISCO system to provide the title compound **12** (361 mg, 71%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.50 (s, 1H), 7.44–7.22 (m, 6H), 4.61 (d,  $J = 6.9$  Hz, 2H), 3.34 (s, 5H), 1.91 (d,  $J = 11.9$  Hz, 2H), 1.69–1.60 (m, 3H), 1.41 (s, 9H), 1.34–1.29 (m, 3H), 1.20–1.08 (m, 4H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  156.3, 151.9, 147.3, 138.8, 136.3, 128.5, 128.2, 127.5, 127.4, 125.8, 110.0, 78.6, 59.5, 52.1, 49.3, 48.7, 44.5, 37.1, 31.8, 28.1, 27.6; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{27}\text{H}_{37}\text{N}_6\text{O}_4$ , 509.2876; found 509.2869.

***tert*-Butyl ((1*r*,4*r*)-4-((7-amino-4-(benzyl(ethyl)amino)-5H-pyrrolo[3,2-*d*]pyrimidin-5-yl)methyl)cyclohexyl)carbamate (13).** To a suspension of 10% palladium on active carbon (13 mg, 0.013 mmol) in methanol (8.0 mL) was added **12** (125 mg, 0.25 mmol) at room temperature. The resulting mixture was stirred under hydrogen atmosphere for 3.0 h. Then the mixture was diluted with EtOAc and filtered through a pad of Celite and condensed. The residue was purified by column chromatography with ISCO system to provide the title compound **13** (100 mg, 84%) as a green oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.35 (s, 1H), 7.41 (d,  $J = 4.6$  Hz, 1H), 7.37–7.26 (m, 5H), 5.17 (s, 2H),

4.44 (d,  $J = 7.3$  Hz, 2H), 3.93–3.84 (m, 2H), 3.34 (s, 3H), 1.93 (d,  $J = 10.7$  Hz, 2H), 1.69 (d,  $J = 11.9$  Hz, 2H), 1.42 (s, 9H), 1.32 (t,  $J = 7.2$  Hz, 3H), 1.25–1.13 (m, 4H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  156.4, 150.3, 145.7, 136.1, 128.8, 128.5, 127.5, 127.2, 123.9, 118.8, 111.1, 78.5, 56.2, 52.0, 49.4, 48.4, 44.5, 39.5, 37.8, 31.7, 28.2, 27.4, 12.4; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{27}\text{H}_{39}\text{N}_6\text{O}_2$ , 479.3134; found 479.3128.

***tert*-Butyl ((1*r*,4*r*)-4-((4-(Ethyl(phenethyl)amino)-7-(4-methoxybenzamido)-5H-pyrrolo[3,2-*d*]pyrimidin-5-yl)methyl)cyclohexyl)carbamate (14).** To a solution of **13** (80 mg, 0.17 mmol), triethyl amine (25.3 mg, 0.25 mmol), catalytic amount of DMAP in anhydrous THF (3.0 mL) was added 4-methoxybenzoyl chloride (28.4 mg, 0.17 mmol) at room temperature. The resulting mixture was stirred for 5.0 h and quenched with water and diluted with EtOAc (20 mL). The aqueous phase was extracted with EtOAc (3 $\times$ ). The combined organic phase was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and condensed. The residue was purified by column chromatography with ISCO system to give the title compound **14** (53 mg, 51% yield) as a yellow solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.90 (d,  $J = 7.8$  Hz, 2H), 7.65 (s, 1H), 7.56 (s, 1H), 7.35–7.19 (m, 5H), 6.94 (d,  $J = 8.7$  Hz, 2H), 5.29 (s, 2H), 5.13–5.00 (m, 1H), 4.44 (s, 1H), 3.96 (d,  $J = 6.9$  Hz, 2H), 3.86 (s, 3H), 3.37–3.25 (m, 1H), 1.94–1.81 (m, 3H), 1.51–1.45 (m, 2H), 1.41 (s, 9H), 1.35–1.18 (m, 3H), 0.96 (t,  $J = 9.7$  Hz, 4H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  169.1, 163.0, 156.4, 151.3, 142.8, 137.9, 131.0, 129.2, 128.1, 127.3, 127.0, 125.6, 113.5, 105.7, 78.5, 56.3, 54.6, 36.7, 31.6, 28.6, 27.3; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{35}\text{H}_{45}\text{N}_6\text{O}_4$ , 613.3502; found 613.3498.

***N*-(5-(((1*r*,4*r*)-4-Aminocyclohexyl)methyl)-4-(ethylamino)-5H-pyrrolo[3,2-*d*]pyrimidin-7-yl)-4-methoxybenzamide (15).** To a solution of compound **14** (46 mg, 0.075 mmol) in methanol (5.0 mL) was added 10% palladium on activated carbon (5 mg, 0.0047 mmol) at room temperature. The resulting mixture was added 3 drops of acetic acid and was then heat at 60 °C under  $\text{H}_2$  atmosphere for 6.0 h. The reaction mixture was then diluted with EtOAc (20 mL) and filtered through a pad of Celite. The solvent was removed and the residue was purified by column chromatography with ISCO system to give the debenzylated intermediate (22 mg, 56%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.47 (s, 1H), 8.02 (d,  $J = 8.9$  Hz, 2H), 7.82 (s, 1H), 7.09 (d,  $J = 8.9$  Hz, 2H), 4.28 (d,  $J = 7.2$  Hz, 2H), 3.89 (s, 3H), 3.80 (q,  $J = 7.2$  Hz, 2H), 2.97 (s, 1H), 1.79–1.76 (m, 4H), 1.50 (s, 2H), 1.37 (t,  $J = 7.3$  Hz, 3H), 1.24–1.07 (m, 5H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  169.1, 163.5, 150.7, 148.1, 129.8, 129.3, 124.6, 113.7, 112.8, 109.1, 56.6, 54.7, 37.1, 36.1, 31.5, 28.4, 27.3, 13.0. To a solution of the debenzylated intermediate (15 mg, 0.029 mmol) in dichloromethane (5.0 mL) was added trifluoroacetic acid (32.7 mg, 0.29 mmol) at room temperature. The resulting mixture was heated at 60 °C for 4.0 h. Then the reaction was diluted with dichloromethane (22.0 mL) and washed with  $\text{NaHCO}_3$  (sat.), water and brine sequentially. The organic phase was dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and condensed. The residue was purified through preparative HPLC to give the title compound **15** (9.2 mg, 75% yield) as a clear oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.47 (s, 1H), 8.02 (d,  $J = 8.9$  Hz, 2H), 7.82 (s, 1H), 7.09 (d,  $J = 8.9$  Hz, 2H), 4.28 (d,  $J = 7.2$  Hz, 2H), 3.89 (s, 3H), 3.80 (q,  $J = 7.2$  Hz, 2H), 2.97 (s, 1H), 1.95–1.75 (m, 4H), 1.50 (s, 2H), 1.53–1.46 (m, 3H), 1.24–1.07 (m, 5H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  168.9, 163.5, 150.8, 148.0, 130.1, 129.8, 129.3, 124.5, 113.8, 112.8, 108.9, 56.1, 54.7, 49.5, 36.5, 36.1, 29.3, 27.4, 13.0; HRMS (TOF, ESI+)  $m/z$   $[\text{M} + \text{H}]^+$  calculated for  $\text{C}_{23}\text{H}_{31}\text{N}_6\text{O}_2$ , 423.2508; found 423.2518.



**Selectivity Profiling.** The selected compounds were screened against 48 kinases using the Profiler Pro Kit (Caliper Life Sciences). Briefly, preplated enzyme stocks were reconstituted with 15  $\mu\text{L}$  of supplied reconstitution buffer containing DTT and Protease Inhibitor Cocktail. One microliter of compound at a concentration of 260  $\mu\text{M}$  in 100% DMSO was transferred to the plate and the reaction was initiated by the addition of 10  $\mu\text{L}$  substrate solution containing ATP and cofactors specific for each enzyme. The reaction was incubated at room temperature for 90 min and 45  $\mu\text{L}$  of termination buffer was then added to stop the reaction. The plates were read in a Caliper EZ Reader.

**Microfluidic Capillary Electrophoresis (MCE) Assay.** The activity assays were performed in a 384 well, polypropylene microtiterplate, using 0.3 nM FLT-3 (Life Technologies catalog PHC9415), in a final volume of 50  $\mu\text{L}$  of 50 mM HEPES pH 7.4 containing 10 mM  $\text{MgCl}_2$ , 1 mM DTT, 0.01% Triton X-100, 0.1% bovine serum albumin (BSA), 1  $\mu\text{M}$  fluorescent peptide substrate (5-FAM-KKKKEEYFFF-CONH<sub>2</sub>) and 275  $\mu\text{M}$  ATP (at  $K_m$  for Flt3). All reactions were terminated following a 60-min incubation, by addition of 20  $\mu\text{L}$  of 70 mM EDTA. Phosphorylated and unphosphorylated substrate peptides were separated on a LabChip EZ Reader equipped with a 12-sipper chip in separation buffer supplemented with CR-8 and analyzed using EZ Reader software.

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

Experimental procedures, characterization of all compounds and biological methods, and copies of <sup>1</sup>H and <sup>13</sup>C spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*Tel: 919-843-8456. Fax: 919-843-8465. E-mail: [xiaodonw@unc.edu](mailto:xiaodonw@unc.edu)

### Author Contributions

X.W., W.Z., and J.L. conceived and designed the experiments, W.Z., J.L., and M.S. performed the experiments, X.W., W.Z., and J.L. wrote the manuscript and Supporting Information.

### Funding

This work was supported by the University Cancer Research Fund, School of Pharmacy, and Federal Funds from the National Cancer institute, National Institute of Health, under Contract No. HHSN261200800001E. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank Professor Stephen Frye for comments on manuscript and Professor K. H. Lee's lab for HRMS support.

## ■ REFERENCES

(1) Imamura, S.; Oguro, Y. Preparation of fused heterocyclic derivatives, particularly pyrrolo[3,2-d]pyrimidines and pyrrolo[3,2-b]pyridines, and their use for treating cancer. Int. Patent WO2007004749A1, 2007.

(2) Oguro, Y.; Miyamoto, N.; Takagi, T.; Okada, K.; Awazu, Y.; Miki, H.; Hori, A.; Kamiyama, K.; Imamura, S. N-Phenyl-N'-[4-(5H-pyrrolo[3,2-d]pyrimidin-4-yloxy)phenyl]ureas as novel inhibitors of VEGFR and FGFR kinases. *Bioorg. Med. Chem.* **2010**, *18* (20), 7150–7163.

(3) Saavedra, O. M.; Claridge, S. W.; Zhan, L.; Raepfel, F.; Vaisburg, A.; Raepfel, S.; Deziel, R.; Mannion, M.; Zhou, N. Z.; Isakovic, L. Thienopyridine and thienopyrimidine derivatives and their preparation, pharmaceutical compositions, and use as inhibitors of VEGF receptor and HGF receptor signaling for treatment of proliferative diseases. Int. Patent WO2007054831A2, 2007.

(4) Blake, J. F.; Kallan, N. C.; Xiao, D.; Xu, R.; Bencsik, J. R.; Skelton, N. J.; Spencer, K. L.; Mitchell, I. S.; Woessner, R. D.; Gloor, S. L.; Risom, T.; Gross, S. D.; Martinson, M.; Morales, T. H.; Vigers, G. P. A.; Brandhuber, B. J. Discovery of pyrrolopyrimidine inhibitors of Akt. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 5607–5612.

(5) Bluhm, H.; Hochguertel, M.; Kroth, H.; Essers, M.; Gege, C.; Richter, F.; Taveras, A. Heterobicyclic matrix metalloprotease inhibitors, their preparation, pharmaceutical compositions, and use in therapy. Int. Patent WO2008063669A1, 2008.

(6) Flanagan, M. E.; Blumenkopf, T. A.; Brissette, W. H.; Brown, M. F.; Casavant, J. M.; Chang, S.-P.; Doty, J. L.; Elliott, E. A.; Fisher, M. B.; Hines, M.; Kent, C.; Kudlacz, E. M.; Lillie, B. M.; Magnuson, K. S.; McCurdy, S. P.; Munchhof, M. J.; Perry, B. D.; Sawyer, P. S.; Strelevitz, T. J.; Subramanyam, C.; Sun, J.; Whipple, D. A.; Changelian, P. S. Discovery of CP-690,550: A potent and selective janus kinase (JAK) inhibitor for the treatment of autoimmune diseases and organ transplant rejection. *J. Med. Chem.* **2010**, *53*, 8468–8484.

(7) Calderwood, D. J.; Johnston, D. N.; Munschauer, R.; Rafferty, P. Pyrrolo[2,3-d]pyrimidines containing diverse N-7 substituents as potent inhibitors of Lck. *Bioorg. Med. Chem. Lett.* **2002**, *12* (12), 1683–6.

(8) Evans, G. B.; Furneaux, R. H.; Lenz, D. H.; Painter, G. F.; Schramm, V. L.; Singh, V.; Tyler, P. C. Second generation transition state analogue inhibitors of human 5'-methylthioadenosine phosphorylase. *J. Med. Chem.* **2005**, *48*, 4679–4689.

(9) Morris, P. E., Jr.; Elliott, A. J.; Walton, S. P.; Williams, C. H.; Montgomery, J. A. Synthesis and biological activity of a novel class of purine nucleoside phosphorylase inhibitors. *Nucleosides, Nucleotides* **2000**, *19*, 379–404.

(10) Girardet, J. L.; Koh, Y.-H.; Shaw, S.; Kin, H. W. Preparation of purines, azapurines, and deazapurines as non-nucleoside reverse transcriptase inhibitors for treatment of HIV infection. Int. Patent WO2006122003A2, 2006.

(11) (a) Zhao, L.; Tao, K.; Li, H.; Zhang, J. Practical one-pot protocol for the syntheses of 2-chloro-pyrrolo[3,2-d]pyrimidines. *Tetrahedron* **2011**, *67*, 2803–2806. (b) Norman, M. H.; Chen, N.; Chen, Z.; Fotsch, C.; Hale, C.; Han, N.; Hurt, R.; Jenkins, T.; Kincaid, J.; Liu, L.; Lu, Y.; Moreno, O.; Santora, V. J.; Sonnenberg, J. D.; Karbon, W. Structure–activity relationships of a series of pyrrolo[3,2-d]pyrimidine derivatives and related compounds as neuropeptide Y5 receptor antagonists. *J. Med. Chem.* **2000**, *43*, 4288–4312.

(12) (a) Cherng, Y.-J. Synthesis of substituted pyridines by the reactions of halopyridines with sulfur, oxygen and carbon nucleophiles under focused microwave irradiation. *Tetrahedron* **2002**, *58* (24), 4931–4935. (b) Luo, G.; Chen, L.; Poindexter, G. S. Microwave-assisted synthesis of aminopyrimidines. *Tetrahedron Lett.* **2002**, *43* (33), 5739–5742.

(13) Miyaura, N.; Yamada, K.; Suzuki, A. A new stereospecific cross-coupling by the palladium-catalyzed reaction of 1-alkenylboranes with 1-alkenyl or 1-alkynyl halides. *Tetrahedron Lett.* **1979**, 3437–3440.

(14) King, A. O.; Okukado, N.; Negishi, E. Highly general stereo-, regio-, and chemo-selective synthesis of terminal and internal conjugated enynes by the palladium-catalyzed reaction of alkynylzinc reagents with alkenyl halides. *J. Chem. Soc., Chem. Commun.* **1977**, 683–684.

(15) Bambuch, V.; Otmár, M.; Pohl, R.; Masojdková, M.; Holy, A. C-Functionalization of 9-deazapurines by cross-coupling reactions. *Tetrahedron* **2007**, *63*, 1589–1601.

(16) Wiernik, P. H. FLT3 inhibitors for the treatment of acute myeloid leukemia. *Clin. Adv. Hematol. Oncol.* **2010**, *8* (6), 429-36–444.

(17) (a) Pommereau, A.; Pap, E.; Kannt, A. Two simple and generic antibody-independent kinase assays: comparison of a bioluminescent and a microfluidic assay format. *J. Biomol. Screening* **2004**, *9* (5), 409–16.

(b) Dunne, J.; Reardon, H.; Trinh, V.; Li, E.; Farinas, J. Comparison of on-chip and off-chip microfluidic kinase assay formats. *Assay Drug Dev. Technol.* **2004**, *2* (2), 121–9. (c) Bernasconi, P.; Chen, M.; Galasinski, S.; Popa-Burke, I.; Bobasheva, A.; Coudurier, L.; Birkos, S.; Hallam, R.; Janzen, W. P. A chemogenomic analysis of the human proteome: Application to enzyme families. *J. Biomol. Screening* **2007**, *12* (7), 972–82.