

Combining the Petasis 3-Component Reaction with Multiple Modes of Cyclization: A Build/Couple/Pair Strategy for the Synthesis of Densely Functionalized Small Molecules

Thomas Flagstad,[†] Mette R. Hansen,[†] Sebastian T. Le Quement,[†] Michael Givskov,^{‡,§} and Thomas E. Nielsen^{*,†,§}

[†]Department of Chemistry, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark


[‡]Department of International Health, Immunology and Microbiology, Faculty of Health and Medical Sciences, University of Copenhagen, DK-2200 Copenhagen, Denmark

[§]Singapore Centre on Environmental Life Sciences Engineering, Nanyang Technological University, Singapore 637551, Singapore

Supporting Information

ABSTRACT: A *build/couple/pair* strategy for the synthesis of complex and densely functionalized small molecules is presented. The strategy relies on synthetically tractable building blocks (*build*), that is, diversely substituted hydrazides, α -hydroxy aldehydes, and boronic acids, which undergo Petasis 3-component reactions (*couple*) to afford densely functionalized *anti*-hydrazido alcohols. The resulting scaffolds can subsequently be converted via chemoselective cyclization reactions (*pair*), including intramolecular Diels–Alder or Ru-alkylidene catalyzed ring-closing metathesis, into sets of structurally diverse heterocycles in good yields in only 3–4 steps.

KEYWORDS: Petasis 3-component reaction, intramolecular Diels–Alder reaction, ring-closing metathesis, *build/couple/pair*, diversity-oriented synthesis



In recent years, new synthetic concepts and methodologies have been proposed for the design and synthesis of molecular screening libraries.¹ Challenged by a tendency of traditional vendor libraries to incorporate simple and relatively “flat” molecules, as defined by their high content of sp^2 -hybridized carbon,² evidence suggests that many available compound collections are not meeting the structural requirements of more demanding targets, such as protein–protein interactions, transcription factors and nucleic acid macromolecules.^{1b} Some particularly successful strategies have heralded molecular diversity as a design parameter to access target-relevant chemical space. These strategies typically rely on short synthetic pathways yielding skeletally and stereochemically diverse molecular scaffolds, frequently drawing upon structural inspiration from natural products.³ By maximizing the number of stereogenic carbon atoms in these scaffolds, emerging trends emphasize that spatially well-defined molecules more smoothly may be elaborated into selective modulators of biological targets.⁴ In an effort to meet some of these challenging goals, diversity-oriented synthesis (DOS) tactics referred to as *build/couple/pair* (B/C/P) strategies have recently been proposed.^{1e,5}

We herein report our efforts toward the development of a synthetic pathway that starts with readily available building

blocks (*build*), comprising diverse mono-, di-, and trisubstituted hydrazides, α -hydroxy aldehydes and boronic acids, and combines the use of robust organic reactions, namely, Petasis 3-component (*couple*),⁶ intramolecular Diels–Alder (IMDA),⁷ and Ru-alkylidene catalyzed ring-closing metathesis (RCM) reactions (*pair*) (Figure 1).⁸ We have previously reported the applicability of hydrazides in the Petasis 3-component reaction (3-CR), and the efficient and selective elaboration of hydrazido alcohols (I, Figure 1) into oxazolidinones and oxadiazolones by controlled carbonylation processes (VI and VIII, Figure 1).⁹ We have also successfully combined Petasis 3-CRs with Ru-catalyzed cyclizations of simple amines to construct diverse 5- and 7-membered ring-systems.^{5f}

In a continuous search for screening-relevant molecular entities, we desired to investigate if an elaboration of these strategies could further expand the array of accessible heterocycles from the Petasis 3-CR template, specifically via strategic positioning of functional groups, such as vinyl and furyl moieties, which would participate in RCM and IMDA reactions (II–V and VII, Figure 1).

Received: June 11, 2014

Revised: October 28, 2014

Published: December 3, 2014




Figure 1. Build/couple/pair strategy utilizing the Petasis 3-component (couple) and cyclization (pair) reactions for the synthesis of densely functionalized compounds.

In accordance with our previous work,⁹ the Petasis 3-CR of hydrazides (1), α -hydroxy aldehydes (2), and both vinyl and aryl boronic acids (3) proceeded smoothly in MeOH to provide an array of densely functionalized diastereomerically pure *anti*-hydrzido alcohols in good to excellent yields (4{1–25}, Scheme 1). When using electron-rich 2-furyl boronic acid yields were typically higher compared to the less electron-rich vinyl and phenyl boronic acids, and the reactions were generally promoted using 1,1,1,3,3,3-hexafluoro-propan-2-ol (HFIP) as a cosolvent.¹⁰


Interestingly, when *N'*-allylated hydrazides (5) were subjected to the Petasis 3-CR with 2-furyl boronic acid in refluxing MeOH, the expected Petasis 3-CR products (6) rapidly formed before a clean conversion (*albeit* more slowly) provided the diastereomerically pure IMDA adducts 7{1} and 7{2} (Scheme 2).

Having confirmed the viability of furyl-functionalized Petasis 3-CR products as IMDA precursors, we next decided to investigate if the *N'*-linked double bond could be introduced at a later synthetic stage, thereby providing a more divergent approach to the IMDA products. This would also allow for the use of unsubstituted hydrazides in the Petasis 3-CR. Gratifyingly, Petasis 3-CR product 4{12} could be easily allylated with allyl bromide followed by refluxing in toluene to give IMDA product 7{1} in acceptable yield (41% over two steps, Scheme 3). Acrylic variants of the scaffold were obtained analogously by subjecting 4{12} and 4{4} to *O*-TBS-protection, *N*-acryloylation with acryloyl chloride and heating (refluxing toluene), providing IMDA adducts 11{1} and 11{2} in good yields over 3 steps (51% and 47%, respectively).

The scope of the acryloylation/IMDA sequence was investigated for a series of furyl-containing Petasis 3-CR products 4{1–13} and 4{15}. In general, the desired products (12{1–6} and 12{10–13}) were obtained in decent to good yields (47–76%). The slightly lower yields observed for the formation of 12{7–9} (25–44%) presumably reflect the low solubility of their corresponding starting materials during the acryloylation step.

Interestingly, when switching to methacryloyl chloride as the acryloylating agent (Scheme 5), the reaction of compound 9{2} provided a mixture of anti diastereomers 13{1} and 13{2} after IMDA reaction (58% and 14% over 3 steps, respectively).


Scheme 1. Petasis 3-CR: Synthesis of Densely Functionalized *Anti*-Hydrzido Alcohols (Couple)^a




^aReagents and conditions: (a) TBSCl, imidazole, DMF, rt; (b) acryloyl chloride, Et₃N, THF, –78 °C. ^b10% HFIP added as a cosolvent.

Having established efficient protocols for IMDA reactions of vinyl and furyl moieties, pairwise reactive diene combinations were subjected to Ru-catalyzed RCM. For example, when subjecting the Petasis 3-CR products 4{21–23} and 4{25} to RCM conditions, three different ring systems (compounds 14{1}, 14{3–4}, and 14{5}) were obtained in good yields (58–80%, Scheme 6). Unfortunately, despite the use of a range of different RCM conditions, Petasis 3-CR product 4{20} could never be converted to the desired 7-membered ring 14{6}. However, a related ring system could be obtained by

Scheme 2. Tandem Petasis 3-CR and Intramolecular IMDA Reaction (Couple–Pair)



Scheme 3. Functional Group Pairing via IMDA Reaction (Pair)^a



^aReagents and conditions: (a) allyl bromide, Et₃N, DMF, 60 °C; (b) toluene, reflux; (c) TBSCl, imidazole, DMF, rt; (d) acryloyl chloride, Et₃N, CH₂Cl₂, -78 °C.


conversion of acryloylated compound **4**{19} to compound **14**{2} in good yield (70%), suggesting that the basic *N* of **4**{20} inactivates the metathesis catalyst. Attempts made to synthesize larger ring systems by RCM of the Petasis 3-CR products **4**{17} and **4**{24} were all unfruitful.

Finally, we envisioned that the exocyclic alcohol-moiety in the IMDA products (originating from the α -hydroxy aldehydes) could serve as an entry for further skeletalization reactions. Accordingly, compounds **12**{12} and **12**{6} were mesylated and subjected to NaH in a set of commonly used solvents at room temperature. However, under these conditions the mesylated compounds **15**{1–2} appeared completely unreactive. This issue was resolved by elevating the temperature, which resulted in the formation of different products depending on the nature of the solvent. For instance, complex reaction mixtures were observed when using THF and CH₂Cl₂, whereas reactions in toluene or DMF at 80 °C resulted in clean conversions to either eliminated product **16** (toluene) or cyclized product **17**{1} (DMF). In the same manner, subjection of mesylate **15**{2} to NaH in toluene provided compound **17**{2} as a single enantiomer in good yield (76%).


The stereochemistry of compound **17**{2} was assigned via X-ray crystallography (Figure 2), which provided unambiguous confirmation of the exo-anti selectivity of the IMDA reaction and proved that the NaH-mediated cyclization proceeds via S_N2-inversion.

In conclusion, we have shown the feasibility of forming a collection of structurally diverse heterocycles through a build/couple/pair pathway that combines the Petasis 3-CR reaction

Scheme 4. Sequential Acryloylation and IMDA Reactions (Pair)




Scheme 5. Intramolecular Diels–Alder Reactions (Pair)



of diverse hydrazides with regio- and stereoselective cyclization reactions, including ring-closing metathesis and intramolecular Diels–Alder reactions. Using this strategy, 10 structurally distinct scaffolds were synthesized in good yields in only 3–4 steps. Functionalized hydrazido alcohols accessed through the Petasis 3-CR are synthetically versatile templates for subsequent cyclization reactions at several positions, thus constituting valuable compounds of modular and synthetically programmable pathways. The presented key cyclization steps of the pathway take advantage of pairwise reactive functional groups of a densely functionalized hydrazido alcohol template,

Scheme 6. Ring-Closing Metathesis of Diene-Containing Hydrazido Alcohols (Pair)^a



^aReagents and conditions: (a) Grubbs II (10 mol %), toluene, 65 °C, 2 h; (b) Grubbs II (10 mol %), CH₂Cl₂, rt, overnight.

Scheme 7. Solvent-Dependent Cyclization and Elimination of Mesylates




Figure 2. Crystal Structure of Compound **17{2}**.

leaving any remaining handle amenable to additional diversification, including secondary cyclization processes, and appendage decoration.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and compound characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: ten@kemi.dtu.dk.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the Technical University of Denmark, DSF Center for Antimicrobial Research, Danish Council for Strategic Research, Danish Council for Independent Research, and Lundbeck Foundation.

■ REFERENCES

- (1) (a) Burke, M. D.; Schreiber, S. L. A planning strategy for diversity-oriented synthesis. *Angew. Chem., Int. Ed.* **2004**, *43*, 46–58. (b) Tan, D. S. Diversity-oriented synthesis: Exploring the intersections between chemistry and biology. *Nat. Chem. Biol.* **2005**, *1*, 74–84. (c) Spiegel, D. A.; Schroeder, F. C.; Duvall, J. R.; Schreiber, S. L. An oligomer-based approach to skeletal diversity in small-molecule synthesis. *J. Am. Chem. Soc.* **2006**, *128*, 14766–14767. (d) Comer, E.; Rohan, E.; Deng, L.; Porco, J. A., Jr. An approach to skeletal diversity using functional group pairing of multifunctional scaffolds. *Org. Lett.* **2007**, *9*, 2123–2126. (e) Nielsen, T. E.; Schreiber, S. L. Towards the optimal screening collection: A synthesis strategy. *Angew. Chem., Int. Ed.* **2008**, *47*, 48–56. (f) Galloway, W. R. J. D.; Isidro-Llobet, A.; Spring, D. R. Diversity-oriented synthesis as a tool for the discovery of novel biologically active small molecules. *Nat. Commun.* **2010**, *1*:80, 1–13. (g) Dandapani, S.; Marcaurelle, L. A. Current strategies for diversity-oriented synthesis. *Curr. Opin. Chem. Biol.* **2010**, *14*, 362–370. (h) Wetzel, S.; Bon, R. S.; Kumar, K.; Waldmann, H. Biology-oriented synthesis. *Angew. Chem., Int. Ed.* **2011**, *50*, 10800–10826. (i) *Diversity-Oriented Synthesis: Basics and Applications in Organic Synthesis, Drug Discovery, and Chemical Biology*; Trabocchi, A., Ed.; Wiley: Hoboken, NJ, 2013. (j) O'Connor, C. J.; Beckmann, H. S. G.; Spring, D. R. Diversity-oriented synthesis: Producing chemical tools for dissecting biology. *Chem. Soc. Rev.* **2012**, *41*, 4444–4456.
- (2) (a) Lovering, F.; Bikker, J.; Humblet, C. Escape from flatland: Increasing saturation as an approach to improving clinical success. *J. Med. Chem.* **2009**, *52*, 6752–6756. (b) Lovering, F. Escape from flatland 2: Complexity and promiscuity. *Med. Chem. Commun.* **2013**, *4*, 515–519.
- (3) (a) Cordier, C.; Morton, D.; Murrison, S.; Nelson, A.; O'Leary-Steele, C. Natural products as an inspiration in the diversity-oriented synthesis of bioactive compound libraries. *Nat. Prod. Rep.* **2008**, *25*, 719–737. (b) Morton, D.; Leach, S.; Cordier, C.; Warriner, S.; Nelson, A. Synthesis of natural-product like molecules with over eighty distinct scaffolds. *Angew. Chem., Int. Ed.* **2009**, *48*, 104–109. (c) Marcaurelle, L. A.; Comer, E.; Dandapani, S.; Duvall, J. R.; Gerard, B.; Kesavan, S.; Lee, M. D.; Liu, H.; Lowe, J. T.; Marie, J. C.; Mulrooney, C. A.; Pandya, B. A.; Rowley, A.; Ryba, T. D.; Suh, B. C.; Wei, J.; Young, D. W.; Akella, L. B.; Ross, N. T.; Zhang, Y. L.; Fass, D. M.; Reis, S. A.; Zhao, W. N.; Haggarty, S. J.; Palmer, M.; Foley, M. A. An aldol-based build/couple/pair strategy for the synthesis of medium- and large-sized rings: Discovery of macrocyclic histone deacetylase inhibitors. *J. Am. Chem. Soc.* **2010**, *132*, 16962–16976. (d) Moura-Lettsa, G.; DiBlasia, C. M.; Bauer, R. A.; Tan, D. S. Solid-phase synthesis and chemical space analysis of a 190-membered alkaloid/terpenoid-like library. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 6745–6750. (e) Basu, S.; Ellinger, B.; Rizzo, S.; Deraeve, C.; Schürmann, M.; Preut, H.; Arndt, H. D.; Waldmann, H. Biology-oriented synthesis of a natural-product inspired oxepane collection yields a small-molecule activator of the wnt-pathway. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 6805–6810.
- (4) Clemons, P. A.; Wilson, J. A.; Dančik, V.; Muller, S.; Carrinski, H. A.; Wagner, B. K.; Koehler, A. N.; Schreiber, S. L. Quantifying structure and performance diversity for sets of small molecules

comprising small-molecule screening collections. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 6817–6822.

(5) (a) Kumagai, N.; Muncipinto, G.; Schreiber, S. L. Short synthesis of skeletally and stereochemically diverse small molecules by coupling Petasis condensation reactions to cyclization reactions. *Angew. Chem., Int. Ed.* **2006**, *45*, 3635–3638. (b) Zhou, A.; Rayabarapu, D.; Hanson, P. R. “Click, click, cyclize”: A DOS approach to sultams utilizing vinyl sulfonamide linchpins. *Org. Lett.* **2009**, *11*, 531–534. (c) Uchida, T.; Rodriguez, M.; Schreiber, S. L. Skeletally diverse small molecules using a build/couple/pair strategy. *Org. Lett.* **2009**, *11*, 1559–1562. (d) Hung, A. W.; Ramek, A.; Wang, Y.; Kaya, T.; Wilson, J. A.; Clemons, P. A.; Young, D. W. Route to three-dimensional fragments using diversity-oriented synthesis. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 6799–6804. (e) Fitzgerald, M. E.; Mulrooney, C. A.; Duvall, J. R.; Wei, J.; Suh, B. C.; Akella, L. B.; Vrcic, A.; Marcaurelle, L. A. Build/couple/pair strategy for the synthesis of stereochemically diverse macrolactams via head-to-tail cyclization. *ACS Comb. Sci.* **2012**, *14*, 89–96. (f) Ascic, E.; Le Quement, S. T.; Ishoey, M.; Daugaard, M.; Nielsen, T. E. Build/couple/pair strategy combining the Petasis 3-component reaction with Ru-catalyzed ring-closing metathesis and isomerization. *ACS Comb. Sci.* **2012**, *14*, 253–257.

(6) For early accounts, see: (a) Petasis, N. A.; Akritopoulou, I. The boronic-acid Mannich reaction: A new method for the synthesis of geometrically pure allylamines. *Tetrahedron Lett.* **1993**, *34*, 583–586. (b) Dyker, G. Amino acid derivatives by multicomponent reactions. *Angew. Chem., Int. Ed.* **1997**, *36*, 1700–1702. For reviews on the Petasis 3-CR, see: (c) Petasis, N. A. In *Multicomponent Reactions*; Zhu, J., Bienaymé, H., Eds.; Wiley-VCH: Weinheim, Germany, 2005; pp 199–223. (d) Batey, R. A. In *Boronic Acids*; Hall, D. G., Ed.; Wiley-VCH: Weinheim, Germany, 2005; pp 279–304. (e) Candeias, N. R.; Montalbano, F.; Cal, P. M. S. D.; Gois, P. M. P. Boronic acid and esters in the Petasis-borono Mannich multicomponent reaction. *Chem. Rev.* **2010**, *110*, 6169–6193.

(7) For review on IMDA, see: Takao, K.; Munakata, R.; Tadano, K. Recent advances in natural product synthesis by using intramolecular Diels–Alder reactions. *Chem. Rev.* **2005**, *105*, 4779–4807.

(8) For a recent review on metathesis reactions, see: Vougioukalakis, G. C.; Grubbs, R. H. Ruthenium-based heterocyclic carbene-coordinated olefin metathesis catalysts. *Chem. Rev.* **2010**, *110*, 1746–1787.

(9) Le Quement, S. T.; Flagstad, T.; Mikkelsen, R. J. T.; Hansen, M. R.; Givskov, M. C.; Nielsen, T. E. Petasis three-component coupling reactions of hydrazides for the synthesis of oxadiazolones and oxazolidinones. *Org. Lett.* **2012**, *14*, 640–643.

(10) Nanda, K. K.; Trotter, B. W. Diastereoselective Petasis Mannich reactions accelerated by hexafluoro-isopropanol: A pyrrolidine-derived arylglycine synthesis. *Tetrahedron Lett.* **2005**, *46*, 2025–2028.