


Pathophysiology of Ebola Virus Infection: Current Challenges and Future Hopes

Andrea Rivera and Ilhem Messaoudi*

Division of Biomedical Sciences, University of California, Riverside, Riverside, California 92521, United States

ABSTRACT: The filoviruses, Ebola virus (EBOV) and Marburg virus (MARV), are among the deadliest viruses that cause disease in humans, with reported case fatality rates of up to 90% in some outbreaks. The high virulence of EBOV and MARV is largely attributed to the ability of these viruses to interfere with the host immune response. Currently, there are no approved vaccines or postexposure therapeutics, and treatment options for patients infected with EBOV are limited to supportive care. In this review, we discuss mechanisms of EBOV pathogenesis and its ability to subvert host immunity as well as several vaccines and therapeutics with respect to their evaluation in small animal models, nonhuman primates, and human clinical trials.

KEYWORDS: Ebola virus, viral hemorrhagic fever, innate immunity, adaptive immunity, immunoevasion, antivirals, Ebola vaccines, Ebola therapeutics



Ebola virus (EBOV) is a filamentous enveloped virus containing a negative strand RNA genome 19 kb in length that encodes for a nucleoprotein (NP), glycoprotein (GP), RNA-dependent RNA polymerase (L), and four structural proteins termed VP24, VP30, VP35, and VP40. Viral replication is carried out by NP, VP35, and L; the active polymerase complex is composed of VP35 (a polymerase cofactor) and L (polymerase) while NP drives RNA encapsidation. VP30 is a transcriptional activator and is also involved in nucleocapsid formation and assembly.¹ VP24 is a matrix protein that contributes to nucleocapsid formation whereas matrix protein VP40 facilitates the budding of progeny virion from infected cells.^{1b,2} Glycoprotein (GP) covers the surface of the virion and is the sole host attachment factor for EBOV.³ Moreover, EBOV expresses two soluble forms of GP, sGP, and ssGP through RNA editing.⁴

Infection with EBOV results in severe viral hemorrhagic fever with fatality rates that can reach 90% in humans depending on the species. Infection with Zaire Ebola virus (EBOV), Sudan Ebola virus (SUDV), and Bundibugyo Ebola virus (BDBV) is associated with 70–90, 50, and 40% mortality rates, respectively.⁵ In contrast, infection with either Tai Forest Ebola virus (TAFV) or Reston Ebola virus (RESTV) has not been associated with human fatalities.⁵ EBOV is transmitted by direct contact with infected bodily fluid. The incubation period ranges from 2 to 21 days with an average incubation period of 5–7 days. Early symptoms are general and nonspecific including fever, general malaise, and myalgia. Following early symptoms, cases can show maculo-papular rash, petichiae, conjunctival hemorrhage, melena, hematemesis, shock, and encephalopathy. During the terminal stage of disease, there is an increase in vascular permeability, massive tissue injury,

dysregulation of the coagulation cascade, and hemorrhage. Multiorgan failure and shock are usually the main causes of death.⁶

Previous EBOV outbreaks have occurred mostly in remote regions of central Africa including the Democratic Republic of Congo, Sudan, Gabon, and Uganda. However, in March 2014, the World Health Organization (WHO) reported an outbreak of Ebola virus disease originating in the Guinean capital, Conakry, marking the first large urban setting for EBOV transmission. This has led to the first known epidemic in West Africa and is the largest and longest outbreak since the virus was first discovered in 1976.⁷ Genome sequencing of the Guinea strain shows 97% identity to EBOV.⁸ Sequencing studies of 99 EBOV genomes from 78 patients in Sierra Leone reveal an accumulation of inter- and intrahost genetic variation. Patterns of viral transmission suggest that this variant diverged from central African lineages around 2004.⁹ As of March 8, 2015, the CDC reports a total 24 247 confirmed, probable, and suspected cases of EBOV disease and 9961 reported deaths in 8 countries (Guinea, Liberia, Sierra Leone, Mali, Nigeria, Senegal, Spain, and United States). Currently, there are no FDA-approved treatment strategies. Supportive care, consisting of oral fluid rehydration and nutritional supplementation, is the mainstay in treating Ebola hemorrhagic fever (EHF). The high mortality rate has caused concern to public health officials worldwide, prompting a sense of urgency to accelerate clinical trials to identify effective vaccines and postexposure therapeutics.

Received: December 15, 2014

Published: March 17, 2015




Figure 1. Dysregulation of the immune system by the Ebola virus. Monocytes, macrophages, and dendritic cells are the preferred sites of filovirus replication. Infection of monocytes and macrophages triggers the robust expression of inflammatory mediators. Inflammatory mediators, reactive oxygen species, and nitric oxide can induce apoptosis leading to lymphocyte death. The infection of dendritic cells impairs their maturation and suppresses type I (IFN) responses, thereby preventing T cell activation. Production of EBOV-soluble glycoprotein (sGP) usurps GP-specific antibodies.

Nonhuman primates, particularly cynomolgus macaques and rhesus macaques, are the gold standard animal models used in filovirus study because they are susceptible to infection by the same strains that cause disease in humans and exhibit a strong similarity of viral hemorrhagic fever to humans.¹⁰ Small animal models, such as inbred mice and guinea pigs, have also been used; however, these animal models require the use of adapted EBOV strains and do not exhibit the characteristic hemorrhagic manifestations seen with human and nonhuman primate infection.¹¹ In a recent study using Collaborative Cross mice, animals exhibited a spectrum of disease outcomes following infection with mouse-adapted EBOV (MA-EBOV), including hemorrhagic fever.¹² However, MA-EBOV has diminished virulence in nonhuman primates. Of three rhesus macaques injected with a large challenge dose of 5000 pfu of MA-EBOV, two showed mild illness and survived infection.¹³ A single amino acid mutation in both NP and VP24 are the determinants of the virulence of MA-EBOV in mice.¹⁴ Although MA-EBOV infection in mice correlated with the evasion of IFN-induced responses,¹⁴ the determinants for the virulence of EBOV may differ between mouse and primate models. The purpose of this review is to summarize our current understanding of the pathogenesis of EBOV and discuss the

various vaccine platforms and therapeutics currently being investigated.

■ DYSREGULATION OF INNATE IMMUNITY BY EBOV INFECTION

Although EBOV targets a wide range of cell lineages, *in vivo* studies in nonhuman primates using immunohistochemistry and *in situ* hybridization suggest that EBOV replicates preferentially in monocytes, macrophages, and dendritic cells.^{10b,15} The infection of monocytes and macrophages triggers the robust expression of inflammatory mediators¹⁶ (Figure 1). Indeed, several studies have shown that fatal infections were associated with the uncontrolled secretion of proinflammatory cytokines, chemokines, and growth factors such as IL-1 β , IL-6, IL-8, IL-10, MCP-1, MIP-1 α , MIP-1 β , and TNF α as well as nitric oxide and reactive oxygen species, which, at the time of death, can reach 5 to 1000 times the levels detected in survivors and healthy individuals.¹⁷ In contrast, survivors show a transient and moderate upregulation in levels of IL-1 β , IL-6, TNF α , MIP1 α , and MIP1 β early in the disease.^{17a} These studies suggest that protection from fatal EBOV infection may depend on an early yet regulated inflammatory response.

Ebola virus infection is also characterized by a suppressed type I interferon (IFN) response, which is an integral part of the innate immune response against viral infections. This suppression is mediated by VP24, which prevents the nuclear transport of tyrosine phosphorylated STAT1 (pSTAT1), a key downstream outcome of IFN signaling, thereby preventing IFN-induced gene expression.¹⁸ VP24 binds the C-terminus of karyopherin α proteins (karyopherin $\alpha 1$, $\alpha 5$, $\alpha 6$), which normally transport pSTAT1 through the nuclear pore, therefore competing with pSTAT1 to bind the nuclear transporter.¹⁹ In addition to VP24, VP35 was also found to block the activation of transcription factor IFN regulatory factor 3 (IRF-3), thus decreasing IFN production.²⁰ VP35 interacts with SUMO E2 enzyme Ubc9 and E3 ligase PIAS1, which in turn leads to increased SUMOylation and degradation of transcription factor IRF-7.²¹ VP35 can also inhibit retinoic acid-inducible gene-I (RIG-I) helicase signaling by binding to dsRNA²² or binding to PKR activating protein (PACT), which activates RIG-I.²³ Moreover, VP35 can decrease IFN production by impairing IKK ϵ and TBK-1 kinase function.²⁴ Interference with RIG-I signaling can also inhibit the upregulation of a number of costimulatory molecules on dendritic cells (CD40, CD80, CD86, and MHC class II) needed to activate T cells.²⁵ Additionally, *in vitro* studies have shown that EBOV infection inhibits dendritic cells maturation into mobile, antigen-presenting cells and impairs their ability to stimulate antigen-specific T cell responses²⁶ (Figure 1).

■ DYSREGULATION OF LYMPHOCYTE FUNCTION BY EBOV INFECTION

One major consequence of EBOV infection is severe lymphopenia.^{10a} EBOV infection leads to the loss of both human CD4⁺ and CD8⁺ T cells after 4 days of *in vitro* culture.²⁷ Similarly, EBOV infection resulted in a loss of peripheral CD4⁺ and CD8⁺ T cells in both mouse models at 2 to 3 dpi²⁸ and in nonhuman primates at 4 dpi.²⁹ A significant loss of circulating natural killer (NK) cells was also described in mice²⁸ and in cynomolgus macaques.^{10b,29} B cell loss is more controversial, with some studies reporting the loss of B lymphocytes in mice²⁸ and macaques¹⁵ and other studies reporting no changes in B cell counts in nonhuman primates.^{29,30} Flow cytometric analysis of peripheral blood mononuclear cells (PBMCs) from humans and cynomolgus macaques infected with EBOV revealed an increase in the percentage of CD4⁺ and CD8⁺ T cells expressing cell death receptor Fas (CD95), suggesting that apoptosis is the primary mechanism of lymphocyte death.^{17f,29} Transmission electron microscopy and TUNEL staining confirmed lymphopenia *in vivo* and *in vitro* occurred by apoptosis.¹⁵ Increased plasma levels of apoptosis mediators such as soluble Fas (sFas) and 41/7 nuclear matrix protein, upregulation of Fas and FasL mRNA in PBMCs, and dramatic DNA fragmentation in leukocytes were detected during the terminal stage of human EBOV infection.³¹ Furthermore, PBMCs from survivors showed an upregulation of antiapoptotic Bcl-2 mRNA whereas fatalities showed a significant decrease in Bcl-2 mRNA as well as a decrease in CD3, CD8, and TCR-V β mRNA.^{15,31b}

The fact that EBOV does not replicate within lymphocytes during infection suggests that lymphocyte apoptosis is an indirect result of viral replication.^{15,31a} Some of the proinflammatory mediators released by EBOV-infected monocytes such as TNF α , reactive oxygen species, and nitric oxide have been shown to induce apoptosis³² (Figure 1).

Furthermore, immunohistochemistry, flow cytometry, and RNA analysis showed a high expression of TNF-related apoptosis inducing ligand (TRAIL) in 90% of EBOV-infected adherent human monocytes/macrophages.^{17c} In addition to lymphopenia, EBOV infection interferes with T cell activation as evident by decreased expression of activation marker CD44.^{26,29} The absence of EBOV-specific T cell responses is consistent with the lack of T cell cytokines (IL-2, IL-4, IL-5, IL-12, and IFN γ) in the plasma of EBOV-infected patients.^{17f,33}

EBOV also subverts the host humoral immune response, as fatalities are associated with a lack of EBOV-specific IgG antibodies.^{31b} The loss of CD4⁺ T cells, which are required for isotype class switching, may explain the lack of Ebola-specific IgM and IgG antibodies observed in patients who succumb to infection. In contrast, asymptomatic EBOV-infected patients developed IgM responses between 10 and 18 days and IgG responses between 17 and 25 days specific to GP, NP, and VP40.³⁴ EBOV also subverts humoral responses through the production of sGP,³⁰ which sequesters GP-specific antibodies needed to control viral replication³⁵ (Figure 1). The importance of antibodies in protection against EBOV infection is highlighted by passive immunization studies. During one of the EBOV outbreaks in the Democratic Republic of Congo, 7/8 patients treated with blood transfusions from five convalescent patients who generated EBOV-specific IgG antibodies survived.³⁶ Similarly, 100% of naïve mice were protected against infection when treated with serum from vaccinated mice that survived challenge with mouse-adapted EBOV.³⁷

■ VACCINES AGAINST EBOLA

Several vaccine platforms against EBOV have been developed. Replication-deficient vaccines include DNA-based vaccines, virus-like particles (VLPs), and recombinant adenovirus vectors (rAd). Replication competent constructs include recombinant human parainfluenza virus 3 (rHPIV3), recombinant vesicular stomatitis virus (rVSV), and more recently recombinant rabies virus (RABV) and recombinant cytomegalovirus (CMV).³⁸ Two promising vaccine candidates, rAd and rVSV vectors expressing GP (Table 1), have entered clinical trials.

Table 1. Vaccines against Ebola Virus

vaccine	rAdS/ChAd3	rVSV/ Δ G/GP EBOV
replication potential	replication deficient	replication competent
mechanism of protection	CD8 ⁺ T cell mediated	antibody mediated
booster requirement	modified vaccinia Ankara	
durability	10 months with booster vaccination in nonhuman primates	12–18 months in rodents 14 months in nonhuman primates (MARV)
clinical trial	phase II/III	phase II/III

The vaccination of cynomolgus macaques with the replication-deficient rAd serotype 5 vector expressing EBOV-GP (rAd5-EBOVGP) resulted in the generation of EBOV-GP-specific antibodies and GP-specific CD8⁺ T cell responses within 3 weeks of immunization and protected 100% of the animals against challenge.³⁹ Passive transfer of EBOV-specific IgG purified from nonhuman primates vaccinated with DNA

and rAd5 vectors expressing EBOV GP to naïve animals protected only 25% of naïve animals 6 or 16 h before EBOV challenge, whereas the depletion of CD8⁺ T cells in vaccinated animals 4 days before challenge resulted in 4/5 nonhuman primates succumbing to infection, suggesting that CD8⁺ T cells play a more important role in protection compared to antibodies in this vaccine platform.⁴⁰ Despite these promising results, humans have preexisting immunity against Ad5, which may interfere with the efficacy of the rAd5-EBOVGP vaccine.³⁸ On the other hand, adenoviruses isolated from chimpanzees (ChAds) provide better vaccine vectors because of their low seroprevalence in humans. The immunization of cynomolgus macaques with a single inoculation of recombinant ChAd3 expressing EBOV GP (rChAd3-EBOVGP) resulted in complete protection when challenged with EBOV 5 weeks after vaccination but conferred only 50% protection when the animals were challenged 10 months after immunization. To improve long-term efficacy of this vaccine platform, a booster vaccination using recombinant-modified vaccinia Ankara expressing EBOV-GP was added 8 weeks after the rChAd3-EBOVGP vaccination, which resulted in complete protection at 10 months.⁴¹ Phase I clinical trials of rChAd3 began in September 2014 by the National Institute of Allergy and Infectious Diseases (NIAID) to evaluate the vaccine's safety and immunogenicity. Twenty adults were vaccinated with either 2×10^{10} or 2×10^{11} particles. Reactivity to rChAd3 was dose-dependent as two participants developed fever 1 day after vaccination with the 2×10^{11} dose. Similarly, immunogenicity was also dose-dependent, with higher GP-specific antibodies and T-cell responses detected in the group that received the 2×10^{11} dose 4 weeks after vaccination.⁴² In parallel, the NIH partnered with the United Kingdom-based international consortium to test the safety and efficacy of rChAd3, at a dose of either 2×10^{10} or 2×10^{11} , among 60 volunteers at the University of Oxford in England and 40 volunteers in Mali and has observed similar results with regard to safety and immune response.⁴³ The vaccine will enter a phase II/III trial called Partnership for Research on Ebola Vaccines in Liberia (PREVAIL). This study will enroll 27 000 healthy men and women to investigate rChAd3 and rVSV/ΔG/GP and is estimated to be completed in June 2016.⁴³

Vesicular stomatitis virus (VSV) is a nonsegmented, negative-stranded RNA virus and a member of the Rhabdoviridae family. Recombinant VSV, in which the VSV G protein is replaced with EBOV GP (rVSV/ΔG/GP EBOV), confers 100% protection in nonhuman primates and mice challenged 28 days after immunization with EBOV and mouse-adapted EBOV, respectively.⁴⁵ Antibodies play a critical role in rVSV/ΔG/GP-mediated protection against EBOV. In a nonhuman primate study, only the animals depleted of CD4⁺ T cells during vaccination, which lacked GP-specific antibodies, succumbed to infection, suggesting that antibodies were required for protection.⁴⁶ Because rVSV/ΔG/GP is a live-attenuated virus, several studies have investigated its safety. No toxicity was observed in over 80 nonhuman primates given rVSV/ΔG/GP MARV or rVSV/ΔG/GP EBOV vaccines.^{45a,47} Moreover, this vaccine was safe in a mouse model of severe combined immunodeficiency (NOD-SCID) and in simian human immunodeficiency (SHIV)-infected rhesus macaques.^{45b,47b} Administration of a single dose of rVSV/ΔG/GP confers 100% protection against EBOV challenge for up to 6 months in macaques and 18 months in mice and guinea pigs, and this protection correlates with GP-specific IgG titers.^{48,49}

This vaccine platform has also provided protection in macaques against MARV for up to 14 months after vaccination, further demonstrating the durability of VSV-based vaccines against filoviruses.⁵⁰ Administration of a mixture vaccine containing rVSV/ΔG/GP from MARV, EBOV, and SUDV to cynomolgus macaques resulted in their survival following challenge with MARV, EBOV, SUDV, and TAFV, demonstrating that a single injection of a multivalent vaccine is as efficacious as the administration of a single specificity vaccine.⁵¹ Finally, rVSV/ΔG/GP has also been shown to be effective postexposure. Administration of rVSV/ΔG expressing EBOV or SUDV GP to rhesus macaques 20–30 min after challenge resulted in 50 and 100% protection, respectively.⁵² In March 2009 in Germany, a virologist working in a biosafety level 4 sustained a needle stick injury with a syringe that contained EBOV mixed with Freund's adjuvant. A single dose of 5×10^7 pfu rVSV/ΔG/GP EBOV was administered to her 48 h after the injury. Following postexposure vaccination, the virologist developed a fever 12 h later in addition to rVSV viremia detected via PCR for 2 days but remained healthy during the 3 week observation period.⁵³ During the 2014 outbreak, rVSV/ΔG/GP EBOV was used as an emergency postexposure vaccination and was administered 43 h after a physician experienced a needle stick injury while working in an Ebola treatment unit in Sierra Leone.⁵⁴ Although it is unknown if rVSV/ΔG/GP EBOV was effective as a postexposure vaccination, the patient had a self-limited febrile syndrome and cytokine- and Ebola glycoprotein-specific adaptive immune response after vaccination.⁵⁴ The NIH and WHO have carried out phase I clinical trials to test the safety and efficacy of rVSV/ΔG/GP EBOV, with a dose ranging from 3×10^6 to 1×10^8 pfu, in various locations in the U.S., Europe, and Africa. Although safety data from phase I studies have not yet been published, there is sufficient safety and efficacy information to push the vaccine forward into the PREVAIL phase II/III trial.⁴³ The CDC in collaboration with Sierra Leone has launched another phase II/III trial called Sierra Leone Trial to Introduce a Vaccine against Ebola (STRIVE). The study will test a dose of 2×10^7 pfu of rVSV/ΔG/GP on 6000 participants.⁵⁵

■ THERAPEUTICS AGAINST EBOLA

Various therapeutic candidates that either directly inhibit viral replication, modulate clinical symptoms, or prolong survival time have been evaluated either as sole or adjunctive postexposure therapies in rodent and nonhuman primates. A complete list of Ebola therapeutics is summarized in Table 2. Two anticoagulant recombinant proteins, recombinant nematode anticoagulant protein c2 (rNAPc2) and recombinant human activated protein C (rhAPC), have been evaluated in preventing increased coagulation during filovirus infection.⁵⁹ However, both anticoagulants provided only partial protection in nonhuman primates challenged with EBOV.⁵⁹ Administration of rNAPc2 to six rhesus macaques 10 min after EBOV challenge prolonged the survival time from a mean time to death of 8.3 days to 11.7 days with a 33% survival rate.^{59a} Treatment with rhAPC in 11 rhesus macaques starting 30–60 min after EBOV challenge and daily for 7 days increased the mean time to death from 8.3 to 12.6 days with 2 of 11 rhesus macaques surviving.^{59b} Although rNAPc2 has gone through phase II trials for use in the prevention of venous thromboembolism after orthopedic surgery⁶⁰ and coronary revascularization,⁶¹ it has not been assessed in clinical trials for the treatment of EBOV exposure. Additionally, rhAPC has not

Table 2. Therapeutics for EBOV Infection

molecule	mechanism of action	development phase	reference
Favipiravir	a nucleotide analog that inhibits RNA polymerase	has been shown to protect mice against EBOV; administered to one EBOV-infected patient who survived currently in phase II trial	63
PMOs AVI-6002 (AVI-7537 and AVI-7539)	AVI 7537 targets EBOV VP24; AVI-7539 targets EBOV VP35	AVI-6002 has been shown to provide partial protection of rhesus macaques against EBOV; AVI-7537 confers sole protection observed; AVI-6003 has been shown to completely protect cynomolgus macaques from MARV infection	71
AVI-6003 (AVI-7287 and AVI-7288)	AVI-7287 targets MARV VP24; AVI-7288 targets MARV NP	completed phase I clinical trial	
ZMapp	monoclonal cocktail of antibodies against different epitopes of EBOV-GP	has been shown to completely protect cynomolgus macaques; protected 5/7 patients infected with EBOV; currently in phase I clinical trial	83, 85
BXC4430	an adenosine analog that inhibits RNA transcription	shown to protect cynomolgus macaques against Marburg virus; in phase I clinical trial	66
Brincidofovir	inhibits DNA synthesis; mechanism against RNA viruses is unknown	shows efficacy against EBOV in vitro; protected one of two U.S. patients infected with EBOV; withdrawn from phase II clinical trial	68, 69
TKM-Ebola	treatment with SNALP carrying siRNAs against EBOV L polymerase, VP24, and VP35	protected macaques following EBOV challenge; phase I trial halted; FDA expanded access to TKM-Ebola on an emergency basis	74,75
rhAPC	anticoagulant	recombinant human activated protein C is an anticoagulant that prolongs survival time and provides partial protection in nonhuman primates.	59b
rNAPc2	anticoagulant	recombinant nematode anticoagulant protein c2 prolongs survival time and provides partial protection in nonhuman primates	59a
IFN β	interferon	prolongs mean time to death in rhesus macaques infected with EBOV; may be possible adjunctive postexposure therapy	62
chloroquine	the antiviral mechanism is unknown	FDA approved as an antimalarial agent; through a screening study evaluating 1012 FDA-approved drugs, chloroquine was found to possess in vitro activity against EBOV; found to protect mice against EBOV infection in vivo	87
selective estrogen receptor modulators: clomiphene and toremifene	inhibits EBOV entry	clomiphene and toremifene inhibit EBOV infection in vitro and show partial protection in mice challenged with mouse-adapted EBOV	88
FGI compounds: FGI-103, FGI-104, FGI-106	FGI compounds are broad-spectrum viral inhibitors; antiviral mechanism against EBOV is unknown	exhibited efficacy against EBOV in vitro and in mouse models	89–91
SAH hydrolase inhibitor Dezanepplanocin A	targets SAH hydrolase, indirectly resulting in reduced methylation of 5' cap of viral mRNA; may also increase IFN α production	early dose protected mice against EBOV challenge	92
glucosidase inhibitors: CM-10–18, IHVRI1029, IHVRI7028 and IHVRI9029	glucosidase inhibitors inhibit endoplasmic reticulum α -glucosidases I and II and prevent maturation of viral glycosylated envelope and inhibit viral entry	CM-10–18, IHVRI1029, IHVRI7028, and IHVRI9029 has been shown to suppress the mortality of MARV and EBOV infection in mice	93
antioxidant NSC62914	acts as a scavenger of reactive oxygen species and upregulates oxidative stress-induced genes	exhibits antiviral activity in vitro and in vivo in mice infected with EBOV or Marburg virus	94
HSPAS inhibitors	targets HSPAS chaperone (EBOV-associated host factor)	PMOs targeting HSPAS reduced viral replication and protected mice against lethal EBOV infection	95
SAH hydrolase inhibitor Neplanocin A, 3-	targets SAH hydrolase, indirectly resulting in reduced methylation of 5' cap of viral mRNA	efficacy of Neplanocin A against EBOV has been shown in vitro	96, 97
benzylpiperazine adamantane diamides LJ-001	inhibits EBOV entry via cholesterol transporter Niemann-Pick C1 in vitro a rhodamine derivative which prevents the fusion of viral and cellular membranes and prevents viral entry	shown to inhibit EBOV entry in vitro shown to inhibit EBOV entry in vitro	98, 100
CMLDBU3402	broad-spectrum antiviral inhibits EBOV RNA transcription	shown to inhibit EBOV entry in vitro	101
ion channel blockers: amiodarone, dronedarone, verapamil	inhibit EBOV entry	shown to inhibit EBOV entry in vitro	102
heme oxygenase-1 (HO-1)	inhibit EBOV replication; mechanism unknown	HO-1 induction significantly suppressed EBOV replication in vitro	103
Lamivudine	cytosine analog approved for treatment of HIV and hepatitis B virus; unknown mechanism against EBOV	not yet evaluated in context of EBOV; studies against EBOV in vitro are currently being conducted	104
duY11	a rigid amphipathic fusion inhibitor (RAFI) targets the entry of enveloped viruses and may potentially inhibit the entry of EBOV	not yet assessed in in vitro or in vivo studies against EBOV	105
inhibitors of cathepsin L cleavage	prevention of cathepsin L cleavage of viral glycoproteins may inhibit EBOV entry	not yet assessed in in vitro or in vivo studies	106

Table 2. continued

reference	development phase	mechanism of action	molecule	statins
107	no in vitro or in vivo studies in efficacy against Ebola; however, may be used to treat viral sepsis during EBOV infection; FDA approved	anti-inflammatory		

been evaluated in clinical trials for use against EBOV infection. Treatment with interferon beta (IFN β) for five EBOV-infected rhesus macaques 18 h and 1, 3, 5, 7, and 9 days postinfection did not alter mortality but significantly increased the mean time to death from 8.3 to 13.8 days⁶² and may be used as an adjunctive therapeutic.

Antivirals that have been evaluated as therapeutics against EBOV include nucleotide analogs (Favipiravir and BCX4430), Brincidofovir, and antisense therapeutics (phosphorodiamidate morpholino oligomers (PMOs) and siRNAs) and are summarized in Table 2. Favipiravir (T-705) is an oral nucleotide analog that, when converted into its active metabolite ribofuranosyl triphosphate, inhibits the viral RNA-dependent RNA polymerase by directly competing with GTP. Currently licensed for influenza outbreaks, Favipiravir (T-705) was shown to suppress EBOV replication by 4 log₁₀ units in Vero E6 cells 1 h after infection.⁶³ In a mouse model in which mice lack the type I IFN α/β receptor and are susceptible to wild-type EBOV, treatment with Favipiravir 6 h after EBOV challenge resulted in complete survival of all five mice.⁶³ To date, Favipiravir has been used to treat one French nurse infected with Ebola who recovered.⁶⁴ Favipiravir entered phase II evaluations in December 2014 in Guinea and is sponsored by Institut National de la Sante Et de la Recherche Medicale, France. Nucleoside analogue inhibitor BCX4430 is an adenosine analog with a C-nucleoside instead of the N-glycoside and a 1,4 imino group instead of the 1,4 oxygen. Once metabolized to the active triphosphate nucleotide form and after pyrophosphate cleavage, it is incorporated into the nascent viral RNA chain and terminates transcription. It is not incorporated into mammalian RNA or DNA. Administration of BCX4430 in cynomolgus macaque as late as 48 h following infection with Marburg virus resulted in 100% protection.⁶⁶ BioCryst Pharmaceuticals and NIAID have initiated a phase I clinical trial of BCX4430 in the United Kingdom.⁶⁷

PMOs are synthetic antisense molecules that are able to target mRNA in a sequence-specific fashion and suppress translation through steric hindrance. AVI-6002 consists of PMO AVI-7537, which targets VP24, and AVI-7539, which targets VP35. AVI-6003 consists of PMO AVI-7287, which targets MARV VP24, and AVI-7288, which targets MARV NP. AVI-6002 protected 62.5% of rhesus monkeys against EBOV infection when given 30–60 min postinfection and daily for 14 days. AVI-6003 protected 100% of cynomolgus macaques from MARV infection when given 30–60 min postinfection and daily for 14 days.⁷⁰ A follow-up study discovered AVI-7537 alone protected 75% (6/8) cynomolgus macaques against EBOV when administered 1 h \pm 30 min after challenge and daily for 14 days whereas treatment solely with AVI-7539 did not result in any survival past 10 days postinfection.⁷¹ Results from a phase I clinical trial evaluating the safety and pharmacokinetic profiles of AVI-6002 and AVI-6003 in two groups of 30 individuals revealed that both AVI-6002 and AVI-6003 were safe and well tolerated with doses ranging from 0.005 to 4.5 mg/kg per component.⁷² Small-interfering RNA (siRNA) can inhibit the translation of mRNA and has been shown to specifically downregulate MARV transcription of nucleocapsid protein in Vero E6 cells, significantly decreasing viral protein production and viral release.⁷³ The design of stable nucleic acid–lipid particles (SNALP) to efficiently deliver siRNAs in vivo to target cells facilitated therapeutic evaluation. SNALP-encapsulated siRNAs targeting the EBOV polymerase L gene completely protected five guinea pigs from viremia and

death when administered in a dose of 0.75 mg/kg 1 h after challenge and daily on days 1–6 postinfection.⁷⁴ Treatment with a SNALP carrying siRNAs targeting EBOV L polymerase, VP24, and VP35, called TKM-Ebola, resulted in the survival of 2/3 macaques that were given a 2 mg/kg dose after 30 min and on days 1, 3, and 5 after EBOV challenge and 4/4 macaques that were given a 2 mg/kg dose after 30 min and on days 1, 2, 3, 4, 5, and 6 postchallenge.⁷⁵ However, phase I trials of TKM-Ebola in early 2014 were halted after elevated cytokine levels were detected in healthy participants. In October 2014, the FDA expanded access to TKM-Ebola on an emergency basis.⁷⁶ Brincidofovir (CMX001) is a lipid conjugate of cidofovir that can be converted intracellularly into cidofovir diphosphate, which has been shown to inhibit DNA polymerase. Although brincidofovir is in clinical trials for diseases caused by DNA viruses, in vitro studies carried out at the CDC and NIH revealed that brincidofovir is active against the Ebola virus.⁶⁸ While the mechanism by which brincidofovir acts against Ebola is unknown, brincidofovir has been used to treat two patients infected with Ebola in the United States (one survived and one died). Although Chimerix initiated a phase II clinical trial in October 2014,⁶⁹ they withdrew the study in January 2015.

Given the large body of both experimental and clinical data supporting the role of antibodies in protection against EBOV infections, several monoclonal treatment modalities were tested. Human monoclonal antibody (mAb) KZ52, which targeted one epitope in EBOV-GP, was able to protect guinea pigs⁷⁷ but failed to protect rhesus macaques against lethal challenge with EBOV strain Kikwit when administered 1 day before challenge and 4 days after challenge.⁷⁸ On the other hand, the administration of a combination of two human–mouse chimeric neutralizing mAbs to three nonhuman primates 1 day prior to as well as 1 and 3 days after lethal EBOV challenge resulted in the protection of 1/3 nonhuman primates and a prolonged time to death in a second animal.⁷⁹ Passive transfer of polyclonal IgG purified from vaccinated NHPs that survived challenge with either EBOV or MARV as late as 48 h after virus challenge protected naïve NHPs against both MARV and EBOV lethal challenge.⁸⁰ Together with the previously failed trials of monoclonal antibodies, these observations strongly suggested the need to target multiple epitopes on GP in order to achieve protection against EBOV.

To develop a multispecificity antibody cocktail, several EBOV GP-specific mAbs generated using the rVSV/ΔG/GP EBOV vaccine were evaluated in immunocompetent mice and guinea pigs individually or as pools of 3 to 4 mAbs. In contrast to individually administered mAbs, which were ineffective, pools of 3 mAbs were found to give complete protection in guinea pigs when administered 2 days postinfection.⁸¹ These studies paved the way for the development of two cocktails of monoclonal antibodies (either chimeric (c) or human (h)) termed MB-003 (clones c13C6, h13F6, and c6D8) and ZMab (clones m1H3, m2G4, and m4G7) that mediated the complete protection of nonhuman primates when administered 1–3 days post-EBOV challenge.⁸² Further refinement of these cocktails led to the generation of ZMapp containing monoclonal antibodies with the highest efficacy (c13C6, c2G4, and c4G7). This combination is a highly effective postexposure therapeutic that can protect nonhuman primates even when administered 5 days postexposure to Ebola to symptomatic animals⁸³ (Table 2). ZMapp has been used to treat seven patients infected with EBOV, resulting in five patients surviving.⁸⁴ In February 2015, the NIAID initiated phase I

clinical trial evaluations of ZMapp in Liberia and the United States. The study will be conducted on 200 patients positive for Ebola virus infection who will be randomly assigned to one of two groups: the control group will receive the current standard of care, and the second group will receive three infusions of ZMapp administered 3 days apart.⁸⁵

CONCLUSIONS AND FUTURE OUTLOOK

EBOV is one of the most feared pathogens due to its high lethality rates. Subversion of the innate immune response by Ebola virus occurs at various levels from suppression of the IFN response to the generation of a robust cytokine storm to the inhibition of dendritic cell maturation. Dysregulation of innate immune pathways together with the characteristic lymphopenia of Ebola prevents the development of cellular and humoral responses as evident by the lack of EBOV-specific T and B cell responses and the sequestration of neutralizing antibodies. The current Ebola outbreak in West Africa has spurred a public health emergency and highlighted the need for licensed vaccines and postexposure therapeutics. With the substantial amount of data pointing to the critical role of antibody-mediated protection against Ebola, it is imperative that research efforts focus on evaluating the safety and efficacy of monoclonal antibody cocktails. Highly promising antibody cocktail ZMapp is grown in tobacco plants, which is thought to take less time and money to produce than in rodents. However, clinical evaluations of ZMapp were delayed until early of 2015 due to the slow production of ZMapp. The production of one course of treatment (14 g) requires 78 tobacco plants to produce.⁸⁶ Consequently, efforts must be allocated to developing novel methods to increase yields of the monoclonal antibody cocktail. Although using more traditional methods of generating monoclonal antibodies in rodents may be a slower route than tobacco plant production, this established technique might enable increased production in order to ensure that enough of the treatment is available for further clinical evaluations and future outbreaks.

In addition to monoclonal cocktails, efforts to evaluate and license the rVSV/ΔG/GP EBOV vaccine should be increased. Compared to the rAd vaccine platform, advantages of the rVSV/ΔG/GP EBOV platform include limited pre-existing immunity, the ability to administer it mucosally, the long duration of immunity after one dose without booster requirement, and its potential use as a postexposure vaccination. With the Ebola outbreak slowing down, the opportunity to obtain efficacy data through human trials may decrease. In this case, the FDA animal rule should be used, in which laboratory animal data is used to show efficacy, in order to move vaccine candidates through the approval process and start stockpiling the most promising candidate. This will ensure that we have immediate immunization available to protect frontline healthcare workers and contain the spread of a future outbreak.

The 2014 outbreak highlights that Ebola virus disease is not limited to central Africa and is a global problem that requires global cooperation. The World Health Organization's role in the event of an outbreak should be to effectively coordinate regional collaboration where data and expertise relevant to the outbreak in addition to critical information about disease surveillance is shared. Efforts to improve the infrastructure of healthcare in African nations should be undertaken, such as developing inexpensive systems to equip hospitals to be able to manage routine healthcare needs as well as outbreaks.

Educating the public is also equally important, as many individuals in Ebola outbreak countries are fearful of modern health care, preventing those who have come into contact with the disease to seek health care and contributing to the spread of the virus. Therefore, increased training of the local health care force to build trust and effectively communicate with the public in times of crisis is urgently required. Community engagement is absolutely required, especially when preventive measures are at odds with cultural beliefs and religious practices. Public health education regarding the dynamics of Ebola transmission is also urgently needed, especially with traditional burial practices, which involve washing and touching of the deceased. It is clear that our approaches in combatting Ebola virus disease will engage many areas ranging from vaccine and therapeutic discovery to improving the healthcare infrastructure, training, and community participation.

AUTHOR INFORMATION

Corresponding Author

*Phone: 951-827-7774. Email: messaoud@ucr.edu.

Notes

The authors declare no competing financial interest.

REFERENCES

- (1) (a) Muhlbinger, E., Lotfering, B., Klenk, H. D., and Becker, S. (1998) Three of the four nucleocapsid proteins of Marburg virus, NP, VP35, and L, are sufficient to mediate replication and transcription of Marburg virus-specific monocistronic minigenomes. *J. Virol.* 72, 8756–8764. (b) Dolnik, O., Kolesnikova, L., and Becker, S. (2008) Filoviruses: Interactions with the host cell. *Cell. Mol. Life Sci.* 65, 756–776.
- (2) (a) Ruigrok, R. W., Schoehn, G., Dessen, A., Forest, E., Volchkov, V., Dolnik, O., Klenk, H. D., and Weissenhorn, W. (2000) Structural characterization and membrane binding properties of the matrix protein VP40 of Ebola virus. *J. Mol. Biol.* 300, 103–112. (b) Huang, Y., Xu, L., Sun, Y., and Nabel, G. J. (2002) The assembly of Ebola virus nucleocapsid requires virion-associated proteins 35 and 24 and posttranslational modification of nucleoprotein. *Mol. Cell* 10, 307–316. (c) Hartlieb, B., and Weissenhorn, W. (2006) Filovirus assembly and budding. *Virology* 344, 64–70.
- (3) Kawaoka, Y. (2005) How Ebola virus infects cells. *New Engl. J. Med.* 352, 2645–2646.
- (4) Volchkov, V. E., Volchkova, V. A., Slenczka, W., Klenk, H. D., and Feldmann, H. (1998) Release of viral glycoproteins during Ebola virus infection. *Virology* 245, 110–119.
- (5) (a) Kuhn, J. H., Bao, Y., Bavari, S., Becker, S., Bradfute, S., Brauburger, K., Rodney Brister, J., Bukreyev, A. A., Cai, Y., Chandran, K., Davey, R. A., Dolnik, O., Dye, J. M., Enterlein, S., Gonzalez, J. P., Formenty, P., Freiberg, A. N., Hensley, L. E., Hoenen, T., Honko, A. N., Ignatyev, G. M., Jahrling, P. B., Johnson, K. M., Klenk, H. D., Kobinger, G., Lackmeyer, M. G., Leroy, E. M., Lever, M. S., Muhlbinger, E., Netesov, S. V., Olinger, G. G., Palacios, G., Patterson, J. L., Paweska, J. T., Pitt, L., Radoshitzky, S. R., Ryabchikova, E. I., Saphire, E. O., Shestopalov, A. M., Smither, S. J., Sullivan, N. J., Swanepoel, R., Takada, A., Towner, J. S., van der Groen, G., Volchkov, V. E., Volchkova, V. A., Wahl-Jensen, V., Warren, T. K., Warfield, K. L., Weidmann, M., and Nichol, S. T. (2014) Virus nomenclature below the species level: a standardized nomenclature for filovirus strains and variants rescued from cDNA. *Arch. Virol.* 159, 1229–1237. (b) Lefebvre, A., Fiet, C., Belpois-Duchamp, C., Tiv, M., Astruc, K., and Aho Glele, L. S. (2014) Case fatality rates of Ebola virus diseases: a meta-analysis of World Health Organization data. *Med. Mal. Infect.* 44, 412–416.
- (6) (a) Feldmann, H., and Geisbert, T. W. (2011) Ebola haemorrhagic fever. *Lancet* 377, 849–862. (b) Bwaka, M. A., Bonnet, M. J., Calain, P., Colebunders, R., De Roo, A., Guimard, Y.,

- Katwiki, K. R., Kibadi, K., Kipasa, M. A., Kuvula, K. J., Mapanda, B. B., Massamba, M., Mupapa, K. D., Muyembe-Tamfum, J. J., Ndaberey, E., Peters, C. J., Rollin, P. E., Van den Enden, E., and Van den Enden, E. (1999) Ebola hemorrhagic fever in Kikwit, Democratic Republic of the Congo: clinical observations in 103 patients. *J. Infect. Dis.* 179, S1–S7.
- (7) (a) System Ebola Epidemiology Team, I. M., Update: Ebola Virus Disease Epidemic - West Africa, November 2014. *Morbidity and Mortality Weekly Report* 2014, 63, 1064–1066. (b) Team, W. H. O. E. R. (2014) Ebola virus disease in West Africa—the first 9 months of the epidemic and forward projections. *N. Engl. J. Med.* 371, 1481–1495.
- (8) Baize, S., Pannetier, D., Oestereich, L., Rieger, T., Koivogui, L., Magassouba, N., Soropogui, B., Sow, M. S., Keita, S., De Clerck, H., Tiffany, A., Dominguez, G., Loua, M., Traore, A., Kolie, M., Malano, E. R., Heleze, E., Bocquin, A., Mely, S., Raoul, H., Caro, V., Cadar, D., Gabriel, M., Pahlmann, M., Tappe, D., Schmidt-Chanasit, J., Impouma, B., Diallo, A. K., Formenty, P., Van Herp, M., and Gunther, S. (2014) Emergence of Zaire Ebola virus disease in Guinea. *New Engl. J. Med.* 371, 1418–1425.
- (9) Gire, S. K., Goba, A., Andersen, K. G., Sealfon, R. S., Park, D. J., Kanneh, L., Jalloh, S., Momoh, M., Fullah, M., Dudas, G., Wohl, S., Moses, L. M., Yozwiak, N. L., Winnicki, S., Matranga, C. B., Malboeuf, C. M., Qu, J., Gladden, A. D., Schaffner, S. F., Yang, X., Jiang, P. P., Nekoui, M., Colubri, A., Coomber, M. R., Fonnies, M., Moigboi, A., Gbakie, M., Kamara, F. K., Tucker, V., Konuwa, E., Saffa, S., Sellu, J., Jalloh, A. A., Kovoma, A., Koninga, J., Mustapha, I., Kargbo, K., Foday, M., Yillah, M., Kanneh, F., Robert, W., Massally, J. L., Chapman, S. B., Bochicchio, J., Murphy, C., Nusbaum, C., Young, S., Birren, B. W., Grant, D. S., Scheiffelin, J. S., Lander, E. S., Hapfi, C., Gevao, S. M., Gnirke, A., Rambaut, A., Garry, R. F., Khan, S. H., and Sabeti, P. C. (2014) Genomic surveillance elucidates Ebola virus origin and transmission during the 2014 outbreak. *Science* 345, 1369–1372.
- (10) (a) Jaax, N. K., Davis, K. J., Geisbert, T. J., Vogel, P., Jaax, G. P., Topper, M., and Jahrling, P. B. (1996) Lethal experimental infection of rhesus monkeys with Ebola-Zaire (Mayinga) virus by the oral and conjunctival route of exposure. *Arch. Pathol. Lab. Med.* 120, 140–155. (b) Geisbert, T. W., Hensley, L. E., Larsen, T., Young, H. A., Reed, D. S., Geisbert, J. B., Scott, D. P., Kagan, E., Jahrling, P. B., and Davis, K. J. (2003) Pathogenesis of Ebola hemorrhagic fever in cynomolgus macaques: evidence that dendritic cells are early and sustained targets of infection. *Am. J. Pathol.* 163, 2347–2370. (c) Geisbert, T. W., Young, H. A., Jahrling, P. B., Davis, K. J., Larsen, T., Kagan, E., and Hensley, L. E. (2003) Pathogenesis of Ebola hemorrhagic fever in primate models: evidence that hemorrhage is not a direct effect of virus-induced cytolysis of endothelial cells. *Am. J. Pathol.* 163, 2371–2382.
- (11) (a) Bray, M., Davis, K., Geisbert, T., Schmaljohn, C., and Huggins, J. (1998) A mouse model for evaluation of prophylaxis and therapy of Ebola hemorrhagic fever. *J. Infect. Dis.* 178, 651–661. (b) Connolly, B. M., Steele, K. E., Davis, K. J., Geisbert, T. W., Kell, W. M., Jaax, N. K., and Jahrling, P. B. (1999) Pathogenesis of experimental Ebola virus infection in guinea pigs. *J. Infect. Dis.* 179, S203–S217.
- (12) Rasmussen, A. L., Okumura, A., Ferris, M. T., Green, R., Feldmann, F., Kelly, S. M., Scott, D. P., Safronetz, D., Haddock, E., LaCasse, R., Thomas, M. J., Sova, P., Carter, V. S., Weiss, J. M., Miller, D. R., Shaw, G. D., Korth, M. J., Heise, M. T., Baric, R. S., de Villena, F. P., Feldmann, H., and Katze, M. G. (2014) Host genetic diversity enables Ebola hemorrhagic fever pathogenesis and resistance. *Science* 346, 987–991.
- (13) Bray, M., Hatfill, S., Hensley, L., and Huggins, J. W. (2001) Haematological, biochemical and coagulation changes in mice, guinea-pigs and monkeys infected with a mouse-adapted variant of Ebola Zaire virus. *J. Comp. Pathol.* 125, 243–253.
- (14) Ebihara, H., Takada, A., Kobasa, D., Jones, S., Neumann, G., Theriault, S., Bray, M., Feldmann, H., and Kawaoka, Y. (2006) Molecular determinants of Ebola virus virulence in mice. *PLoS Pathog.* 2, e73.
- (15) Geisbert, T. W., Hensley, L. E., Gibb, T. R., Steele, K. E., Jaax, N. K., and Jahrling, P. B. (2000) Apoptosis induced in vitro and in vivo

during infection by Ebola and Marburg viruses. *Lab. Invest.* 80, 171–186.

(16) Stroher, U., West, E., Bugany, H., Klenk, H. D., Schnittler, H. J., and Feldmann, H. (2001) Infection and activation of monocytes by Marburg and Ebola viruses. *J. Virol.* 75, 11025–11033.

(17) (a) Baize, S., Leroy, E. M., Georges, A. J., Georges-Courbot, M. C., Capron, M., Bedjabaga, I., Lansoud-Soukate, J., and Mavoungou, E. (2002) Inflammatory responses in Ebola virus-infected patients. *Clin. Exp. Immunol.* 128, 163–168. (b) Ebihara, H., Rockx, B., Marzi, A., Feldmann, F., Haddock, E., Brining, D., LaCasse, R. A., Gardner, D., and Feldmann, H. (2011) Host response dynamics following lethal infection of rhesus macaques with Zaire ebolavirus. *J. Infect. Dis.* 204, S991–S999. (c) Hensley, L. E., Young, H. A., Jahrling, P. B., and Geisbert, T. W. (2002) Proinflammatory response during Ebola virus infection of primate models: possible involvement of the tumor necrosis factor receptor superfamily. *Immunol. Lett.* 80, 169–179. (d) Sanchez, A., Lukwiya, M., Bausch, D., Mahanty, S., Sanchez, A. J., Wagoner, K. D., and Rollin, P. E. (2004) Analysis of human peripheral blood samples from fatal and nonfatal cases of Ebola (Sudan) hemorrhagic fever: cellular responses, virus load, and nitric oxide levels. *J. Virol.* 78, 10370–10377. (e) Villinger, F., Rollin, P. E., Brar, S. S., Chikkala, N. F., Winter, J., Sundstrom, J. B., Zaki, S. R., Swanepoel, R., Ansari, A. A., and Peters, C. J. (1999) Markedly elevated levels of interferon (IFN)-gamma, IFN-alpha, interleukin (IL)-2, IL-10, and tumor necrosis factor-alpha associated with fatal Ebola virus infection. *J. Infect. Dis.* 179, S188–S191. (f) Wauquier, N., Becquart, P., Padilla, C., Baize, S., and Leroy, E. M. (2010) Human fatal zaire ebola virus infection is associated with an aberrant innate immunity and with massive lymphocyte apoptosis. *PLoS Negl. Trop. Dis.* 4, e837.

(18) (a) Reid, S. P., Valmas, C., Martinez, O., Sanchez, F. M., and Basler, C. F. (2007) Ebola virus VP24 proteins inhibit the interaction of NPI-1 subfamily karyopherin alpha proteins with activated STAT1. *J. Virol.* 81, 13469–13477. (b) Reid, S. P., Leung, L. W., Hartman, A. L., Martinez, O., Shaw, M. L., Carbone, C., Volchkov, V. E., Nichol, S. T., and Basler, C. F. (2006) Ebola virus VP24 binds karyopherin alpha1 and blocks STAT1 nuclear accumulation. *J. Virol.* 80, 5156–5167. (c) Mateo, M., Reid, S. P., Leung, L. W., Basler, C. F., and Volchkov, V. E. (2010) Ebolavirus VP24 binding to karyopherins is required for inhibition of interferon signaling. *J. Virol.* 84, 1169–1175.

(19) Xu, W., Edwards, M. R., Borek, D. M., Feagins, A. R., Mittal, A., Alinger, J. B., Berry, K. N., Yen, B., Hamilton, J., Brett, T. J., Pappu, R. V., Leung, D. W., Basler, C. F., and Amarasinghe, G. K. (2014) Ebola virus VP24 targets a unique NLS binding site on karyopherin alpha 5 to selectively compete with nuclear import of phosphorylated STAT1. *Cell Host Microbe* 16, 187–200.

(20) Basler, C. F., Mikulasova, A., Martinez-Sobrido, L., Paragas, J., Muhlberger, E., Bray, M., Klenk, H. D., Palese, P., and Garcia-Sastre, A. (2003) The Ebola virus VP35 protein inhibits activation of interferon regulatory factor 3. *J. Virol.* 77, 7945–7956.

(21) Chang, T. H., Kubota, T., Matsuoka, M., Jones, S., Bradfute, S. B., Bray, M., and Ozato, K. (2009) Ebola Zaire virus blocks type I interferon production by exploiting the host SUMO modification machinery. *PLoS Pathog.* 5, e1000493.

(22) Cardenas, W. B., Loo, Y. M., Gale, M., Jr., Hartman, A. L., Kimberlin, C. R., Martinez-Sobrido, L., Saphire, E. O., and Basler, C. F. (2006) Ebola virus VP35 protein binds double-stranded RNA and inhibits alpha/beta interferon production induced by RIG-I signaling. *J. Virol.* 80, 5168–5178.

(23) Luthra, P., Ramanan, P., Mire, C. E., Weisend, C., Tsuda, Y., Yen, B., Liu, G., Leung, D. W., Geisbert, T. W., Ebihara, H., Amarasinghe, G. K., and Basler, C. F. (2013) Mutual antagonism between the Ebola virus VP35 protein and the RIG-I activator PACT determines infection outcome. *Cell Host Microbe* 14, 74–84.

(24) Prins, K. C., Cardenas, W. B., and Basler, C. F. (2009) Ebola virus protein VP35 impairs the function of interferon regulatory factor-activating kinases IKKepsilon and TBK-1. *J. Virol.* 83, 3069–3077.

(25) Lubaki, N. M., Ilinykh, P., Pietzsch, C., Tigabu, B., Freiberg, A. N., Koup, R. A., and Bukreyev, A. (2013) The lack of maturation of

Ebola virus-infected dendritic cells results from the cooperative effect of at least two viral domains. *J. Virol.* 87, 7471–7485.

(26) Mahanty, S., Hutchinson, K., Agarwal, S., McRae, M., Rollin, P. E., and Pulendran, B. (2003) Cutting edge: impairment of dendritic cells and adaptive immunity by Ebola and Lassa viruses. *J. Immunol.* 170, 2797–2801.

(27) Gupta, M., Spiropoulou, C., and Rollin, P. E. (2007) Ebola virus infection of human PBMCs causes massive death of macrophages, CD4 and CD8 T cell sub-populations in vitro. *Virology* 364, 45–54.

(28) Bradfute, S. B., Braun, D. R., Shamblin, J. D., Geisbert, J. B., Paragas, J., Garrison, A., Hensley, L. E., and Geisbert, T. W. (2007) Lymphocyte death in a mouse model of Ebola virus infection. *J. Infect. Dis.* 196, S296–S304.

(29) Reed, D. S., Hensley, L. E., Geisbert, J. B., Jahrling, P. B., and Geisbert, T. W. (2004) Depletion of peripheral blood T lymphocytes and NK cells during the course of ebola hemorrhagic fever in cynomolgus macaques. *Viral Immunol.* 17, 390–400.

(30) Mehedi, M., Falzarano, D., Seebach, J., Hu, X., Carpenter, M. S., Schnittler, H. J., and Feldmann, H. (2011) A new Ebola virus nonstructural glycoprotein expressed through RNA editing. *J. Virol.* 85, 5406–5414.

(31) (a) Baize, S., Leroy, E. M., Mavoungou, E., and Fisher-Hoch, S. P. (2000) Apoptosis in fatal Ebola infection. Does the virus toll the bell for immune system? *Apoptosis* 5, 5–7. (b) Baize, S., Leroy, E. M., Georges-Courbot, M. C., Capron, M., Lansoud-Soukate, J., Debre, P., Fisher-Hoch, S. P., McCormick, J. B., and Georges, A. J. (1999) Defective humoral responses and extensive intravascular apoptosis are associated with fatal outcome in Ebola virus-infected patients. *Nat. Med.* 5, 423–426.

(32) (a) Rath, P. C., and Aggarwal, B. B. (1999) TNF-induced signaling in apoptosis. *J. Clin. Immunol.* 19, 350–364. (b) Simon, H. U., Haj-Yehia, A., and Levi-Schaffer, F. (2000) Role of reactive oxygen species (ROS) in apoptosis induction. *Apoptosis* 5, 415–418. (c) Snyder, C. M., Shroff, E. H., Liu, J., and Chandel, N. S. (2009) Nitric oxide induces cell death by regulating anti-apoptotic BCL-2 family members. *PLoS One* 4, e7059.

(33) Leroy, E. M., Baize, S., Debre, P., Lansoud-Soukate, J., and Mavoungou, E. (2001) Early immune responses accompanying human asymptomatic Ebola infections. *Clin. Exp. Immunol.* 124, 453–460.

(34) Leroy, E. M., Baize, S., Volchkov, V. E., Fisher-Hoch, S. P., Georges-Courbot, M. C., Lansoud-Soukate, J., Capron, M., Debre, P., McCormick, J. B., and Georges, A. J. (2000) Human asymptomatic Ebola infection and strong inflammatory response. *Lancet* 355, 2210–2215.

(35) Mohan, G. S., Li, W., Ye, L., Compans, R. W., and Yang, C. (2012) Antigenic subversion: a novel mechanism of host immune evasion by Ebola virus. *PLoS Pathog.* 8, e1003065.

(36) Mupapa, K., Massamba, M., Kibadi, K., Kuvula, K., Bwaka, A., Kipasa, M., Colebunders, R., and Muyembe-Tamfum, J. J. (1999) Treatment of Ebola hemorrhagic fever with blood transfusions from convalescent patients. International Scientific and Technical Committee. *J. Infect. Dis.* 179, S18–S23.

(37) Gupta, M., Mahanty, S., Bray, M., Ahmed, R., and Rollin, P. E. (2001) Passive transfer of antibodies protects immunocompetent and immunodeficient mice against lethal Ebola virus infection without complete inhibition of viral replication. *J. Virol.* 75, 4649–4654.

(38) Friedrich, B. M., Trefry, J. C., Biggins, J. E., Hensley, L. E., Honko, A. N., Smith, D. R., and Olinger, G. G. (2012) Potential vaccines and post-exposure treatments for filovirus infections. *Viruses* 4, 1619–50.

(39) Sullivan, N. J., Geisbert, T. W., Geisbert, J. B., Shedlock, D. J., Xu, L., Lamoreaux, L., Custers, J. H., Popernack, P. M., Yang, Z. Y., Pau, M. G., Roederer, M., Koup, R. A., Goudsmit, J., Jahrling, P. B., and Nabel, G. J. (2006) Immune protection of nonhuman primates against Ebola virus with single low-dose adenovirus vectors encoding modified GPs. *PLoS Med.* 3, e177.

(40) Sullivan, N. J., Hensley, L., Asiedu, C., Geisbert, T. W., Stanley, D., Johnson, J., Honko, A., Olinger, G., Bailey, M., Geisbert, J. B., Reimann, K. A., Bao, S., Rao, S., Roederer, M., Jahrling, P. B., Koup, R.

A., and Nabel, G. J. (2011) CD8+ cellular immunity mediates rAd5 vaccine protection against Ebola virus infection of nonhuman primates. *Nat. Med.* 17, 1128–1131.

(41) Stanley, D. A., Honko, A. N., Asiedu, C., Trefry, J. C., Lau-Kilby, A. W., Johnson, J. C., Hensley, L., Ammendola, V., Abbate, A., Grazioli, F., Foulds, K. E., Cheng, C., Wang, L., Donaldson, M. M., Colloca, S., Folgori, A., Roederer, M., Nabel, G. J., Mascola, J., Nicosia, A., Cortese, R., Koup, R. A., and Sullivan, N. J. (2014) Chimpanzee adenovirus vaccine generates acute and durable protective immunity against ebolavirus challenge. *Nat. Med.* 20, 1126–1129.

(42) Ledgerwood, J. E., DeZure, A. D., Stanley, D. A., Novik, L., Enama, M. E., Berkowitz, N. M., Hu, Z., Joshi, G., Ploquin, A., Sitar, S., Gordon, I. J., Plummer, S. A., Holman, L. A., Hendel, C. S., Yamshchikov, G., Roman, F., Nicosia, A., Colloca, S., Cortese, R., Bailer, R. T., Schwartz, R. M., Roederer, M., Mascola, J. R., Koup, R. A., Sullivan, N. J., and Graham, B. S. (2014) Chimpanzee Adenovirus Vector Ebola Vaccine—Preliminary Report. *N. Engl. J. Med.*, DOI: 10.1056/NEJMoal410863.

(43) Diseases, N. I. o. A. a. I. PREVAIL Phase2/3 Clinical Trial of Investigational Ebola Vaccines. <http://www.niaid.nih.gov/news/QA/Pages/EbolaVaxResultsQA.aspx> (accessed March 7, 2015).

(44) Stover, K. NIH to Launch Human Safety Study of Ebola Vaccine Candidate. <http://www.niaid.nih.gov/news/newsreleases/2014/Pages/EbolaVaxCandidate.aspx> (accessed December 12, 2014).

(45) (a) Jones, S. M., Feldmann, H., Stroher, U., Geisbert, J. B., Fernando, L., Grolla, A., Klenk, H. D., Sullivan, N. J., Volchkov, V. E., Fritz, E. A., Daddario, K. M., Hensley, L. E., Jahrling, P. B., and Geisbert, T. W. (2005) Live attenuated recombinant vaccine protects nonhuman primates against Ebola and Marburg viruses. *Nat. Med.* 11, 786–790. (b) Jones, S. M., Stroher, U., Fernando, L., Qiu, X., Alimonti, J., Melito, P., Bray, M., Klenk, H. D., and Feldmann, H. (2007) Assessment of a vesicular stomatitis virus-based vaccine by use of the mouse model of Ebola virus hemorrhagic fever. *J. Infect. Dis.* 196, S404–S412.

(46) Marzi, A., Engelmann, F., Feldmann, F., Habberth, K., Shupert, W. L., Brining, D., Scott, D. P., Geisbert, T. W., Kawaoaka, Y., Katze, M. G., Feldmann, H., and Messaoudi, I. (2013) Antibodies are necessary for rVSV/ZEBOV-GP-mediated protection against lethal Ebola virus challenge in nonhuman primates. *Proc. Natl. Acad. Sci. U.S.A.* 110, 1893–1898.

(47) (a) Geisbert, T. W., Daddario-Dicaprio, K. M., Geisbert, J. B., Reed, D. S., Feldmann, F., Grolla, A., Stroher, U., Fritz, E. A., Hensley, L. E., Jones, S. M., and Feldmann, H. (2008) Vesicular stomatitis virus-based vaccines protect nonhuman primates against aerosol challenge with Ebola and Marburg viruses. *Vaccine* 26, 6894–6900. (b) Geisbert, T. W., Daddario-Dicaprio, K. M., Lewis, M. G., Geisbert, J. B., Grolla, A., Leung, A., Paragas, J., Matthias, L., Smith, M. A., Jones, S. M., Hensley, L. E., Feldmann, H., and Jahrling, P. B. (2008) Vesicular stomatitis virus-based ebola vaccine is well-tolerated and protects immunocompromised nonhuman primates. *PLoS Pathog.* 4, e1000225.

(48) Qiu, X., Fernando, L., Alimonti, J. B., Melito, P. L., Feldmann, F., Dick, D., Stroher, U., Feldmann, H., and Jones, S. M. (2009) Mucosal immunization of cynomolgus macaques with the VSVDeltaG/ZEBOVGP vaccine stimulates strong ebola GP-specific immune responses. *PLoS One* 4, e5547.

(49) Wong, G., Audet, J., Fernando, L., Fausther-Bovendo, H., Alimonti, J. B., Kobinger, G. P., and Qiu, X. (2014) Immunization with vesicular stomatitis virus vaccine expressing the Ebola glycoprotein provides sustained long-term protection in rodents. *Vaccine* 32, S722–S729.

(50) Mire, C. E., Geisbert, J. B., Agans, K. N., Satterfield, B. A., Versteeg, K. M., Fritz, E. A., Feldmann, H., Hensley, L. E., and Geisbert, T. W. (2014) Durability of a vesicular stomatitis virus-based marburg virus vaccine in nonhuman primates. *PLoS One* 9, e94355.

(51) Geisbert, T. W., Geisbert, J. B., Leung, A., Daddario-DiCaprio, K. M., Hensley, L. E., Grolla, A., and Feldmann, H. (2009) Single-injection vaccine protects nonhuman primates against infection with marburg virus and three species of ebola virus. *J. Virol.* 83, 7296–7304.

(52) (a) Feldmann, H., Jones, S. M., Daddario-DiCaprio, K. M., Geisbert, J. B., Stroher, U., Grolla, A., Bray, M., Fritz, E. A., Fernando, L., Feldmann, F., Hensley, L. E., and Geisbert, T. W. (2007) Effective post-exposure treatment of Ebola infection. *PLoS Pathog.* 3, e2. (b) Geisbert, T. W., Daddario-DiCaprio, K. M., Williams, K. J., Geisbert, J. B., Leung, A., Feldmann, F., Hensley, L. E., Feldmann, H., and Jones, S. M. (2008) Recombinant vesicular stomatitis virus vector mediates postexposure protection against Sudan Ebola hemorrhagic fever in nonhuman primates. *J. Virol.* 82, S664–S668.

(53) Gunther, S., Feldmann, H., Geisbert, T. W., Hensley, L. E., Rollin, P. E., Nichol, S. T., Stroher, U., Artsob, H., Peters, C. J., Ksiazek, T. G., Becker, S., ter Meulen, J., Olschlager, S., Schmidt-Chanasit, J., Sudeck, H., Burchard, G. D., and Schmiedel, S. (2011) Management of accidental exposure to Ebola virus in the biosafety level 4 laboratory, Hamburg, Germany. *J. Infect. Dis.* 204, S785–S790.

(54) Lai, L., Davey, R., Beck, A., Xu, Y., Suffredini, A. F., Palmore, T., Kabbani, S., Rogers, S., Kobinger, G., Alimonti, J., Link, C. J., Jr., Rubinson, L., Stroher, U., Wolcott, M., Dorman, W., Uyeki, T. M., Feldmann, H., Lane, H. C., and Mulligan, M. J. (2015) Emergency Postexposure Vaccination With Vesicular Stomatitis Virus-Vectored Ebola Vaccine After Needlestick. *JAMA*, DOI: 10.1001/jama.2015.1995.

(55) NIH STRIVE (Sierra Leone Trial to Introduce a Vaccine Against Ebola). <https://clinicaltrials.gov/ct2/show/record/NCT02378753> (accessed March 7, 2015).

(56) Chaib, F.; Jasarevic, T.; Harris, M. WHO welcomes approval of a second Ebola vaccine trial in Switzerland. <http://www.who.int/mediacentre/news/statements/2014/second-ebola-vaccine/en/> (accessed December 12, 2014).

(57) Fox, M. Joint Pain Interrupts Ebola Vaccine Trial. <http://www.nbcnews.com/storyline/ebola-virus-outbreak/joint-pain-interrupts-ebola-vaccine-trial-n266516> (accessed December 12, 2014).

(58) Stover, K. NIH Begins Early Human Clinical Trial of VSV Ebola Vaccine. <http://www.niaid.nih.gov/news/newsreleases/2014/Pages/VSEbolaVaccine.aspx> (accessed December 12, 2014).

(59) (a) Geisbert, T. W., Hensley, L. E., Jahrling, P. B., Larsen, T., Geisbert, J. B., Paragas, J., Young, H. A., Fredeking, T. M., Rote, W. E., and Vlasuk, G. P. (2003) Treatment of Ebola virus infection with a recombinant inhibitor of factor VIIa/tissue factor: a study in rhesus monkeys. *Lancet* 362, 1953–1958. (b) Hensley, L. E., Stevens, E. L., Yan, S. B., Geisbert, J. B., Macias, W. L., Larsen, T., Daddario-DiCaprio, K. M., Cassell, G. H., Jahrling, P. B., and Geisbert, T. W. (2007) Recombinant human activated protein C for the postexposure treatment of Ebola hemorrhagic fever. *J. Infect. Dis.* 196, S390–S399.

(60) Lee, A., Agnelli, G., Buller, H., Ginsberg, J., Heit, J., Rote, W., Vlasuk, G., Costantini, L., Julian, J., Comp, P., van Der Meer, J., Piovela, F., Raskob, G., and Gent, M. (2001) Dose-response study of recombinant factor VIIa/tissue factor inhibitor recombinant nematode anticoagulant protein c2 in prevention of postoperative venous thromboembolism in patients undergoing total knee replacement. *Circulation* 104, 74–78.

(61) Moons, A. H., Peters, R. J., Bijsterveld, N. R., Piek, J. J., Prins, M. H., Vlasuk, G. P., Rote, W. E., and Buller, H. R. (2003) Recombinant nematode anticoagulant protein c2, an inhibitor of the tissue factor/factor VIIa complex, in patients undergoing elective coronary angioplasty. *J. Am. College Cardiol.* 41, 2147–2153.

(62) Smith, L. M., Hensley, L. E., Geisbert, T. W., Johnson, J., Stossel, A., Honko, A., Yen, J. Y., Geisbert, J., Paragas, J., Fritz, E., Olinger, G., Young, H. A., Rubins, K. H., and Karp, C. L. (2013) Interferon-beta therapy prolongs survival in rhesus macaque models of Ebola and Marburg hemorrhagic fever. *J. Infect. Dis.* 208, 310–318.

(63) Oestereich, L., Ludtke, A., Wurr, S., Rieger, T., Munoz-Fontela, C., and Gunther, S. (2014) Successful treatment of advanced Ebola virus infection with T-705 (favipiravir) in a small animal model. *Antiviral Res.* 105, 17–21.

(64) Trompiz, G.; Lawson, H.; Powell, S. First French Ebola patient leaves hospital. <http://www.reuters.com/article/2014/10/04/us-health-ebola-france-idUSKCN0HTOD720141004>.

- (65) (a) WHO Ebola vaccines, therapies, and diagnostics. http://www.who.int/medicines/emp Ebola_q_as/en/. (b) NIH Efficacy of Favipiravir Against Ebola (IJKI). <https://clinicaltrials.gov/ct2/show/record/NCT02329054?term=favipiravir+ebola> (accessed March 7, 2015).
- (66) Warren, T. K., Wells, J., Panchal, R. G., Stuthman, K. S., Garza, N. L., Van Tongeren, S. A., Dong, L., Retterer, C. J., Eaton, B. P., Pegoraro, G., Honnold, S., Bantia, S., Kotian, P., Chen, X., Taubenheim, B. R., Welch, L. S., Minning, D. M., Babu, Y. S., Sheridan, W. P., and Bavari, S. (2014) Protection against filovirus diseases by a novel broad-spectrum nucleoside analogue BCX4430. *Nature* 508, 402–405.
- (67) NIH A Phase 1 Study to Evaluate the Safety, Tolerability and Pharmacokinetics of BCX4430. <https://clinicaltrials.gov/ct2/show/NCT02319772?term=BCX4430> (accessed March 7, 2015).
- (68) Schepers, J. T. *Chimerix Announces Emergency Investigational New Drug Applications for Brincidofovir Authorized by FDA for Patients with Ebola Virus Disease*. <http://ir.chimerix.com/releasedetail.cfm?releaseid=874647> (accessed December 12, 2014).
- (69) Health, N. I. o. An Open-Label, Multicenter Study of the Safety and Anti Viral Activity of Brincidofovir (BCV, CMX001) for Ebola Virus Disease. <https://clinicaltrials.gov/ct2/show/NCT02271347?term=chimerix> (accessed December 13, 2014).
- (70) Warren, T. K., Warfield, K. L., Wells, J., Swenson, D. L., Donner, K. S., Van Tongeren, S. A., Garza, N. L., Dong, L., Mourich, D. V., Crumley, S., Nichols, D. K., Iversen, P. L., and Bavari, S. (2010) Advanced antisense therapies for postexposure protection against lethal filovirus infections. *Nat. Med.* 16, 991–994.
- (71) Warren, T. K., Whitehouse, C. A., Wells, J., Welch, L., Heald, A. E., Charleston, J. S., Sazani, P., Reid, S. P., Iversen, P. L., and Bavari, S. (2015) A Single Phosphorodiamidate Morpholino Oligomer Targeting VP24 Protects Rhesus Monkeys against Lethal Ebola Virus Infection. *mBio* 6, e02344-14.
- (72) Heald, A. E., Iversen, P. L., Saoud, J. B., Sazani, P., Charleston, J. S., Axtelle, T., Wong, M., Smith, W. B., Vutikullird, A., and Kaye, E. (2014) Safety and pharmacokinetic profiles of phosphorodiamidate morpholino oligomers with activity against ebola virus and marburg virus: results of two single-ascending-dose studies. *Antimicrob. Agents Chemother.* 58, 6639–6647.
- (73) Fowler, T., Bamberg, S., M^ller, P., Klenk, H. D., Meyer, T. F., Becker, S., and Rudel, T. (2005) Inhibition of Marburg virus protein expression and viral release by RNA interference. *J. Gen. Virol.* 86, 1181–1188.
- (74) Geisbert, T. W., Hensley, L. E., Kagan, E., Yu, E. Z., Geisbert, J. B., Daddario-DiCaprio, K., Fritz, E. A., Jahrling, P. B., McClintock, K., Phelps, J. R., Lee, A. C., Judge, A., Jeffs, L. B., and MacLachlan, I. (2006) Postexposure protection of guinea pigs against a lethal ebola virus challenge is conferred by RNA interference. *J. Infect. Dis.* 193, 1650–1657.
- (75) Geisbert, T. W., Lee, A. C., Robbins, M., Geisbert, J. B., Honko, A. N., Sood, V., Johnson, J. C., de Jong, S., Tavakoli, I., Judge, A., Hensley, L. E., and MacLachlan, I. (2010) Postexposure protection of non-human primates against a lethal Ebola virus challenge with RNA interference: a proof-of-concept study. *Lancet* 375, 1896–1905.
- (76) Young, S. *FDA allows expanded use of experimental Ebola drug*. <http://www.cnn.com/2014/09/22/health/ebola-drug-tekmira/> (accessed December 12, 2014).
- (77) Parren, P. W., Geisbert, T. W., Maruyama, T., Jahrling, P. B., and Burton, D. R. (2002) Pre- and postexposure prophylaxis of Ebola virus infection in an animal model by passive transfer of a neutralizing human antibody. *J. Virol.* 76, 6408–6412.
- (78) Oswald, W. B., Geisbert, T. W., Davis, K. J., Geisbert, J. B., Sullivan, N. J., Jahrling, P. B., Parren, P. W., and Burton, D. R. (2007) Neutralizing antibody fails to impact the course of Ebola virus infection in monkeys. *PLoS Pathog.* 3, e9.
- (79) Marzi, A., Yoshida, R., Miyamoto, H., Ishijima, M., Suzuki, Y., Higuchi, M., Matsuyama, Y., Igarashi, M., Nakayama, E., Kuroda, M., Saijo, M., Feldmann, F., Brining, D., Feldmann, H., and Takada, A. (2012) Protective Efficacy of Neutralizing Monoclonal Antibodies in a Nonhuman Primate Model of Ebola Hemorrhagic Fever. *PLoS One* 7, e36192.
- (80) Dye, J. M., Herbert, A. S., Kuehne, A. I., Barth, J. F., Muhammad, M. A., Zak, S. E., Ortiz, R. A., Prugar, L. I., and Pratt, W. D. (2012) Postexposure antibody prophylaxis protects nonhuman primates from filovirus disease. *Proc. Natl. Acad. Sci. U.S.A.* 109, 5034–5039.
- (81) Qiu, X., Fernando, L., Melito, P. L., Audet, J., Feldmann, H., Kobinger, G., Alimonti, J. B., and Jones, S. M. (2012) Ebola GP-specific monoclonal antibodies protect mice and guinea pigs from lethal Ebola virus infection. *PLoS Negl. Trop. Dis.* 6, e1575.
- (82) (a) Qiu, X., Audet, J., Wong, G., Pillet, S., Bello, A., Cabral, T., Strong, J. E., Plummer, F., Corbett, C. R., Alimonti, J. B., and Kobinger, G. P. (2012) Successful treatment of ebola virus-infected cynomolgus macaques with monoclonal antibodies. *Sci. Transl. Med.* 4, 138ra81. (b) Olinger, G. G., Jr., Pettitt, J., Kim, D., Working, C., Bohorov, O., Bratcher, B., Hiatt, E., Hume, S. D., Johnson, A. K., Morton, J., Pauly, M., Whaley, K. J., Lear, C. M., Biggins, J. E., Scully, C., Hensley, L., and Zeitlin, L. (2012) Delayed treatment of Ebola virus infection with plant-derived monoclonal antibodies provides protection in rhesus macaques. *Proc. Natl. Acad. Sci. U.S.A.* 109, 18030–18035.
- (83) Qiu, X., Wong, G., Audet, J., Bello, A., Fernando, L., Alimonti, J. B., Fausther-Bovendo, H., Wei, H., Aviles, J., Hiatt, E., Johnson, A., Morton, J., Swope, K., Bohorov, O., Bohorova, N., Goodman, C., Kim, D., Pauly, M. H., Velasco, J., Pettitt, J., Olinger, G. G., Whaley, K., Xu, B., Strong, J. E., Zeitlin, L., and Kobinger, G. P. (2014) Reversion of advanced Ebola virus disease in nonhuman primates with ZMapp. *Nature* 514, 47–53.
- (84) McCarthy, M. (2014) US signs contract with ZMapp maker to accelerate development of the Ebola drug. *BMJ* 349, g5488.
- (85) Health, N. I. o. *Libera-US. clinical research partnership opens trial to test Ebola treatments*. 2015.
- (86) Park, A. *See How Ebola Drugs Grow in Tobacco Leaves*. <http://time.com/3457472/see-how-ebola-drugs-grow-in-tobacco-leaves/> (accessed March 7, 2015).
- (87) Madrid, P. B., Chopra, S., Manger, I. D., Gilfillan, L., Keepers, T. R., Shurtleff, A. C., Green, C. E., Iyer, L. V., Dilks, H. H., Davey, R. A., Kolokoltsov, A. A., Carrion, R., Jr., Patterson, J. L., Bavari, S., Panchal, R. G., Warren, T. K., Wells, J. B., Moos, W. H., Burke, R. L., and Tanga, M. J. (2013) A systematic screen of FDA-approved drugs for inhibitors of biological threat agents. *PLoS One* 8, e60579.
- (88) Johansen, L. M., Brannan, J. M., Delos, S. E., Shoemaker, C. J., Stossel, A., Lear, C., Hoffstrom, B. G., Dewald, L. E., Schornberg, K. L., Scully, C., Lehar, J., Hensley, L. E., White, J. M., and Olinger, G. G. (2013) FDA-approved selective estrogen receptor modulators inhibit Ebola virus infection. *Sci. Trans. Med.* 5, 190ra79.
- (89) Warren, T. K., Warfield, K. L., Wells, J., Enterlein, S., Smith, M., Ruthel, G., Yunus, A. S., Kinch, M. S., Goldblatt, M., Aman, M. J., and Bavari, S. (2010) Antiviral activity of a small-molecule inhibitor of filovirus infection. *Antimicrob. Agents Chemother.* 54, 2152–2159.
- (90) Kinch, M. S., Yunus, A. S., Lear, C., Mao, H., Chen, H., Fesseha, Z., Luo, G., Nelson, E. A., Li, L., Huang, Z., Murray, M., Ellis, W. Y., Hensley, L., Christopher-Hennings, J., Olinger, G. G., and Goldblatt, M. (2009) FGI-104: a broad-spectrum small molecule inhibitor of viral infection. *Am. J. Trans. Res.* 1, 87–98.
- (91) Aman, M. J., Kinch, M. S., Warfield, K., Warren, T., Yunus, A., Enterlein, S., Stavale, E., Wang, P., Chang, S., Tang, Q., Porter, K., Goldblatt, M., and Bavari, S. (2009) Development of a broad-spectrum antiviral with activity against Ebola virus. *Antiviral Res.* 83, 245–251.
- (92) Bray, M., Raymond, J. L., Geisbert, T., and Baker, R. O. (2002) 3-deazaneplanocin A induces massively increased interferon-alpha production in Ebola virus-infected mice. *Antiviral Res.* 55, 151–159.
- (93) Chang, J., Warren, T. K., Zhao, X., Gill, T., Guo, F., Wang, L., Comunale, M. A., Du, Y., Alonzi, D. S., Yu, W., Ye, H., Liu, F., Guo, J. T., Mehta, A., Cuconati, A., Butters, T. D., Bavari, S., Xu, X., and Block, T. M. (2013) Small molecule inhibitors of ER alpha-glucosidases are active against multiple hemorrhagic fever viruses. *Antiviral Res.* 98, 432–440.

(94) Panchal, R. G., Reid, S. P., Tran, J. P., Bergeron, A. A., Wells, J., Kota, K. P., Aman, J., and Bavari, S. (2012) Identification of an antioxidant small-molecule with broad-spectrum antiviral activity. *Antiviral Res.* 93, 23–29.

(95) Reid, S. P., Shurtleff, A. C., Costantino, J. A., Tritsch, S. R., Retterer, C., Spurgers, K. B., and Bavari, S. (2014) HSPA5 is an essential host factor for Ebola virus infection. *Antiviral Res.* 109, 171–174.

(96) Züst, R., Cervantes-Barragan, L., Habjan, M., Maier, R., Neuman, B. W., Ziebuhr, J., Szretter, K. J., Baker, S. C., Barchet, W., Diamond, M. S., Siddell, S. G., Ludewig, B., and Thiel, V. (2011) Ribose 2'-O-methylation provides a molecular signature for the distinction of self and non-self mRNA dependent on the RNA sensor Mda5. *Nature Immunol.* 12, 137–143.

(97) Ye, W., and Schnell, S. W. (2014) The enantiomers of the 1',6'-isomer of neplanocin A: synthesis and antiviral properties. *Bioorg. Med. Chem.* 22, 5315–5319.

(98) Cote, M., Misasi, J., Ren, T., Bruchez, A., Lee, K., Filone, C. M., Hensley, L., Li, Q., Ory, D., Chandran, K., and Cunningham, J. (2011) Small molecule inhibitors reveal Niemann-Pick C1 is essential for Ebola virus infection. *Nature* 477, 344–348.

(99) Basu, A., Li, B., Mills, D. M., Panchal, R. G., Cardinale, S. C., Butler, M. M., Peet, N. P., Majgier-Baranowska, H., Williams, J. D., Patel, I., Moir, D. T., Bavari, S., Ray, R., Farzan, M. R., Rong, L., and Bowlin, T. L. (2011) Identification of a small-molecule entry inhibitor for filoviruses. *J. Virol.* 85, 3106–3119.

(100) Wolf, M. C., Freiberg, A. N., Zhang, T., Akyol-Ataman, Z., Grock, A., Hong, P. W., Li, J., Watson, N. F., Fang, A. Q., Aguilar, H. C., Porotto, M., Honko, A. N., Damoiseaux, R., Miller, J. P., Woodson, S. E., Chantasirivisal, S., Fontanes, V., Negrete, O. A., Krogstad, P., Dasgupta, A., Moscona, A., Hensley, L. E., Whelan, S. P., Faull, K. F., Holbrook, M. R., Jung, M. E., and Lee, B. (2010) A broad-spectrum antiviral targeting entry of enveloped viruses. *Proc. Natl. Acad. Sci. U.S.A.* 107, 3157–3162.

(101) Filone, C. M., Hodges, E. N., Honeyman, B., Bushkin, G. G., Boyd, K., Platt, A., Ni, F., Strom, K., Hensley, L., Snyder, J. K., and Connor, J. H. (2013) Identification of a broad-spectrum inhibitor of viral RNA synthesis: validation of a prototype virus-based approach. *Chem. Biol.* 20, 424–433.

(102) Gehring, G., Rohrmann, K., Atenchong, N., Mittler, E., Becker, S., Dahlmann, F., Pohlmann, S., Vondran, F. W., David, S., Manns, M. P., Ciesek, S., and von Hahn, T. (2014) The clinically approved drugs amiodarone, dronedarone and verapamil inhibit filovirus cell entry. *J. Antimicrob. Chemother.* 69, 2123–2131.

(103) Hill-Batorski, L., Halfmann, P., Neumann, G., and Kawaoka, Y. (2013) The cytoprotective enzyme heme oxygenase-1 suppresses Ebola virus replication. *J. Virol.* 87, 13795–137802.

(104) Fauci, A. S.; Collins, F. S. *NIH Ebola Update: Working Toward Treatments and Vaccines*. <http://directorsblog.nih.gov/2014/10/14/nih-ebola-update-working-toward-treatments-and-vaccines/>.

(105) St. Vincent, M. R., Colpitts, C. C., Ustinov, A. V., Muqadas, M., Joyce, M. A., Barsby, N. L., Epand, R. F., Epand, R. M., Khramyshev, S. A., Valueva, O. A., Korshun, V. A., Tyrrell, D. L., and Schang, L. M. (2010) Rigid amphipathic fusion inhibitors, small molecule antiviral compounds against enveloped viruses. *Proc. Natl. Acad. Sci. U.S.A.* 107, 17339–17344.

(106) Elshabrawy, H. A., Fan, J., Haddad, C. S., Ratia, K., Broder, C. C., Caffrey, M., and Prabhakar, B. S. (2014) Identification of a broad-spectrum antiviral small molecule against severe acute respiratory syndrome coronavirus and Ebola, Hendra, and Nipah viruses by using a novel high-throughput screening assay. *J. Virol.* 88, 4353–4365.

(107) Fedson, D. S. (2015) A practical treatment for patients with ebola virus disease. *J. Infect. Dis.* 211, 661–662.