

CJ-1639: A Potent and Highly Selective Dopamine D3 Receptor Full Agonist

Jianyong Chen,[†] Gregory T. Collins,[‡] Beth Levant,^{||} James Woods,[‡] Jeffrey R. Deschamps,[⊥] and Shaomeng Wang^{*,†,§}

Departments of [†]Internal Medicine, [‡]Pharmacology, and [§]Medicinal Chemistry, University of Michigan, 1500 East Medical Center Drive, Ann Arbor, Michigan 48109, United States


^{||}Department of Pharmacology, Toxicology and Therapeutics, University of Kansas Medical Center, Kansas City, Kansas 66160, United States

[⊥]Center for Biomolecular Science and Engineering, Naval Research Laboratory, 4555 Overlook Avenue, Washington, DC 20375, United States

S Supporting Information

ABSTRACT: We have identified several ligands with high binding affinities to the dopamine D3 receptor and excellent selectivity over the D2 and D1 receptors. CJ-1639 (**17**) binds to the D3 receptor with a K_i value of 0.50 nM and displays a selectivity of >5000 times over D2 and D1 receptors in binding assays using dopamine receptors expressed in the native rat brain tissues. CJ-1639 binds to human D3 receptor with a K_i value of 3.61 nM and displays over >1000-fold selectivity over human D1 and D2 receptors. CJ-1639 is active at 0.01 mg/kg at the dopamine D3 receptor in the rat and only starts to show a modest D2 activity at doses as high as 10 mg/kg. CJ-1639 is the most potent and selective D3 full agonist reported to date.

KEYWORDS: Dopamine receptors, ligands, agonists, drug abuse



Dopamine is an endogenous neurotransmitter in the central nervous system and exerts its effects through activation of five distinct dopamine receptor subtypes that belong to the G protein-coupled receptor superfamily. The receptors are grouped into the D1-like (D1 and D5) and D2-like (D2, D3, and D4) receptor subtypes. Numerous studies have provided strong evidence that the D3 receptor is a promising therapeutic target for a variety of conditions, including drug abuse, restless legs syndrome, schizophrenia, Parkinson's disease, and depression.^{1–3} Extensive efforts have been devoted to the discovery and development of potent and selective D3 ligands, including agonists, partial agonists, and antagonists.^{5,6}

Because of the high degree of sequence identity within the transmembrane helices between D2 and D3 receptors and the near-identity of the residues inferred to form the binding site in these receptors,⁴ it has been a challenge to design highly selective D3 ligands with excellent aqueous solubility and bioavailability.^{5,6} Pramipexole (**1**), quinpirole (**2**), and 7-hydroxy dipropylamino-tetralin (7-OH-DPAT, **3**), three widely used D3 agonists (Figure 1), all have modest selectivity (<100-fold) over the D2 receptor based upon their in vitro binding data⁴ and a narrow range of selectivity at the D3 receptor over the D2 receptor in vivo.^{7,8} SB-277011A (**4**),⁹ SB-414796 (**5**),¹⁰ NGB 2904 (**6**),¹¹ and BP 897(**7**)¹² are four commonly used D3 antagonists (Figure 1), but they also have a selectivity of 100-fold or less for D3 over D2. Furthermore, these four ligands have limited aqueous solubility. R-PG648 (**8**) (Figure 1), whose design is based upon the core structure of NGB 2904, is a potent D3 antagonist with a selectivity of >400-fold over D2 and also has a much improved

aqueous solubility as compared to NGB 2904.¹³ Recently, compound **9** (Figure 1) was reported to be a potent and orally active D3 antagonist with a 500-fold selectivity over D2 in in vitro functional assays.¹⁴ The in vivo selectivity for compounds **8** and **9** at the D3 receptor over the D2 receptor has not been reported.

In our effort to identify D3 ligands with high selectivity and good solubility, we have previously reported the design of a set of new compounds based upon pramipexole (**1**),¹⁵ which included CJ-1037 (**10**) and CJ-998 (**11**) as the two best compounds (Figure 2). Both **10** and **11** bind to D3 with high affinities and have a good selectivity for D3 over D2 and excellent aqueous solubility (>50 mg/mL). Unlike pramipexole, which is a D3 full agonist, compounds **10** and **11** appear to behave as partial agonists with low intrinsic agonist activity in rats.¹⁵ However, despite their high binding affinities to D3, both compounds **10** and **11** only show in vivo activity at a very high dose (32 mg/kg), indicating their poor bioavailability in the brain. Here, we report further modifications based upon compound **11**. These efforts have now yielded a set of new D3 ligands (Figure 2 and Table 1) that are highly potent and selective, not only in vitro, but also in vivo.

Newman's group showed that introduction of a hydroxyl group in the linker region in their D3 ligands improves D3 selectivity and aqueous solubility.^{13,16} Accordingly, we designed and synthesized compounds **12** and **13** by placing a hydroxyl group in the cyclohexane ring of compound **11** to improve its

Received: April 19, 2011

Accepted: June 19, 2011

Published: June 28, 2011




Figure 1. Chemical structures of representative known D3 agonists (1–3) and antagonists (4–9).

Table 1. Binding Affinities at the D₁-Like, D₂-Like, and D₃ Receptors in Binding Assays Using Rat Brain^a

ligand	K _i ± SEM (nM)			selectivity	
	D ₃ ([³ H]-R(+)-7-OH-DPAT)	D ₂ -like [³ H]spiperone	D ₁ -like [³ H]SCH23390	D ₂ -like/D ₃	D ₁ -like/D ₃
1	0.45 ± 0.034	39 ± 4	>10000	87	>10000
11	0.80 ± 0.05	92 ± 9	>5000	115	>5000
12	1.6 ± 0.25	1628 ± 234	11557 ± 996	1039	7373
13	0.93 ± 0.094	3549 ± 421	14057 ± 1095	3816	15115
14	0.41 ± 0.038	584 ± 72	6125 ± 271	1416	14848
15	0.35 ± 0.019	564 ± 52	3058 ± 198	1627	8822
16	0.40 ± 0.087	725 ± 45	1616 ± 167	1827	4074
17	0.50 ± 0.080	2568 ± 302	10127 ± 1002	5136	20254

^a Binding affinities for all of the compounds in this table were evaluated under the assay conditions as described in the Supporting Information.

solubility and also to explore the effect of this hydroxyl group for binding and selectivity at the D₃ receptor (Figure 2). Compound **12** with a *cis*-hydroxycyclohexyl group and compound **13**, the corresponding *trans* isomer, have K_i values of 1.6 and 0.9 nM to D₃, respectively, in our in vitro binding assays using dopamine receptors expressed in the native rat brain tissues. Thus, they are as potent as compound **11**, indicating that introduction of a hydroxyl group at the bridge carbon atom of the cyclohexyl group is not detrimental to D₃ binding. Both compounds **12** and **13**, however, have a reduced binding affinity to D₂ as compared to pramipexole. As a result, **12** and **13** display a selectivity of >1000 times and >3000 times, respectively, for D₃ over D₂.

We next synthesized compounds **14** and **15** with a four-membered ring in the linker region to replace the six-membered ring in compound **11**. Compound **14**, with a *trans*-cyclobutyl group, has a K_i value of 0.41 nM to D₃ and is thus two times more potent than **11**. Compound **14** binds to D₂ with a K_i value of 584 nM and is six times less potent than **11**. Hence, compound **14** is a highly potent D₃ ligand and displays >1000-fold selectivity for D₃ over D₂. Compound **15**

with a *cis*-cyclobutyl group in the linker region binds to D₃ with a K_i value of 0.35 nM, very similar to that for compound **14**. Compound **15** also shows a >1000-fold selectivity for D₃ over D₂. Hence, replacement of the *trans*-cyclohexyl group in compound **11** with either a *trans*- or a *cis*-cyclobutyl group improves both the binding affinity for D₃ and the selectivity over D₂.

Encouraged by the promising binding and selectivity data observed for compounds **12**–**15**, we next synthesized compounds **16** and **17**, in which a hydroxyl group is introduced to the bridge carbon atom of the four-membered ring in compounds **14** and **15**, respectively. Compound **16** binds to D₃ with a K_i value of 0.40 nM and displays a selectivity of 1827 times over D₂. Compound **17** binds to D₃ with a K_i of 0.50 nM and displays a selectivity of >5000 times over D₂. In addition to their high binding affinities to D₃ and excellent selectivity over D₂, compounds **16** and **17** also have excellent aqueous solubility (>100 mg/mL) in their HCl salt form.

The syntheses of compounds **12** and **13** were outlined in Scheme 1. Basically, 2-naphthoyl chloride was reacted with




Figure 2. Chemical structures of two previously reported D₃ ligands, **10** and **11**, and designed new analogues **12–17**.

trans-4-aminocyclohexanol in the presence of triethylamine, followed by oxidation with PCC in DCM to give ketone **18**. Ketone **18** was treated with allylmagnesium bromide at $-78\text{ }^{\circ}\text{C}$ to afford *cis*-cyclohexanol **20** and *trans*-cyclohexanol **21**. Compounds **20** and **21** were separated by silica gel chromatography (hexane:ethyl acetate = 1:1). *cis*-Aldehyde **22** was obtained by the treatment of *cis*-cyclohexanol **20** with osmium tetroxide followed by sodium periodate. *trans*-Aldehyde **23** was obtained in a similar manner. Reductive amination of pramipexole **1** with **22** and **23** gave designed compounds **12** and **13**, respectively. Details of the synthesis of compounds **12–17** are provided in the Supporting Information.


The stereochemistry of *cis*-conformation of compound **12** was confirmed by X-ray crystallographic analysis of key intermediate **20** (Figure 3). The stereochemistry of *cis*-conformation of compound **16** was confirmed by X-ray crystallographic analysis of a key intermediate (see the Supporting Information).

Our previous studies have shown that the induction of yawning by D₂-like agonists is a sensitive and well-validated measure of agonist activity at the dopamine D₃ receptor,^{7,8,17} whereas the inhibition of yawning and induction of hypothermia by D₂-like agonists are indicative of an agonist activation at the D₂ receptor.^{17–19} We have evaluated compounds **16** and **17** for their *in vivo* function in the yawning and hypothermia assays in the rat and included pramipexole as the reference compound in the evaluations. The results are shown in Figure 4.

Pramipexole produced dose-dependent increases in yawning at low doses and inhibition of yawning and induction of hypothermia

at higher doses (Figure 4A,D). The data are consistent with its full agonist activity profile at the D₃ and D₂ receptors and a narrow *in vivo* selectivity at the D₃ receptor.^{7,8} Like pramipexole, both **16** and **17** produced a dose-dependent increase in yawning (Figure 4A). Unlike pramipexole, which induced yawning over a very narrow dose range, compound **16** and **17** produced increases in yawning over a very wide range of doses, with significant increases in yawning observed at a dose range of 0.1–10.0 mg/kg for compound **16** (CJ-1638) and 0.01–10.0 mg/kg for compound **17** (CJ-1639) (Figure 4A). Although both compounds have very similar binding affinities to D₃ based upon our *in vitro* binding data, our yawning assay showed that compound **17** is more active than **16** in activation of the D₃ receptor in the rat and has a significant yawning activity at a dose of 0.01 mg/kg. These data suggest that compound **17** has a better bioavailability in the brain than compound **16**. Both compounds produced peak levels of yawning at a dose of 1.0 mg/kg (Figure 4A).

Because **17** is more potent *in vivo* in the yawning assay, we focused our further evaluation on this compound. Consistent with previous reports, significant decreases in core body temperature were observed with doses of 0.32 and 1.0 mg/kg pramipexole (Figure 4D). In comparison, compound **17** had no significant effect on core body temperature at 3.2 mg/kg and had a significant but a modest effect at 10 mg/kg (Figure 4D). These data suggest that compound **17** readily crosses the blood–brain barrier and functions as a highly selective, full efficacy agonist at the D₃ receptor.

Scheme 1. Synthesis of Compounds 12 and 13^a

^a Conditions and reagents: (i) 2-Naphthoyl chloride, *trans*-4-aminocyclohexanol, triethylamine, DCM. (ii) PCC, DCM, room temperature, 12 h. (iii) Allylmagnesium bromide, THF, -78°C , 2 h. (iv) OsO_4 , NaIO_4 , THF– H_2O , room temperature, 30 min. (v) Compound 22, $\text{NaBH}(\text{OAc})_3$, $\text{CH}_3\text{CO}_2\text{H}$, DCM. (vi) Compound 23, $\text{NaBH}(\text{OAc})_3$, $\text{CH}_3\text{CO}_2\text{H}$, DCM.




Figure 3. Crystal structure of intermediate 20.

The *in vivo* profiles of activity for compound 17 and pramipexole were assessed by the antagonist interaction studies with the known D3 antagonist SB-277011A⁹ and the D2 antagonist L-741,626²⁰ in both yawning and hypothermia assays. The selective D3 antagonist SB-277011A effectively inhibited the induction of yawning by both 0.1 and 0.32 mg/kg of pramipexole and 1 and 10 mg/kg of 17 (Figure 4B,C). In comparison, although the selective D2 antagonist L-741,626 failed to modulate the yawning effect induced by the lower doses of pramipexole

and 17 (0.1 and 1.0 mg/kg, respectively), a significant increase in the yawning effect was observed when L-741,626 was administered prior to the higher doses of pramipexole and 17 (0.32 and 10.0 mg/kg, respectively; Figure 4B,C). These data further confirm the agonist activity for pramipexole and compound 17 at the D3 receptor and suggest that D2 agonist effects may be observed with high doses of pramipexole (~ 1.0 mg/kg) and 17 (~ 10.0 mg/kg). In the hypothermia assay, although the D2 antagonist L-741,626 significantly reversed the hypothermia




Figure 4. Functional evaluations of the D3 and D2 activity of pramipexole and compounds **16** and **17** in yawning and hypothermia assays in rats. *, $P < 0.05$; **, $P < 0.005$; and ***, $P < 0.001$.

effect induced by 0.32 mg/kg of pramipexole, it did not have a significant effect on the modest hypothermia induced by 10 mg/kg of compound **17** (Figure 4E). Of note, both pramipexole at 0.32 mg/kg and compound **17** at 10 mg/kg produced similar modest decreases in core body temperature (approximately 0.75 °C) with pretreatment of L-741,626 at 1 mg/kg (Figure 4E). Taken together, these in vivo data show that **17** functions as a potent, highly selective, full efficacy agonist at the D3 receptor, with the onset of modest D2 agonist activity observed at 10 mg/kg for compound **17** and at 0.32 mg/kg for pramipexole. Pramipexole was one of the most selective D3 agonists identified to date⁷ and has been extensively used in in vitro and in vivo studies. The profiles of activity obtained in the current studies have shown that compound **17** functions as a full agonist at the D3 receptor at a dose of 0.01 mg/kg but has minimal D2 activity at a dose of 10.0 mg/kg, displaying not only higher potency but also much better selectivity than pramipexole.

To further assess the binding affinity, selectivity, and functional activity of compound **17**, we evaluated this compound in the Addiction Treatment Discovery Programs of the National Institute on Drug Abuse (NIDA) using cloned human dopamine receptors (Supporting Information). Compound **17** binds to human D3 receptor with a K_i value of 3.61 ± 0.53 nM and to human D2 receptor with a K_i value of >4000 nM. Furthermore, **17** shows no detectable binding to human D1 receptor at 10000 nM. Functional testing using a mitogenesis functional assay for the D3 receptor in CHO cells showed that **17** is a potent and full agonist with an EC_{50} value of 1.70 ± 0.53 nM and 103.4% of maximum stimulation of quinpirole as the standard D3 agonist control. Therefore, compound **17** is also a highly potent D3 full agonist in the functional assays using the cloned human D3 receptor and displays a >1000 -fold selectivity for D3 over D2

and D1 in the binding assays using cloned human dopamine receptors.

In summary, through exploration of the linker region of our previously reported D3 ligand **11**, we have identified a number of new D₃ ligands with high binding affinities to the D₃ receptor and a selectivity of >1000 times over the D₂ and D₁ receptors based upon our in vitro binding data. Compound **17** binds to D₃ with a K_i value of 0.50 nM and shows a selectivity of >5000 times over D₂ and >10000 over D₁ in our binding assays using rat brain. The high binding affinity and selectivity of **17** for the D₃ receptor were further confirmed using cells transfected with cloned human dopamine receptors. Our in vivo functional evaluation demonstrated that compound **17** is a highly potent and selective D₃ full agonist. While it is active at the D₃ receptor at doses as low as 0.01 mg/kg, it has a minimal activity at the D₂ receptor in the rat at doses as high as 10 mg/kg. Its full agonist profile at the D₃ receptor was also confirmed using cells transfected with cloned human D₃ receptor in the mitogenesis assay. To the best of our knowledge, compound **17** is the most active and selective D₃ full agonist reported to date. Furthermore, compound **17** also has good aqueous solubility. Taken together, our data show that compound **17** (CJ-1639) is an excellent pharmacological tool with which to elucidate the role of the D₃ receptor in different neurological conditions in animal models. Extensive in vivo studies are in progress to evaluate its therapeutic potential for the treatment of drug abuse and other neurological conditions in which the D₃ receptor may play a role, and the results will be reported in due course.

■ ASSOCIATED CONTENT

S Supporting Information. Experimental details of chemical synthesis, chemical data, X-ray crystallographic analysis, and in

vitro and in vivo characterizations of the ligands. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*Tel: 734-615-0362. Fax: 734-647-9647. E-mail: shaomeng@umich.edu.

Funding Sources

This work was supported by a grant from the NIDA, NIH (R01DA020669). Crystallographic studies were supported by a grant from NIDA under contract Y1-DA-1101. We are grateful to the Addiction Treatment Discovery Programs at the NIDA, NIH, for evaluation of CJ-1639 for its binding affinities and functional activity in cells transfected with cloned human dopamine receptors under the contract NIDA Y1-DA-0101, performed by Dr. Aaron Janowsky at the Oregon Health & Science University (Portland, Oregon).

REFERENCES

- Joyce, J. N. Dopamine D3 receptor as a therapeutic target for antipsychotic and antiparkinsonian drugs. *Pharmacol. Ther.* **2001**, *90*, 231–259.
- Crocq, M. A.; Mant, R.; Asherson, P.; Williams, J.; Hode, Y.; Shimohama, S.; Sawada, H.; Kitamura, Y.; Taniguchi, T. Disease model: Parkinson's disease. *Trends Mol. Med.* **2003**, *9*, 360–365.
- Volkow, N. D.; Fowler, J. S.; Wang, G. J. Role of dopamine in drug reinforcement and addiction in humans: Results from imaging studies. *Behav. Pharmacol.* **2002**, *13*, 355–366.
- Levant, B. The D3 dopamine receptor: Neurobiology and potential clinical relevance. *Pharmacol. Rev.* **1997**, *49*, 231–252.
- Newman, A. H.; Grundt, P.; Nader, M. A. Dopamine D3 receptor partial agonists and antagonists as potential drug abuse therapeutic agents. *J. Med. Chem.* **2005**, *48*, 3663–3679.
- Heidbreder, C. A.; Newman, A. H. Current perspectives on selective dopamine D(3) receptor antagonists as pharmacotherapeutics for addictions and related disorders. *Ann. N.Y. Acad. Sci.* **2010**, *1187*, 4–34.
- Collins, G. T.; Newman, A. H.; Grundt, P.; Rice, K. C.; Husbands, S. M.; Chauvignac, C.; Chen, J.; Wang, S.; Woods, J. H. Yawning and hypothermia in rats: Effects of dopamine D3 and D2 agonists and antagonists. *Psychopharmacology (Berlin)* **2007**, *193*, 159–170.
- Collins, G. T.; Witkin, J. M.; Newman, A. H.; Svensson, K. A.; Grundt, P.; Cao, J.; Woods, J. H. Dopamine Agonist-Induced Yawning in Rats: A Dopamine D3 Receptor-Mediated Behavior. *J. Pharmacol. Exp. Ther.* **2005**, *314*, 310–319.
- Reavill, C.; Taylor, S. G.; Wood, M. D.; Ashmeade, T.; Austin, N. E.; Avenell, K. Y.; Boyfield, I.; Branch, C. L.; Cilia, J.; Coldwell, M. C. Pharmacological actions of a novel, high-affinity, and selective human dopamine D3 receptor antagonist, SB-277011-A. *J. Pharmacol. Exp. Ther.* **2000**, *294*, 1154–1165.
- Macdonald, G. J.; Branch, C. L.; Hadley, M. S.; Johnson, C. N.; Nash, D. J.; Smith, A. B.; Stemp, G.; Thewlis, K. M.; Vong, A. K.; Austin, N. E.; Jeffrey, P.; Winborn, K. Y.; Boyfield, I.; Hagan, J. J.; Middlemiss, D. N.; Reavill, C.; Riley, G. J.; Watson, J. M.; Wood, M.; Parker, S. G.; Ashby, C. R. Design and synthesis of trans-3-(2-(4-((3-(3-(S-methyl-1,2,4-oxadiazolyl))-phenyl)carboxamido)cyclohexyl)ethyl)-7-methylsulfonyl-2,3,4,5-tetrahydro-1H-3-benzazepine (SB-414796): A potent and selective dopamine D3 receptor antagonist. *J. Med. Chem.* **2003**, *46*, 4952–4964.
- Yuan, J.; Chen, X.; Brodbeck, R.; Primus, R.; Braun, J.; Wasley, J. W. F.; Thurkauf, A. NGB 2904 and NGB 2849: Two highly selective dopamine D3 receptor antagonists. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 2715–2718.

- Pilla, M.; Perachon, S.; Sautel, F.; Garrido, F.; Mann, A.; Wermuth, C. G.; Schwartz, J. C.; Everitt, B. J.; Sokoloff, P. Selective inhibition of cocaine-seeking behaviour by a partial dopamine D3 receptor agonist. *Nature* **1999**, *400*, 371–375.

- Newman, A. H.; Grundt, P.; Cyriac, G.; Deschamps, J. R.; Taylor, M.; Kumar, R.; Ho, D.; Luedtke, R. R. N-(4-(4-(2,3-Dichloro- or 2-methoxyphenyl)piperazin-1-yl)butyl)heterobiarylcarboxamides with Functionalized Linking Chains as High Affinity and Enantioselective D3 Receptor Antagonists. *J. Med. Chem.* **2009**, *52*, 2559–2570.

- Micheli, F.; Arista, L.; Bonanomi, G.; Blaney, F. E.; Braggio, S.; Capelli, A. M.; Checchia, A.; Damiani, F.; Di-Fabio, R.; Fontana, S.; Gentile, G.; Griffante, C.; Hamprecht, D.; Marchioro, C.; Mugnaini, M.; Piner, J.; Ratti, E.; Tedesco, G.; Tarsi, L.; Terreni, S.; Worby, A.; Ashby, C. R.; Heidbreder, C. 1,2,4-Triazolyl azabicyclo[3.1.0]hexanes: A new series of potent and selective dopamine D(3) receptor antagonists. *J. Med. Chem.* **2010**, *53*, 374–391.

- Chen, J.; Collins, G. T.; Zhang, J.; Yang, C.-Y.; Levant, B.; Woods, J.; Wang, S. Design, synthesis and evaluation of potent and highly selective dopamine 3 receptor ligands with a novel profile of pharmacologic and behavioral activity. *J. Med. Chem.* **2008**, *51*, 5905–5908.

- Grundt, P.; Prevatt, K. M.; Cao, J.; Taylor, M.; Floresca, C. Z.; Choi, J.-K.; Jenkins, B. G.; Luedtke, R. R.; Newman, A. H. Heterocyclic Analogues of N-(4-(4-(2,3-Dichlorophenyl)piperazin-1-yl)butyl)arylcarboxamides with Functionalized Linking Chains as Novel Dopamine D3 Receptor Ligands: Potential Substance Abuse Therapeutic Agents. *J. Med. Chem.* **2007**, *50*, 4135–4146.

- Collins, G. T.; Calinski, D. M.; Newman, A. H.; Grundt, P.; Woods, J. H. Food restriction alters N'-propyl-4,5,6,7-tetrahydrobenzothiazole-2,6-diamine dihydrochloride (pramipexole)-induced yawning, hypothermia, and locomotor activity in rats: Evidence for sensitization of dopamine D2 receptor-mediated effects. *J. Pharmacol. Exp. Ther.* **2008**, *325*, 691–697.

- Boulay, D.; Depoortere, R.; Perrault, G.; Borrelli, E.; Sanger, D. J. Dopamine D2 receptor knock-out mice are insensitive to the hypolocomotor and hypothermic effects of dopamine D2/D3 receptor agonists. *Neuropharmacology* **1999**, *38*, 1389–1396.

- Chaperon, F.; Tricklebank, M. D.; Unger, L.; Neijt, H. C. Evidence for regulation of body temperature in rats by dopamine D2 receptor and possible influence of D1 but not D3 and D4 receptors. *Neuropharmacology* **2003**, *44*, 1047–1053.

- Kulagowski, J. J.; Broughton, H. B.; Curtis, N. R.; Mawer, I. M.; Ridgill, M. P.; Baker, R.; Emms, F.; Freedman, S. B.; Marwood, R.; Patel, S.; Ragan, C. I.; Leeson, P. D. 3-[[4-(4-Chlorophenyl)piperazin-1-yl]-methyl]-1H-pyrrolo[2,3-b]pyridine: An Antagonist with High Affinity and Selectivity for the Human Dopamine D4 Receptor. *J. Med. Chem.* **1996**, *39*, 1941–1942.

NOTE ADDED AFTER ASAP PUBLICATION

This paper was published on the Web on June 28, 2011. Details of X-ray crystallographic analysis were added to the Supporting Information and the corrected version was reposted on July 6, 2011.