

Discovery and Characterization of ML398, a Potent and Selective Antagonist of the D₄ Receptor with *in Vivo* Activity

Cynthia B. Berry,^{†,‡,§} Michael Bubser,^{‡,||} Carrie K. Jones,^{‡,||} John P. Hayes,^{†,‡,⊥} James A. Wepy,^{†,‡,⊥} Charles W. Locuson,^{‡,§,||} J. Scott Daniels,^{‡,§,||} Craig W. Lindsley,^{†,‡,§,||} and Corey R. Hopkins^{*,†,‡,§,||}

[†]Department of Chemistry, Vanderbilt University, Nashville, Tennessee 37232, United States


[‡]Department of Pharmacology, Vanderbilt University Medical Center, Nashville, Tennessee 37232, United States

[§]Vanderbilt Specialized Chemistry Center for Probe Development (MLPCN), Nashville, Tennessee 37232, United States

^{||}Vanderbilt Center for Neuroscience Drug Discovery, Vanderbilt University Medical Center, Nashville, Tennessee 37232, United States

S Supporting Information

ABSTRACT: Herein, we report the structure–activity relationship of a chiral morpholine-based scaffold, which led to the identification of a potent and selective dopamine 4 (D₄) receptor antagonist. The 4-chlorobenzyl moiety was identified, and the compound was designated an MLPCN probe molecule, ML398. ML398 is potent against the D₄ receptor with IC₅₀ = 130 nM and K_i = 36 nM and shows no activity against the other dopamine receptors tested (>20 μM against D₁, D_{2S}, D_{2L}, D₃, and D₅). Further *in vivo* studies showed that ML398 reversed cocaine-induced hyperlocomotion at 10 mg/kg.



ML398

D₄ IC₅₀ = 130 nM

D₄ K_i = 36 nM

Selectivity D₁, D_{2S}, D_{2L},

D₃, D₅ > 20 μM

B:P = 2.0

KEYWORDS: Dopamine 4 receptor antagonist, ML398, addiction, MLPCN

Dopamine receptors are members of the Class A G-protein coupled receptors (GPCRs) superfamily. GPCRs, also known as seven-transmembrane domain receptors (7TM receptors), are protein receptors that mediate most of the physiological responses to many hormones, neurotransmitters, etc., and constitute the largest class of drug targets. The dopamine receptors are further divided into five subtypes, included in two families. The D₁-like family of receptors contains D₁ and D₅, while the D₂-like family contains D₂, D₃, and D₄.^{1,2} The dopamine receptors are associated with numerous neurological processes including memory, learning, motivation, pleasure, and cognition; and as such, they are familiar drug targets.² Included in the list of disorders linked to dysfunction of dopaminergic signaling is schizophrenia,^{3–5} attention-deficit hyperactivity disorder (ADHD),^{6–8} Parkinson's disease,^{9,10} and drug^{2,11} and alcohol^{12,13} dependence. Recently there has been mounting evidence linking elevations in synaptic dopamine levels with the reinforcing effects of cocaine and hence its abuse potential.^{14–16}

Cocaine is a powerful stimulant made from the leaves of the coca plant and produces short-term euphoria and energy bursts. According to the National Survey on Drug Use and Health (NSDUH), there are ~2 M current cocaine users in the US with young adults (18–25 years old) representing the largest population of users. Unfortunately, there are no approved treatments for cocaine dependence.¹⁷ Cocaine does not directly bind to the D₁ and D₂ receptors, but rather binds to the dopamine transporter, thereby increasing synaptic levels of dopamine and its downstream effects on D₁ and D₂ receptors


that enables the cocaine reinforcement effects. The involvement of D₄ receptors as another potential target for cocaine reinforcement/dependence is due to its tissue distribution in the limbic and cortical brain regions implicated in cocaine addiction.^{3,18} In fact, the dopamine D₄ receptor has been coined the “adventure gene” due to the higher novelty seeking scores in individuals grouped by the long, polymorphic repeat region in exon III of the D4DR (L-D4DR) gene.^{19,20} Although the notion of the adventure gene has been questioned,^{21,22} further data has suggested that D4 has a role in severity of dependence.²³ While there have been numerous studies on the role of dopamine D4, the field has been hampered by the lack of selective D₄ receptor antagonists.² Herein, we report the discovery and characterization of a potent and selective dopamine D₄ receptor antagonist, ML398.

Recently we reported an enantioselective synthesis of a chiral morpholine analogue via an organocatalytic α -chlorination of aldehydes followed by cyclization to form the morpholine scaffold.²⁴ As part of the previous report, we synthesized a known dopamine D₄ antagonist, **1**, in order to confirm the stereochemistry of the active enantiomer (Figure 1).²⁵ Biological evaluation of **1** revealed the (*R*)-enantiomer was the active isomer with a D₄ IC₅₀ = 180 nM and K_i = 70 nM. (*R*)-**1** was inactive against dopamine D₁ and D₂ (>100 μM) and weakly active against D₃ (IC₅₀ = 46.2 μM and K_i = 15.7 μM).

Received: June 27, 2014

Accepted: July 9, 2014

Published: July 9, 2014



1


Receptor		(±)-1	(S)-1	(R)-1
D ₁	K _i	>100	>100	>100
	IC ₅₀	>100	>100	>100
D ₂	K _i	>100	>100	>100
	IC ₅₀	>100	>100	>100
D ₃	K _i	10.8	25.9	15.7
	IC ₅₀	31.8	76.4	46.2
D ₄	K _i	0.14	>100	0.07
	IC ₅₀	0.36	>100	0.18

Figure 1. Structure and activity of initial hit, (R)-1. K_i and IC₅₀ values are in μM.

The (S)-1 was inactive against D₄ (>100 μM), and the racemic-1 was less active (IC₅₀ = 360 nM and K_i = 140 nM). On the basis of the activity and binding of (R)-1, we embarked on a medicinal chemistry campaign to explore the structure–activity relationship (SAR) around the benzimidazole portion of the molecule.

In an effort to evaluate a number of compounds, the racemic phenethyl morpholine was synthesized and then rapidly analogued and assayed at D₄. Active compounds would then be assayed in enantiopure form. The racemic synthesis of the initial analogues for the SAR evaluation is outlined in Scheme 1

Scheme 1. Synthesis of Racemic Morpholine D₄ Receptor Antagonists^{a,24}




















^aReagents and conditions: (a) DL-proline (10 mol %), NCS, CH₂Cl₂, 0 °C; (b) NaBH₄, MeOH; 86%; (c) Tf₂O, lutidine, DCM, −78 °C; (d) BnNHCH₂CH₂OH, DCM, −78 °C → rt; 78%; (e) KOtBu, CH₃CN, −20 °C; 54%; (f) H₂, Pd/C, MeOH; (g) R-Br, K₂CO₃, CH₃CN, rt; (h) DIPEA, acid chloride, DMF, rt, 1 h; (i) Pd₂(dba)₃, XANTPHOS, Cs₂CO₃, aryl bromide, 1,4-dioxane, 100 °C, 18 h; (j) aryl isocyanate, THF, rt, 1 h; (k) DIPEA, sulfonyl chloride, DMF, 40 °C, 1 h

and is identical to the previous work, except for the organocatalyst.²⁴ The racemic synthesis utilizes DL-proline as the catalyst to obtain the racemic α-chloro aldehyde, 3. Next, the alcohol was activated (Tf₂O, lutidine) and displaced with the amino alcohol to yield 4. Next, the morpholine ring was formed via an intramolecular cyclization and then the benzyl protecting group was removed to yield the common

intermediate, 5. The benzylic analogues in Table 1 were synthesized by alkylation of the morpholine nitrogen (R-Br,

Table 1. SAR Evaluation of the N-Morpholine Substituent^a



Entry	R	D ₄ (% inh. @ 10 μM)
6a		94
6b		98
6c		97
6d		54
6e		88
6f		88
6g		74
6h		51
6i		23
6j		35
6k		13
6l		17
6m		5
6n		17
6o		5
6p		2

^aAll assays were performed on the human D₄ receptor.²⁶

K₂CO₃). The amide analogues were synthesized via the corresponding acid chloride (DIPEA, DMF, rt, 1 h), the direct aryl compounds were synthesized via a palladium catalyzed cross-coupling (Pd₂(dba)₃, XANTPHOS, Cs₂CO₃, and 1,4-dioxane), the urea analogues were made via the aryl isocyanate (THF, rt, 1 h), and the sulfonamides were made via reaction with the appropriate sulfonyl chloride (DIPEA, DMF, 40 °C, 1 h).

The first set of analogues were substituted benzyl analogues (6a–6h), which all showed varying degrees of dopamine D₄ receptor inhibition (51–98% at 10 μM) (Table 1). Interestingly, the pyridyl analogues (2-, 3-, and 4-pyridylmethyl) were all inactive (6i–6k). Introduction of an amide group (thus changing the linker group from a methylene to a carbonyl) was not tolerated and led to inactive compounds (6l). Deletion of the linker group (direct arylation of the morpholine) was also not tolerated (6m and 6n). Lastly, the

urea analogue (**6o**) and sulfonamides (**6p**) were also inactive, potentially suggesting the need for a flexible or rotatable bond between the morpholine and the right-hand substituent (See Supplemental Table 1 for full SAR).

Next, the active benzyl analogues were progressed into the IC₅₀ and K_i determinations, and clear SAR trends emerged (Table 2). First, ortho substituents were less tolerated than the

Table 2. IC₅₀ and K_i Evaluation of the Racemic and Enantiomerically Pure Benzyl Analogues

compd	D ₄ (% inh. @ 10 μM) ^a	IC ₅₀ (μM)	K _i (μM)
(±)- 6a	94	0.16	0.043
(R)- 6a	96	0.23	0.065
(±)- 6b	98	0.17	0.046
(R)- 6b	86	0.10	0.028
Selectivity Profile ^a : >20 μM against D ₁ , D _{2S} , and D ₅			
D _{2L} , IC ₅₀ = 16.5 μM; K _i = 5.5 μM			
D ₃ , IC ₅₀ = 8.17 μM; K _i = 2.77 μM			
(±)- 6c	97	0.29	0.081
(R)- 6c	96	0.13	0.036
Selectivity Profile ^a : >20 μM against D ₁ , D _{2S} , D _{2L} , D ₃ , and D ₅			
(±)- 6d	54	3.68	1.02
(±)- 6e	88	0.39	0.11
(±)- 6f	88	1.14	0.32
(±)- 6g	74	1.48	0.41
(±)- 6h	51	3.88	1.07

^aAll assays were performed on the human receptor.²⁶

meta- and *para*-substituents (even the small fluorine analogues) (**6d** and **6h**). Second, the *para*-fluoro (**6f**) and unsubstituted benzyl (**6g**) were less active. Lastly, the most potent compounds contained a *meta*- or *para*-trifluoromethoxybenzyl group (**6a**, **6e**), a *para*-methoxybenzyl group (**6b**), or a *para*-chlorobenzyl group (**6c**). On the basis of these results, both enantiomers of **6a–c** were synthesized using the previously published route.²⁴ Both the (R)- and (S)-enantiomers were evaluated, and as previously determined, the (S)-enantiomers were inactive. All three of the compounds tested were potent antagonists of the dopamine D₄ receptor with high binding affinities ((R)-**6a**, IC₅₀ = 230 nM, K_i = 65 nM; (R)-**6b**, IC₅₀ = 100 nM, K_i = 28 nM; (R)-**6c**, IC₅₀ = 130 nM, K_i = 36 nM). On the basis of these results, we further profiled (R)-**6b** and (R)-**6c** for their selectivity against the other dopamine receptors, and (R)-**6c** was inactive (>20 μM) against all of the dopamine receptors tested. Thus, on the basis of potency, binding, and selectivity profile, (R)-**6c** was declared an MLPCN probe (ML398).

ML398 was further profiled in a battery of Tier 1 *in vitro* DMPK assays (Table 3). The intrinsic clearance was assessed in hepatic microsomes (rat and human), and ML398 was shown to be unstable to oxidative metabolism and predicted to display high clearance in both species. In addition, using an equilibrium dialysis approach, the protein binding of ML398 was evaluated, and it was shown to have good free fraction in both species (6.1% in human and 3.9% in rat). ML398 was evaluated for its inhibition of the cytochrome P450 (CYP) enzymes using a cocktail approach in human liver microsomes as a screen for potential drug–drug interaction liability. ML398 displayed no significant activity against the panel of CYPs. ML398 was also profiled in an *in vivo* tissue distribution study (plasma and brain levels). Because of the predicted high clearance, ML398 was dosed at a single dose via intraperitoneal (IP) route of

Table 3. *In Vitro* and *In Vivo* PK Properties of ML398

	ML398	(R)-1
<i>In Vitro</i> PK Properties		
microsome predicted hepatic clearance (mL/min/kg)		
rat CL _{HEP}	67.5	65.1
human CL _{HEP}	15.7	17.9
plasma unbound fraction (F _u)		
human	0.061	0.012
rat	0.039	0.133
CYP inhibition (IC ₅₀ , μM)		
1A2	>30	13.1
2C9	>30	4.8
2D6	18.0	12.0
3A4	>30	15.6
plasma exposure in SD Rat		
10 mg/kg, intraperitoneal, 0.25 hr sample		
plasma (nM)	482	1935
brain (nM)	987	3558
Brain:Plasma	2.0	1.8

administration with a suspension formulation and then evaluated at a single time point. ML398 readily crosses the blood–brain barrier with a B/P ratio of ~2 and total brain concentrations of ~1 μM. Lastly, ML398 was tested using EuroFins Lead Profiling screen (radioligand binding assay panel of 68 GPCRs, ion channels, and transporters screened at 10 μM). ML398 was found to not significantly interact with 63 of the 68 assays performed (<50% binding at 10 μM) (See Supporting Information). ML398 did have activity against five targets (adrenergic, α₁A (77%); histamine, H₁ (93%); sigma, σ₁ (99%); dopamine transporter, DAT (72%); norepinephrine transporter, NET (68%).

Having identified a potent, selective, and brain penetrant D₄ antagonist, we wanted to test its ability to reverse hyperlocomotion induced by cocaine. It is believed that cocaine increases locomotor activity by increasing synaptic concentrations of dopamine by blocking dopamine reuptake or by enhancing the release of dopamine. This, in turn, increases stimulation of postsynaptic dopamine receptors. The increased locomotor activity can be blocked by selective dopamine antagonists as well as haloperidol.²⁷ The cocaine-induced hyperlocomotion assay is used to establish PK/PD relationship. The effects of both (R)-1 and ML398 were evaluated in this assay, and the results are shown in Figure 2. Cocaine was




Figure 2. Effects of ML398 and (R)-1 on reversing cocaine-induced hyperlocomotion in rats.

shown to significantly induce hyperactivity in rats, which is characterized by an increase in the number of beam breaks using the SmartFrame open field activity chambers. Both test compounds were dosed via IP administration at doses of 3 mg/kg and 10 mg/kg. Although (*R*)-1 showed a linear trend of reversal, the data was not significant. However, ML398 did show a statistically significant reversal of cocaine-induced hyperlocomotion at the highest dose tested (10 mg/kg).

In conclusion, we have identified a new, potent, and selective dopamine D₄ antagonist based on a chiral morpholine scaffold. As many of the previous D₄ antagonists contain a piperidine moiety, the reduced basicity of the morpholine may help contribute to the unprecedented selectivity. The SAR studies showed that the benzylic substitution is optimal as the amides, ureas, and arylamion analogues were all inactive. In addition, the (*R*)-enantiomer was confirmed as the active isomer within this series. ML398 is >100-fold selective versus the other dopamine receptors and is highly brain penetrant. ML398 also was shown to reverse cocaine-induced hyperlocomotion in rats. Further optimization studies are ongoing in an effort to discover an improved molecular probe for biological study as well and development of a radioligand for the D₄ receptor.

■ ASSOCIATED CONTENT

5 Supporting Information

General methods for the synthesis and characterization of all compounds. General methods for the *in vitro* and *in vivo* DMPK protocols and *in vivo* pharmacology. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*(C.R.H.) Phone: 1 615-936-6892. Fax: 1 615-936-4381. E-mail: corey.r.hopkins@vanderbilt.edu.

Author Contributions

[†]J.P.H. and J.A.W. contributed equally to this work. C.B.B., M.B., C.K.J., C.W.L., J.S.D., and C.R.H. drafted/corrected the manuscript. C.B.B., J.P.H., and J.A.W. performed the chemical synthesis. C.W.L. and C.R.H. oversaw the target selection and interpreted the biological data. C.W.L. and J.S.D. performed the *in vitro* DMPK experiments. M.B. and C.K.J. performed the *in vivo* experiments. All authors have given approval to the final version of the manuscript.

Funding

Vanderbilt is a member of the MLPCN and houses the Vanderbilt Specialized Chemistry Center for Accelerated Probe Development. This work was generously supported by the NIH/MLPCN Grant U54 MH084659 (to C.W.L.).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors would like to thank Mr. Frank Byers for technical assistance with the *in vivo* PK experiment and Mr. David Myers for help with the statistical evaluations.

■ ABBREVIATIONS

DIPEA, diisopropyl ethyl amine; DMF, dimethylformamide; THF, tetrahydrofuran; NCS, *N*-chlorosuccinimide; DCM, dichloromethane

■ REFERENCES

- (1) Girault, J.-A.; Greengard, P. The neurobiology of dopamine signaling. *Arch. Neurol.* **2004**, *61*, 641–644.
- (2) Ye, N.; Neumeyer, J. L.; Baldessarini, R. J.; Zhen, X.; Zhang, A. Update 1 of: Recent progress in development of dopamine receptor subtype-selective agents: potential therapeutics for neurological and psychiatric disorders. *Chem. Rev.* **2013**, *113*, PR123–PR178.
- (3) Lahti, R. A.; Roberts, R. C.; Cochrane, E. V.; Primus, R. J.; Gallager, D. W.; Conley, R. R.; Tamminga, C. A. Direct determination of dopamine D₄ receptors in normal and schizophrenic postmortem brain tissue: a [³H]NGD-94-1 study. *Mol. Psychiatry* **1998**, *3*, 528–533.
- (4) Sanyal, S.; Van Tol, H. H. M. Review the role of dopamine D₄ receptors in schizophrenia and antipsychotic action. *J. Psychiat. Res.* **1997**, *31*, 219–232.
- (5) Bristow, L. J.; Kramer, M. S.; Kulagowski, J.; Patel, S.; Ragan, C. I.; Seabrook, G. R. Schizophrenia and L-745,870, a novel dopamine D₄ receptor antagonist. *Trends Pharmacol. Sci.* **1997**, *18*, 186–188.
- (6) Woolley, M. L.; Waters, K. A.; Reavill, C.; Bull, S.; Lacroix, L. P.; Martyn, A. J.; Hutcheson, D. M.; Valerio, E.; Bate, S.; Jones, D. N. C.; Dawson, L. A. Selective dopamine D₄ receptor agonist (A-412997) improves cognitive performance and stimulates motor activity without influencing reward-related behaviour in rat. *Behav. Pharmacol.* **2008**, *19*, 765–776.
- (7) Tarazi, F. I.; Zhang, K.; Baldessarini, R. J. Dopamine D₄ receptors: beyond schizophrenia. *J. Recept. Signal Transduction Res.* **2004**, *24*, 131–147.
- (8) Rubinstein, M.; Cepeda, C.; Hurst, R. S.; Flores-Hernandez, J.; Ariano, M. A.; Falzone, T. L.; Kozell, L. B.; Meshul, C. K.; Bunzow, J. R.; Low, M. J.; Levine, M. S.; Grandy, D. K. Dopamine D₄ receptor-deficient mice display cortical hyperexcitability. *J. Neurosci.* **2001**, *21*, 3756–3763.
- (9) Huot, P.; Johnston, T. H.; Koprach, J. B.; Aman, A.; Fox, S. H.; Brotchie, J. M. L-745,870 reduces L-DOPA-induced dyskinesia in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-lesioned macaque model of Parkinson's disease. *J. Pharmacol. Exp. Ther.* **2012**, *342*, 576–585.
- (10) Marin, C.; Aguilar, E.; Rodriguez-Oroz, M. C.; Bartoszyk, G. D.; Obeso, J. A. Local administration of serizotan into the subthalamic nucleus attenuates levodopa-induced dyskinesias in 6-OHDA-lesioned rats. *Psychopharmacology* **2009**, *204*, 241–250.
- (11) Kotler, M.; Cohen, H.; Segman, R.; Gritsenko, I.; Nemanov, L.; Lerer, B.; Kramer, I.; Zer-Zion, M.; Kletzl, I.; Ebstein, R. P. Excess dopamine D₄ receptor (D₄DR) exon III seven repeat allele in opioid-dependent subjects. *Mol. Psychiatry* **1997**, *2*, 251–254.
- (12) George, S. R.; Cheng, R.; Nguyen, T.; Israel, Y.; O'Dowd, B. F. Polymorphisms of the D₄ dopamine receptor alleles in chronic alcoholism. *Biochem. Biophys. Res. Commun.* **1993**, *196*, 107–114.
- (13) Rubinstein, M.; Phillips, T. J.; Bunzow, J. R.; Falzone, T. L.; Dziejczapolski, G.; Zhang, G.; Fang, Y.; Larson, J. L.; McDougall, J. A.; Chester, J. A.; Saez, C.; Pugsley, T. A.; Gershanik, O.; Low, M. J.; Grandy, D. K. Mice lacking dopamine D₄ receptors are supersensitive to ethanol, cocaine, and methamphetamine. *Cell* **1997**, *90*, 991–1001.
- (14) Bergman, J.; Roof, R. A.; Furman, C. A.; Conroy, J. L.; Mello, N. K.; Sibley, D. R.; Skolnick, P. Modification of cocaine self-administration by buspirone (buspar): potential involvement of D₃ and D₄ dopamine receptors. *Int. J. Neuropsychopharm.* **2013**, *16*, 445–458.
- (15) Costanza, R. M.; Terry, P. The dopamine D₄ receptor antagonist L-745,870: effects in rats discriminating cocaine from saline. *Eur. J. Pharmacol.* **1998**, *345*, 129–132.
- (16) Caine, S. B.; Negus, S. S.; Mello, N. K.; Patel, S.; Bristow, L.; Kulagowski, J.; Vallone, D.; Saiardi, A.; Borrelli, E. Role of dopamine D₂-like receptors in cocaine self-administration: studies with D₂ receptor mutant mice and novel D₂ receptor antagonists. *J. Neurosci.* **2002**, *22*, 2977–2988.
- (17) What Is the Scope of Cocaine Use in the United States? <http://www.drugabuse.gov/publications/research-reports/cocaine/what-scope-cocaine-use-in-united-states>.

(18) Primus, R. J.; Thurkauf, A.; Xu, J.; Yevich, E.; McInerney, S.; Shaw, K.; Tallman, J. F.; Gallager, D. W. II. Localization and characterization of dopamine D₄ binding sites in rats and human brain by use of the novel, D₄ receptor-selective ligand [³H]NGD 94-1. *J. Pharmacol. Exp. Ther.* **1997**, *282*, 1020–1027.

(19) Ebstein, R. P.; Novick, O.; Umansky, R.; Priel, B.; Osher, Y.; Blaine, D.; Bennett, E. R.; Nemanov, L.; Katz, M.; Belmaker, R. H. Dopamine D4 receptor (D4DR) exon III polymorphism associated with the human personality trait of novelty seeking. *Nat. Genet.* **1996**, *12*, 78–80.

(20) Ebstein, R. P.; Herzog, S. Saga of an adventure gene: novelty seeking, substance abuse and the dopamine D4 receptor (D4DR) exon III repeat polymorphism. *Mol. Psychiatry* **1997**, *2*, 381–384.

(21) Baron, M. Mapping genes for personality: is the saga sagging? *Mol. Psychiatry* **1998**, *3*, 106–108.

(22) Lusher, J. M.; Chandler, C.; Ball, D. Dopamine D4 receptor gene (DRD4) is associated with novelty seeking (NS) and substance abuse: the saga continues. *Mol. Psychiatry* **2001**, *6*, 497–499.

(23) Lusher, J.; Ebersole, L.; Ball, D. Dopamine D4 receptor gene and severity of dependence. *Addict. Biol.* **2000**, *5*, 469–472.

(24) O'Reilly, M. C.; Lindsley, C. W. A general, enantioselective synthesis of protected morpholines and piperazines. *Org. Lett.* **2012**, *14*, 2910–2913.

(25) Showell, G. A.; Emms, F.; Marwood, R.; O'Connor, D.; Patel, S.; Leeson, P. D. Binding of 2,4-disubstituted morpholines at human D₄ dopamine receptors. *Bioorg. Med. Chem.* **1998**, *6*, 1–8.

(26) Eurofins Scientific. <http://www.eurofins.com/en.aspx>.

(27) O'Neill, M. F.; Shaw, G. S. Comparison of dopamine receptor antagonists on hyperlocomotion induced by cocaine, amphetamine, MK-801 and the dopamine D₁ agonist C-APB in mice. *Psychopharmacology* **1999**, *145*, 237–250.

■ NOTE ADDED AFTER ASAP PUBLICATION

This paper was published ASAP on July 15, 2014. The abstract graphic was corrected and the revised version was reposted on July 16, 2014.