

ACCOUNTS OF CHEMICAL RESEARCH®

APRIL, 1988

Registered in U.S. Patent and Trademark Office; Copyright 1988 by the American Chemical Society

Generation of Simple Enols in Solution

BRIAN CAPON,* BO-ZHANG GUO, FU CHIU KWOK, ARUP K. SIDDHANTA, and CÉSAR ZUCCO

Chemistry Department, University of Glasgow, Glasgow G12 8QQ, Scotland, and Chemistry Department, University of Hong Kong, Pokfulam Road, Hong Kong

Received July 15, 1987 (Revised Manuscript Received October 20, 1987)

Since the subject of simple enols was reviewed by Hart in 1979,¹ notable advances have been made by Rappoport² and by Kresge³ and their co-workers. We have also carried out investigation in this field,⁴⁻¹⁰ and the purpose of this Account is to describe the progress that we have made.

As the following quotations illustrate, many textbooks of organic chemistry tell their readers that vinyl alcohol and other simple enols are very unstable species:

"But all attempts to prepare vinyl alcohol result—like hydration of acetylene—in formation of acetaldehyde."¹¹

"Vinyl alcohol is an unstable compound that rearranges spontaneously to acetaldehyde."¹²

"Since long before the nature of tautomerization was understood, chemists have been fascinated with the challenge of isolating the simplest of the enols, vinyl alcohol; to date this feat has not been accomplished. The equilibrium heavily favors the aldehyde."¹³

Brian Capon left the University of Glasgow in 1982 to join the Scientific Staff of the Beilstein Institute, Frankfurt, BRD. In 1983 he assumed his present position as Head of the Chemistry Department, University of Hong Kong. He is the author of a previous Account (*Acc. Chem. Res.* 1981, 14, 306).

Bo-Zhang Guo was born in Indonesia. He obtained his B.Sc. degree at Jnan University, Canton, PRC, and has been working toward his Ph.D. degree at Hong Kong University since 1984.

Fu Chiu Kwok was born in Hong Kong and obtained a B.Sc. degree from Hong Kong University in 1983 and a Ph.D. degree in 1987. He works at present as a chemist in the Hong Kong Government Laboratory.

Arup K. Siddhanta obtained his B.Sc., M.Sc., and Ph.D. (1981) degrees from the University of Calcutta, India. From 1980 to 1983 he was a postdoctoral fellow at the University of Glasgow. After a further spell (1983–1985) of postdoctoral research at the University of Texas, he returned to India, where he now works as a Pool Scientist in the Central Salt and Marine Chemical Research Institute (CSIR Laboratory), Bhavnagar.

César Zucco was born in Brusque, Santa Catarina, Brazil. He obtained his B.Sc. degree from the Fundação Universidade Regional de Blumenau, Santa Catarina, Brazil, his M.Sc. degree from the Universidade Federal de Santa Catarina, Florianópolis, Santa Catarina, Brazil, and his Ph.D. degree from the University of Glasgow. He is presently Professor Adjunto in the Chemistry Department of the Universidade Federal de Santa Catarina, Santa Catarina, Brazil.

In the spring of 1979 it occurred independently to Hart¹ and to one of us⁴ that perhaps simple enols were not so unstable as generally thought and that by the use of the right techniques it might be possible to generate them in solution or even isolate them. In our own work we initially concentrated on vinyl alcohol, and the design of our first experiments was based on some earlier work of Lienhard and Wang.¹⁴ These workers concluded that the rate constants for the H₃O⁺-catalyzed ketonization of 1-hydroxycyclohexene and for the hydrolysis of 1-methoxycyclohexene are approximately the same. If this also applied to the ketonization of vinyl alcohol and the hydrolysis of methyl vinyl ether, then k_{H^+} for the former reaction should only be about 1 M⁻¹ s⁻¹ at 25 °C.¹⁵ This suggested to us that if a precursor could be found that generated vinyl alcohol with a rate constant several powers of 10 greater than this, it should be possible to obtain a metastable solution of the latter. In fact, it is now known^{7,9} that k_{H^+} for the ketonization

* To whom correspondence should be addressed at the University of Hong Kong.

(1) Hart, H. *Chem. Rev.* 1979, 79, 515.

(2) See: Nadler, E. B.; Rappoport, Z. *J. Am. Chem. Soc.* 1987, 109, 2112 and references cited therein.

(3) (a) See: Kresge, A. J. *CHEMTECH* 1986, 250. (b) Chiang, Y.; Hojatti, M.; Keefe, J. R.; Kresge, A. J.; Schepp, N. P.; Wirz, J. *J. Am. Chem. Soc.* 1987, 109, 4000 and references cited therein. (c) Chiang, Y.; Kresge, A. J.; Walsh, P. A. *J. Am. Chem. Soc.* 1986, 108, 6314.

(4) Capon, B.; Rycroft, D. S.; Watson, T. W. *J. Chem. Soc., Chem. Commun.* 1979, 724.

(5) Capon, B.; Rycroft, D. S.; Watson, T. W.; Zucco, C. *J. Am. Chem. Soc.* 1981, 103, 1761.

(6) Capon, B.; Siddhanta, A. K. *Tetrahedron Lett.* 1982, 23, 3199.

(7) Capon, B.; Zucco, C. *J. Am. Chem. Soc.* 1982, 104, 7567.

(8) Capon, B.; Siddhanta, A. K. *J. Org. Chem.* 1984, 49, 255.


(9) Capon, B.; Siddhanta, A. K.; Zucco, C. *J. Org. Chem.* 1985, 50, 3580.

(10) Capon, B.; Kwok, F. C. *Tetrahedron Lett.* 1986, 27, 3275.

(11) Morrison, R. T.; Boyd, R. W. *Organic Chemistry*, 5th ed.; Allyn and Bacon: Boston, 1987; p 426.

(12) Solomons, T. W. G. *Organic Chemistry*, 2nd ed.; Wiley: New York, 1980; p 336.


(13) Allinger, N. L.; Cava, M. P.; De Jongh, D. C.; Johnson, C. R.; LeBel, A. N.; Stevens, C. L. *Organic Chemistry*, 2nd ed.; Worth Publishers: New York, 1976; p 174.



of enols in aqueous solution is $10\text{--}10^2$ times greater than that for the hydrolysis of the corresponding enol ethers but nevertheless as described below, vinyl alcohol and some other simple enols are still sufficiently unreactive for this argument to apply.

Generation of Simple Enols in Solution

An examination of previous work¹⁶ showed that the hydronium ion catalyzed hydration of ketene acetals (e.g., 1) frequently has rate constants ca. $10^6 \text{ M}^{-1} \text{ s}^{-1}$ and



that the hydronium ion catalyzed hydrolyses of ortho esters (e.g., 2 and 3) frequently have rate constants $10^3\text{--}10^4 \text{ M}^{-1} \text{ s}^{-1}$ at 25°C . A ketene acetal or ortho ester might therefore be a suitable precursor for the generation of vinyl alcohol in solution.

The first precursor to be chosen was therefore methoxy(vinyloxy)methyl chloroacetate (4), a particularly reactive orthoformate derivative. It was thought that this ester might be hydrolyzed in D_2O as shown in Scheme I to give O-deuteriated vinyl alcohol. If in this sequence of reactions steps 1 and 2 were faster than the conversion of O-deuteriated vinyl alcohol into deuteriated acetaldehyde (step 3), then a solution of O-deuteriated vinyl alcohol would be obtained. This is in fact what happens. In CD_3CN at -20°C the signals of the vinyl group of 4 are approximately a first-order ABC system with $\delta_A = 6.53$, $\delta_B = 4.58$, and $\delta_C = 4.32$ and $J_{AB} = 14.7$, $J_{AC} = 4.9$, and $J_{BC} = 1.7$ Hz. On addition of D_2O (10% (v/v)), the spectrum of 4 slowly disappears and after about 45 min at -20°C is replaced by the spectra of chloroacetic acid, methyl formate, and a new species with a vinyl group whose ^1H NMR spectrum is again approximately a first-order ABC system with $\delta_A = 6.45$, $\delta_B = 4.18$, and $\delta_C = 3.82$ and $J_{AB} = 14.2$, $J_{AC} = 6.4$ Hz, and $J_{BC} = 1$ Hz. Thus in this new species the signal of the α -proton (H_A) has almost the same chemical shift as that of the precursor 4 but the signals of the β -protons (H_B and H_C) are moved upfield by 0.4–0.5 ppm. The new species was stable in this acidic solution at -20°C for several hours but if sodium deuterioxide was added or if the temperature was raised to $+20^\circ \text{C}$, it was rapidly converted into CH_2DCHO .

(14) Lienhard, G. E.; Wang, T. C. *J. Am. Chem. Soc.* **1969**, *91*, 1146.

(15) Kresge, A. J.; Sagatys, D. S.; Chen, H. L. *J. Am. Chem. Soc.* **1977**, *99*, 7228.

(16) Kankaanpera, A.; Tuominen, H. *Suom. Kemistil. B* **1967**, *40*, 271.





Figure 1. ^1H NMR spectrum of vinyl alcohol under conditions of slow exchange.



The same species could be generated from six other precursors (5–10; Chart I). The best of these are 7 and 10, which may be regarded as vinyl alcohol with a very labile ortho ester (7) or ketene acetal (10) protecting group. These can be removed under very mild acidic conditions (see also ref 3b), and proof that the new vinylic species is in fact vinyl alcohol was obtained when 10 was allowed to react in CD_3COCD_3 (99% (v/v)) and H_2O (1% (v/v)) which contained 10^{-4} M HCl . Under these conditions the spectrum shown in Figure 1 was obtained. This spectrum of O–H vinyl alcohol differs from that of O–D vinyl alcohol in that now there is a signal of the OH group at $\delta = 7.97$ and the signal of the α -proton of the vinyl group is a doublet quartet. The coupling between the OH and α -CH is 9.98 Hz. Working with a solvent that contains a little more water enables the transformation from fast to slow exchange to be observed by changing the temperature (cf. Figure 2).

This method provides a good, and possibly the best, way of preparing solutions of vinyl alcohol. The precursors 7 and 10 are made quite easily (eq 1 and 2; Scheme II), and with a little care it is possible to convert them into vinyl alcohol without any detectable amounts of acetaldehyde being present. Attempts to isolate vinyl alcohol from such solutions and to achieve the goal set by Allinger et al.¹³ have, however, so far been unsuccessful. The purest sample that has been obtained up to now was isolated by low-temperature trap-to-trap distillation of a solution of vinyl alcohol prepared from 10 in slightly aqueous acetone and was approximately 70 mol % pure (^1H NMR). The major impurities were acetone and methyl acetate.¹⁷

The use of ortho ester of ketene acetal precursors to generate simple enols is of fairly general application and

(17) Zucco, C., unpublished observations.







Figure 2. Temperature variation of the OH and C_{α} -H signals of the ^1H NMR spectrum of vinyl alcohol in CD_3COCD_3 (94.5%)– H_2O (5.5%). Note the presence of small amounts of acetaldehyde ($\delta = 9.71$) and vinyl acetate ($\delta = 7.27$).




has been used to generate enols 11–18 (Chart II). In addition to acyclic orthoacetates, cyclic orthoacetates such as 20 and 21 can be used. These esters have the advantage of being more reactive than their acyclic analogues.¹⁸ They are prepared very easily⁹ by adding a chloro or β,γ -unsaturated alcohol to 2-methylene-

(18) Chiang, Y.; Kresge, A. J.; Salmona, P.; Young, C. I. *J. Am. Chem. Soc.* 1974, 96, 4494.



1,3-dioxolane (19) followed by elimination of HCl or migration of a double bond (see eq 3 and 4, Scheme III). This class of precursor could probably be used for the generation of other enols. With all of the enols studied except the acetone enol 14, the precursor can be converted to the enol without detectable (by ^1H NMR) amounts of the keto form. To generate acetone enol 14 the best precursor is 22, but even this reacts too slowly in CD_3COCD_3 – D_2O (8:2 (v/v)) to form 100% 14 before some of the latter is converted into $[^2\text{H}_1]$ acetone.



The other limitation of ketene acetal and ortho ester precursors is that for more complex enols their synthesis is laborious. For this reason we examined the possibility of using trimethylsilyl derivatives. The rate constant for the H_3O^+ -catalyzed hydrolysis of (vinyloxy)trimethylsilane (23) was determined by Tidwell and co-workers to be $50.4 \text{ M}^{-1} \text{ s}^{-1}$ in 50% H_2O –acetonitrile at 25°C .¹⁹ This compares with a value of $45.9 \text{ M}^{-1} \text{ s}^{-1}$ for $k_{\text{H}_3\text{O}^+}$ for the ketonization of vinyl alcohol in H_2O at 25°C .⁷ These data suggest that it is unlikely that 23 would be a suitable precursor for vinyl alcohol, and in our hands this has proved to be so. However, we have used trimethylsilyl ethers to generate a variety of more stable enols such as the heterocyclic enols 24–27,^{20,21} the homocyclic enols 28–32,²¹ 2-methylprop-1-en-1-ol (13),²² (*E*)- and (*Z*)-1-hydroxybuta-1,3-diene (33 and 34),²² and enols 35, 36, and 37 (Chart III), in which the double


(19) Novice, M. H.; Seikaly, H. R.; Seiz, A. D.; Tidwell, T. T. *J. Am. Chem. Soc.* 1980, 102, 5835.

(20) Kwok, F. C. Ph.D. Thesis, University of Hong Kong, 1987.

(21) Capon, B.; Guo, B. Z.; Kwok, F. C.; Wu, Z. P. *Pure Appl. Chem.* 1987, 59, 1577.

(22) Guo, B. Z., unpublished observations.

Chart III




bond is conjugated with a cyclopropane ring or a phenyl group.²²

In a typical experiment^{10,20} 3-((trimethylsilyl)oxy)-benzofuran was allowed to hydrolyze at 30 °C in a mixture of acetone-*d*₆ and D₂O (9:1 (v/v)) that contained DCl (5 × 10⁻⁴ M). Removal of the trimethylsilyl group could be followed by the disappearance of the signal at $\delta = 0.27$ (s) and formation of the signal at $\delta = 0.05$ due to trimethylsilanol. There was very little change in the rest of the spectrum except that the singlet of H-2 moved upfield from $\delta = 7.45$ to $\delta = 7.3$. The 3-hydroxybenzofuran was stable for several hours under these conditions but on addition of more acid, the spectrum changed rapidly to that of 2-deuterio-benzofuranone. This method, with slight variations, was used for the other enols listed above.

Stereochemistry of Enols

As described above the HO-C _{α} H coupling constant for vinyl alcohol is 9.98 Hz. This value decreases slightly with increasing temperature in the range -80 to +40 °C. It was thought that, by analogy to what is found with HC-CH bonds, this coupling constant should be related to the dihedral angle between the OH and C _{α} H bonds. Two planar conformations are possible for vinyl alcohol, *s*-cis (38) and *s*-trans (39), for which



the dihedral angles are 180° and 0°, respectively. There are also an infinite number of nonplanar gauche conformations possible. When a methyl group is introduced cis to the hydroxy group of vinyl alcohol as in (*Z*)-prop-1-en-1-ol (11, L = H) and 2-methylprop-1-en-1-ol (13, L = H), $J(\text{OH}-\text{C}_{\alpha}\text{H}) < 6$ Hz and increases with increasing temperature. With (*E*)-prop-1-enol (12, L

Table I
Equilibrium Constants for Enolization in Water at 25 °C

equilibrium	$K_{\text{enol}}(\text{E}/\text{K})$	ref
1. acetaldehyde = vinyl alcohol ^a	2.7 × 10 ⁻⁷ or 3.6 × 10 ⁻⁷	7
2. acetone = 2-hydroxypropene	6.0 × 10 ⁻⁹	26 ^b
3. isobutyraldehyde = 1-hydroxy-2-methylpropene	1.37 × 10 ⁻⁴	3c ^c
4. acetophenone = 1-hydroxystyrene	1.25 × 10 ⁻⁸	27 ^b
5. 1-indanone = 3-indenol	1.85 × 10 ⁻⁸	20
6. 2-indanone = 2-indenol	8.19 × 10 ⁻⁵	20
7. cyclohepta-3,5-dien-1-one = 1-hydroxycycloheptatriene	1.19 × 10 ⁻³	22

^a 15 °C. The different values are derived from different literature values for the rate constant for the enolization of acetaldehyde (see ref 7). ^b The enol was generated photochemically. ^c The enol was generated by protonation of the enolate ion.

= H), however, this coupling constant is >9 Hz. These results are interpreted as indicating that for vinyl alcohol and 12, which have hydrogen cis to the hydroxyl, the predominant conformation is *s*-cis but with enols 11 and 13, which have a methyl group cis to the hydroxyl, the predominant conformation is *s*-trans. As the temperature is raised, the higher energy conformation becomes more highly populated so the coupling constants of vinyl alcohol and 12 decrease whereas those of 11 and 13 increase.³ The long-range coupling constant between the OH and the trans β -CH, which for vinyl alcohol is 0.4 Hz and for (*Z*)-prop-1-en-1-ol is 0.95 Hz, also supports this conclusion. In the *s*-trans conformation (40) proposed for the latter, there is a *W* relationship between the OH and the trans C β -H, which is the most favorable for long-range allylic coupling and hence should lead to a larger coupling constant.

These conclusions are in agreement with MO calculations of Bouma and Radom, which showed the *s*-cis conformation of vinyl alcohol to be 2.2 kcal mol⁻¹ more stable than the *s*-trans conformation and which indicated "no evidence" for a nonplanar conformation.²³

Thermodynamic Stability of Enols

As discussed by Toullec,²⁴ many of the earlier measurements of equilibrium constants for enol formation, $K_{\text{enol}}(\text{E}/\text{K})$, based on halogen titration, are too high, although the estimate ($K_{\text{enol}} = 10^{-8}$) made by Gutsche and Pasto on the basis of bond energy considerations is approximately correct.²⁵ The generation of simple enols as described above allows their ketonization rate constants to be determined, and as it is usually possible to determine the rate constants for enolization of the keto form by halogen trapping or deuterium exchange, the equilibrium constants can be determined as the ratio of the rate constants of the forward and backward reactions (see Table I).

When the carbon-carbon double bond of the enol is conjugated with a benzene ring but the carbon-oxygen double bond of the keto form is not, as expected, the enol form is stabilized (compare Table I, entry 6, with Table I, entry 5). Conjugation with additional car-

(23) Bouma, W. J.; Radom, L. *J. Mol. Struct.* 1978, 43, 267.

(24) Toullec, J. *Adv. Phys. Org. Chem.* 1982, 18, 43.

(25) Gutsche, C. D.; Pasto, D. J. *Fundamentals of Organic Chemistry*; Prentice-Hall: Englewood Cliffs, NJ, 1975; p 253.


(26) Chiang, Y.; Kresge, A. J.; Tang, Y. S.; Wirz, J. *J. Am. Chem. Soc.* 1984, 106, 460.

(27) Chiang, Y.; Kresge, A. J.; Wirz, J. *J. Am. Chem. Soc.* 1984, 106, 6392.

Table II
Equilibrium Constants for the Enolization of Some
Heterocyclic Compounds in Water at 25 °C

equilibrium	$K_{\text{enol}}(\text{E}/\text{K})$
1. 3-ketobenzofuran \rightleftharpoons 3-hydroxybenzofuran	8.70×10^{-4}
2. 3-ketobenzothiophene \rightleftharpoons 3-hydroxybenzothiophene	6.7
3. 3-ketoindole \rightleftharpoons 3-hydroxyindole	0.086
4. 3-keto-1-methylindole \rightleftharpoons 3-hydroxy-1-methylindole	0.303
5. 3-ketofuran \rightleftharpoons 3-hydroxyfuran	$<10^{-2}$
6. 3-ketothiophene \rightleftharpoons 3-hydroxythiophene	2.96
7. 3-ketopyrrole \rightleftharpoons 3-hydroxypyrrrole	0.13
8. 3-keto-1-methylpyrrole \rightleftharpoons 3-hydroxy-1-methylpyrrole	0.18

bon-carbon double bonds also stabilizes the enol form, although whether any of the high stability of 1-hydroxycycloheptatriene can be ascribed to homoaromaticity is not clear. The introduction of β -methyl groups (see Table I, entries 1 and 3) also stabilizes the enol. On the other hand, the introduction of an α -methyl or α -phenyl substituent, to give the enol of a ketone, causes a decrease in K_{enol} (see Table I, entries 1, 2, and 4). This result presumably arises from a stabilization of the keto form, as an α -methyl or α -phenyl substituent should stabilize the mesomeric zwitterionic structure (see 41 \leftrightarrow 42).



The heterocyclic enols 24 are all more stable than their homocyclic analogue 3-indenol (cf. Table I, entry 5, with Table II, entries 1–4). Even 3-hydroxybenzofuran is much more stable than 3-indenol despite the low aromaticity of the five-membered ring of benzofuran.²⁸ When the heterocyclic ring is more aromatic, as with pyrrole, thiophene, and their benzo derivatives, the 3-hydroxy tautomers are sufficiently stable for there to be detectable amounts of them present at equilibrium.

Heterocyclic enols with the hydroxy group in the 2-position could only be generated for the thiophene and benzothiophene series (27, 26). These are formally enols of thioesters and while the enolic form is stabilized by the aromaticity of the heterocyclic ring, the keto form is stabilized by the thioester resonance.²⁹ The result is that 2-hydroxybenzothiophene, with $K_{\text{enol}} = 2.29 \times 10^{-5}$ in 90% CH_3CN –10% H_2O (v/v) (25 °C), is much less stable than 3-hydroxybenzothiophene, for which $K_{\text{enol}} = 0.31$, measured under the same conditions, and is only slightly more stable than its homocyclic analogue 2-hydroxyindene, for which $K_{\text{enol}} = 3.42 \times 10^{-6}$.

Kinetic Stability of Simple Enols

The pH-rate profile for the ketonization of vinyl alcohol is a U-shaped curve with H_3O^+ , HO^- , and H_2O -catalyzed reactions. In the pH range 3.5–5.0, the H_2O -catalyzed reaction predominates and the rate of ketonization is almost constant, with $k = (1.5\text{--}2.0) \times 10^{-2} \text{ s}^{-1}$ at 15 °C. This corresponds to a half-life of about 35–45 s. At pHs lower than 3.5 and higher than 5.0, the

(28) Dewar, M. J. S.; Harget, A. J.; Trinajstić, N.; Worley, S. D. *Tetrahedron* 1970, 26, 4505.

(29) The delocalization energies of simple thioesters have been estimated to be ca. 11 kcal mol⁻¹. Pastrez, J. J. *Am. Chem. Soc.* 1977, 99, 7004.

Table III
Rate Constants for the Ketonization of Simple Enols in
Water ($I = 1.0 \text{ M}$) at 15 °C

	$k_{\text{H}^+}/$ ($\text{M}^{-1} \text{ s}^{-1}$)	$k_{\text{HO}^-}/$ ($\text{M}^{-1} \text{ s}^{-1}$)	$k_{\text{H}_2\text{O}}/\text{s}^{-1}$
1. vinyl alcohol	20.2	1.5×10^{17} ^a	1.38×10^{-2}
2. (Z)-propenol	2.80	5.03×10^4	
3. 2-methylprop-1-en-1-ol	0.370	1.38×10^3	
4. 3-chloro-2-hydroxypropene	6.18	4.74×10^7	6.0×10^{-2}
5. hydroxypropadiene	5.67	1.11×10^9	7.61×10^{-3}

^a A value of $1.99 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ for k_{HO^-} can be calculated for the results given in ref 3b ($I = 0.1 \text{ M}$, 25 °C).


rate increases as the H_3O^+ - and HO^- -catalyzed reactions become more important.⁷ The rate constants for these reactions and those for the ketonization of a number of other enols measured by ourselves and by Kresge and co-workers are summarized in Tables III–V.

As might be predicted from their expected effects on the stability of cationic and anionic transition states, an α -methyl group causes a large rate enhancement of the hydronium ion catalyzed ketonization but has only a small effect on the hydroxide ion catalyzed ketonization (cf. Table IV, entry 2, with Table IV, entry 1, and Table III, entry 1), whereas a β -methyl substituent has a much larger rate-retarding effect on the HO^- -catalyzed ketonization than on the H_3O^+ -catalyzed ketonization (cf. Table III, entries 2 and 3, with Table III, entry 1). β -Methyl substituents also cause a decrease in the rate of the H_2O -catalyzed reaction so that the half-life of (Z)-propenol at the minimum of the pH-rate profile (pH 3.5–6) at 15 °C is approximately 10 min and that of 2-methylprop-1-en-1-ol is approximately 1 h (pH 3–5.5).

The introduction of a chloro substituent into the methyl group of 2-hydroxypropene has a large rate-decreasing effect on the H_3O^+ -catalyzed ketonization but only a small effect on the HO^- -catalyzed ketonization (cf. Table IV, entry 2, with Table III, entry 4). On the other hand, the HO^- -catalyzed ketonization of hydroxypropadiene is much faster than that of vinyl alcohol (see Table III), which can be explained on the basis of the vinyl anion character of the transition state of the former reaction.⁸

Groups that are conjugated with the carbon-carbon double bond of the enol but not with the carbon-oxygen double bond of the keto tautomer of course stabilize the enol kinetically. This is illustrated by entries 6–10 of Table IV.

The monocyclic heterocyclic enols (25) undergo H_3O^+ -catalyzed ketonization faster than their benzo analogues (24) (see Table V), which can be explained by a mesomeric effect as symbolized by 43, which is



particularly effective when $\text{X} = \text{NH}$ or NMe . 3-Hydroxybenzofuran and 3-hydroxybenzothiophene undergo hydroxide ion catalyzed ketonization much faster than the 3-hydroxyindoles. This result can be explained partly as arising from the smaller $\text{p}K_{\text{a}}$'s of the two former compounds (see Table V), which result in a higher concentration of the anion, which is presumably the reactive species.

Table IV
Rate Constants for the Ketonization of Simple Enols in Water ($I = 1.0 \text{ M}$) at 25°C

	$k_{\text{H}^+}/(\text{M}^{-1} \text{ s}^{-1})$	$k_{\text{HO}^-}/(\text{M}^{-1} \text{ s}^{-1})$	$k_{\text{H}_2\text{O}}/\text{s}^{-1}$	ref
1. vinyl alcohol	45.9			7
2. 2-hydroxypropene ^a	4.7×10^3 ^b	3.43×10^7		26
3. 1-hydroxystyrene ^a	1250	2.07×10^7	1.9	27
4. 2-hydroxyindene	3.48		5.5×10^{-2}	20
5. 3-hydroxyindene	903 ^c			20
6. (<i>Z</i>)-1-hydroxybuta-1,3-diene ^c	9.55×10^{-2}	1.28×10^5	3.99×10^{-3}	22
7. (<i>E</i>)-1-hydroxybuta-1,3-diene ^c	0.53	1.14×10^5	3.03×10^{-3}	22
8. (<i>Z</i>)-2-cyclopropylvinyl alcohol	1.25	4.48×10^4	2.54×10^{-3}	22
9. (<i>Z</i>)-2-(2,2-dichlorocyclopropyl)vinyl alcohol	0.36	1.79×10^6	6.29×10^{-3}	22
10. (<i>Z</i>)-2-hydroxystyrene	0.16	2.59×10^6	6.7×10^{-3}	22

^a $I = 0.1 \text{ M}$. ^bEstimated from k_{H^+} for the enolization of acetone, $2.8 \times 10^{-5} \text{ M}^{-1} \text{ s}^{-1}$ (Albery, W. J.; Gelles, J. S. *J. Chem. Soc., Faraday Trans. 1982*, **78**, 1569) and $K_{\text{enol}} = 6 \times 10^{-9}$ (ref 26). The value agrees well with a recent direct measurement, $5.38 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ (ref 3b). ^cExtrapolated from values in acetonitrile-water mixtures. ^dTo yield 2-butenal and 3-butenal.

Table V
Rate Constants for the Ketonization of Heterocyclic Enols in Water ($I = 1.0 \text{ M}$) at 25.0°C ^a

	$k_{\text{H}^+}/(\text{M}^{-1} \text{ s}^{-1})$	$10^{-5}k_{\text{HO}^-}/(\text{M}^{-1} \text{ s}^{-1})$	$10^3k_{\text{H}_2\text{O}}/\text{s}^{-1}$	$\text{p}K_{\text{a}}$
1. 3-hydroxybenzofuran	0.592	4.05	6.41	9.16
2. 3-hydroxybenzothiophene	0.579	1.14	5.59	8.86
3. 3-hydroxyindole	3.43	0.00994	1.05	10.46
4. 3-hydroxy-1-methylindole	5.82	0.134	1.00	10.76
5. 3-hydroxyfuran	50.1	27.9	8.60	
6. 3-hydroxythiophene	5.83	0.187	2.21	
7. 3-hydroxypyrrole	2.49×10^4	0.346	17.0	
8. 3-hydroxy-1-methylpyrrole	1.17×10^4	0.154	8.87	

^aReferences 10, 20, and 21.

2-Hydroxythiophene ($k_{\text{H}^+} = 11.5 \text{ M}^{-1} \text{ s}^{-1}$) undergoes acid-catalyzed ketonization faster than 3-hydroxythiophene ($k_{\text{H}^+} = 1.78$) in 50% CH_3CN -50% H_2O (v/v) at 25°C , and 2-hydroxybenzothiophene ($k_{\text{H}^+} = 12.4 \text{ M}^{-1} \text{ s}^{-1}$) undergoes acid-catalyzed ketonization faster than 3-hydroxybenzothiophene ($k_{\text{H}^+} = 0.314$) in 90% CD_3CN -10% H_2O (v/v) at 25°C . The thermodynamic instability of these 2-hydroxy heterocyclic compounds (see above), which are formally enols at the carboxylic acid level of oxidation, is therefore paralleled by kinetic instability. 2-Hydroxybenzothiophene, which has its hydroxyl group attached to the five-membered aromatic ring, is also kinetically less stable than its homocyclic analogue 2-hydroxyindene ($k_{\text{H}^+} = 2.79 \text{ M}^{-1} \text{ s}^{-1}$ in 90% CH_3CN -10% H_2O). This is in complete contrast to

what is found in the 3-hydroxy series.

Conclusion

The results described in this Account show that many simple enols are species that, if generated in the right way, can be investigated by the standard spectroscopic techniques of organic chemistry. They, like the tetrahedral intermediates of *O,O*-acyl-transfer reactions,³⁰ were once hypothetical species, but now the door is open for a full and detailed investigation of their properties.

(30) See: Capon, B.; Ghosh, A. K.; Grieve, D. McL. A. *Acc. Chem. Res.* **1981**, *14*, 306. Capon, B.; Dosunmu, M. I.; Sanchez, M. de N. de M. *Adv. Phys. Org. Chem.* **1985**, *21*, 37. McClelland, R. A.; Santry, L. J. *Acc. Chem. Res.* **1983**, *16*, 394.