

ACCOUNTS OF CHEMICAL RESEARCH®

JANUARY 1989

Registered in U.S. Patent and Trademark Office; Copyright 1989 by the American Chemical Society

Free Radical Chain Reactions Involving Alkyl- and Alkenylmercurials

GLEN A. RUSSELL

Department of Chemistry, Iowa State University, Ames, Iowa 50011

Received July 28, 1987 (Revised Manuscript Received July 29, 1988)

Alkylmercurials are readily available organometallic reagents possessing moderate reactivity in electrophilic substitution and low reactivity in nucleophilic attack at carbon.¹ Pyrolysis of organomercurials has been utilized as a method to generate alkyl radicals useful in homolytic aromatic substitution processes² and in defining a set of σ -constants for substituent stabilization of benzylic radicals.³ One of the most widely studied chain reactions of an organomercurial has been the alkaline NaBH_4 reduction of alkylmercury halides or carboxylates to yield the alkane⁴ or products derived from alkyl radical attack upon a suitable coreactant.⁵ Both the pyrolysis of R_2Hg and the alkaline NaBH_4 reduction of RHgCl are reactions that involve unstable mercury(I) intermediates.

During the past several years, we have developed a series of free radical reactions in which RHgX or R_2Hg participates in the propagation step of a chain process that does not usually involve RHg^+ as an intermediate other than in the initiation step.⁶⁻²⁶ One group of these reactions involves the homolytic displacement of an alkyl radical from a mercury atom by an electron-accepting carbon- or heteroatom-centered radical. This type of homolytic displacement is not unexpected since it has been recognized for some time in the cleavage of R_2Hg or RHgX by molecular halogens.²⁷ Competing electrophilic substitution by halogen can occur, but it has been possible to define conditions where either the free radical or ionic substitution will predominate.²⁸ Another group of reactions involves electron transfer

to RHgX from an electron-donating free radical or radical ion and leads directly to the alkyl radical, mercury metal, and X^- .⁶ The unusual reactivity of carbon-mercury bonds toward both electron-accepting and electron-donating paramagnetic species is the focus

- (1) Larock, R. C. *Organomercury Compounds in Organic Synthesis*; Springer-Verlag: Heidelberg and New York, 1985.
- (2) Bass, K. C.; Nababsing, P. *J. Chem. Soc. C* **1966**, 1184, 2019.
- (3) Dinctürk, S.; Jackson, R. A. *J. Chem. Soc., Chem. Commun.* **1979**, 172.
- (4) Whitesides, G. M.; San Filippo, J. *J. Am. Chem. Soc.* **1970**, *92*, 6611.
- (5) Giese, B. *Radicals in Organic Synthesis: Formation of Carbon-Carbon Bonds*; Pergamon: Oxford, 1986.
- (6) Russell, G. A.; Hershberger, J.; Owens, K. *J. Am. Chem. Soc.* **1979**, *101*, 1312.
- (7) Hershberger, J.; Russell, G. A. *Synthesis* **1980**, 475.
- (8) Russell, G. A.; Hershberger, J. *J. Am. Chem. Soc.* **1980**, *102*, 7603.
- (9) Russell, G. A.; Hershberger, J.; Owens, K. *J. Organomet. Chem.* **1982**, *225*, 43.
- (10) Russell, G. A.; Tashtoush, H. *J. Am. Chem. Soc.* **1983**, *105*, 1398.
- (11) Russell, G. A.; Tashtoush, H.; Ngoviwatchai, P. *J. Am. Chem. Soc.* **1984**, *106*, 4622.
- (12) Russell, G. A.; Guo, D. *Tetrahedron Lett.* **1984**, *25*, 5239.
- (13) Russell, G. A.; Khanna, R. K. *J. Am. Chem. Soc.* **1985**, *107*, 1450.
- (14) Russell, G. A.; Khanna, R. K. *Tetrahedron* **1985**, *41*, 4133.
- (15) Russell, G. A.; Guo, D.; Khanna, R. K. *J. Org. Chem.* **1985**, *50*, 3423.
- (16) Russell, G. A.; Ngoviwatchai, P. *Tetrahedron Lett.* **1985**, *26*, 4975.
- (17) Russell, G. A.; Khanna, R. K.; Guo, D. *J. Chem. Soc., Chem. Commun.* **1986**, 632.
- (18) Russell, G. A.; Ngoviwatchai, P. *Tetrahedron Lett.* **1986**, *27*, 3479.
- (19) Russell, G. A.; Jiang, W.; Hu, S. S.; Khanna, R. K. *J. Org. Chem.* **1986**, *51*, 5498.
- (20) Russell, G. A.; Khanna, R. K. *Adv. Chem. Ser.* **1987**, *No. 215*, 355.
- (21) Russell, G. A.; Ngoviwatchai, P.; Tashtoush, H.; Hershberger, J. *Organometallics* **1987**, *6*, 1414.
- (22) Russell, G. A.; Khanna, R. K. *Phosphorus Sulfur* **1987**, *29*, 271.
- (23) Russell, G. A.; Ngoviwatchai, P. *Tetrahedron Lett.* **1987**, *28*, 6113.
- (24) Russell, G. A.; Ngoviwatchai, P.; Tashtoush, H. I. *Organometallics* **1988**, *7*, 696.
- (25) Russell, G. A.; Ngoviwatchai, P.; Tashtoush, H. I.; Pla-Dalmau, A.; Khanna, R. K. *J. Am. Chem. Soc.* **1988**, *110*, 3530.
- (26) Russell, G. A.; Hu, S.; Herron, S.; Baik, W.; Ngoviwatchai, P.; Jiang, W.; Nebgen, M.; Wu, Y.-W. *J. Phys. Org. Chem.*, in press.
- (27) Jensen, F. R.; Gale, L. H. *J. Am. Chem. Soc.* **1960**, *82*, 148.
- (28) Winstein, S.; Traylor, T. G. *J. Am. Chem. Soc.* **1956**, *78*, 2597.

Glen A. Russell, Distinguished Professor of the College of Sciences and Humanities of Iowa State University, was born in New York State in 1925. Following degrees from Rensselaer Polytechnic Institute (B.Ch.E. 1947, M.S. 1948) and Purdue (Ph.D. 1951), he spent 7 years at the General Electric Research Laboratory (Schenectady) before joining Iowa State University. His research interests have centered on free radicals and in particular upon the formation and reactions of radicals and radical ions by electron transfer.


of this Account.

Formation of Alkyl Radicals from Alkylmercurials

Alkylmercury(I) species have only transitory existence because of their facile fragmentation to alkyl radicals and mercury metal.²⁹ Alkylmercury(II) compounds also generate alkyl radicals upon thermolysis or photolysis. It appears that the rate of spontaneous (thermal) formation of radicals from *tert*-butylmercurials decreases from the unisolable *t*-BuHgH to (*t*-Bu)₂Hg to *t*-BuHgX (X = Cl, Br, I, OAc). Formation of *t*-Bu[•] can be measured by spin trapping with *N*-*t*-butylphenylnitron.³⁰ In Me₂SO at 100 °C this technique leads to $d[t\text{-Bu}^\bullet]/dt = k[t\text{-BuHgX}]$, where *k* increases from 1.4×10^{-6} with X = Cl to 1.4×10^{-5} for X = I and $4.4 \times 10^{-3} \text{ s}^{-1}$ with X = *t*-Bu.²⁶ The secondary alkyl- and primary alkylmercury derivatives are, of course, considerably more stable in thermal or photochemical radical-forming processes. With irradiation from a 275-W fluorescent sunlamp through Pyrex, typical decomposition rates for 0.1 M *t*-BuHgCl of a few percent per hour are observed as measured by the consumption of the radical trap (*t*-Bu)₂NO[•].


Alkylmercury hydrides, formed by the reaction of RHgX with alkaline NaBH₄, react with alkyl radicals by hydrogen atom transfer to form RHg[•].¹² A free radical chain process leads to the formation of the alkane (RH) or, in the presence of an olefinic radical trap, to the reduced addition product (Scheme I).³¹ For such processes involving RHg[•], the structure of the alkyl group has little effect on the reactivity of its precursor; competitive reactions involving *tert*-butyl- and *n*-butylmercury chlorides with NaBH₄ lead to approximately equal amounts of *tert*-butylation and *n*-butylation products, particularly when a reactive radicalophile (e.g., CH₂=C(Cl)CN) is present to trap R[•].¹⁹

Scheme I



Dialkylmercurials react cleanly with heteroatom-centered acceptor radicals such as RS[•], PhSe[•], PhTe[•], RSO₂[•], or halogen atoms to form the Hg(II) salt and R[•] (Scheme IIa).¹⁰ With carbon-centered radicals (e.g., CCl₃[•]), hydrogen atom abstraction and β-elimination of RHg[•] becomes an important process, and a chain reaction leading to alkene and Hg⁰ has been reported (Scheme IIb).³²

Scheme II



(29) Skinner, H. A. *Adv. Organomet. Chem.* 1964, 2, 49.

(30) Blaukat, U.; Neuman, W. P. *J. Organomet. Chem.* 1973, 49, 323.



(31) Giese, B. *Angew. Chem., Int. Ed. Engl.* 1985, 24, 553.

Table I
Relative Reactivities of Alkylmercury Chlorides toward Free Radicals at 35–40 °C



radical ^a	solvent	<i>t</i> -BuHgCl: <i>i</i> -PrHgCl: <i>n</i> -BuHgCl
RCMe ₂ NO ₂ ^{••}	Me ₂ SO	1.0 ^b :0.07:<0.005
Bu ₃ Sn [•]	PhH	1.0:0.025 ^c :0.005
ClHg [•]	PhH	1.0:0.011 ^c :0.0001
PhS [•]	Me ₂ SO	1.0:0.08:<0.003
PhSe [•]	PhH	1.0:-:<0.004
RCH ₂ CHP(O)(OEt) ₂	Me ₂ SO	1.0:-:<0.003
I [•]	Me ₂ SO	1.0:0.006 ^c :<0.0001

^a Radical precursors were Me₂C=NO₂[•], (*E*)-PhCH=CHSnBu₃, (*E*)-PhCH=C(H)HgCl, PhSSPh, PhSeSePh, CH₂=CHP(O)(OEt)₂, and (*E*)-PhCH=CHI or Ph₂C=CHI. ^b PhCH₂HgCl is 4.7 times as reactive. ^c *c*-C₆H₁₁[•].

Monoalkylmercurials (RHgX) also react readily with acceptor radicals (A[•]). For example, SO₄^{•-}, generated by reaction of S₂O₈²⁻ with I⁻ or Ag⁺, readily displaces R[•]; in Me₂SO at 25 °C strong CIDNP multiplet signals (A/E) for RH, R(-H) and R-R can be observed from the diffusive encounter of primary, secondary, or tertiary alkyl radicals. Alkylmercury salts will also undergo electron transfer with easily oxidized donor radicals (D[•]) or radical anions (D^{•-}). Competing elimination reactions are seldom observed. Thus, reactions 1 and 2 can serve as propagation steps in a number of



chain reactions. In a similar fashion, monomeric HgCl formed in a β-elimination process will regenerate R[•] (reaction 3).¹¹ This process contributes to the efficiency



of RHgX in thermal or photochemical initiation reactions. Monomeric [•]HgX also reacts in a radical manner with reagents Y-A (reaction 4) such as the dichalcogenides or sulfonyl halides, sulfides, and selenides to generate a heteroatom-centered radical (A[•]), which serves as the acceptor radical in reaction 2.¹⁰

In direct-competition experiments with an excess of a mixture of two alkylmercury chlorides, it is observed that in reactions 1–3 the stability of the incipient R[•] has a large effect upon the rate of its production, with *k*_{*t*-Bu}/*k*_{*n*-Bu} as great as 10³.¹⁹ We conclude that reactions 1–3 are processes in which the radical is formed in the rate-determining step; the reactions are probably concerted and quite certainly do not involve RHg[•] intermediates. Table I lists relative reactivity data observed when excesses of mixtures of two alkylmercury chlorides are attacked by acceptor or donor radicals in free radical chain processes.

Reactions 1–3 occur to yield substituted alkyl radicals, e.g., PhCOCH₂[•] or 2-methoxycyclohexyl radical. However, the reactions are very inefficient or do not occur at all when R[•] = CF₃[•], 1-alkenyl, 1-alkynyl, or aryl, presumably because of the stability of the radical that would be formed. Heteroatom-centered acceptor radicals such as PhS[•], PhSO₂[•], (EtO)₂PO[•], and RCO₂[•]

(32) Nugent, W. A.; Kochi, J. K. *J. Organomet. Chem.* 1977, 124, 327, 349, 371.

Table II
Substitutions Occurring by a Homolytic Addition-Elimination Chain Process



reactants (equiv)	conditions ^a	product	% yield, (E/Z)	ref
(E)-PhCH=C(H)HgCl and RHgCl (5), R = <i>t</i> -BuCH ₂ ; <i>c</i> -C ₆ H ₁₁ ; <i>t</i> -Bu	Me ₂ SO, R, 12 h; 12 h; 4 h	PhCH=CHR	23; 78 (6); 97 (31)	11
Ph ₂ C=C(H)HgBr and <i>i</i> -PrMgI (2); <i>t</i> -BuMgCl (2); (<i>c</i> -C ₆ H ₁₁) ₃ B (5)	PhH, R, 12 h; R, 5 h; S, 1 h	Ph ₂ C=CHR	35; 38; 42	24
PhCH=C(H)HgR, R = <i>n</i> -Bu; <i>c</i> -C ₆ H ₁₁ ; <i>t</i> -Bu	PhH, R, 24 h	PhCH=CHR	12; 60; 73	24
(E)- <i>t</i> -BuCH=C(H)HgSR, R = Ph; <i>n</i> -Bu	PhH, S (PhH, S, 3 h; 12 h) 3 h; 12 h	<i>t</i> -BuCH=CHSR	100 (>50); 100 (>50)	8
(E)- <i>t</i> -BuCH=C(H)HgCl and PhYPh, Y = S; Se; Te	PhH, S, 6 h; 2 h; 18 h	<i>t</i> -BuCH=CHYPh	100 (>50); 95 (>50); 89 (>50)	8
(E)- <i>t</i> -BuCH=C(H)HgCl and PhSO ₂ Cl (1); (PhSO ₂) ₂ Hg (5); PhSO ₂ Na (1.2)	Me ₂ SO, S, 3 h; 12 h; <i>t</i> -BuOH/H ₂ O, S, 30 h	<i>t</i> -BuCH=CHSO ₂ Ph	99 (>50); 42 (>50); 81 (>50)	10
(E)- <i>t</i> -BuCH=C(H)HgCl and A ⁻ (1.2), A = (EtO) ₂ POK; PhP(OBu)OK; <i>n</i> -PrSO ₂ Na	Me ₂ SO, S, 29 h; 3.5 h; <i>t</i> -BuOH/H ₂ O, S, 30 h	<i>t</i> -BuCH=CHA	76; 84; 75	8
CH ₂ =C(H)HgCl and A ₂ Hg, A = PhS (1); PhSe (1); PhSO ₂ (5)	Me ₂ SO, R, 20 h; 20 h; 12 h	CH ₂ =CHA	76; 39; 43	21
(E)-PhCH=CHSnBu ₃ and RHgCl (1.2), R = <i>n</i> -Bu; <i>i</i> -Pr; <i>t</i> -Bu	PhH, R, 40 h; S, 18 h; S, 14 h	PhCH=CHR	46 (6); 86 (12); 83 (49)	11
(E)-PhCH=CHI and RHgCl, R = <i>n</i> -Bu (2); <i>i</i> -Pr (2); <i>t</i> -Bu (5)	Me ₂ SO, R, 48 h; 48 h; 1 h	PhCH=CHR	22; 70 (9); 100 (25)	11
(Z)-PhCH=CHI and <i>t</i> -BuHgCl (5)	Me ₂ SO, S, 6 h	PhCH=CHBu- <i>t</i>	90 (0.04)	25
(E)-PhCH=C(H)Q, Q = PhS; PhSO; PhSO ₂ and <i>t</i> -BuHgCl (5; 5; 2)	Me ₂ SO, R, 24 h	PhCH=CHBu- <i>t</i>	36; 32 (21); 43 (81)	11
Ph ₂ C=C(H)Q, Q = Bu ₃ Sn; HgBr; PhSO ₂ and <i>t</i> -BuHgCl (2; 5; 5)	PhH, R, 18 h; Me ₂ SO, R, 12 h; 12 h	Ph ₂ C=CHBu- <i>t</i>	78; 100; 88	11
Ph ₂ C=C(H)Q, Q = HgBr; I; SPh and <i>i</i> -PrHgCl (5)	Me ₂ SO, R, 12 h; 8 h; 96 h	Ph ₂ C=CHPr- <i>i</i>	96; 89; 55	11
Ph ₂ C=C(H)I and A ₂ Hg (1), A = PhS; PhSO ₂ ; (EtO) ₂ PO	Me ₂ SO, R, 12 h	Ph ₂ C=CHA	100; 93; 86	11
Ph ₂ C=C(H)I and (RCO ₂) ₂ Hg, R = Et (1); <i>i</i> -Pr (2); <i>t</i> -Bu (1)	Me ₂ SO, R, 24 h	Ph ₂ C=CHR	48; 39; 28	16
PhC≡CX, X = Bu ₃ Sn; SO ₂ Ph; SPh and <i>t</i> -BuHgCl (5; 5; 0.2)	Me ₂ SO, R, 24 h	PhC≡CBu- <i>t</i>	61; 57; 44	18
PhC≡CI and RHgCl (1.5), R = <i>n</i> -Bu; <i>c</i> -C ₆ H ₁₁ ; <i>t</i> -Bu	Me ₂ SO, S, 7 h	PhC≡CR	48; 93; 100	18
CH ₂ =CHCH ₂ X, X = Bu ₃ Sn; Br; SO ₂ Ph and <i>t</i> -BuHgCl (5)	PhH, S, 4 h; Me ₂ SO, S, 2 h; 4 h	<i>t</i> -BuCH ₂ CH=CH ₂	55; 91; 85	26
HC≡CCH ₂ Cl and <i>t</i> -BuHgCl (5), NaI (5)	Me ₂ SO, S, 7 h	<i>t</i> -BuCH=C=CH ₂	71	26

^aR = 350-nm Rayonet photoreactor; S = 275-W sunlamp ca. 20 cm from Pyrex reaction vessel.

(which decarboxylates to R[•] + CO₂) are also readily formed in reaction 2 or 3 by radical attack upon the appropriate Hg(II) salt.¹¹



Substitution Processes Occurring by a Homolytic Addition-Elimination Mechanism

1-Alkenyl- and 1-alkynylmercurials open up a radical reaction channel in which substitution can occur by an addition-elimination process (Scheme III).^{8,11} Monomeric HgCl thus formed regenerates an alkyl radical by reaction with RHgCl (reaction 3) or by the analogous reaction with RMgCl.²⁴ Heteroatom-centered radicals, which can take the place of R[•] in Scheme III, can be formed by reaction with Y-A reagents (reaction 4),⁸ with HgA₂ (reaction 5),¹¹ or in certain cases by electron





transfer with A⁻ = RSO₂⁻, (EtO)₂PO⁻, RP(OR')O⁻, or RS⁻ (reaction 6).⁸ Examples of photostimulated radical chain substitution reactions of vinylmercurials are given in Scheme IV and in Table II.

Scheme III




The radical addition-elimination sequence occurs with a variety of leaving groups in addition to Hg(I) species. Thus, reaction 7a occurs by a radical chain

process where the leaving group Q can be not only HgCl but also Bu₃Sn, PhS, PhSO₂, and halogen.¹¹ Mercury(II) salts participate in the free radical exchange of one group for another (reaction 7b). Whether exchange




occurs readily depends upon the leaving-group abilities of Q[•] and A[•]. Thus, in the β-styrenyl system, (EtO)₂PO[•] displaces I[•], Bu₃Sn[•], or XHg[•] readily but Br[•], PhS[•], or PhSO₂[•] ineffectively or not at all.

With PhSSPh reaction 8 is observed with R¹ = R² = H not only for Q = HgCl but also for Bu₃Sn, R₃Rb, or Zr(Cp)₂Cl.²¹ When the attacking radical is PhS[•] or




PhSO₂[•], reversal of the addition step is possible, and in reaction 8 only products from α-addition are observed even though the preferred site of radical attack is at the β-carbon atom. Thus, reaction of CH₂=CHSnPh₃ with *p*-MeC₆H₄SH proceeds in a chain fashion to yield mainly *p*-MeC₆H₄SCH₂CH₂SnPh₃ as expected for preferred terminal attack followed by a facile hydrogen atom transfer.³³ However, in reaction with the less reactive PhSSPh or (PhS)₂Hg, only the products from α-substitution (CH₂=CHSPh) are observed for CH₂=CHSnBu₃.²¹

Scheme IV




1-Alkenyl- or 1-alkynylmercurials such as PhCH=C(H)HgSR, PhCH=C(H)HgR, or PhC≡CHgR undergo a radical chain decomposition involving the addition-elimination sequence followed by decomposition of the eliminated Hg(I) species to form Hg⁰ and PhS[•] and PhS[•] (reaction 9).⁸ The photostimulated decomposi-




tion of a 1:1 mixture of *t*-BuHgCH=CHPh and *n*-BuHgCH=CHPh initially forms *t*-BuCH=CHPh and *n*-BuCH=CHPh in an approximately equal ratio ($k_{n\text{-Bu}}/k_{t\text{-Bu}} = 0.7$) since there is little effect of the structure of R on the reactivity of PhCH=C(H)HgR toward alkyl radicals.²⁴


Homolytic aromatic substitution by an addition-elimination mechanism occurs less efficiently than alkenyl substitutions. However, with activated aromatic systems, such as 2-benzothiazole derivatives, a chain reaction with RHgCl has been observed (reaction 10, Q = I, PhSO₂).¹¹



Substitution by alkyl radical attack in allylic, propargylic, and allenic systems by the S_H2' addition-elimination mechanism (reaction 11) is well-known for substrates with Q = Bu₃Sn,³⁴ PhS,³⁵ or cobaloximes.³⁶



With alkylmercurials as the radical precursor, such substitutions also occur readily in a chain fashion with halogen as the leaving groups (e.g., reaction 12). With



poor leaving groups that are also electron withdrawing, e.g., Q = O₂CR, CN, OTos, or OP(O)(OEt)₂, the intermediate radicals (RCH₂ĊHCH₂Q, RCH=ĊHCH₂Q) react with RHgX as an acceptor species to form RCH₂CH(HgX)CH₂Q or RCH=CH(HgX)CH₂Q. When Q is the tosyl or diethyl phosphate group, these intermediates readily undergo elimination to form the allyl or allenyl products, particularly in the presence of iodide ions.

Addition-elimination in 3-butenyl and 5-hexenyl

systems can lead to cyclized products containing 3- and 5-membered rings.³⁷ One example of such a process involving an alkylmercurial as the alkylating agent is given in reaction 13, a process previously described for solvent-derived radicals.³⁸




Table II presents some representative examples of homolytic substitutions involving mercury reagents that proceed by the addition-elimination mechanism. In those substitutions where reactions 1-3 are involved, the processes are usually more efficient with R = *t*-Bu than for *i*-Pr or *n*-Bu. As demonstrated in Table II, homolytic substitution in 1-alkenyl derivatives can occur in a stereospecific manner with retention of configuration.¹⁶ The degree of stereospecificity depends upon the nature of the leaving group, the size of the attacking radical, and presumably the lifetime of the intermediate radical undergoing β-elimination. Thus, C₆H₁₁[•] gives retention of configuration with (*E*)- or (*Z*)-MeO₂CCH=CHSnBu₃, MeO₂CCH=CHI, PhCH=CHI, or MeO₂CCH=CHCl, but *t*-Bu[•] reacts in a stereospecific manner only with the iodides. Apparently, in the β-elimination of Scheme III, I[•] is a better leaving group than Bu₃Sn[•] or Cl[•]. With Q = Bu₃Sn or Cl and R = *t*-Bu in Scheme V, the β-elimination is apparently slow enough and the concentration of the high-energy conformation 1 so low, that both *E* and *Z* substrates react via the low-energy conformation 2 leading to the *E* product. With a smaller R group and Q = Bu₃Sn or Cl or with Q = I and R = *t*-Bu, elimination from conformation 1 occurs more rapidly than the interconversion of 1 to 2. Stereospecificity is also observed in the reactions of *t*-BuHgCl with (*E*)- or (*Z*)-ClCH=CHHgCl.

Substitution Processes Involving S_H2 Reactions of Alkyl Radicals

The addition-elimination sequence of Scheme III is but one example of a process in which an alkyl radical is recycled to yield a species capable of participating in reactions 1 and 2. Scheme VI presents another general chain sequence involving the formation of donor or acceptor radicals. An example of this process has already been given in Scheme IIa. Atom-transfer reactions of R[•] followed by reaction 2 can be involved in the homolytic cleavage of alkylmercury bonds by the halogens or HBr. Other Y-Q reagents that have been observed to participate in Scheme VI are HSnR₃,^{39,40}

(33) Taylor, R. D.; Wardell, J. L. *J. Organomet. Chem.* 1974, 77, 311.

(34) Kosugi, M.; Kurino, K.; Takayama, K.; Migita, T. *J. Organomet. Chem.* 1973, 56, C11.

(35) Keck, G. E.; Byers, J. H. *J. Org. Chem.* 1985, 50, 5442. Keck, G. E.; Burnett, D. A. *J. Org. Chem.* 1987, 52, 2959.

(36) Crease, A. E.; Gupta, B. D.; Johnson, M. D.; Bialkowska, E.; Duong, K. N. V.; Gaudemer, A. *J. Chem. Soc., Perkin Trans. 1* 1979, 2611.

(37) Ashcroft, M. R.; Bury, A.; Cooksey, C. J.; Davies, A. G.; Gupta, B. D.; Johnson, M. D.; Morris, H. *J. Organomet. Chem.* 1980, 195, 89.

(38) Kharrat, A.; Gardrat, C.; Maillard, B. *Can. J. Chem.* 1984, 62, 2385.

(39) Quirk, R. P. *J. Org. Chem.* 1972, 37, 3554.

Scheme V






Figure 1. Reaction of 1:1 mixture of *t*-BuHgCl and *n*-BuHgCl with substituted alkenes (initially 0.10 M) in the presence of an excess of NaBH₄ or Bu₃SnH at 25 °C: (A) CH₂=C(Cl)CN and NaBH₄ in CH₂Cl₂; (B) CH₂=CHCO₂Et and NaBH₄ in CH₂Cl₂; (C) CH₃=C(Cl)CN and Bu₃SnH in PhH.

N-benzyl-1,4-dihydrinicotinamide,⁴¹ HSPH, RSSR, PhSeSePh, PhTeTePh, ClSO₂Ar, and PhSeSO₂Ar.¹⁰

Scheme VI



The reaction of RHgCl with Bu₃SnH has been formulated to involve RHgH as an intermediate³⁹ or to involve halogen atom abstraction by Bu₃Sn•.⁴⁰ As shown in Table I, the reactivity of RHgCl toward Bu₃Sn• increases with the stability of the incipient R•, an observation that we believe excludes RHg• as an intermediate. When applied in the Giese reaction with CH₂=C(Cl)CN as the substrate (reaction 14, Figure 1)



and with an excess of a 1:1 mixture of *t*-BuHgCl and *n*-BuHgCl, the use of Bu₃SnH in PhH or alkaline NaBH₄ in CH₂Cl₂ gives dramatically different results.⁴² With NaBH₄ essentially a 1:1 mixture of *t*-BuCH₂CH(Cl)CN and *n*-BuCH₂CH(Cl)CN is formed, particularly

(40) Bloodworth, A. J.; Courtneidge, J. L. *J. Chem. Soc., Perkin Trans. 1* 1982, 1797.

(41) Kurosawa, H.; Okada, H.; Yasuda, M. *Tetrahedron Lett.* 1980, 21, 959.

(42) Russell, G. A. *Prepr.—Am. Chem. Soc., Div. Petrol. Chem.* 1986, 31, 891.

at high concentrations of CH₂=C(Cl)CN. With Bu₃SnH the initial reaction products are more than 100:1 in favor of the *tert*-butylated product. The borohydride reaction involves the intermediates RHgH and RHg•, which appear to be excluded at least in the initial reaction sequence involving Bu₃SnH.

The reaction of an alkyl radical with BH₄⁻ could in principle form RH and the donor species BH₃⁻. However, BH₄⁻ is unreactive toward alkyl radical attack. When (5-hexenyl)mercury chloride is reduced by alkaline NaBH₄, the cyclized (methylcyclopentane) and uncyclized (1-hexene) products are formed. With excess BH₄⁻ the uncyclized/cyclized ratio is a linear function of the concentration of RHgCl (presumably rapidly converted to RHgH) employed. However, excess BH₄⁻ itself has no effect on the amount of cyclization, indicating a low reactivity of BH₄⁻ with the 5-hexenyl radical.¹² Reactions of RHgX/BH₄⁻ thus involves Hg(I) intermediates formed by radical attack upon RHgH. On the other hand, the RHgX/Bu₃SnH system involves mainly a chain process in which R• is displaced by the attack of Bu₃Sn• at mercury.

Formation of Donor and Acceptor Radicals by Addition of R• to Unsaturated Systems

Unsaturated systems with polar substituents yield carbon-centered adduct radicals that react as donors or acceptors in reactions 1 and 2. Reactions of 1,1-diarylethylenes with *t*-BuHgCl are illustrative.¹⁷ With CH₂=CPh₂, photolysis of *t*-BuHgCl yields *t*-BuCH₂CPh₂•, which does not attack *t*-BuHgCl at 40 °C. The radicals disproportionate to yield *t*-BuCH₂CHPh₂ and *t*-BuCH=CPh₂ in a 1:1 ratio by a nonchain process. With CH₂=C(C₆H₄OMe-*p*)₂ the resulting adduct radical is a donor species that undergoes electron transfer with *t*-BuHgCl, thereby establishing a chain process. Loss of a proton from the dianisylcarbinyl cation gives the substitution product, *t*-BuCH=C(C₆H₄OMe-*p*)₂, in good yield. 1,1-Bis(*p*-nitrophenyl)ethylene also reacts with *t*-BuHgCl by a chain process, but now the product after protonation is *t*-BuCH₂CH(C₆H₄NO₂-*p*)₂. The intermediate bis(*p*-nitrophenyl)carbinyl radical is an acceptor species that reacts with *t*-BuHgCl to form the anion (whether *t*-BuCH₂CH(HgCl)(C₆H₄NO₂-*p*)₂ is an intermediate is unknown; with powerful electron acceptors reaction 2 may proceed directly to A⁻ + HgX⁺ + R•).

With the unsaturated radicaphile is an anion, the resulting adduct is a radical anion (reaction 15), which

Table IV
Kinetic Chain Lengths (kcl) in Photostimulated Reaction of *t*-BuHgCl at 35–40 °C in Me₂SO^a

substrate (equiv)	product	init kcl
PhSSPh (0.2)	<i>t</i> -BuSPh	400 ^b
Me ₂ C=NO ₂ ⁻ (1)	<i>t</i> -BuCMe ₂ NO ₂	50 ^c
CH ₂ =CHP(O)(OEt) ₂ (0.25)	<i>t</i> -BuCH ₂ C(H)(HgCl)P(O)(OEt) ₂	105 (95) ^d
CH ₂ =CHSO ₂ Ph (0.4)	<i>t</i> -BuCH ₂ CH(HgCl)SO ₂ Ph	103
HC≡CCOCH ₃ (0.3)	<i>t</i> -BuCH=C(HgCl)COCH ₃	32 (26) ^d
HC≡CPh (0.3)	<i>t</i> -BuCH=C(HgCl)Ph	15 (14) ^d
Ph ₂ C=CHI (0.3)	<i>t</i> -BuCH=CPh ₂	100
(<i>E</i>)-PhCH=CHSnBu ₃ (0.3)	(<i>E</i>)- <i>t</i> -BuCH=CHPh	71
(<i>E</i>)-PhCH=CHHgCl (0.3)	(<i>E</i>)- <i>t</i> -BuCH=CHPh	56
CH ₂ =C(C ₆ H ₄ OMe- <i>p</i>) ₂ (0.3)	<i>t</i> -BuCH=C(C ₆ H ₄ OMe- <i>p</i>) ₂	18


^a [*t*-BuHgCl]₀ = 0.5 M; irradiation by a 275-W fluorescent sunlamp in 6-mm NMR tubes. Irradiation conditions were only approximately the same for different substrates. ^b 420 in the presence of 1 equiv of *n*-BuHgCl. ^c In the presence of 18-crown-6; longer kinetic chain lengths were observed in its absence. ^d PhH solvent.

The products of addition of R₂HgX to an alkene (RCH₂CH(HgX)(EWG)) or alkyne (RCH=C(HgX)Y) can be converted by reaction with BH₄⁻ to RCH₂CH₂(EWG) or RCH=CHY or with I₂ to RCH₂CH(I)(EWG) or RCH=C(I)Y. The saturated adducts with electron-withdrawing groups are also susceptible to hydrolytic electrophilic cleavage to form RCH₂CH₂(EWG). This product can be formed in one step in reaction 14 but only if the alkene or alkyne has a high reactivity towards R[•]. Otherwise, the alkyl radical is trapped by the intermediate R₂HgH or MH to form RH, and the alkene or alkyne fails to participate significantly in the chain reaction of Scheme I.¹⁹

Kinetic Chain Lengths and Relative Reactivities

Free radical chain reactions of alkylmercurials with mixtures of substrates is a convenient technique for measuring their relative reactivities (e.g., R'CH=CHQ, CH₂=CHCH₂Q, Y-Q, CH₂=CH(EWG)) toward an alkyl radical. This technique has been applied to electronegatively substituted alkenes using NaBH₄ in reaction 14^{43,44} and to a variety of unsaturated anions in reaction 15.^{13,14,20} The technique requires that the relative yields of the substitution or addition products are a measure of the ratio k_A/k_B in Scheme IX. If the chain reactions have a significant kinetic chain length (kcl) with both reactant A and reactant B, the values of k'_A and k'_B in Scheme IX do not influence the ratio of the products P_A and P_B formed in a competition because a significant kcl requires that essentially all adduct radicals (RA[•], RB[•]) will react to continue the chain. Under such conditions, $k_A/k_B = [P_A][B]/[P_B][A]$ or if the ratio of [A]/[B] changes during the reaction, $k_A/k_B = \log \{(1 - [P_A])/[A]_0\} / \log \{(1 - [P_B])/[B]_0\}$.

Scheme IX



The initial kcl's of the photochemical reactions of alkylmercurials are conveniently measured by the nitroxide inhibition method using ¹H NMR to follow the course of the reaction (Figure 2).¹⁹ If (*t*-Bu)₂NO[•] traps the radicals that would have otherwise initiated a chain reaction, the initial kcl can be obtained from the initial




Figure 2. Reaction of 0.12 M CH₂=CHP(O)(OEt)₂ with 4 equiv of *t*-BuHgCl in Me₂SO at 25 °C (dark) or 40 °C (*hν*): (A) dark; (B) sunlamp irradiation; (C) sunlamp irradiation, 0.012 M (*t*-Bu)₂NO[•], initial kcl ~ 100; (D) dark, 6 equiv of NaI; initial kcl ~ 500; (E) sunlamp irradiation, 6 equiv of NaI. Product after NaBH₄ workup was *t*-BuCH₂CH₂P(O)(OEt)₂.

reaction rate divided by the rate of chain initiation, provided the rate of initiation is maintained at a constant value, for example, in a photostimulated process. The kcl is a function of the rate of initiation, and as the reaction proceeds, the kcl decreases as the reactants are consumed. Table IV gives the measured values of the initial kcl observed in some photostimulated reactions of *t*-BuHgCl. The reactions are of moderate kcl but are suitable for relative reactivity studies at least under the initial reaction conditions.

Alkylmercurials react with a variety of substrates by reactions that are clearly radical in nature and that do not occur in the absence of an initiation reaction. The reactions of *t*-BuHgX thus define the chemo-, regio-, and stereospecificity of *t*-Bu[•] addition or substitution. The resulting reactivity data can be used as a diagnostic test for free radical intervention in the reactions of other *tert*-butylating agents. For example, the reaction of *t*-Bu[•] with (*E*)- or (*Z*)-PhCH=CHI gives substitution with 96 ± 1% retention of configuration, a stereospecificity quantitatively different from that observed for substitution with *t*-BuMgCl.²³ On the other hand, (*E*)-PhCH=CHSO₂Ph reacts with *t*-BuHgCl by competing chain reactions to yield (*E*)-PhCH=CHBu-*t* (73%) and PhCH(Bu-*t*)CH(HgCl)SO₂Ph (27%), a regiochemistry essentially the same as observed for reactions with *t*-BuLi or *t*-BuMgCl.^{23,26} The relative reactivities for substitution of PhS or PhSO₂ via reaction 7a for the series PhC≡CSPH, PhC≡CSO₂Ph, and PhCH=CHSO₂Ph observed for *t*-Bu[•] at 40 °C in Me₂SO are 0.3:1.6:1.0, while with *t*-BuLi at 0 °C in THF the reactivity series is

(43) Giese, B. *Angew. Chem., Int. Ed. Engl.* 1983, 22, 753.

(44) Giese, B.; Kretschmar, G.; Meixner, J. *Chem. Ber.* 1980, 113, 2787.

Table V
Relative Reactivities toward *tert*-Butyl Radical at 35-40 °C

substrate	final product	rel react
CH ₂ =C(Cl)CN	<i>t</i> -BuCH ₂ CH(Cl)CN ^a	160
PhSO ₂ Cl	<i>t</i> -BuCl	11
CH ₂ =CHSO ₂ Ph	<i>t</i> -BuCH ₂ CH(HgCl)SO ₂ Ph	7.4
PhSSPh	<i>t</i> -BuSPh	4.3
CH ₂ =CHCO ₂ Et	<i>t</i> -BuCH ₂ CH(HgCl)CO ₂ Et	3.0
CH ₂ =CHP(O)(OEt) ₂	<i>t</i> -BuCH ₂ CH(HgCl)P(O)(OEt) ₂	1.0 ^b
HC≡CCO ₂ Et	<i>t</i> -BuCH=C(HgCl)CO ₂ Et	0.6
CH ₂ =CPh ₂	<i>t</i> -BuCH ₂ CHPh ₂ + <i>t</i> -BuCH=CPh ₂	0.4
MeSSMe	<i>t</i> -BuSMe	0.3
CH ₂ =CHSOPh	<i>t</i> -BuCH ₂ CH(HgCl)SOPh	0.2
CH ₂ =CHSPh	<i>t</i> -BuCH ₂ CH ₂ SPh ^a	0.1
Ph ₂ C=CHI	<i>t</i> -BuCH=CPh ₂	0.1
(<i>E</i>)-PhCH=CHI	(<i>E</i>)- <i>t</i> -BuCH=CHPh	0.28
(<i>Z</i>)-PhCH=CHI	(<i>Z</i>)- <i>t</i> -BuCH=CHPh	0.18
(<i>E</i>)-PhCH=CHBr	(<i>E</i>)- <i>t</i> -BuCH=CHPh	0.35
(<i>E</i>)-PhCH=CHCl	(<i>E</i>)- <i>t</i> -BuCH=CHPh	0.22
(<i>Z</i>)-PhCH=CHCl	(<i>E</i>)- <i>t</i> -BuCH=CHPh	0.03
PhC≡CH	<i>t</i> -BuCH=C(HgCl)Ph	0.04
Me ₂ C=NO ₂ Li	<i>t</i> -BuCMe ₂ NO ₂	0.02
<i>i</i> -PrSSPr- <i>i</i>	<i>t</i> -BuSPr- <i>i</i>	4 × 10 ⁻⁸
<i>t</i> -BuSSBu- <i>t</i>	(<i>t</i> -Bu) ₂ S	5 × 10 ⁻⁴

^a Observed with NaBH₄/CH₂Cl₂. ^b Data of ref 45 with *E*_a = 4 kcal/mol yields 4.8 × 10⁶ M⁻¹ s⁻¹ for the rate of attack of *t*-Bu[•] at 35 °C.

0.3:1.8:1.0.²⁶ Evidence for *t*-Bu[•] attack is also provided in a comparison of the reactivity of 2-cyclopentenone and 2-cyclohexanone with *t*-BuHgI/I⁻ (*hν*/40 °C, Me₂SO), (*t*-Bu)₂CuLi (-30 °C, Et₂O), and (*t*-Bu)₃ZnLi/TMEDA (0 °C, Et₂O). The C₅/C₆ relative reactivities are 3.8 for *t*-Bu[•], 4.2 for (*t*-Bu)₂CuLi, and 3.9 for (*t*-Bu)₃ZnLi.²⁶ Reaction with *n*-Bu[•] gives C₅/C₆

= 2.4 but now the *n*-butylcuprate or -zincate reacts preferentially with 2-cyclohexenone (C₅/C₆ ~ 0.2) by a mechanism obviously not involving attack by the butyl radical. Table V lists some additional relative reactivities toward *t*-Bu[•] from which the absolute rate constants can be calculated from the known rate constant for addition of *t*-Bu[•] to CH₂=CHP(O)(OEt)₂.⁴⁵ Relative reactivities and regioselectivities of attack of *c*-C₆H₁₁[•] and PhS[•] upon 1,2-disubstituted alkenes or alkynes are reported elsewhere.^{24,25}

Conclusions

Alkylmercurials are excellent sources of alkyl radicals that can be utilized in a variety of alkylation reactions proceeding by a chain process. *tert*-Butylmercury halides are more reactive than primary alkylmercurials in many of these processes not only in chain initiation but also in the chain propagation reactions 1-3 with a variety of donor or acceptor radicals formed by addition, addition-elimination, or S_H2 reactions of the *tert*-butyl radical.

It is a pleasure to acknowledge the contributions, many of which are unpublished, of my collaborators, J. Hershberger, H. Tashitouch, P. Ngoviwatchi, W. Jiang, S. Hu, R. K. Khanna, S. Herron, D. Guo, Y.-W. Wu, and A. Pla Dalmau. Financial support has been provided by the National Science Foundation and the Petroleum Research Fund.

(45) Baban, J. A.; Roberts, B. P. *J. Chem. Soc., Perkin Trans. 2* 1981, 161.

Probing Ring Conformations with EPR Spectroscopy¹

K. U. INGOLD^{2a} and J. C. WALTON^{*,2b}

Division of Chemistry, National Research Council of Canada, Ottawa, Ontario, Canada K1A 0R6, and Department of Chemistry, University of St. Andrews, St. Andrews, Scotland KY16 9ST

Received March 28, 1988

The conformational options open to alicyclic molecules have far-reaching chemical consequences. NMR spectroscopy has proved to be an extremely valuable tool for studying ring conformations. The equatorial preferences of substituents and, particularly, the dynamic stereochemistry of cyclohexanes and of medium-ring compounds have been investigated in this way.³⁻⁵ The analogous technique of EPR spectroscopy has been used to study the conformations of some persistent radicals such as mono- and bicyclic semidiones⁶ and nitroxides.⁷ This technique has also been used to study ring inversions for a few transient cyclic radicals such as cyclohexyl and other radicals in which the unpaired electron is localized on one or more ring carbon atoms.⁸⁻¹³ Unfortunately, such radicals contain the

planar radical center in the ring, which drastically alters the ring's conformation in comparison with that of the

(1) Issued as NRCC No. 29627.

(2) (a) NRCC. (b) University of St. Andrews.

(3) Anet, F. A. L.; Anet, R. In *Dynamic Nuclear Magnetic Resonance Spectroscopy*; Jackman, L. M., Cotton, F. A., Eds.; Academic: New York, 1975; Chapter 14, pp 543-619.

(4) Dale, J. *Top. Stereochem.* 1976, 9, 199-270.

(5) Burkert, U.; Allinger, N. L. *Molecular Mechanics*; ACS Monograph 177; American Chemical Society: Washington, DC, 1982; pp 89-108.

(6) Russell, G. A. In *Radical Ions*; Kaiser, E. T., Kevan, L., Eds.; Wiley-Interscience: New York, 1968; Chapter 3, pp 87-150.

(7) See, e.g.: Brière, R.; Lemaire, H.; Rassat, A. *Bull. Soc. Chim. Fr.* 1965, 3273-3283. Rolfe, R. E.; Sales, K. D.; Utley, J. H. P. *J. Chem. Soc., Perkin Trans. 2* 1973, 1171-1177. Chiarelli, R.; Rassat, A. *Tetrahedron* 1973, 29, 3639-3647.

(8) Fessenden, R. W.; Schuler, R. H. *J. Chem. Phys.* 1963, 39, 2147-2195.

(9) Corvaja, C.; Giacometti, G.; Sartori, G. *J. Chem. Soc., Faraday Trans. 2* 1974, 70, 709-718.

(10) Bonazzola, L.; Leray, N.; Marx, R. *Chem. Phys. Lett.* 1974, 24, 88-90.

(11) Gilbert, B. C.; Trenwith, M. *J. Chem. Soc., Perkin Trans. 2* 1975, 1083-1090.

(12) Hori, Y.; Shimada, S.; Kashiwabara, H. *J. Phys. Chem.* 1986, 90, 3073-3079.

(13) Berson, J. A.; Griller, D.; Owens, K.; Wayner, D. D. M. *J. Org. Chem.* 1987, 52, 3316-3319.

Keith U. Ingold received his B.Sc. degree from University College London and his D.Phil. from Oxford. He is Associate Director of the Division of Chemistry of the National Research Council of Canada, and his research has concentrated on the chemistry of free radicals in solution.

John C. Walton received his B.Sc. and D.Sc. degrees from Sheffield University and his Ph.D. from St. Andrews. He is now a Reader in organic chemistry at St. Andrews. His research interests are mainly in the area of organic free radicals.