

Touching All the Bases: Synthesis of Inositol Polyphosphate and Phosphoinositide Affinity Probes from Glucose

GLENN D. PRESTWICH*

Departments of Chemistry and Biochemistry & Cell Biology, University at Stony Brook,
Stony Brook, New York 11794-3400

Received April 17, 1996

Introduction

Receptor-activated cleavage of phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂, or simply PIP₂) by phospholipase C (PLC) (Figure 1) releases the second messenger D-*myo*-inositol 1,4,5-trisphosphate (Ins-(1,4,5)P₃).¹ In turn, Ins(1,4,5)P₃ (IP₃) interacts stereospecifically with membrane receptors to promote the release of Ca²⁺, a key event in cellular signal transduction.^{2,3} This process mediates cellular responses to hormones, neurotransmitters, and other cellular signals including odorants^{4–6} and bitter taste.⁷ The interaction of individual inositol polyphosphates (IP_{*n*}s) with cellular targets has been intensely studied in the past eight years, and both the chemistry^{8,9} and biochemistry of IP_{*n*}s^{10–12} have been extensively reviewed.

However, the phosphoinositide (PI) pathway is not restricted to the inositol polyphosphates alone. A PI 3-kinase^{13,14} is linked to protein tyrosine kinases activated by a number of peptide hormones,^{15,16} and the principal reaction product appears to be PtdIns-(3,4,5)P₃, or simply PIP₃, arising from phosphorylation of PIP₂ (Figure 1). Phosphoinositide polyphosphates are important in recruitment of signaling proteins to cell membranes,¹⁷ in modulating actin polymerization,¹⁸ and in regulating membrane traffic.¹⁹ Finally, the PI pathway generates (or responds to) a myriad of other inositol polyphosphates.²⁰ Affinity chromatography methods have been employed to separate IP_{*n*} kinase, phosphatase, and binding activities.^{21–24} A selection of the most important IP_{*n*}s is shown in Figure 2, with the number of phosphates increasing from left to right.

About the IP₃ Receptor. A 260 kDa receptor protein (IP₃R) that specifically recognizes Ins(1,4,5)-P₃ and mediates its role in calcium release²⁵ was first shown to be localized to the endoplasmic reticulum in cerebellar neurons.²⁶ The Ca²⁺ channel is formed by a homotetramer of IP₃R subunits, which exist in at least three main subtypes, each with alternatively spliced forms.^{10,27} Heterotetramers also form naturally and increase receptor diversity.^{28,29} IP₃R undergoes phosphorylation of two serines during activa-

tion,³⁰ and an ATP binding site has been identified by photoaffinity labeling.³¹ Ca²⁺-dependent calmodulin binding to the receptor has been observed,³² and the immunophilin FKBP modulates Ca²⁺ flux through the IP₃R channel.^{33,34}

* Present address: Department of Medicinal Chemistry, School of Pharmacy, The University of Utah, Salt Lake City, Utah 84112. Phone: (801) 581-7063. Fax: (801) 581-7087. E-mail: gprestwich@deans.pharm.utah.edu.

- (1) Berridge, M. J. *Nature* **1993**, *361*, 315–325.
- (2) Berridge, M. J. *Mol. Cell. Endocrinol.* **1992**, *98*, 119–124.
- (3) Clapham, D. E. *Cell* **1995**, *80*, 259–268.
- (4) Boekhoff, I.; Tareilus, E.; Strotmann, J.; Breer, H. *EMBO J.* **1990**, *9*, 2543–2458.
- (5) Fadool, D. A.; Ache, B. W. *Neuron* **1992**, *9*, 1–20.
- (6) (a) Ronnett, G. V.; Snyder, S. H. *Trends Neurosci.* **1992**, *15*, 508–513. (b) Bruch, R. C. *Comp. Biochem. Physiol., B* **1996**, *113*, 451–459.
- (7) Hwang, P. M.; Verma, A.; Bredt, D.; Snyder, S. H. *Proc. Natl. Acad. Sci. U.S.A.* **1990**, *87*, 7395–7399.
- (8) Billington, D. C. *The Inositol Phosphates: Chemical Synthesis and Biological Significance*, VCH Verlagsgesellschaft: Weinheim, Germany, 1993.
- (9) Potter, B. V. L.; Lampe, D. *Angew. Chem., Int. Ed. Eng.* **1995**, *34*, 1933–1972.
- (10) Furuichi, T.; Mikoshiba, K. *J. Neurochem.* **1995**, *64*, 953–960.
- (11) Bezprozvanny, I.; Ehrlich, B. E. *J. Membr. Biol.* **1995**, *145*, 205–216.
- (12) Potter, B. V. L.; Nahorski, S. R. In *Drug Design for Neuroscience*, Kozikowski, A. P., Ed.; Raven Press: New York, 1993; pp 383–416.
- (13) (a) Carpenter, C. L.; Cantley, L. C. *Biochemistry* **1990**, *29*, 11147–11160. (b) Carpenter, C. L.; Cantley, L. C. *Curr. Opin. Cell. Biol.* **1996**, *8*, 153–158.
- (14) Kapeller, R.; Toker, A.; Cantley, L. C.; Carpenter, C. L. *J. Biol. Chem.* **1995**, *270*, 25985–25991.
- (15) Hawkins, P. T.; Jackson, T. R.; Stephens, L. R. *Nature* **1992**, *358*, 157–159.
- (16) Wennstrom, S.; Hawkins, P.; Cooke, F.; Hara, K.; Yonezawa, K.; Kasuga, M.; Jackson, T.; Claesson-Welsh, L.; Stephens, L. *Curr. Biol.* **1994**, *4*, 385–393.
- (17) Hyvonen, M.; Macias, M. J.; Nilges, M.; Oschkinat, H.; Saraste, M.; Wilmanns, M. *EMBO J.* **1995**, *14*, 4676–4685.
- (18) Sohn, R. H.; Chen, J.; Koblan, K. S.; Bray, P. F.; Goldschmidt-Clermont, P. J. *J. Biol. Chem.* **1995**, *270*, 21114–21120.
- (19) (a) Janmey, P. *Chem. Biol.* **1995**, *2*, 61–65. (b) DeCamilli, P.; Emr, S. D.; McPherson, P. S.; Novick, P. *Science* **1996**, *271*, 1533–1539.
- (20) Shears, S. B. *Pharmacol. Ther.* **1991**, *49*, 70–104.
- (21) Prestwich, G. D.; Marecek, J. F.; Mourey, R. J.; Theibert, A. B.; Ferris, C. D.; Danoff, S. K.; Snyder, S. H. *J. Am. Chem. Soc.* **1991**, *113*, 1822–1825.
- (22) Theibert, A. B.; Estevez, V. A.; Ferris, C. D.; Danoff, S. K.; Barrow, R. K.; Prestwich, G. D.; S. H. Snyder *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 3165–3169.
- (23) Abdullah, M.; Hughes, P. J.; Craxton, A.; Gigg, R.; Marecek, J. F.; Prestwich, G. D.; S. B. Shears *J. Biol. Chem.* **1992**, *267*, 22340–22345.
- (24) Hirata, M.; Watanabe, Y.; Ishimatsu, T.; Yanaga, F.; Koga, T.; Ozaki, S. *Biochem. Biophys. Res. Commun.* **1990**, *168*, 379–386.
- (25) Irvine, R. F. *Biochem. Soc. Trans.* **1995**, *23*, 27–35.
- (26) Ross, C. A.; Meldolesi, J.; Milner, T. A.; Satoh, T.; Supattapone, S.; Snyder, S. H. *Nature* **1989**, *339*, 468–470.
- (27) Blondel, O.; Takeda, J.; Janssen, H.; Seino, S.; Bell, G. I. *J. Biol. Chem.* **1993**, *268*, 11356–11363.
- (28) Joseph, S. K.; Lin, C.; Pierson, S.; Thomas, A. P.; Maranto, A. R. *J. Biol. Chem.* **1995**, *270*, 23310–23316.
- (29) Monkawa, T.; Miyawaki, A.; Sugiyama, T.; Yoneshima, H.; Yamamoto-Hino, M.; Furuichi, T.; Saruta, T.; Hasegawa, M.; Mikoshiba, K. *J. Biol. Chem.* **1995**, *270*, 14700–14704.
- (30) Ferris, C. D.; Cameron, A. M.; Bredt, D. S.; Haganir, R. L.; Snyder, S. H. *Biochem. Biophys. Res. Commun.* **1991**, *175*, 192–198.
- (31) Maeda, N.; Kawasaki, T.; Nakade, S.; Yokota, N.; Taguchi, T.; Kasai, M.; Mikoshiba, K. *J. Biol. Chem.* **1991**, *266*, 1109–1116.

Glenn D. Prestwich, a *bona fide* boomer from 1948, grew up in California, upstate New York, and Maryland. Lured West, he graduated from CalTech (B.Sc., 1970) and Stanford University (Ph.D., 1974) and then headed back East to Cornell University and then to Nairobi, Kenya (NIH postdoc, 1974–1976). From organic synthesis, he moved first into insect natural products chemistry and biochemistry, and then to the applications of photoaffinity labeling in ligand–receptor interactions. He was on the faculty in the Department of Chemistry and the Department of Biochemistry & Cell Biology at the State University of New York at Stony Brook from 1977 to 1996. Recently, the metastatic proliferation of his interests in biochemistry, cell biology, and biotechnology has led him back West; he is now the Presidential Professor and Chair of Medicinal Chemistry at the University of Utah. This Account is a celebration of why organic chemistry is still the central science for much of biology.




Figure 1. Phosphatidylinositol 4,5-bisphosphate is converted to three additional cell signaling molecules by phosphatidylinositol 3-kinase and phospholipase C. Note: For all phosphate monoesters, the symbol “=” signifies the presence of one or two anionic charges, protons, sodiums, or other appropriate monovalent cations, depending on the isolation procedure used and the buffer employed.




Figure 2. A selection of naturally-occurring and synthetic inositol polyphosphates.

The primary structure of the 2749-amino acid IP₃R^{35,36} shows six³⁷ transmembrane domains at the C-terminus. A construct lacking the N-terminal 418

(32) Danoff, S. K.; Ferris, C. D.; Donath, C.; Fischer, G. A.; Munemitsu, S.; Ullrich, A.; Snyder, S. H.; Ross, C. A. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 2951–2955.

(33) Cameron, A. M.; Steiner, J. P.; Sabatini, D. M.; Kaplin, A. I.; Walensky, L. D.; Snyder, S. H. *Proc. Natl. Acad. Sci. U.S.A.* **1995**, *92*, 1784–1788.

(34) Cameron, A. M.; Steiner, J. P.; Roskams, A. J.; Ali, S. M.; Ronnett, G. V.; Snyder, S. H. *Cell* **1995**, *83*, 463–472.

(35) Miyawaki, A.; Furuichi, T.; Ryou, Y.; Yoshikawa, S.; Nakagawa, T.; Saitoh, T.; Mikoshiba, K. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 4911–4915.

(36) Mignery, G. A.; Newton, C. L.; Archer, B. T., III; Südhof, T. C. *J. Biol. Chem.* **1990**, *265*, 12679–12685.

amino acids failed to bind IP₃, and an IP₃R (1–788) construct possessed full IP₃ binding activity.³⁸ The binding region for IP₃ in different IP₃R proteins was localized by neutralizing antibodies, by truncation studies,^{35,38} and by photoaffinity labeling³⁹ to an Arg- and Lys-rich stretch near residue 475.

Rationally synthesized affinity probes for Ins(1,4,5)P₃ were independently developed in Japan, Germany,

(37) Michikawa, T.; Hamanaka, H.; Otsu, H.; Yamamoto, A.; Miyawaki, A.; Furuichi, T.; Tashiro, Y.; Mikoshiba, K. *J. Biol. Chem.* **1994**, *269*, 9184–9189.

(38) Mignery, G. A.; Südhof, T. C. *EMBO J.* **1990**, *9*, 3893–3898.

(39) Mourey, R. J.; Estevez, V. A.; Marecek, J. F.; Barrow, R. K.; Prestwich, G. D.; Snyder, S. H. *Biochemistry* **1993**, *32*, 1719–1726.




Figure 3. Design features for myo -Ins(1,4,5) P_3 receptor probes.

and the USA.^{21,40–42} Compounds prepared by Ozaki and co-workers feature derivatization of the C-2 hydroxyl group of IP_3 ,⁴³ and these affinity ligands allowed facile purification of soluble binding proteins for IP_3 .⁴⁴ Our group employed selective modification of the P-1 phosphate⁴⁵ (Figure 3) based on two precedents: (i) semisynthetic IP_3 analogs⁴⁶ that stimulated calcium release *in vitro* and (ii) photolabile P-1-modified phosphate esters of IP_3 that showed activity *prior* to photochemical deprotection.⁴⁷ The importance of the 3-OH and 6-OH groups of IP_3 in IP_3R activation of Ca^{2+} flux has been demonstrated.^{48,49} The aminoethyl-tethered probes⁴¹ showed K_D values in rat pancreatic cells 9-fold lower than that of IP_3 , and a photoaffinity derivative labeled three small proteins (<49 kDa). In contrast, the 1-*O*-aminopropyl-tethered [¹²⁵I]-azidosalicylamide- IP_3 ([¹²⁵I]ASA- IP_3) and [³H](4-benzoyldihydrocinnamide- IP_3 ([³H]BZDC- IP_3) developed in our labs specifically labeled a >220 kDa protein, the rat brain IP_3R .³⁹

Receptors for IP_4 and IP_6 . Several proteins specifically recognize other inositol polyphosphates; in particular, high-affinity binding of Ins(1,3,4,5) P_4 (hereafter, IP_4) is distinct from IP_3 binding in brain tissues.^{50–52} Several putative IP_4R s have been purified by affinity chromatography (rat brain²²) or by conventional means (pig brain⁵³). The metabolism and functional roles for IP_5 or IP_6 were recently summarized⁵⁴ and include specific high-affinity binding to

cerebellar membranes,²² to pituitary gland membranes,^{55,56} to specific domains in synaptic vesicle proteins,⁵⁷ to the assembly proteins^{58,59} AP-2 and AP-3 important in endocytosis, to the coated vesicles in the Golgi complex,⁶⁰ and to brain and liver cytosolic⁶¹ proteins. IP_5 and IP_6 ⁶² can regulate IP_4 3-phosphatase activity and enhance Ca^{2+} uptake in primary cultures of brain cells.

The IP_6R isolated from the IP_4 affinity column was efficiently photoaffinity labeled by [¹²⁵I]ASA- IP_4 in racemic^{63,64} and chiral forms.⁶⁵ The partial amino acid sequencing of this IP_6R and the sequencing of a cDNA clone showed that this protein was identical to AP-2,^{59,66} one of several protein complexes associated with the formation of clathrin-coated vesicles.⁶⁷ AP-2 consists of two larger (114 and 106 kDa) and two smaller (50 and 17 kDa) subunits; AP-2 binds ATP and PI intermediates, leading to self-association.^{68,69}

Development of the Ferrier Rearrangement Route to Chiral IP_n Derivatives

A poster presented by Steve Bender at a 1990 American Chemical Society meeting showed the use of a Ferrier rearrangement⁷⁰ to prepare selectively-protected Ins(1,3,4,5) P_4 derivatives from α -D-glucose. Bender's route readily allowed differentiation of the 1-phosphate from the 3,4,5-trisphosphates, thus enabling facile introduction of a linker group. Interest-

(40) Hirata, M.; Yanaga, F.; Koga, T.; Ogasawara, T.; Watanabe, Y.; Ozaki, S. *J. Biol. Chem.* **1990**, *265*, 8404–8407.

(41) Schäfer, R.; Nehls-Sahabandu, M.; Grabowski, B.; Dehlinger-Kremer, M.; Schulz, I.; Mayr, G. *Biochem. J.* **1990**, *272*, 817–825.

(42) Ozaki, S.; Watanabe, Y.; Hirata, M.; Ogasawara, T.; Kanematsu, T.; Yoshida, M. In *Drug Design for Neuroscience*; Kozikowski, A. P., Ed.; Raven Press: New York, 1993; pp 417–434.

(43) Ozaki, S.; Watanabe, Y.; Ogasawara, T.; Hirata, M.; Kanematsu, T. *Carbohydr. Res.* **1992**, *234*, 189–206.

(44) Kanematsu, T.; Takeya, H.; Watanabe, Y.; Ozaki, S.; Yoshida, M.; Koga, T.; Iwanaga, S.; Hirata, M. *J. Biol. Chem.* **1992**, *267*, 6518–6525.

(45) Prestwich, G. D.; Marecek, J. F. In *Inositol Phosphates and Related Compounds: Synthesis and Therapeutic Potential*; Reitz, A., Ed.; American Chemical Society: Washington, DC, 1991; pp 122–131.

(46) Henne, V.; Mayr, G. W.; Grabowski, B.; Koppitz, B.; Soling, H.-D. *Eur. J. Biochem.* **1988**, *174*, 95–101.

(47) Walker, J. W.; Feeney, J.; Trentham, D. R. *Biochemistry* **1989**, *28*, 3272–3280.

(48) Kozikowski, A. P.; Ognyanov, V. I.; Fauq, A. H.; Nahorski, S. R.; Wilcox, R. A. *J. Am. Chem. Soc.* **1993**, *115*, 4429–4434.

(49) Hirata, M.; Watanabe, Y.; Yoshida, M.; Koga, T.; Ozaki, S. *J. Biol. Chem.* **1993**, *268*, 19260–19266.

(50) Challiss, R. A. J.; Willcocks, A. L.; Mulloy, B.; Potter, B. V. L.; Nahorski, S. R. *Biochem. J.* **1991**, *274*, 861–867.

(51) Theibert, A. B.; Supattapone, S.; Ferris, C. D.; Danoff, S. K.; Evans, R. K.; Snyder, S. H. *Biochem. J.* **1990**, *267*, 441–445.

(52) Reiser, G.; Schäfer, R.; Donié, F.; Hülser, E.; Nehls-Sahabandu, M.; Mayr, G. W. *Biochem. J.* **1991**, *280*, 533–539.

(53) Donié, F.; Reiser, G. *Biochem. J.* **1991**, *275*, 453–457.

(54) Sasakawa, N.; Sharif, M.; Hanley, M. R. *Biochem. Pharmacol.* **1995**, *50*, 137–146.

(55) Nicoletti, F.; Bruno, V.; Cavallaro, S.; Copani, A.; Sortino, M. A.; Canonico, P. L. *Mol. Pharmacol.* **1990**, *37*, 689–693.

(56) Hawkins, P. T.; Reynolds, D. J. M.; Poyner, D. R.; Hanley, M. R. *Biochem. Biophys. Res. Commun.* **1990**, *167*, 819–827.

(57) Fukuda, M.; Aruga, J.; Niinobe, M.; Aimoto, S.; Mikoshiba, K. *J. Biol. Chem.* **1994**, *269*, 29206–29211.

(58) Ye, W. L.; Ali, N.; Bembek, M. E.; Shears, S. B.; Lafer, E. M. *J. Biol. Chem.* **1995**, *270*, 1564–1568.

(59) Voglmaier, S. M.; Keen, J. H.; Murphy, J. E.; Ferris, C. D.; Prestwich, G. D.; Snyder, S. H.; Theibert, A. B. *Biochem. Biophys. Res. Commun.* **1992**, *187*, 158–163.

(60) Fleischer, B.; Xie, J. P.; Mayrleitner, M.; Shears, S. B.; Palmer, D. J.; Fleischer, S. *J. Biol. Chem.* **1994**, *269*, 17826–17832.

(61) Xie, J. P.; Fleischer, B.; Fleischer, S. *Biochemistry* **1994**, *33*, 7908–7916.

(62) Stephens, L. R.; Hawkins, P. T.; Stanley, A. F.; Moore, T.; Poyner, D. R.; Morris, P. J.; Hanley, M. R.; Kay, R. R.; Irvine, R. F. *Biochem. J.* **1991**, *275*, 485–499.

(63) Estevez, V. A.; Prestwich, G. D. *Tetrahedron Lett.* **1991**, *32*, 1623–1626.

(64) Theibert, A. B.; Estevez, V. A.; Mourey, R. J.; Marecek, J. F.; Barrow, R. K.; G. D. Prestwich; Snyder, S. H. *J. Biol. Chem.* **1992**, *267*, 9071–9079.

(65) Estevez, V. A.; Prestwich, G. D. *J. Am. Chem. Soc.* **1991**, *113*, 9885–9887.

(66) Timerman, A. P.; Mayrleitner, M. M.; Lukas, T. J.; Chadwick, C. C.; Saito, A.; Watterson, D. M.; Schindler, H.; Fleischer, S. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, *89*, 8976–8980.

(67) Kirchhausen, T. *Curr. Opin. Struct. Biol.* **1993**, *3*, 182–188.

(68) Beck, K. A.; Keen, J. H. *J. Biol. Chem.* **1991**, *266*, 4442–4447.

(69) Keen, J. *Annu. Rev. Biochem.* **1990**, *59*, 415–438.

(70) Ferrier, R. J.; Middleton, S. *Chem. Rev.* **1993**, *93*, 2779–2831.




Figure 4. Retrosynthetic strategy for the Ferrier rearrangement route to *myo*-Ins(1,3,4,5)P₄.

ingly, the enzyme *D-myo*-inositol 3-phosphate synthase converts glucose-6-phosphate to the inositol 1-phosphate by a biochemical ring opening⁷¹ followed by a stereospecific intramolecular aldol reaction⁷² that is in fact the biological counterpart of the Ferrier rearrangement. We therefore switched from our racemic synthetic approaches based on inositol as a cheap starting material to a chiral synthesis based on glucose as the starting material (Figure 4). In this Account, we show how the Ferrier route can provide access to many different IP_{*n*} and PIP_{*n*} derivatives bearing tethered reporter groups.

P-1-Tethered Ins(1,3,4,5)P₄. Racemic P-1-*O*-(aminopropyl)-Ins(1,3,4,5)P₄ was first synthesized from *myo*-inositol⁶³ by modification of a route²¹ to the corresponding Ins(1,4,5)P₃ derivative using selective protection–deprotection strategies.⁸ To obtain the *D-myo* enantiomer, we employed a Ferrier-based synthesis starting with methyl α-D-glucopyranoside (**1**)⁶⁵ (Figure 5). The use of *p*-methoxybenzyl (PMB) ethers (e.g., in **2**) allowed deprotection under oxidative conditions without affecting hydrogenolytically labile groups. Thus, the key precursor **3** for the Ferrier rearrangement was prepared by tritylation, PMB etherification, detritylation, and oxidation/enol acetate formation. The Ferrier rearrangement of **3** with mercuric acetate gave inosose **4** with >90% axial 2-OH, and stereoselective reduction of this β-hydroxyketone with NaBH(OAc)₃ provided the key differentially-protected chiral intermediate **5** in 17% overall yield. This route has become a staple of our research effort, and has been scaled up to the multigram quantities of **5** for preparation of PIP₃ analogs (see below). The aminopropyl-IP₄ analog **7a**, obtained via hydrogenolysis of **6** followed by ion exchange purification, has now been employed to prepare biotinylated (**7c**) and fluorescent probes in addition to IP₄ affinity columns and IP₄ photoaffinity labels (**7b**).

Bender presented mechanistic studies and optimized the stereoselectivity of the Ferrier rearrangement leading to unmodified *D-myo*-Ins(1,3,4,5)P₄.^{73a} The Ferrier reaction proceeded via oxymercuration to stable organomercurials that did not cyclize to product

inososes until excess chloride ion was added to drive the reaction to completion. When workup was performed immediately after chloride addition, diastereomeric α-mercurio ketones were formed from the (*Z*)- and (*E*)-enol acetates. With excess chloride, both enol acetates appeared to give the axial C-2 OH (*myo*-inositol numbering) and equatorial C-1 OAc with strong diastereoselectivity. A comprehensive review of this reaction was recently published,⁷⁰ and a detailed mechanistic study using deuterated glucose was described.^{73b} Several additional selectively P-1-modified Ins(1,3,4,5)P₄ derivatives were prepared by our route.⁷⁴

Extension to Other IP₄ and IP₃ Derivatives

Selective manipulation of protecting groups in our original Ferrier route to Ins(1,3,4,5)P₄ has provided access to several unnatural IP₄ regioisomers. The strategy devised by Dr. G. Dormán involved first blocking the C-4 and C-6 hydroxyls of methyl α-D-glucopyranoside as the 4-methoxybenzylidene acetal (Figure 6). After protection of the two remaining OH groups, a regioselective reductive cleavage of the benzylidene acetal gave the primary alcohol at C-6 for further processing.

The developmental drug α-Trinositol, *D-myo*-Ins-(1,2,6)P₃, was produced by selective degradation of phytic acid with phytase.⁷⁵ Until 1996, this compound was under development by Perstorp Pharma as a novel anti-inflammatory⁷⁶ and anti-vasoconstrictive therapeutic agent for analgesia.⁷⁷ Two chemically-modified photoaffinity probes were prepared to determine the molecular target of α-Trinositol.

Figure 7 shows the synthesis⁷⁸ of P-5-(aminopropyl)-Ins(1,2,5,6)P₄, selected on the basis of similar biological activity and binding affinity⁷⁹ of Ins(1,2,5,6)P₄ relative to α-Trinositol. Primary alcohol **8** was converted by the sequence described above to give inosose **9**. Introduction of the 1,2,6-tris(phosphomonoester) groups, PMB removal, installation of the P-5-aminopropyl linker **10**, and hydrogenolysis followed by amidation gave [³H]BZDC-P-5-*O*-(aminopropyl)-*D-myo*-Ins(1,2,5,6)P₄ (**11**). Proteins in rat brain, pig aorta, or heart muscle and human umbilical vascular smooth muscle cells were labeled using photoprobe **11**. Although [³H]-α-Trinositol bound to specific sites in epithelial cells of umbilical vasculature,⁸⁰ only non-specific photolabeling was observed with **11** (A. Chaudhary, unpublished results). Binding assays with pig aorta membranes⁷⁹ using **11** as a ligand showed only low-affinity sites.

On the basis of these results and the knowledge that 3,4,5-triesters appeared to be active forms of α-Trinositol,⁷⁵ we reasoned that a 4-acylated Ins(1,2,6)P₃ might be a more suitable photoprobe. To this end, the

(74) Sawada, T.; Shirai, R.; Matsuo, Y.; Kabuyama, Y.; Kimura, K.; Fukui, Y.; Hashimoto, Y.; Iwasaki, S. *BioMed. Chem. Lett.* **1995**, *5*, 2263–2266.

(75) Blum, C.; Karlsson, S.; Schlewer, G.; Spiess, B.; Rehnberg, N. *Tetrahedron Lett.* **1995**, *36*, 7239–7242.

(76) Woie, K.; Reed, R. K. *Am. J. Respir. Crit. Care Med.* **1994**, *150*, 924–928.

(77) Wahlestedt, C.; Reis, D. J.; Yoo, H.; Adamsson, M.; Andersson, D.; Edvinsson, L. *Neurosci. Lett.* **1992**, *143*, 123–126.

(78) Chaudhary, A.; Dormán, G.; Prestwich, G. D. *Tetrahedron Lett.* **1994**, *35*, 7521–7524.

(79) Yoo, H. Y.; Fallgren, B.; Lindahl, A.; Wahlestedt, C. *Eur. J. Pharmacol., Mol. Pharmacol. Sect.* **1994**, *268*, 55–63.

(80) Walsh, D. A.; Mapp, P. I.; Polak, J. M.; Blake, D. R. *J. Pharmacol. Exp. Ther.* **1995**, *273*, 461–469.

(71) Migaud, M. E.; Frost, J. W. *J. Am. Chem. Soc.* **1995**, *118*, 495–501.

(72) Wong, Y.-H.; Sherman, W. R. *J. Biol. Chem.* **1985**, *261*, 11083–11092.

(73) (a) Bender, S. L.; Budhu, R. J. *J. Am. Chem. Soc.* **1991**, *113*, 9883–9885. (b) Yamauchi, N.; Terachi, T.; Eguchi, T.; Kakinuma, K. *Tetrahedron* **1994**, *50*, 4125–4136.




Figure 5. Ferrier route to *D-myo*-P-1-*O*-(aminopropyl)-Ins(1,3,4,5)P₄. Reagents: (a) Ph₃CCl, DMAP, Et₃N, DMF; (b) NaH, *p*-MB-Cl, DMF; (c) 5% H₂SO₄, CH₃OH; (d) (COCl)₂, DMSO, Et₃N, CH₂Cl₂; (e) Ac₂O, K₂CO₃; (f) Hg(OAc)₂, Me₂CO-H₂O; then saturated NaCl; (g) NaBH(OAc)₃, AcOH; (h) BOM-Cl, Bu₄NBr, H⁺ sponge, CH₃CN; (i) NaOH, MeOH; (j) phosphite linker, tetrazole, CH₂Cl₂; (k) *m*-CPBA; (l) DDQ, wet CH₃CN; (m) (BnO)₂PN*i*Pr₂, tetrazole, CH₂Cl₂; (n) *m*-CPBA; (o) H₂ (4 atm), Pd/C, EtOH; (p) Chelex column (Na⁺ form), Et₃NH⁺HCO₃⁻ (TEAB) buffer, pH 8.3.




Figure 6. Benzylidene ketal approach for preparation of derivatives of Ins(1,2,5,6)P₄, Ins(1,2,4,5)P₄, and Ins(1,4,5)P₃.

4,6-benzylidene ketal of methyl α-D-glucopyranoside was opened with (*i*Bu)₂AlH to glucose derivative **12** (Figure 8); Ferrier rearrangement and selective reduction gave intermediate **13**, and manipulation of protecting groups and introduction of phosphates gave **14**. The C-4 OH was acylated to a 5-aminopentanoyl derivative, which was amidated to photoaffinity label **15**. Photoaffinity labeling experiments with the 4-acyl-[³H]BZDC derivative **15** revealed selectively-labeled proteins in human platelet membranes and in human umbilical smooth muscle epithelial cells (A. Chaudhary, unpublished results).

Efforts to characterize the rat olfactory IP₃R using P-1-tethered Ins(1,4,5)P₃ affinity matrix and photoaffinity labels had been unsuccessful, despite promising results with catfish olfactory IP₃R.⁸¹ Curiously, Ins-(2,4,5)P₃ showed 50- to 100-fold higher potency relative to Ins(1,4,5)P₃ in rat olfactory tissues in two physiological assays. In addition, several cell types exhib-

ited Ca²⁺ release in response to *D-myo*-Ins(1,2,4,5)P₄ with the same profile expected for stimulation of the IP₃R by Ins(1,4,5)P₃.^{82,83} A hybrid probe was prepared⁸⁴ from intermediate **16** (Figure 9). Protecting groups were selectively repositioned (**17**, **18**) and the required phosphate esters were installed (**19**) to give the P-2-linked Ins(1,2,4,5)P₄ (**20**).

Since our earlier work on tethered IP₃ had yielded racemic material, a new route (Figure 10) was developed to *D-myo*-P-1-*O*-(aminopropyl)-Ins(1,4,5)P₃ affinity probes based on the scheme in Figure 9.⁸⁵ Thus, manipulation of acetal **21** via enol acetate **22** and protection of C-3 as a benzyl ether led to rearranged intermediate **23**. The chiral IP₃ derivative **24** was converted with [³H]BZDC-NHS ester to photoprobe **25**, which has been employed to photoaffinity label the PIP₂-binding pleckstrin homology (PH) domain of phospholipase C-δ₁.⁸⁶

Onward to IP₆ Derivatives

Our original synthesis of *meso*-P-2-*O*-(aminohexyl)-Ins(1,2,3,4,5,6)P₆ is shown in Figure 11.^{87a} Ketal **26** was converted to key intermediate **27**, the linker was

(81) Restrepo, D.; Teeter, J. H.; Honda, E.; Boyle, A. G.; Marecek, J. F.; Prestwich, G. D.; Kalinoski, D. L. *Am. J. Physiol. (Cell Physiol.)* **1992**, *263*, C667–C673.

(82) Hirata, M.; Narumoto, N.; Watanabe, Y.; Kanematsu, T.; Koga, T.; Ozaki, S. *Mol. Pharmacol.* **1994**, *45*, 271–276.

(83) Mills, S. J.; Safrany, S. T.; Wilcox, R. A.; Nahorski, S. R.; Potter, B. V. L. *BioMed. Chem. Lett.* **1993**, *3*, 1505–1510.

(84) Chen, J.; Dormán, G.; Prestwich, G. D. *J. Org. Chem.* **1996**, *61*, 393–397.

(85) Dormán, G.; Chen, J.; Prestwich, G. D. *Tetrahedron Lett.* **1995**, *36*, 8719–8722.

(86) (a) Tall, E.; Dormán, G.; Garcia, P.; Shah, S.; Prestwich, G. D.; Rebecchi, M. J. *J. Biol. Chem.*, submitted for publication. (b) Ferguson, K. M.; Lemmon, M. A.; Schlessinger, J.; Sigler, P. B. *Cell* **1996**, *83*, 1037–1046 reports a crystal structure of the PLC-δ₁ PH domain.

(87) (a) Marecek, J. F.; Prestwich, G. D. *Tetrahedron Lett.* **1991**, *32*, 1863–1866. (b) Chen, J.; Prestwich, G. D. *J. Labelled Compd. Radiopharm.* **1996**, in press.




Figure 7. Synthesis of P-5-linked Ins(1,2,5,6)P₄. Reagents: (a) *p*-methoxybenzaldehyde dimethyl acetal, pTsOH, DMF; (b) BnBr, NaH, DMF; (c) TMSCl, NaBH₃CN; (d)–(f) “Ferrier sequence”, steps (d)–(f) in Figure 5; (g) NaBH(OAc)₃, AcOH; (h) NaOH, MeOH; (i) “phosphorylation”, steps (m) and (n) in Figure 5; (j) (NH₄)₂Ce(NO₃)₆; (k) “linker attachment”, steps (j) and (k) in Figure 5; (l) “deprotection sequence”, steps (o) and (p) in Figure 5; (m) [³H]BZDC-NHS, DMF, 0.25 M TEAB, pH 8.3; then DEAE-cellulose (HCO₃⁻ form).




Figure 8. Synthesis of 4-acylated D-*myo*-Ins(1,2,6)P₃. Reagents: (a) Bu₄N⁺HSO₄⁻-BnBr, CH₂Cl₂, aqueous NaOH; (b) PMB-Cl, NaH, DMF; (c) DIBAL, CH₂Cl₂; (d)–(g) Ferrier sequence; (h) NaOH, MeOH; (i) phosphorylation; (j) (NH₄)₂Ce(NO₃)₆, MeCN-H₂O (9:1); (k) CbzNH(CH₂)₄COOH, DCC, DMAP; (l) deprotection sequence; (m) “photolabel attachment”, step (m) in Figure 7.

attached (**28**), and protecting groups were cleaved (**29**). The IP₆ affinity column **30b** allowed purification of Ins(1,3,4)P₃ 5/6-kinase²³ and IP₆ kinase,⁸⁸ and the [³H]-BZDC-IP₆ probe **30a** has proven to be a powerful tool for labeling the active sites of proteins important in intracellular protein trafficking (J. D. Olszewski, A. Chaudhary, B. Mehrotra, unpublished results). For this material, the asymmetric route from glucose made

(88) Voglmaier, S. M.; Bembenek, M. E.; Kaplin, A. I.; Dormán, G.; Olszewski, J. D.; Prestwich, G. D.; Snyder, S. H. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, *93*, 4305–4310.




Figure 9. Synthesis of D-*myo*-P-2-*O*-(aminopropyl)-Ins(1,2,4,5)-P₄. Reagents: (a) NaOH, MeOH; (b) TsOH, 2,2-dimethoxypropane, Me₂CO; (c) NaH, BnBr, DMF; (d) TsOH, Me₂CO-H₂O; (e) Bu₂SnO, *n*-Bu₄NI, toluene; then PMB-Cl; (f) linker attachment; (g) (NH₄)₂Ce(NO₃)₆, MeCN-H₂O; (h) phosphorylation; (i) deprotection sequence.




Figure 10. Synthesis of D-*myo*-P-1-*O*-(aminopropyl)-Ins(1,4,5)-P₃. Reagents: (a) *p*-methoxybenzaldehyde dimethyl acetal, pTsOH, DMF; (b) BnBr, CH₂Cl₂, *n*Bu₄NHSO₄, 10% aqueous NaOH; (c) NaH, PMB-Cl, DMF; (d)–(h) Ferrier sequence; (i) BOM-Cl, Bu₄NBr, H⁺ sponge, MeCN; (j) NaOH, MeOH; (k) linker attachment; (l) DDQ, wet CH₂Cl₂; (m) phosphorylation; (n) deprotection; (o) photolabel attachment.

little sense. On the other hand, since receptors could have chiral recognition sites, a P-1-tethered IP₆ (Figure 11, bottom) was initially prepared in racemic





Figure 11. Synthesis of *meso*-P-2-*O*-(aminohexyl)-Ins(1,2,3,4,5,6)-P₆. Reagents: (a) NaH, AllBr, DMF; (b) H₃O⁺, CH₃OH; (c) Bu₂-SnO; (d) AllBr, DMF; (e)–(f) linker attachment; (g) (Ph₃P)₃RhCl, C₂H₅OH; (h) H₃O⁺, CH₃OH; (i) phosphorylation; (j) *m*-CPBA; (k)–(l) deprotection; (m) BZDC-NHS, TEAB, pH 8.5 for R = BZDC; Affigel-10, NaHCO₃ buffer for R = Affigel-10 resin.

form^{87b} by conversion of **26** to protected intermediate **31**. [³H]BZDC-P-1-(aminohexyl)-linked IP₆ **32** has shown different (and mostly lower) affinities for several IP₆ binding proteins (coatomer, AP-2, synaptotagmin, and cytosolic IP₆BP) relative to **30a** (A. Chaudhary, A. A. Profit, J. Chen, unpublished results). Preparation of the chiral derivative via a Ferrier route awaits justification from biological data.

Phosphoinositide Polyphosphates

Lipid-Modified Analogs of PtdIns(4,5)P₂ and PtdIns(3,4,5)P₃. A number of syntheses of phosphoinositide polyphosphates have recently been presented.^{89–92} However, none of the published routes

allowed access to tritium-labeled materials or permitted site-specific incorporation of a reporter group. Thus, we followed two strategies for introducing reporter groups: modification of the lipid moiety and insertion of a linker on the phosphodiester phosphate. Figure 12 illustrates the structures of several common phospholipids and a variety of fluorophore-linked and photophore-linked PIP₂-type probes.

The acyl-modification strategy is described first. Figure 13 compares the chemical structures of the P-1-tethered photoaffinity labels used as probes for Ins(1,4,5)P₃ and Ins(1,3,4,5)P₄ binding sites with both the native phosphatidylinositol polyphosphates PtdIns(4,5)P₂ (PIP₂) and PtdIns(3,4,5)P₃ (PIP₃) and with the target acyl-modified PIP₂ and PIP₃ photoaffinity labels. Lipid-modified analogs of PIP₂ and PIP₃ were prepared in which either a 6-aminohexanoyl group or a 12-aminododecanoyl group was installed in the *sn*-1 position.⁹³ This strategy allowed attachment of a reporter group (e.g., BZDC photophore, NBD fluorophore) that could either reside in the lipid bilayer or interact with lipid-binding sites on a macromolecular target.

The synthesis (Figure 14) followed a convergent approach, beginning with selective *sn*-1-*O*-acylation⁹⁴ of a 3-PMB-protected chiral glycerol synthon, followed by acylation of the *sn*-2 position and oxidative deprotection with DDQ to give the desired 1,2-*O*-diacylglycerol derivative. Reaction with benzyl(*N,N*-diisopropylamino)chlorophosphine afforded rather labile phosphoramidites **33**, which were then condensed with the appropriately protected *D*-*myo*-inositol derivative. Intermediate **23**⁸⁵ from the P-1-tethered Ins(1,4,5)P₃ route was used in making PIP₂ analogs, while intermediate **5**⁶⁵ from the P-1-tethered Ins(1,3,4,5)P₄ route was used for making PIP₃ analogs. In these intermediates (e.g., **23**), benzyl (Bn) or benzylloxymethyl (BOM) groups protected final hydroxyl groups and PMB groups masked future phosphomonoesters. Deprotection of the PMB groups, phosphorylation, and finally hydrogenolysis and ion exchange chromatography afforded the aminoacyl-tethered analogs of PIP₂ and PIP₃. Attachment of the photophore using [³H]-BZDC-NHS ester⁹⁵ gave the desired photolabile probes **35**. In addition, we prepared an extremely high specific activity, nonhydrolyzable ether analog of PIP₂ (**36**) for binding assays (J. Chen, unpublished results). Thus, condensation of a 1,2-*O*-di-10-undecenyl-*sn*-glycerol-derived phosphoramidite with the IP₃ intermediate **24** led to a protected diether PIP₂ analog. Hydrogenolysis with carrier-free tritium (1 atm) followed by hydrogen (4 atm) gave the tritium-labeled diether **36** with >200 Ci/mmol specific activity.

Triester Analogs of PIP₂ and PIP₃. A novel concept for photolabeling of PIP₂ and PIP₃ binding proteins was conceived by Dr. Q.-M. Gu in which the reporter group could be positioned at the water/

(89) Watanabe, Y.; Hirofujii, H.; Ozaki, S. *Tetrahedron Lett.* **1994**, *35*, 123–124.

(90) Gou, D.-M.; Chen, C.-S. *J. Chem. Soc., Chem. Commun.* **1994**, 2125–2126.

(91) Reddy, K. K.; Saady, M.; Falck, J. R.; Whited, G. *J. Org. Chem.* **1995**, *60*, 3385–3390.

(92) Bruzik, K. S.; Kubiak, R. *J. Tetrahedron Lett.* **1995**, *36*, 2415–2418.

(93) Chen, J.; Profit, A. A.; Prestwich, G. D. *J. Org. Chem.* **1996**, *61*, 6305–6312.

(94) Martin, S. F.; Josey, J. A.; Wong, Y.-L.; Dean, D. W. *J. Org. Chem.* **1994**, *59*, 4805–4820.

(95) Olszewski, J. D.; Dormán, G.; Elliott, J. T.; Hong, Y.; Ahern, D. G.; Prestwich, G. D. *Bioconjugate Chem.* **1995**, *6*, 395–400.




Figure 12. Naturally-occurring phospholipids (PIP₂, PtdIns(4,5)P₂; PS, phosphatidylserine; PE, phosphatidylethanolamine; PC, phosphatidylcholine) and selected PIP₂ analogs with photophoric (BZDC) or fluorophoric (NBD) reporter groups.

phospholipid head group interface.⁹⁶ The PIP₂ analog containing this modification is depicted in Figure 12 at the bottom right. In this strategy, both the diacylglycerol bilayer anchor and the 4,5-bisphosphate (or 3,4,5-trisphosphate) recognition groups were retained. The synthetic route (Figure 15) to the triesters hinges on the use of the 2-cyanoethyl protecting group for the inositol P-1 phosphate that also forms the phosphodiester linkage to the diacylglycerol moiety.

The phosphatidylinositol derivatives were assembled convergently. For the PIP₃ analog, dipalmitoylglycerol was converted to the 2-cyanoethyl *N,N*-diisopropylphosphoramidite derivative and coupled to the inositol **5**⁶⁵ to give the adduct **37a**. The 2-cyanoethyl phosphotriester was stable during the subsequent deprotection and phosphorylation steps, and the phosphorylated product **38a** was converted to the 3-aminopropyl-tethered phosphotriester **39a** during hydrolysis. The aminopropyl group of **39a** was readily converted to the [³H]BZDC-PIP₃ derivative **40a**. The

(96) Gu, Q.-M.; Prestwich, G. D. *J. Org. Chem.*, in press.




Figure 13. Comparison of P-1-tethered photoaffinity probes for IP₃ and IP₄ with PIP₂, PIP₃, and acyl-modified derived photoaffinity analogs of PIP₂ and PIP₃.




Figure 14. Synthesis of acyl-modified PIP₂ and PIP₃ analogs. Reagents: (a) 1-*H*-tetrazole, *m*-CPBA; (b) DDQ, CH₂Cl₂; (c) phosphorylation; (d) deprotection in *t*BuOH-H₂O, NaHCO₃; (e) photolabel attachment.

PIP₂ analog **40b** was prepared following an analogous route, but using the key intermediate **23**⁸⁵ as the inositol component.

The 2-cyanoethyl group provided three advantages. First, the high reactivity of the 2-cyanoethyl *N,N,N,N*-

Figure 15. Synthesis of triester analogs of PIP₂ and PIP₃. Reagents: (a) diacylglycerol cyanoethyl phosphoramidite, tetrazole, CH₂Cl₂; then *m*-CPBA; (b) DDQ, CH₂Cl₂; (c) phosphorylation; (d) deprotection (and reduction of nitrile) in *t*BuOH-H₂O, NaHCO₃; (e) photolabel attachment.

tetraisopropylphosphordiamidite ensured that a bulky primary alcohol and a sterically-congested secondary alcohol could both form efficient new O-P bonds. Second, the 2-cyanoethyl phosphite derivatives were more stable to chromatography than the corresponding benzyl phosphites used for the PIP₂ and PIP₃ phosphodiester route above. Third, not only could the




Figure 16. Application of IP₄ affinity resin to purification of IP₄, IP₆, and PIP₃ binding proteins. The first eluted fraction IP₆R was identified as AP-2; IP₄R-1 and IP₄R-2 are still unidentified; IP₄R-3 has recently been described as the PIP₃ binding protein centaurin.

2-cyanoethyl group serve as the source of the 3-aminopropyl linker, but it could be removed by β -elimination using a trialkylamine to provide high yields of the phosphodiester PIP₂ and PIP₃ (see Figure 13).

Protein Purification and Active Site Labeling

The synthesis of the probes was only the beginning. We have, Pandora-like, succumbed to the temptation to use our newly made keys to try to unlock the black boxes of cellular signaling and intracellular protein transport. Fortunately, these opened boxes have produced not pestilence but treasures beyond our expectations. Selected recent examples are highlighted below.

Labeling with [³H]BZDC-Ins(1,4,5)P₃. Our first results with purification and labeling the IP₃R using P-1-*O*-(3-aminopropyl)-linked derivatives³⁹ mapped the IP₃ binding site. More recently, we showed that PIP₂ binding proteins could also be studied with BZDC-IP₃.⁸⁶ Thus, the δ_1 -isozyme of PLC contains a PH domain responsible for binding of PLC to PIP₂-rich lipid bilayers;^{86b} catalysis of phosphodiester hydrolysis takes place at a different site. Two PLC- β isozymes lack the PH domain, yet still hydrolyze PIP₂ to IP₃ and DAG. Photoaffinity labeling of recombinant proteins comprising either the full-length PLC- δ_1 or only the PH domain with [³H]BZDC-IP₃ modified the N-terminal base-rich sequence unique to this isozyme.⁸⁶ The two PLC- β isozymes were not labeled by this probe. This is the first evidence for an important hydrophobic interaction site of a PH domain in addition to the 4,5-bisphosphate recognition role.

Purification on an IP₄ Affinity Column and Labeling with [³H]BZDC-Ins(1,3,4,5)P₄. Purification by IP₄ affinity chromatography and photoaffinity labeling of putative IP₄ binding proteins from rat brain^{27,77} led to the characterization of four proteins (Figure 16). The first eluting heterooligomer had the highest affinity for IP₆, and was identified⁵⁹ as assembly protein AP-2, an essential component in the formation of clathrin-coated pits during endocytosis. The second and third eluting proteins are heterooligomers with as yet undescribed functions. The fourth protein eluted at much higher ionic strength from this IP₄ affinity column, and the protein was very ef-

ficiently labeled with [³H]BZDC-IP₄. This protein has been cloned and its cDNA sequenced, revealing an unexpected chimeric structure that includes (in addition to the IP₄ binding region) a protein kinase C zinc finger, and several domains suggesting interactions with actin and other cytoskeletal proteins. This protein, named centaurin, is the first protein clearly demonstrated to show preferential binding to PIP₃ relative to IP₄ or any other PI pathway metabolite.⁹⁷

Photolabeling with [³H]BZDC-Ins(1,2,3,4,5,6)-P₆ and Purifications with an IP₆ Affinity Column. P-2-Linked [³H]BZDC-IP₆ was recently prepared⁹⁸ and used along with P-1-linked [³H]BZDC-IP₄ in labeling synaptotagmin C2B domains,⁵⁷ Golgi coatomer proteins,⁶⁰ liver cytosolic IP₆ binding proteins,⁶¹ and assembly proteins AP-2 and AP-3.^{58,69} The P-2-linked IP₆ resin has also proven to be valuable for enzyme purification. For example, an Ins(1,3,4) 5/6-kinase was purified 20-fold in a final step to give homogeneous protein.²³ More recently, the IP₆ kinase responsible for biosynthesis of IP₇ from IP₆ and ATP was purified.⁸⁸

Studies with Modified PIP_n. The PIP₂ and PIP₃ probes have just been completed.^{93,96} Early studies with binding of BZDC-IP₃ and the triester and acyl-modified PIP₂ photolabels to profilin show strikingly different results for each photoprobe. A fluorescent probe, NBD-PIP₂ (see Figure 12), was synthesized⁹³ for studies⁹⁹ on the interaction of a highly basic peptide with the acidic phospholipids PIP₂ and phosphatidylserine (PS) in bilayers. The basic peptides caused sequestration of PIP₂ and PS into lateral domains, thus preventing hydrolysis of PIP₂ by PLC.⁹⁹ The inhibition was released upon phosphorylation of serines in the basic peptides, thereby ablating the necessary electrostatic interaction. Selective labeling of PIP₃ binding proteins with the triester will be reported elsewhere.

(97) Hammonds-Odie, L. P.; Jackson, T. R.; Profit, A. A.; Blader, I. J.; Turck, C.; Prestwich, G. D.; Theibert, A. B. *J. Biol. Chem.* **1996**, *271*, 18859–18868.

(98) Prestwich, G. D.; Dormán, G.; Elliott, J. T.; Marecak, D. F.; Chaudhary, A. *Photochem. Photobiol.*, in press.

(99) Glaser, M.; Wanaski, S.; Buser, C. A.; Boguslavsky, V.; Rashidzade, W.; Morris, A.; Rebecchi, M.; Scarlata, S.; Runnels; Prestwich, G. D.; Chen, J.; Aderem, A.; Ahn, J.; McLaughlin, S. *J. Biol. Chem.*, in press.

Conclusions

Phosphoinositide polyphosphate and inositol polyphosphate recognition sites in target proteins are rich in basic arginine and lysine residues that are protonated at physiological pH. The binding site for a given arrangement of two or more equatorial, anionic phosphates is determined by the number and 3-dimensional display of these bases. Over the past six years, we have sought to understand these binding sites by "touching all the bases" with PIP_n and IP_n substrates bearing reporter groups. A molecular level understanding of regulation of cellular signaling and protein trafficking can be pursued with the palette of phosphoinositide pathway photoprobes described in this Account.

I thank my past and present co-workers in the "IP₃ Group" at Stony Brook for their diligent efforts: M. F. Boehm, V. A. Estevez, J. F. Marecek, A. A. Profit, A. Chaudhary, G. Dormán, J. D. Olszewski, J. Chen, B. Mehrotra, and Q.-M. Gu. Synthetic and biochemical work was first supported by the Center for Biotechnology (a New York State Center of Advanced Technology), and recently by the NIH (Grant NS 29632), DuPont New England Nuclear, and Perstorp Pharma. The stimulating suggestions of many biological colleagues have inspired our pursuit of new target molecules. Among these colleagues, special thanks are due to Professors Solomon H. Snyder and Anne B. Theibert for their roles in getting me started in this research area and for their enduring friendships and research partnerships.

AR960136V