

Behavioral Patterns of Heterometallic Cuboidal Derivatives of $[\text{M}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$ ($\text{M} = \text{Mo}, \text{W}; \text{Q} = \text{S}, \text{Se}$)

RITA HERNANDEZ-MOLINA,[†]
MAXIM N. SOKOLOV,[‡] AND
A. GEOFFREY SYKES^{*,§}

Departamento de Química Inorgánica, Universidad de la Laguna, 38200 La Laguna, Islas Canarias, Spain, Institute of Inorganic Chemistry, Russian Academy of Sciences, pr Lavrentyeva 3, Novosibirsk 630090, Russia, and Department of Chemistry, The University of Newcastle, Newcastle upon Tyne, NE1 7RU U.K.

Received June 5, 2000

ABSTRACT

This Account reports recent progress in the study of some ~20 heterometallic derivatives of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ with reference also to W and Se analogues. Single cubes (3:1) and corner-shared double cubes (6:1), as well as dimers of the 3:1 single cubes, are considered. A classification of the heterometals as subtypes A, B, and C is introduced.

Introduction

Recent studies on metal–chalcogenide cluster complexes have provided exciting new developments which relate to solid-state chemistry, catalysis, and metalloprotein systems.^{1–4} In the latter, Fe_4S_4 was the first cuboidal cluster to be identified, and Fe–S clusters are a major component of biological electron-transport chains.⁵ The structure and reactivity of synthetic analogues have been successfully investigated,¹ but in aqueous solution these have limited stability. In the case of nitrogenase, not only is the larger Fe, S, and Mo (or V) eight-metal cluster Fe_7MoS_9 difficult to synthesize, but further mechanistic understanding is required.⁶

Rita Hernández-Molina was born in La Palma, Canary Islands, in 1965. She obtained her Ph.D. at the University of La Laguna, Tenerife, in 1995 under the supervision of Professors Alfredo Mederos and Pedro Gili. She did postdoctoral work with Professor Sykes's group in Newcastle, and is now lecturer of Inorganic Chemistry at the University of La Laguna. Her current research interests concern chalcogenide-bridged clusters compounds.

Maxim Sokolov, born in Novosibirsk, Russia, in 1967, obtained his Ph.D. at the Institute of Inorganic Chemistry RAS, Novosibirsk, in 1992 under the supervision of Prof. Vladimir Fedin. He carried out postdoctoral work 1994–1999 in Germany, the UK (with Professor Sykes), and Japan. At present he holds a position as Senior Researcher at the Institute of Inorganic Chemistry in Novosibirsk. His research interests are centered on second- and third-row transition metal chemistry.

Geoff Sykes has Ph.D. (1958) and D.Sc. (1973) degrees in chemistry from the University of Manchester and is FRS. His postdoctoral work was done at Princeton and Adelaide. He was Lecturer and Reader at the University of Leeds before being appointed to the Inorganic Chair at the University of Newcastle in 1980. His scientific activities have centered on the mechanisms of inorganic and bioinorganic (metalloprotein) reactions. He has published extensively and received much international recognition.




FIGURE 1. Single $\text{M}_3\text{M}'\text{S}_4$ and double $\text{M}_6\text{M}'\text{S}_8$ heterometal cube core structures ($\text{M} = \text{Mo}, \text{W}$).

A		B				C			
Cr		Fe	Co	Ni	Cu		Ga	Ge	As
Mo			(Rh)	Pd			In	Sn	Sb
W				(Pt)		Hg	Tl	Pb	Bi

FIGURE 2. Elements for which heterometallic derivatives of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ can be prepared.

It is remarkable that alongside Fe only molybdenum and, to a lesser extent, tungsten provide a similar range of cuboidal clusters. The $\text{M}_4\text{Q}_4^{n+}$ ($\text{M} = \text{Mo}, \text{W}; \text{Q} = \text{S}, \text{Se}, \text{Te}$) cubes have all been prepared, although those containing Te remain comparatively rare.⁷ Both Mo and W are octahedrally coordinated, and the aqueous chemistry is accessible as long as acid conditions ($\text{pH} < 1$) are maintained to avoid conjugate-base ($\text{H}_2\text{O} \rightarrow \text{OH}^-$) ligand effects. The $[\text{Mo}_4\text{Q}_4(\text{H}_2\text{O})_{12}]^{5+}$ ($\text{Q} = \text{S}, \text{Se}$) state is most readily prepared, with the 4+ and 6+ cubes readily accessed by electrochemical or chemical means.⁸ On mild oxidation, one of the molybdenums is lost to give the Mo^{IV}_3 incomplete cube $[\text{Mo}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$. The latter can incorporate ~20 different transition and post-transition metals M' to give single ($\text{Mo}_3\text{M}'\text{S}_4$) and/or double cubes ($\text{Mo}_6\text{M}'\text{S}_8$), which together provide the main focus of this Account (Figure 1). Early examples are described in ref 3. One of the most fascinating aspects is that zero oxidation state heteroatoms can be used as the source of M' . Thus, with $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$, direct incorporation of 14 heteroatoms, Fe, Co, Ni, Cu, Pd, Hg, Ga, In, Tl, Ge, Sn, Pb, Sb, and Bi, has been achieved.^{2–4} The existence of tungsten analogues of $[\text{W}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$ ($\text{Q} = \text{S}, \text{Se}$) has been demonstrated, but there is less tendency to incorporate heteroatoms, and to date only ~7 have been identified.⁹

Classification of Heterometallic Derivatives of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$

In 1986, a report¹⁰ that $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ reacts with metallic iron to give the 3:1 heterometallic cube $[\text{Mo}_3\text{FeS}_4(\text{H}_2\text{O})_{10}]^{4+}$ led to an intense search for similar reactions. The extent of the chemistry of 3:1 and 6:1 adducts is summarized with reference to the Periodic Table (Figure 2). Three subtypes of $\text{Mo}_3\text{S}_4^{4+}$ derivatives, A, B, and C, are defined, and distinctive properties of each are now considered.

In type A with Group 6 metals, the Mo_3CrS_4 , Mo_4S_4 , Mo_3WS_4 and W_3MoS_4 cores are formed. The metal atoms are all octahedral and have variable oxidation states, $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{n+}$, $[\text{Mo}_3\text{WS}_4(\text{H}_2\text{O})_{12}]^{n+}$, and $[\text{W}_3\text{MoS}_4-$

[†] Universidad de la Laguna.

[‡] Russian Academy of Sciences.

[§] The University of Newcastle.





FIGURE 3. UV-vis spectra of $[\text{Mo}_3\text{WS}_4(\text{H}_2\text{O})_{12}]^{5+}$ (—), $[\text{Mo}_2\text{W}_2\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$ (- - -) and $[\text{MoW}_3\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$ (···) in 2 M Hpts. The inset gives reduction potentials (E^0) for $[\text{Mo}_x\text{W}_{4-x}\text{S}_4(\text{H}_2\text{O})_{12}]^{6+/5+}$ and $[\text{Mo}_4\text{W}_{4-x}\text{S}_4(\text{H}_2\text{O})_{12}]^{5+/4+}$ couples.¹¹

$(\text{H}_2\text{O})_{12}]^{n+}$ ($n = 4, 5,$ and 6),^{8,11} with $[\text{Mo}_3\text{CrS}_4(\text{H}_2\text{O})_{12}]^{4+}$ an exception.¹² Reduction potentials (vs NHE) for the $6+/5+$ and $5+/4+$ couples of $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{n+}$ are 860 and 210 mV. Although $[\text{W}_4\text{S}_4(\text{CN})_{12}]^{6-/7-}$ have been prepared,⁷ $[\text{W}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{n+}$ cubes have not, so far, been obtained. Details of the UV-vis spectrum ($n = 5$) can be predicted (Figure 3). Also, by extrapolation of data for $[\text{Mo}_x\text{W}_{4-x}\text{S}_4(\text{H}_2\text{O})_{12}]^{n+}$ ($x = 1-4$), reduction potentials of 39 and -627 mV, respectively, are obtained (inset, Figure 3),¹¹ which are 821 and 833 mV more reducing than those for the corresponding Mo_4S_4 couples. The 6:1 corner-shared double cube $[\text{Mo}_7\text{S}_8(\text{H}_2\text{O})_{18}]^{8+}$ is the only example with a transition metal at the nodal position.¹³

For type B with Groups 8–11, 3:1 single cubes are obtained.³ Of these, only the Cu cube exhibits more than one oxidation state, $\text{Mo}_3\text{CuS}_4^{4+}$ and $\text{Mo}_3\text{CuS}_4^{5+}$.¹⁴ The latter is a derivative formed by addition of Cu^I (d^{10}) in a process similar to that of In^I or Sn^{II} .^{15,16} The single cubes incorporating Fe, Co, Ni, Pd, and Cu ($4+$ products) have tetrahedrally coordinated heteroatoms, which for $M' = \text{Co}, \text{Pd},$ and Cu give edge-linked double cubes, $\{[\text{Mo}_3\text{M}'\text{S}_4(\text{H}_2\text{O})_9]_2\}^{8+}$, (1), confirmed by X-ray crystallography.²⁻⁴ The



two μ_4 -S-to- M' bridges formed give a rhombic $M'_2\text{S}_2$ arrangement. These are particularly well characterized in the Pd case, which is air stable (Figure 4).¹⁷ No edge-linked forms have been detected for $[\text{Mo}_3\text{M}'\text{S}_4(\text{H}_2\text{O})_{10}]^{4+}$ ($M' = \text{Fe}, \text{Ni}$), but they are observed with $[\text{Mo}_x\text{W}_{3-x}\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ ($x = 0, 1$).¹⁸ The Ni of $\text{Mo}_3\text{NiS}_4^{4+}$ is also capable of expanding its coordination to give $[\text{Mo}_3(\text{NiL})\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ with the tridentate ligand $L = 1,4,7$ -triazacyclononane.¹⁹ In the case $M' = \text{Co}$, the high air-sensitivity of the edge-linked double cube has made it difficult to confirm the existence of the single cube.²⁰ With $M' = \text{Rh}$ or Pt , the




FIGURE 4. UV-vis spectra ϵ 's per Mo_3 : (A) purple $\{[\text{Mo}_3\text{PdS}_4(\text{H}_2\text{O})_9]_2\}^{8+}$, (B) blue single cube $[\text{Mo}_3(\text{PdCl})\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ on addition of a 10-fold excess of Cl^- , and (C) on addition of 1 M Cl^- .³²





FIGURE 5. Conversion of a single cube $[\text{Mo}_3\text{SnS}_4(\text{H}_2\text{O})_{12}]^{6+}$ to the corresponding double cube by reductive addition.

heteroatom is inert, and for Rh at least octahedral coordination seems likely.^{2,21}

Finally, for type C (Groups 12–15), single cubes are formed by direct addition of Ga^I , In^I , Ge^{II} , and Sn^{II} to $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$, where the heteroatom is octahedral and products have $5+$ or $6+$ charges. Corner-shared 6:1 double cubes are obtained with Hg, In, Tl, Ge, Sn, Pb, As, Sb, and Bi (but not Ga!), all of which are $8+$ in charge.²² A redox-induced interconversion of single and double cubes is observed, (2), with double-cube formation occurring in a




reductive addition step (Figure 5). The above studies have been extended to include $[\text{Mo}_4\text{Se}_4(\text{H}_2\text{O})_{12}]^{n+}$, the incomplete cube $[\text{Mo}_3\text{Se}_4(\text{H}_2\text{O})_9]^{4+}$, and derivatives of the latter.^{8,23} Whereas $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ is extremely stable, with $[\text{Mo}_3\text{Se}_4(\text{H}_2\text{O})_9]^{4+}$ there is a tendency for Se to deposit within days as a result of air oxidation, $\text{Se}^{2-} \rightarrow \text{Se}$.

Preparation and Properties


The driving force for the incorporation of heteroatoms into $\text{M}_3\text{Q}_4^{4+}$ cores ($\text{Q} = \text{S}, \text{Se}$) stems from two sources. Affinity of the heterometal M' for μ_2 -chalcogen atoms (chalcophilicity) is required. The heterometal must also serve as an

electron donor, which is why incorporation requires either M^0 or a low oxidation state. Nonreducing Pb^{II} , Tl^{I} , and Bi^{III} are not incorporated, despite their chalcophilicity. The $\text{Mo}_3\text{PbS}_4^{6+}$ and $\text{Mo}_3\text{BiS}_4^{7+}$ cubes have been prepared in nonaqueous solutions, with negatively charged thiophosphate ester ligands.²⁴ The reactivity of the host cluster is also important: $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ is the most inert and does not incorporate Hg, but $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$, $[\text{Mo}_3\text{Se}_4(\text{H}_2\text{O})_9]^{4+}$, and $[\text{W}_3\text{Se}_4(\text{H}_2\text{O})_9]^{4+}$ give 6:1 products.²⁵ The reactivity of the heterometal in its elementary form is not always very high, and reactions can take several hours or even days for completion. With M^{II} transition metals ($\text{M}' = \text{V}$, Mn , Fe , Co , Ni , Cu , Zn , Cd , Pd , Pt), or with post-transition metals in their higher oxidation states, e.g., Ga^{III} , In^{III} , Ge^{IV} , and Sn^{IV} , no heterometallic cube formation is observed. However, the $\text{Mo}_3\text{M}'\text{S}_4^{4+}$ clusters in addition of heterometal cations to $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ in the presence of BH_4^- . This method is effective in the preparation of $[\text{Mo}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ in 2 M HCl (3)¹⁹ and complete within a few minutes, as compared to ~ 7 days with the metal.



Other, more specific preparative details are now considered. Purification is by O_2 -free (N_2) Dowex cation-exchange chromatography, at which stage different acids can be introduced. In HCl solution, heterometallic single cubes from $[\text{M}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$ ($\text{M} = \text{Mo}$, W ; $\text{Q} = \text{S}$, Se) undergo Cl^- complexing, M' generally $> \text{M}$. The less complexing perchloric acid and *p*-toluenesulfonic acid (Hpts) can be used, but no elution of 8+ double cubes occurs with HClO_4 . With 4 M Hpts, the pts^- complexes at a low level, elution occurs, and an H-bonded $\text{H}^+/\text{H}_2\text{O}/\text{pts}^-$ network induces crystallization of the aqua ions.

Type A. In the reaction of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ with $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$, the Cr^{II} is able to reduce $\text{Mo}_3\text{S}_4^{4+}$ with Cr^{II} uptake (4).¹² Similarly, by using $\text{K}_4[\text{Mo}_2\text{Cl}_8]$ as a source of Mo^{II} ,



reaction with $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$, $[\text{W}_3\text{Se}_4(\text{H}_2\text{O})_9]^{4+}$, $[\text{Mo}_2\text{WS}_4(\text{H}_2\text{O})_9]^{4+}$, and $[\text{MoW}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ occurs, with Mo^{III} a product (5).¹¹ Again, an excess of reducing agent is



required to give prior reduction of the trinuclear species. In some cases with $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and V , Mg , Al , or $\text{H}_3\text{-PO}_2$ as reductant, fragmentation occurs with reassembly into $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$ and $[\text{Mo}_7\text{S}_8(\text{H}_2\text{O})_{18}]^{8+}$. The yields of these two products with H_3PO_2 are $>30\%$ and $\sim 20\%$, respectively, after 3 days at 20°C .¹³

Type B. Direct reaction of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ with Fe , Co , Ni , Cu , or Pd (as Pd black)^{2,3} takes place in 2 M HCl solutions. The metals are incorporated formally as tetrahedral M^0 , leading to $\text{Mo}_3\text{M}'\text{S}_4^{4+}$ and edge-linked products.²² In the Cu case, Dowex chromatography gives $\text{Mo}_3\text{CuS}_4^{4+}$ as product.¹⁴ An interesting exception is the incorporation of nonreducing $\text{Rh}(\text{III})$ by reaction of Cp^*RhCl_2 (Cp^* is $\eta^5\text{-C}_5\text{Me}_5$) with $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$.²¹ Ad-





FIGURE 6. UV-vis spectra of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ (---), $[\text{Mo}_3\text{InS}_4(\text{H}_2\text{O})_{12}]^{5+}$ (—) (ϵ 's per Mo_3), and $[\text{Mo}_6\text{InS}_8(\text{H}_2\text{O})_{18}]^{8+}$ (-.-) (ϵ 's per Mo_6) in 2 M HCl.¹⁵

ducts are also formed with $[\text{PtCl}_4]^{2-}$ (in 2 M HCl) in the presence of H_3PO_2 , and on heating with RhCl_3 in 4 M HCl for 3 h. Analyses have confirmed 3:1 metal ratios, but more extensive characterization has proved difficult.² In the case of $\text{M}' = \text{Ni}$, Pd water-soluble products incorporating $\text{M}'(\text{CO})$,²⁶ $\text{Ni}(\text{C}_2\text{H}_4)$,¹⁸ and $\text{Pd}(t\text{-BuNC})$,¹⁷ have been prepared.

With $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$, both Cu^{I} and Cu metal give $\text{W}_3\text{-CuS}_4^{5+}$ (6), and the 4+ analogue has not, so far, been



prepared.²⁸ This illustrates the difficulty in preparing W_3 cubes in the more reduced 4+ state. Similarly, W_6 corner-shared double cubes are more difficult to prepare than their Mo_6 analogues.

It remains to be seen whether Re , Ru , Os , Ir , or indeed Ag , Au , and Zn can be incorporated into $[\text{M}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$.

Type C. The reaction of $\text{Mo}_3\text{S}_4^{4+}$ with In^{I} , or $\text{In}^{\text{III}} + \text{BH}_4^-$, to give the single cube $\text{Mo}_3\text{InS}_4^{5+}$ suggests direct addition. Metallic In or Ga also incorporates as Ga^+ or In^+ to form $[\text{Mo}_3\text{M}'\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$,¹⁵ where the presence of M^+ cations on the metal surface is possible. There is no evidence for $[\text{Mo}_3\text{InS}_4(\text{H}_2\text{O})_{12}]^{4+}$, and In metal reacts with $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ in 0.1 M Hpts to give the corner-shared product $[\text{Mo}_6\text{InS}_8(\text{H}_2\text{O})_{18}]^{8+}$ (Figure 6). Similarly, from $[\text{Mo}_3\text{S}_3\text{O}(\text{H}_2\text{O})_9]^{4+}$, the product $[\text{Mo}_6\text{InO}_2\text{S}_6(\text{H}_2\text{O})_{18}]^{8+}$ has been obtained, which is a comparatively rare example of oxosulfido trinuclear participation. Incorporation of Hg , Tl , Sn , and Sb metals into the nodal position of double cube $[\text{Mo}_6\text{M}'\text{S}_8(\text{H}_2\text{O})_{18}]^{8+}$ is also observed,^{2,3,29} but $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ reacts with Sn only, and the single-cube product $[\text{W}_3(\text{SnCl}_3)\text{S}_4(\text{H}_2\text{O})_9]^{3+}$ is formed in HCl.³⁰ The heteroatoms of single cubes react with complexing anions, e.g., Cl^- (Figure 7).

With In^{III} , Tl^{I} , Pb^{II} , and Bi^{III} in the presence of BH_4^- , the double cubes $[\text{Mo}_6\text{M}'\text{S}_8(\text{H}_2\text{O})_{18}]^{8+}$ ($\text{M}' = \text{In}$, Tl , Pb , Bi) are formed. However, with Ga^{III} under similar conditions, only the single cube $[\text{Mo}_3\text{GaS}_4(\text{H}_2\text{O})_{12}]^{5+}$ is obtained,¹⁵ and no double cube has been identified, which is unique for




FIGURE 7. UV-vis spectra of $[\text{Mo}_3\text{SnS}_4(\text{H}_2\text{O})_{12}]^{6+}$ (ϵ 's per Mo_3) in 2.0 M acids Hpts (—), HClO_4 (---), and H_2O (— · —).¹⁶





FIGURE 8. Summary of reactions of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and Sn heterometallic products.²³



type C heteroatoms. A procedure involving conversion of single (In, Ge, Sn) to double cubes in the presence of BH_4^- ($\text{M} = \text{Mo}, \text{W}$) (7) is used as the only route to prepare $[\text{W}_6-$



$\text{SnS}_8(\text{H}_2\text{O})_{18}]^{8+}$ and gives approaching quantitative yields. Similarly, $[\text{W}_6\text{GeS}_8(\text{H}_2\text{O})_{18}]^{8+}$ and $[\text{Mo}_6\text{SnSe}_8(\text{H}_2\text{O})_{18}]^{8+}$ have been prepared, and the synthesis of mixed-metal $[\text{Mo}_3\text{W}_3\text{SnS}_8(\text{H}_2\text{O})_{18}]^{8+}$ and mixed-chalcogen $[\text{Mo}_6\text{SnS}_4\text{Se}_4(\text{H}_2\text{O})_{18}]^{8+}$ products has also been achieved.⁹ It should, however, be noted that (7) occurs much less readily with W than with Mo. A summary of the chemistry of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ with Sn is given in Figure 8.

One drawback to using BH_4^- is the fast decomposition with H^+ ($\text{pH} \leq 1$), leading to a loss of BH_4^- , initially in ~ 100 -fold excess. Vigorous H_2 gas evolution and heating effects are difficult to control, and the heterometal is sometimes precipitated as, e.g., metal boride. The use of hypophosphorous acid instead of BH_4^- gives better results, since H_3PO_2 is stable in acidic conditions. The reduction potential for the reaction of H_3PO_2 is a large negative value, but kinetically the reactions are slower. It has been


used in many cases, notably to prepare $[\text{Mo}_6\text{AsS}_8(\text{H}_2\text{O})_{18}]^{8+}$ with NaAsO_2 or Na_3AsO_4 as the As source.²² In the case of In^{III} , formation of $[\text{Mo}_3\text{InS}_4(\text{H}_2\text{O})_{12}]^{5+}$ precedes $[\text{Mo}_6\text{InS}_8(\text{H}_2\text{O})_{18}]^{8+}$ (8, 9).¹⁵



Several attempts have been made to incorporate early transition metals such as Ti and V as well as lanthanoids and actinoids (e.g., the strongly reducing Eu^{2+} and U^{3+}), so far without success.^{2,13} This appears to corroborate the need for some heterometal chalcogen affinity to stabilize the heterometallic product, especially in water.

Molecular and Electronic Structures

Simple principles help to rationalize the cluster types considered herein. The structures are based on single cubes with an $\text{M}_3\text{M}'\text{Q}_4$ core. The M_3Q_4 unit retains features present in the incomplete cube. Indeed, in most oxidation and heteroatom transfer processes, re-formation of $\text{M}_3\text{Q}_4^{4+}$ is observed. The tendency of both components to retain their initial coordination³⁰ is illustrated in (10). When M'



is type C, both M' and M have octahedral coordination, but M' has a preference for Cl^- over H_2O , and products, e.g., $[\text{W}_3(\text{SnCl}_3)\text{S}_4(\text{H}_2\text{O})_9]^{3+}$, are obtained with $[\text{Cl}^-] \geq 0.05$ M. Hence, it is easy to replace the three ligands to M' by three chalcogen atoms of a second M_3Q_4 , as in (7). In the case of Group 12, the single and double cubes $\text{Mo}_3\text{HgS}_4^{4+}$ and $\text{Mo}_6\text{HgS}_8^{8+}$ should interconvert in the presence of $\text{Mo}_3\text{S}_4^{4+}$ without any need for redox. The single cube $\text{Mo}_3\text{HgS}_4^{4+}$ has not so far been identified, but is very likely the initial product from the reaction of $\text{Mo}_3\text{S}_4^{4+}$ with Hg^0 .

The tetrahedral B heteroatoms Co, Ni, Pd, and Cu have affinity for the softer ligands. When no such ligands are present, dimerization occurs by interaction of the heterometal with a $\mu_3\text{-Q}$ of a second cube, and edge-linked double cubes result. An interesting consequence of dimerization is the close contact, giving $\text{M}'\text{-M}'$ bonding. In the Co case, the Co-Co distance is 2.498 Å.³ The corresponding $\text{M}'\text{-M}'$ distances in the Cu and Pd edge-linked double cubes are 2.426 Å (Cu)³ and 2.790 Å (Pd).¹⁷ For $[\text{Mo}_x\text{W}_{3-x}\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$, distances are 2.549 Å ($x = 1$) and 2.560 Å ($x = 0$).¹⁸

Many heterometal cubes have Mo-Mo distances close to the 2.732 Å value of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9](\text{pts})_4 \cdot 9\text{H}_2\text{O}$,⁸ which is diamagnetic and has a triangular arrangement of three metal-metal bonded Mo^{IV} (d^2) atoms. The type C $[\text{Mo}_6\text{M}'\text{S}_8(\text{H}_2\text{O})_{18}]^{8+}$ cubes have an 8+ charge in all cases, which suggests that two $\text{Mo}_3\text{S}_4^{4+}$ units are retained as a common feature and are bridged by M^0 . The Mo-M' separations are ~ 1 Å longer than Mo-Mo, and no Mo-M' metal-metal bonding is apparent.²² Formally, the double cubes can be written as $(\text{Mo}_3\text{S}_4^{4+})_2\text{M}^0$ and the single cubes (also with long Mo-M') as $\text{Mo}_3\text{S}_4^{4+}\text{M}'$ ($\text{M}' = \text{Ga}, \text{In}$) and




FIGURE 9. Interaction of two cucurbituril $\text{C}_{36}\text{H}_{36}\text{N}_{24}\text{O}_{12}$ molecules with the double cube $[\text{Mo}_6\text{HgSe}_8(\text{H}_2\text{O})_{14}\text{Cl}_4]^{4+}$ in a supramolecular assembly.²⁵

$\text{Mo}_3\text{S}_4^{4+}\text{M}'^{2+}$ ($\text{M}' = \text{Ge}, \text{Sn}$). Furthermore, when M' is a transition metal (e.g., Cr, Fe, Co, Ni, Cu, or Pd),²² single cubes of 4+ charge are obtained, and the heteroatom M' can be designated M^0 . The short $\text{Mo}-\text{M}'$ bond lengths for type A and type B cubes provide evidence for $\text{Mo}-\text{M}'$ bonding. Quantum chemical calculations provide an explanation of this sharp division.³¹ The energy level of the d electrons of the transition metal is close to the LUMO of the M_3Q_4 fragment. This makes formation of covalent $\text{M}'-\text{M}$ bonds possible. In contrast, the post-transition metals have only high-energy s or p orbitals available, and instead of $\text{M}'-\text{M}$ bonding the electron density is transferred to the M_3Q_4 core.

In the case of $[\text{Mo}_7\text{S}_8(\text{H}_2\text{O})_{18}]^{8+}$, each cube has three short $\text{Mo}-\text{Mo}$ (2.770 Å) and three long (3.046 Å) $\text{Mo}-\text{Mo}'$ (nodal) separations.³ Here the $\text{Mo}-\text{Mo}'$ bond appears less strong. A structure approximating to $(\text{Mo}_3\text{S}_4^{4+})_2\text{Mo}^0$, with adjacent Mo 's assigned as Mo^{IV} and Mo^0 , is clearly less satisfactory, and an intermediate oxidation state may apply.

Characterization of the cubes by X-ray crystallography is not always possible, and in many cases crystallization has only been achieved as pts^- salts of the aqua ions $[\text{Mo}_6\text{M}'\text{S}_8(\text{H}_2\text{O})_{18}](\text{pts})_8 \cdot x\text{H}_2\text{O}$.^{3,22} Recently, however, crystals of cucurbituril adducts have been prepared. Cucurbituril ($\text{C}_{36}\text{H}_{36}\text{N}_{24}\text{O}_{12}$) is a macrocyclic cavitand molecule of D_{6h} symmetry, having two identical carbonyl-fringed portals. These can H-bond with coordinated water molecules to give supramolecular aggregates. Crystals of $[\text{M}_6\text{HgQ}_8(\text{H}_2\text{O})_{14}\text{Cl}_4](\text{C}_{36}\text{H}_{36}\text{N}_{24}\text{O}_{12})\text{Cl}_4 \cdot 14\text{H}_2\text{O}$ ($\text{M} = \text{Mo}, \text{W}; \text{Q} = \text{S}, \text{Se}$) have been prepared by reaction of $[\text{M}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$ with metallic Hg in 4 M hydrochloric acid and crystallization in the presence of cucurbituril.²⁵ These have cluster cations $[\text{M}_6\text{HgQ}_8(\text{H}_2\text{O})_{14}\text{Cl}_4]^{4+}$ (with Cl^- replacing H_2O ligands) sandwiched between two cucurbituril units (Figure 9). The $\text{Hg}-\text{S}$ distances are unusually long, with two at 2.768(2) Å and four longer at 2.903(2) Å, in good agreement with the value of 2.84(12) Å reported for $[\text{Mo}_6\text{HgS}_8(\text{H}_2\text{O})_{18}](\text{pts})_8 \cdot 20\text{H}_2\text{O}$.³ The complex $[\text{Hg}(\text{[9]-aneS}_3)_2]^{2+}$ is the closest structural analogue, with six $\text{Hg}-\text{S}$ bonds varying between 2.638(3) and 2.728(3) Å, which seems to be the upper limit for $\text{Hg}^{\text{II}}-\text{S}$ bonds. It is concluded²⁵ that, in the double cube, Hg retains a large part of its formal Hg^0 oxidation state.





FIGURE 10. Structure of $[\text{W}_3\text{Se}_4(\text{CN})_9]^{5-}$ anions showing $\text{Se} \cdots \text{Se}$ interactions between two adjacent clusters.³¹

One of the most striking features of the cucurbituril structures is that they are isostructural with assemblies incorporating a dimeric trinuclear unit $[\text{W}_3\text{Se}_4(\text{H}_2\text{O})_8\text{Cl}]_2 \cdot (\text{C}_{36}\text{H}_{36}\text{N}_{24}\text{O}_{12})\text{Cl}_6 \cdot 12\text{H}_2\text{O}$.²⁵ Here, two W_3Se_4 units are held together by $\text{Se} \cdots \text{Se}$ interactions (3.617–3.740 Å) and are embedded between the cucurbituril molecules ($\text{O} \cdots \text{O}$, 2.719–2.779 Å) in exactly the same way as the heterometallic double cubes. Similar dimeric aggregates have been reported previously for $[\text{W}_3\text{Se}_4(\text{NCS})_9]^{5-}$ and $[\text{W}_3\text{Se}_4(\text{CN})_9]^{5-}$ (Figure 10), as well as for other M_3Q_4 ($\text{M} = \text{Mo}, \text{W}; \text{Q} = \text{S}, \text{Se}$) derivatives.³² Moreover, in binary MQ_2 chalcogenides of the early transition metals, the chalcogen atoms of the neighboring layers interact with each other in the same way. Metal atoms and small molecules can intercalate between these layers. Thus, the structures are formally related to those of a matrix and of an intercalate. This means that the two W_3Se_4 trinuclear units, sandwiched between two cucurbituril molecules, mimic rather closely two neighboring layers $\text{QM}_3 \cdots \text{QM}_3$ in solid MQ_2 (with the M_3 units as building blocks). These results show that solid-state chemistry is related to the chemistry of small molecular clusters in a more intimate way than might have been suspected.

Stability in Air

Single and double cubes are, in almost all cases, O_2 sensitive, and rigorous air-free techniques are required. With O_2 , the cubes re-form the trinuclear incomplete cube. However, both the single $[\text{Mo}_3(\text{PdCl})\text{S}_4(\text{H}_2\text{O})_9]^{3+}$ and double $[\{\text{Mo}_3\text{PdS}_4(\text{H}_2\text{O})_9\}_2]^{8+}$ cubes are air stable over long periods. Also, $[\text{Mo}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ is one of the more stable cubes in air and shows no decay over periods of ~1 h. Heterometallic cubes from $[\text{W}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$ ($\text{Q} = \text{S}, \text{Se}$) are less stable in air. The more tungsten atoms included, the more air-sensitive the cluster becomes. Thus, for the cubes $[\text{Mo}_x\text{W}_{4-x}\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$,¹¹ with $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ (or O_2) as oxidant, the 6+ cube is first obtained, which fragments with loss of W exclusively as a high oxidation state product (11).




Heteroatom-Transfer Reactions

Transfer²² of Sn^{II} is observed in the reaction of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ with $[\text{Mo}_3\text{SnSe}_4(\text{H}_2\text{O})_{12}]^{6+}$, $[\text{W}_3\text{SnSe}_4(\text{H}_2\text{O})_{12}]^{6+}$, and $[\text{W}_3\text{SnS}_4(\text{H}_2\text{O})_{12}]^{6+}$, and the reactions can be written as in (12). No corresponding reactions of $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$



with $[\text{W}_3\text{SnSe}_4(\text{H}_2\text{O})_{12}]^{6+}$ or of $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and $[\text{W}_3\text{Se}_4(\text{H}_2\text{O})_9]^{4+}$ with $[\text{Mo}_3\text{SnSe}_4(\text{H}_2\text{O})_{12}]^{6+}$ are observed. The reaction of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ with $[\text{Mo}_3\text{SnSe}_4(\text{H}_2\text{O})_{12}]^{6+}$ gives an equilibrium mix,²³ with $K = 4.0(9)$ for (13). Affinities of




the different aqua ions for Sn^{II} are $\text{Mo}_3\text{S}_4^{4+} > \text{Mo}_3\text{Se}_4^{4+} > \text{W}_3\text{Se}_4^{4+} > \text{W}_3\text{S}_4^{4+}$.

Displacement of the heteroion In^{I} by Sn^{II} occurs, and the affinity $\text{Sn}^{\text{II}} > \text{In}^{\text{I}}$ for $\text{M}_3\text{Q}_4^{4+}$ ($\text{M} = \text{Mo}, \text{W}; \text{Q} = \text{S}, \text{Se}$) has been established (14). Reactions occur by re-equili-




bration (15) and redistribution of In^{I} between the tri-



nuclear species.

The above mechanism holds for post-transition elements. No reaction between $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$ and $\text{Mo}_3\text{FeS}_4^{4+}$, or $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ and $\text{Mo}_3\text{NiS}_4^{4+}$, is observed, and there is no evidence for equilibria as in (15). However, Cu^{2+} gives a net displacement of heteroatoms from $\text{Mo}_3\text{FeS}_4^{4+}$ and $\text{Mo}_3\text{NiS}_4^{4+}$, which has been explained by a redox mechanism involving intermediate formation of small amounts of Cu^+ (16), followed by (17).




Reactions with H^+

Group 13 (Ga, In, Tl) heterometallic cube¹⁵ derivatives of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ are oxidized by H^+ in HCl, giving H_2 , which can be determined by gas chromatography. The reaction is unique to Group 13 derivatives. Thus, the Tl-containing double cube $[\text{Mo}_6\text{TlS}_8(\text{H}_2\text{O})_{18}]^{8+}$ reacts with H^+ according to (18). However, the In-containing double



cubes $[\text{Mo}_6\text{InS}_8(\text{H}_2\text{O})_{18}]^{8+}$ and $[\text{Mo}_6\text{InO}_2\text{S}_6(\text{H}_2\text{O})_{18}]^{8+}$ react with H^+ with formation of the single cube, e.g., (19). The



reaction of $\text{Mo}_3\text{InS}_4^{5+}$ (and $\text{Mo}_3\text{GaS}_4^{5+}$) with 2 M HCl is slow, and no H_2 was detected by GC after 2 d. However, in 4 M HCl, reaction according to (20) is observed. No H_2 was detected in 2 M Hpts. Chloride coordination to the



heterometal may catalyze the oxidation by H^+ . Kinetic studies (25 °C) on the reactions of H^+ with $[\text{Mo}_6\text{InO}_2\text{S}_6(\text{H}_2\text{O})_{18}]^{8+}$ and $[\text{Mo}_6\text{TlS}_8(\text{H}_2\text{O})_{18}]^{8+}$ give second-order rate constants $k_{\text{H}} = 4.9 \times 10^{-3}$ (In) and $0.25 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$ (Tl),¹⁵ $I = 2.00 \text{ M}$ (pts⁻).


Substitution at Heteroatoms

Reactions involving the substitution of H_2O by L at the Ni of $[\text{Mo}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ have been studied.¹⁹ Formation rate constants for $\text{L} = \text{Cl}^-$, Br^- , I^- , and NCS^- lie in the narrow range $9.4\text{--}44 \text{ M}^{-1} \text{ s}^{-1}$. Such behavior is interpreted in terms of a dissociative interchange I_d mechanism (21). Inclusion of two water-soluble phosphines, 1,3,5-triaza-




7-phosphaadamantane (PTA) and tris(3-sulfonatophenyl)-phosphine ($\text{P}(\text{C}_6\text{H}_4\text{SO}_3)_3^{3-}$), requires a 3-fold extension of this range, but for CO k_{f} is 10 times smaller. It is possible to explain these results while retaining essential features of an I_d process.¹⁹ Rate constants are ca. 10^3 times smaller than those for water exchange on $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$, with an even bigger factor for tetrahedral Ni^{II} . Rate constants for substitution on tetrahedral $[\text{Ni}(\text{CO})_4]$ ($2.0 \times 10^{-2} \text{ M}^{-1} \text{ s}^{-1}$ in toluene) are smaller, and the behavior observed is therefore in the Ni^{0} to Ni^{II} range, with Ni^{0} the more likely if the three $\mu_3\text{-S}$ ligands bring about a stabilizing effect.

The conversion³² of edge-linked $[\{\text{Mo}_3\text{PdS}_4(\text{H}_2\text{O})_9\}_2]^{8+}$ to $[\text{Mo}_3(\text{PdL})\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ is induced by $\text{L} = \text{CO}$, the phosphines PTA and $\text{P}(\text{C}_6\text{H}_4\text{SO}_3)_3^{3-}$, Cl^- , Br^- , and NCS^- , all of which complex at the tetrahedral Pd (22). Cleavage



of an intercube Pd- $\mu_3\text{-S}$ bond occurs in the rate-determining step involving one L. Rate constants $k_{\text{f}}/\text{M}^{-1} \text{ s}^{-1}$ are CO, 1.11×10^5 ; PTA, 27.8×10^5 ; and $\text{P}(\text{C}_6\text{H}_4\text{SO}_3)_3^{3-}$, 9.6×10^5 , where the products have strong colors. The reactions are independent of $[\text{H}^+] = 0.30\text{--}2.0 \text{ M}$. The first stages with $\text{L} = \text{Cl}^-$, Br^- , and NCS^- are too fast to monitor, but equilibrium formation constants K/M^{-1} are Cl^- , 490; Br^- , 8040; and NCS^- , 630. The results indicate a preference for L as compared to H_2O , which is unable to retain the single-cube structure. Subsequent steps have been assigned to substitution at the Mo's, and a $[\text{NCS}^-]$ -independent step has been assigned to the isomerization $\text{Pd-NCS} \rightarrow \text{Pd-SCN}$.³²

Aqueous solution reactions of CO with $[\text{Mo}_3\text{M}'\text{S}_4(\text{H}_2\text{O})_{10}]^{4+}$ ($\text{M}' = \text{Co}, \text{Ni}, \text{Pd}$) have been compared (23).²⁶ Electron counts are 15 for the single cube Mo_3Co and 16




for Mo_3Ni and Mo_3Pd . On bubbling N_2 through the solutions, no reverse reaction is detected. Similar reactions are observed for $[\text{Mo}_3\text{NiSe}_4(\text{H}_2\text{O})_{10}]^{4+}$ and $[\text{W}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$. No reaction is observed for $[\text{Mo}_3\text{FeS}_4(\text{H}_2\text{O})_{10}]^{4+}$ (14 electrons) and $[\text{Mo}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{4+}$ (17 electrons). However, the

higher oxidation state cube $[\text{Mo}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{5+}$ (16 electrons) reacts with CO. The reaction is different from (23) in that $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and $\text{Cu}(\text{CO})^+$ are formed (24). On



bubbling of N_2 through the solutions, the $\text{Cu}(\text{CO})^+$ decomposes with release of CO (25), and the Cu^+ recombines with $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ (26).



Uptake of ethylene by $[\text{M}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ ($\text{M}_3 = \text{Mo}_2\text{W}$, MoW_2 , W_3) in aqueous or organic solvents to give olefin π -complexes $[\text{Mo}_3(\text{NiC}_2\text{H}_4)\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ has been reported and is as expected for Ni^0 .¹⁸ The equilibration of $[\text{Mo}_3\text{-NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ with ethylene is not as extensive as for the tungsten cluster, and in solvent H_2O (for CH_2Cl_2) the reaction is readily reversible.

Substitution of H_2O by NCS^- at the Cr of $[\text{Mo}_3\text{CrS}_4(\text{H}_2\text{O})_{12}]^{4+}$ gives a rate law $k[\text{Mo}_3\text{CrS}_4^{4+}][\text{NCS}^-]$, with $k(25^\circ\text{C}) = 37 \text{ M}^{-1} \text{ s}^{-1}$ in 2.00 M Hpts.¹² The magnitude of k suggests a Cr^{III} rather than Cr^{II} assignment, with the three μ_3 -S ligands to Cr^{III} having a stabilizing effect. The dependence of k on $[\text{H}^+]^{-1}$ indicates a conjugate-base pathway. For such a mechanism, the Cr requires at least two H_2O ligands; hence, a six-coordinate Cr and the formula $[\text{Mo}_3\text{-CrS}_4(\text{H}_2\text{O})_{12}]^{4+}$ are proposed.

At 25°C , the formation constant K for 1:1 Cl^- complexing to $[\text{Mo}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{4+}$ ($K = 3500 \text{ M}^{-1}$) compares with a value ($K = 500 \text{ M}^{-1}$) for the complexing of Cl^- to Cu^+ .¹⁴ The reaction of Cl^- with $[\text{Mo}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{4+}$ is assigned as rapid substitution at the tetrahedral Cu. Similarly, the 1:1 reactions (25°C) of Cl^- with $[\text{Mo}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ ($K = 106 \text{ M}^{-1}$) and $[\text{Mo}_3\text{FeS}_4(\text{H}_2\text{O})_{10}]^{4+}$ ($K = 560 \text{ M}^{-1}$) are fast and in the stopped-flow range. However, in the case of $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{4+}$, 1:1 complexing is much less favorable ($K = 1.98 \text{ M}^{-1}$), and a substantially smaller rate constant of $9.7 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$ is observed.⁸

The only example of kinetic studies on the uptake of a heteroatom is for the reactions of Cu^{I} with $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$.²⁷

Redox Reactions

Studies on the oxidation of single and double cubes with the 2,6-dicarboxylatopyridine (dipicolinate) Co^{III} complex $[\text{Co}(\text{dipic})_2]^-$ (E° for $\text{Co}^{\text{III}}/\text{Co}^{\text{II}}$, 362 mV) or hexaaquairon(III) ($E^\circ \approx 770 \text{ mV}$), stoichiometries (27, 28),



have been carried out. Rate laws first-order in oxidant and cube indicate a rate-controlling first stage followed by fast steps. The reactions of $[\text{Co}(\text{dipic})_2]^-$ are independent of $[\text{H}^+]$ in the range 0.5–2.0 M and are assigned as outer-sphere electron transfer. In the case of the corner-shared

double cubes, intermediate formation of the single cubes can be observed. With stoichiometric amounts of oxidant, the single cube can be isolated in high yield for $\text{M}' = \text{In}$, Sn , and Ge .^{15,16} Relevant stoichiometries for Sn and Ge are as in (29), where the single cube reacts with two further



Co^{III} to give $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$. Oxidation of $[\text{Mo}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{4+}$ with $[\text{Co}(\text{dipic})_2]^-$ has two stages with intermediate formation of $[\text{Mo}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{5+}$ (30, 31).¹⁴



The 5+ oxidation state has recently been confirmed by X-ray crystallography.³³ In pts^- solution, disproportionation of the 5+ ion to give $\text{Mo}_3\text{CuS}_4^{4+}$, $\text{Mo}_3\text{S}_4^{4+}$, and Cu^{II} is observed. No similar process is observed with $\text{W}_3\text{-CuS}_4^{5+}$.²⁷

Virtually all heteroatom derivatives of $[\text{M}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$ react with O_2 . In the absence of electrochemical information on reduction potentials, and considering the difficulty in carrying out mechanistic studies with O_2 , rate constants for the oxidation with $[\text{Co}(\text{dipic})_2]^-$ (k_{Co}) and $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ (k_{Fe}) can be informative. There is little or no $[\text{Co}(\text{dipic})_2]^-$ oxidation of $[\text{Mo}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ and $[\text{Mo}_3(\text{PdCl})\text{S}_4(\text{H}_2\text{O})_9]^{3+}$, which are least reactive with O_2 . Linear free energy plots of $\log k_{\text{Co}}$ vs $\log k_{\text{Fe}}$, where k_{Fe} is for the $[\text{H}^+]$ -independent reaction of $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$, have been reported.²⁰

Future Trends

At present, the formation of heteroatom cubes of $\text{M}_3\text{Q}_4^{4+}$ with Group 6 metals, but nonparticipation of other early transition metals, is quite striking. Cyclopentadienyl (Cp) $[\text{M}'_4\text{S}_4(\text{Cp})_4]$ cubes have been prepared ($\text{M}' = \text{Ti}$, V , Cr , Mo),^{34,35} and incorporation of other $\text{M}'(\text{Cp})$ into $[\text{M}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$ may be possible.²¹ The self-assembly of trinuclear $\{\text{M}_3\text{Q}_7\text{Br}_7\}_x$ polymers by heating together the elements, and ready access to $[\text{M}_3\text{Q}_4(\text{H}_2\text{O})_9]^{4+}$ ($\text{M} = \text{Mo}$, W ; $\text{Q} = \text{S}$, Se),^{8,31} may prove to be a route to other clusters with N, P, As, etc. instead of the chalcogenides. A niobium cluster $[\text{Nb}_4(\mu_4\text{-O})(\mu_3\text{-Te})_4(\text{CN})_{12}]^{6-}$ has recently been prepared with the oxo anion at the center of the cube.³⁶ Determining the synthesis and structure of larger nitrogenase-like clusters remains a challenge. In the lanthanoid area, the face-sharing double cube $[\text{Yb}_6\text{S}_6(\text{SPh})_6(\text{py})_{10}]$ represents a new type of structure.³⁷

References

- Holm, R. H. Trinuclear Cuboidal and Heterometallic Cubane-Type Iron–Sulfur Clusters: New Structural and Reactivity Themes in Chemistry and Biology. *Adv. Inorg. Chem.* **1992**, *38*, 1–71.
- Saysell, D. M.; Sokolov, M. N.; Sykes, A. G. Heterometallic Cuboidal Complexes as Derivatives of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$: Interconversions of single and double cubes and related studies. *ACS Symp. Ser.* **1996**, *653*, 216–239.
- Shibahara, T. Cubane and incomplete cubane-type molybdenum and tungsten oxo/sulfido clusters. *Adv. Inorg. Chem.* **1991**, *37*, 143–173.

- (4) Hidai, M.; Kuwata, S.; Mizobe, Y. Synthesis and Reactivities of Cubane-Type Sulfido Clusters Containing Noble Metals. *Acc. Chem. Res.* **2000**, *33*, 46–52.
- (5) Fontecilla-Camps, J. C.; Ragsdale, S. W. Nickel–Iron–Sulfur Active Sites: Hydrogenase and CO Dehydrogenase. *Adv. Inorg. Chem.* **1999**, *47*, 304.
- (6) Smith, B. E. Structure, Function and Biosynthesis of the Metallo-sulfur Clusters in Nitrogenase. *Adv. Inorg. Chem.* **1999**, *49*, 159–218.
- (7) Fedin, V. P.; Kalinina, I. V.; Samsonenko, D. G.; Mironov, Y. V.; Sokolov, M. N.; Tkachev, S. V.; Virovets, A. V.; Podberezskaya, N. V.; Elsegood, M. R. J.; Clegg, W.; Sykes, A. G. Synthesis, Structure and Properties of Molybdenum and Tungsten Cyano Complexes with Cuboidal $\text{M}_4(\mu_3\text{-E})_4$ (M = Mo, W; E = S, Se, Te) Cores. *Inorg. Chem.* **1999**, *38*, 1956–1965.
- (8) Hernández-Molina, R.; Sykes, A. G. Chalcogenide-Bridged Cuboidal Clusters with M_4Q_4 (M = Mo, W; Q = S, Se, Te) Cores. *J. Chem. Soc., Dalton Trans. (Perspective)* **1999**, 3137–3148.
- (9) Fedin, V. P.; Sokolov, M. N.; Sykes, A. G. Preparation of Heterometallic Single Cubes $[\text{W}_3\text{MS}_4(\text{H}_2\text{O})_{12}]^{n+}$ (M = In, Ge or Sn) and the First Corner-Shared Double Cube $[\text{W}_6\text{SnS}_8(\text{H}_2\text{O})_{18}]^{8+}$ and $[\text{W}_3\text{-Mo}_3\text{SnS}_8(\text{H}_2\text{O})_{18}]^{8+}$ as Derivatives of $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$. *J. Chem. Soc., Dalton Trans.* **1996**, 4089–4092.
- (10) Shibahara, T.; Akashi, H.; Kuroya, H. Cubane-type $\text{Mo}_3\text{FeS}_4^{4+}$ Aqua Ion and X-ray Structure of $[\text{Mo}_3\text{FeS}_4(\text{NH}_3)_9(\text{H}_2\text{O})]\text{Cl}_4$. *J. Am. Chem. Soc.* **1986**, *108*, 1342–1343.
- (11) McLean, I. J.; Hernandez-Molina, R.; Sokolov, M. N.; Seo, M.-S.; Virovets, A. V.; Elsegood, M. R. J.; Clegg, W.; Sykes, A. G. Preparation, Structure and Properties of Three $[\text{Mo}_x\text{W}_{4-x}\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$ (x = 1–3) and $[\text{MoW}_3\text{Se}_4(\text{H}_2\text{O})_{12}]^{5+}$ Cuboidal Complexes Alongside $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$ and $[\text{Mo}_4\text{Se}_4(\text{H}_2\text{O})_{12}]^{5+}$. *J. Chem. Soc., Dalton Trans.* **1998**, 2557–2562.
- (12) Routledge, C. A.; Humanes, M.; Li, Y.-J.; Sykes, A. G. Preparation and Properties of the Heterometallic Cuboidal Complex $[\text{Mo}_3\text{CrS}_4(\text{H}_2\text{O})_{12}]^{4+}$: Comparisons with $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{4+}$. *J. Chem. Soc., Dalton Trans.* **1994**, 1275–1282.
- (13) Sokolov, M. N.; Coichev, N.; Moya, H. D.; Hernandez-Molina, R.; Borman, C. D.; Sykes, A. G. New Procedures for the Preparation of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$, $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$ and $[\text{Mo}_7\text{S}_8(\text{H}_2\text{O})_{18}]^{8+}$ and their Se Analogues: Redox and Substitution Studies on the Double Cube $[\text{Mo}_7\text{S}_8(\text{H}_2\text{O})_{18}]^{8+}$. *J. Chem. Soc., Dalton Trans.* **1997**, 1863–1869.
- (14) Nasreldin, M.; Li, Y.-J.; Mabbs, F. E.; Sykes, A. G. Preparation and Properties of the Heterometallic Cuboidal Cluster $[\text{Mo}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{5+}$ and Comparisons with $[\text{Mo}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{4+}$. *Inorg. Chem.* **1994**, *33*, 4283–4289.
- (15) Hernandez-Molina, R.; Fedin, V. P.; Sokolov, M. N.; Saysell, D. M.; Sykes, A. G. Preparation and Properties of Group 13 (Ga, In, Tl) Heterometallic Single and Corner-Shared Double Cube Derivatives of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and Related Studies. *Inorg. Chem.* **1998**, *37*, 4328–4334.
- (16) Varey, J. E.; Lamprecht, G. J.; Fedin, V. P.; Holder, A.; Clegg, W.; Elsegood, M. R. J.; Sykes, A. G. Interconversion and Reactivity of Two Heterometallic Tin-Containing Cuboidal Clusters from $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$: X-ray Structure of the Single Cube with an $\text{Mo}_3\text{-SnS}_4$ Core. *Inorg. Chem.* **1996**, *35*, 5525–5530.
- (17) Murata, T.; Mizobe, Y.; Gao, H.; Ishii, Y.; Wakabayashi, T.; Nakano, F.; Tanase, T.; Yano, S.; Hidai, M.; Echizen, I.; Nanikawa, H.; Motomura, S. Synthesis of Mixed-Metal Sulfide Cubane-Type Clusters with the Novel PdMo_3S_4 Core and Reactivities of the Pd with Alkenes, CO, tBuNC , and Alkynes. *J. Am. Chem. Soc.* **1994**, *116*, 3389–3398.
- (18) Shibahara, T.; Sakane, G.; Maeyama, M.; Kobashi, H.; Yamamoto, T.; Watase, T. Uptake of Ethylene by Sulfur-bridged Cubane-type Molybdenum/Tungsten–Nickel Clusters $[\text{M}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ ($\text{M}_3 = \text{Mo}_3, \text{Mo}_2\text{W}, \text{MoW}_2, \text{W}_3$): Synthesis, Structures and ^1H NMR Spectra. *Inorg. Chim. Acta* **1996**, *251*, 207–225.
- (19) Saysell, D. M.; Borman, C. D.; Kwak, C.-H.; Sykes, A. G. Ligand Substitution Reactions at the Nickel of $[\text{Mo}_3\text{NiS}_4(\text{H}_2\text{O})_{10}]^{4+}$ with Two Water Soluble Phosphines, CO, Br^- , I^- , and NCS^- and the Inertness of the 1,4,7-Triazacyclononane (L) Complex $[\text{Mo}_3(\text{NiL})\text{-S}_4(\text{H}_2\text{O})_9]^{4+}$. *Inorg. Chem.* **1996**, *35*, 173–178.
- (20) Dimmock, P. W.; Saysell, D. M.; Sykes, A. G. Electron-Transfer reactivity of cuboidal heterometallic Mo_3MS_4 clusters in aqueous acidic solutions. *Inorg. Chim. Acta* **1994**, *225*, 157–162.
- (21) Akashi, H.; Nakano, A.; Shibahara, T. Crystal Structure of Molybdenum–Rhodium Sulfido Cluster Containing $\text{Mo}_3\text{Rh}(\mu_3\text{-S})_4$ Cube. Presented at the XVIIIth IUCr Congress, Glasgow, August 1999, P07.07.041.
- (22) Hernandez-Molina, R.; Edwards, A. J.; Clegg, W.; Sykes, A. G. Preparation, Structure and Properties of the Arsenic-Containing Corner-Shared Double-Cube $[\text{Mo}_6\text{AsS}_8(\text{H}_2\text{O})_{18}]^{8+}$: Metal–Metal Bonding and a Classification of Different Cluster Types. *Inorg. Chem.* **1998**, *37*, 2989–2994.
- (23) Hernandez-Molina, R.; Dybtsev, D. N.; Fedin, V. F.; Elsegood, M. R. J.; Clegg, W.; Sykes, A. G. Preparation, Structure and Reactivity of Heterometallic Sn-Containing Single- and Double-Cube Derivatives of $[\text{Mo}_3\text{Se}_4(\text{H}_2\text{O})_9]^{4+}$ and $[\text{W}_3\text{Se}_4(\text{H}_2\text{O})_9]^{4+}$. *Inorg. Chem.* **1998**, *37*, 2995–3001.
- (24) Lu, S.-F.; Huang, J.-Q.; Wu, Q.-J.; Huang, X.-Y.; Yu, R.-M.; Zheng, Y.; Wu, D.-X. The synthesis and crystal structures of the first species of monocubane type clusters $[\text{Mo}_3\text{PbS}_4]^{6+}$ and $[\text{Mo}_3\text{-BiS}_4]^{7+}$. *Inorg. Chim. Acta* **1997**, *261*, 201–209.
- (25) Sokolov, M. N.; Virovets, A. V.; Dybtsev, D. N.; Gerasko, O. A.; Fedin, V. P.; Hernandez-Molina, R.; Clegg, W.; Sykes, A. G. Metal Incorporation into and Dimerization of M_3E_4 Clusters (M = Mo, W; E = S, Se) in Supramolecular Assemblies with Cucurbituril: A Molecular Model of Intercalation. *Angew. Chem., Int. Ed.* **2000**, *39*, 1659–1661.
- (26) Hernández-Molina, R.; Sykes, A. G. Reactions of the Heterometallic Cuboidal Clusters Mo_3MS_4 (M = Co, Ni, Pd, Cu) and Mo_3NiSe_4 with CO: Electron Counts and Kinetic/Thermodynamic Studies with M = Ni, Pd. *Coord. Chem. Rev.* **1999**, *187*, 291–302.
- (27) Nasreldin, M.; Routledge, C. A.; Sykes, A. G. Preparation and Aqueous Solutions Properties of the Heterometallic Cuboidal Complex $[\text{W}_3\text{CuS}_4(\text{H}_2\text{O})_{10}]^{5+}$. *J. Chem. Soc., Dalton Trans.* **1994**, 2809–2814.
- (28) Sakane, G.; Hashimoto, K.; Takahashi, M.; Takeda, M.; Shibahara, T. Sulfur-Bridged Sandwich Cubane-Type Molybdenum Antimony Cluster. Synthesis, X-ray Structure and ^{121}Sb Mössbauer Spectra of $[\text{Mo}_6\text{SbS}_8(\text{H}_2\text{O})_{18}(\text{pts})_8\cdot 24\text{H}_2\text{O}]$. *Inorg. Chem.* **1998**, *37*, 4231–4234.
- (29) Müller, A.; Fedin, V. P.; Diemann, E.; Bögge, H.; Krickemeyer, E.; Solter, D.; Giuliani, A. M.; Barbieri, R.; Adler, P. Reductive addition at the W_3S_4 core by Sn^{2+} or an unusual supramolecular system: a synergetic reaction leading to the host–guest compound $(\text{Me}_2\text{-NH}_2)_6[\text{W}_3\text{SnCl}_3\text{S}_4(\text{NCS})_9]\cdot 0.5\text{H}_2\text{O}$. *Inorg. Chem.* **1994**, *33*, 2243–2247.
- (30) Bahn, C. C.; Tan, A.; Harris, S. Bonding in $\text{Mo}_3\text{M}'\text{S}_4$ cubane-type clusters. Variations in electronic structure when M' is a main group or transition metal. *Inorg. Chem.* **1998**, *37*, 2770–2778.
- (31) Fedin, V. P.; Lamprecht, G. J.; Kohzuma, T.; Clegg, W.; Elsegood, M. R. J.; Sykes, A. G. Preparation, Structure and Properties of Trinuclear $[\text{M}_3\text{Se}_4(\text{CN})_9]^{5-}$ (M = Mo or W) Complexes Obtained from Mo_3Se_7 . Core Compounds and Related Studies. *J. Chem. Soc., Dalton Trans.* **1997**, 1747–1751.
- (32) Saysell, D. M.; Lamprecht, G. J.; Darkwa, J.; Sykes, A. G. Aqueous Solution Chemistry of the Mo_3PdS_4 Cube: Substitution Reactions and the Double to Single Cube Interconversion Induced by CO, Two Phosphines, Cl^- , Br^- , and NCS^- . *Inorg. Chem.* **1996**, *35*, 5531–5535.
- (33) Akashi, H.; Shibahara, T. Molybdenum–copper–sulfur Mo_3CuS_4 cubes. *Inorg. Chim. Acta* **2000**, *300–303*, 572–580.
- (34) Darkwa, J.; Lockemeyer, J. R.; Boyd, P. D. W.; Rauchfuss, T. B.; Rheingold, A. L. *J. Am. Chem. Soc.* **1988**, *110*, 141–147.
- (35) McGrady J. E. Periodic Trends in Metal–Metal Bonding in Cubane Clusters $[\text{M}_4\text{Q}_4(\text{C}_5\text{H}_5)_4]$ (M = Cr, Mo; Q = O, S). *J. Chem. Soc., Dalton Trans.* **1999**, 1393–1399.
- (36) Fedin, V. P.; Kalimina, I. V.; Virovets, A. V.; Podberezskaya, N. V.; Neretin, I. S.; Slovokholov, Y. L. Synthesis and Structure of a Tetranuclear Niobium Telluride Cuboidal Cluster with a Central $\mu_4\text{-O}$ Ligand. *Chem. Commun.* **1998**, 2579–2580.
- (37) Freedman, D.; Melman, J. H.; Emge, T. J.; Brennan, J. G. Cubane Clusters Containing Lanthanide Ions: $(\text{py})_8\text{Yb}_4\text{Se}_4(\text{SPh})_4$ and $(\text{py})_{10}\text{Yb}_6\text{S}_6(\text{SPh})_6$. *Inorg. Chem.* **1998**, *37*, 4162–4163.

AR960152S