

# PARACEST Agents: Modulating MRI Contrast via Water Proton Exchange

SHANRONG ZHANG,<sup>†,‡</sup> MATTHEW MERRITT,<sup>‡</sup>  
DONALD E. WOESSNER,<sup>‡</sup>  
ROBERT E. LENKINSKI,<sup>§</sup> AND  
A. DEAN SHERRY<sup>\*,†,‡</sup>

*Department of Chemistry, University of Texas at Dallas,  
P.O. Box 830688, Richardson, Texas 75083-0688, The Rogers  
Magnetic Resonance Center, Department of Radiology,  
University of Texas Southwestern Medical Center,  
5801 Forest Park Road, Dallas, Texas 75235-9085, and  
Department of Radiology, Beth Israel Deaconess Medical  
Center and Harvard Medical School, 1 Deaconess Road,  
Boston, Massachusetts 02115*

Received February 21, 2003

## ABSTRACT

Scientific interest in optimizing the properties of gadolinium (III) complexes as MRI contrast agents has led to many new insights into lanthanide ion coordination chemistry in the last two decades. Among these was the surprising observation that water exchange in lanthanide (III) derivatives of DOTA can be modulated dramatically by judicious choice of ligand side chain and Ln<sup>3+</sup> ionic radii. This resulted in the discovery of paramagnetic CEST agents for altering MRI image contrast based upon the chemical exchange saturation transfer mechanism. The goal of this article is to review the factors that govern water molecule and water proton exchange in these complexes and to compare the potential sensitivity of PARACEST agents versus Gd<sup>3+</sup>-based T<sub>1</sub> relaxation agents for altering tissue contrast.

## Introduction

Scientific interest in lanthanide ion coordination chemistry has grown substantially over the past 25 years, driven largely by the emergence of magnetic resonance imaging

Shanrong Zhang was born in Ankang, Shaanxi, China, in 1966. He received his BS in Chemistry from Shaanxi University of Technology (formerly the Hanzhong Teachers College) in 1987 and his PhD in Physical Chemistry from Changchun Institute of Applied Chemistry, Chinese Academy of Sciences in 1995. He joined A. Dean Sherry at the University of Texas at Dallas as a postdoctoral fellow in 1997. Recently, he joined the staff of the Radiology Department of the University of Texas Southwestern Medical Center at Dallas. His major research interests are in the development of new contrast agents for MRI.

Matthew Merritt was born in Raleigh, NC, in 1969. He graduated with a BS in Chemistry from North Carolina State University in 1991. In 1996 he obtained his PhD in Chemistry from Washington University in St. Louis under Dr. Jacob Schaefer. From 1996 to 2000, he held a postdoctoral position with Dr. Gary Drobny at the University of Washington in Seattle. In 2000, he joined the staff of the Radiology Department of the University of Texas Southwestern Medical Center as the NMR facility manager. His other research interests include magic-angle spinning NMR techniques and in vivo <sup>13</sup>C spectroscopy.

Donald E. Woessner received a BA in chemistry and mathematics in 1952 from Carthage College. In 1957 he received a PhD in physical chemistry from the University of Illinois under the direction of Herbert S. Gutowsky on quadrupole resonance relaxation in solids and NMR relaxation in solids and liquids. In 1958, he joined Mobil Research and Development Corporation in Dallas and worked on NMR relaxation in liquids and later on NMR imaging of porous media. He retired from Mobil in 1992 but continues NMR research as an Adjunct Professor of Radiology, University of Texas Southwestern Medical Center at Dallas.

(MRI) as a preeminent tool in clinical medicine.<sup>1</sup> Gadolinium (III) is most often used as a “contrast agent” for MRI because it is the most efficient of all lanthanide ions at relaxing bulk water protons.<sup>2,3</sup> The remaining paramagnetic lanthanide ions are much less efficient at relaxing water protons but can induce sizable NMR hyperfine shifts in nearby NMR nuclei.<sup>4</sup> Hyperfine shifts can provide useful quantitative structural information or may simply be used to “resolve” resonances of NMR active nuclei in biological compartments. This feature has been particularly attractive for separating the NMR signals of intra- and extracellular Na<sup>+</sup> in isolated cells, organs, and tissues in vivo.<sup>5,6</sup> The frequency shifting ability of the remaining lanthanide ions has not found an application in proton imaging because, unlike Gd<sup>3+</sup>-based relaxation systems where rapid water exchange is considered imperative, fast water exchange effectively reduces water proton hyperfine shifts to near zero (weighted average of a small shifted pool and a large bulk pool). Very recently, however, several slow water exchange lanthanide complexes have been reported,<sup>7–15</sup> and in these systems highly hyperfine shifted Ln<sup>3+</sup>-bound water resonances are often seen by high-resolution NMR. As we shall see, this feature once again offers an opportunity to take advantage of the variable hyperfine shifting ability of the lanthanide ion series.<sup>16</sup>

An alternative way to introduce contrast into a MR image would be to alter the total water signal detected in the experiment. This was demonstrated by Balaban and co-workers<sup>17</sup> when they used a presaturation pulse to saturate the broad water signal that lies beneath the sharper bulk water signal in many tissues. This broad resonance is thought to reflect water tightly associated with tissue proteins, lipids, and/or subcellular structures and appears to change with progressive diseases such as multiple sclerosis, ALS, etc. Upon saturation, exchange

\* Corresponding author. Phone: 972-883-2907. Fax: 972-883-2925. E-mail: sherry@utdallas.edu.


<sup>†</sup> University of Texas at Dallas.

<sup>‡</sup> University of Texas Southwestern Medical Center.

<sup>§</sup> Beth Israel Deaconess Medical Center and Harvard Medical School.

Robert E. Lenkinski was born in Mildorf, Germany, in 1947. He received his BS from the University of Toronto (1969) and his PhD from the University of Houston (1972) under the supervision of Professor MR Willcott, III. He spent two years as a postdoctoral fellow with Jacques Reuben at the Weizmann Institute of Science and then returned to Houston for one year before joining the Cancer Center at the University of Alabama, Birmingham. In 1980, he joined the Chemistry Faculty at the University of Guelph and, in 1986, moved to the Department of Radiology, University of Pennsylvania. In 1999, he moved to the Beth Israel Deaconess Medical Center, Harvard Medical School, as Professor of Radiology. His research interests include magnetic resonance imaging and spectroscopy of human diseases and lanthanide chelates as a platform for developing targeted contrast agents for MRI.

A. Dean Sherry was born in Wisconsin in 1945, received a BS in Chemistry from UW-LaCrosse in 1967 and a PhD in Inorganic Chemistry from Kansas State University in 1971, and he was an NIH Postdoctoral Fellow (1971–72) before joining the Chemistry faculty at the University of Texas at Dallas. He served as Chairman of that department from 1979 to 1990 and has held a joint appointment at UT-Southwestern Medical Center as Professor of Radiology since 1990. He works on various aspects of biomedical NMR including the use of <sup>13</sup>C metabolic tracers and multidimensional NMR to follow metabolism in vivo and on novel applications of lanthanide complexes as metabolic imaging agents.



**FIGURE 1.** A scheme of a two pool CEST model. By selective saturation of the small pool (the exchanging  $-OH$  or  $-NH$  site), the large pool (bulk water) will decrease also if the chemical exchange kinetics are appropriate.

occurs between the bound water pool and the larger bulk water pool resulting in a decrease in bulk water signal intensity (Figure 1). This technique is now commonly known as magnetization transfer (MT) imaging.<sup>18</sup> More recently, this same group demonstrated that low molecular weight compounds with slowly exchanging  $-NH$  or  $-OH$  protons may also be used to alter tissue contrast via chemical exchange saturation transfer (CEST) of presaturated spins to bulk water.<sup>19</sup> The molecules first used to demonstrate the CEST effect were diamagnetic and thereby had proton chemical shifts not much different from that of bulk water ( $-NH$  and  $-OH$  groups are typically well within 5 ppm of bulk water). Paramagnetic analogues of such compounds were not known until the discovery of lanthanide complexes having unusually slow water exchange kinetics.<sup>16,20</sup> In this Account, we review the impact of lanthanide ion size and ligand structure on water exchange and demonstrate that large hyperfine shifting lanthanide ions such as  $Dy^{3+}$ ,  $Tb^{3+}$ ,  $Tm^{3+}$ , and  $Yb^{3+}$  offer advantages for introducing MRI contrast via the CEST mechanism. Such slow water exchange paramagnetic complexes are referred to herein as PARACEST agents.

### Tuning the Bound Water Lifetime in $Ln^{3+}$ Complexes

During early development of  $Gd^{3+}$ -based MRI contrast agents, it was commonly assumed that water exchange between  $Ln^{3+}$ -bound, inner-sphere coordination sites and bulk solvent was so fast that it effectively had no impact on the relaxation characteristics of those complexes. However, more recent studies have shown that this is not always the case. For example, it is now known that the bound water lifetime ( $\tau_M$ ) in two clinically used MRI agents,  $GdDTPA^{2-}$  and  $GdDOTA^-$  (both have one inner-sphere bound water,  $q=1$ ), is  $\sim 240$  ns,<sup>21,22</sup>  $\sim 100$ -fold longer than the inner-sphere water lifetime of the  $Gd^{3+}$  aqua ion.<sup>23–25</sup> Solomon–Bloembergen–Morgan theory of paramagnetic relaxation indicates that water exchange in  $GdDTPA^{2-}$  and  $GdDOTA^-$  lowers the relaxivity of these complexes somewhat below their theoretical optimum, but this is irrelevant in terms of their clinical efficacy.  $\tau_M$  is about 10-fold longer ( $\sim 2$   $\mu s$ ) in bis-amide derivatives

of DTPA, like  $GdDTPA-BMA^{26,27}$  and longer still ( $\sim 20$   $\mu s$ ) in tetra-amide derivatives of DOTA such as  $GdDTMA^{3+,7,28}$ . Clearly, complexes with even slower water exchange are detrimental if one's goal is to optimize relaxivity but this same feature does allow one to consider using these complexes as PARACEST agents.

### Factors Affecting $\tau_M$ : Lanthanide Coordination Geometry

Early NMR studies of the  $LnDOTA^-$  complexes showed that two slowly interconverting species are present in aqueous solution and that the populations of these two species are sensitive to the size of the central  $Ln^{3+}$  cation.<sup>29</sup> These two coordination isomers have since been assigned to the square antiprism (commonly referred to as the M isomer) and twisted square antiprism (commonly referred to as the m isomer) structures, related to each other by rotation of the side-chain acetate groups. Interestingly, it was subsequently discovered that these two species have quite different bound water lifetimes, the m isomer showing much faster exchange compared to the M isomer. For example,  $\tau_M$  is 120  $\mu s$  for the M isomer of  $EuDTMA^{3+}$  but only 2  $\mu s$  for the m isomer.<sup>7,8,10</sup>

### Factors Affecting $\tau_M$ : Identity of the Ligand Side Chains

As noted above, the bound water lifetimes for  $LnDOTA^-$  tetra(amide) complexes tend to be considerably longer than for the corresponding  $LnDOTA^-$  complexes. The shapes of  $^{17}O$  curves of Figure 2 immediately show that water exchange is quite different in these complexes and hence sensitive to the number of amide substitutions. Although a detailed analysis is beyond the scope of this review,<sup>30</sup> the maxima in each of these curves (near 328 K for the mono-amide, near 355 K for the bis-amide, and above 370 K for the tetra-amide) show that the bound water lifetimes are in the order  $GdDOTA-4AmP > GdDOTA-2AmP > GdDOTA-1AmP$ . A complete fitting of these data to theory gives  $\tau_M$  values of 26, 6.2, and 1.3  $\mu s$ , respectively, for the M isomers in each complex. The bound water lifetimes in the m isomers are also much shorter ( $\sim 6$ –20 ns) and differ in this series. This illustrates that even one amide side chain appears to have a significant impact on the bound water lifetime.

The chemical characteristics, i.e., charge and polarity, of the extended amide side-chain arms also play a role in determining  $\tau_M$ . For the series of  $EuDOTA^-$  tetraamide complexes shown in Figure 3, it has been found that  $\tau_M$  varies considerably and this has been correlated with a calculated solvent accessibility parameter. Given that the mechanism of water exchange in these tetra-amide complexes is thought to be dissociative,<sup>15</sup> greater solvent accessibility to the  $Eu^{3+}$ -bound water may aid in dissociation of the bound water by hydrogen bonding. In general, for  $EuDOTA^-$  tetra(amide) complexes, the bound water lifetime correlates with polarity of the ligand side chains, i.e., phosphonate ester  $\geq$  carboxylate ester  $\gg$  alkyl groups  $\geq$  simple amides. For example,  $\tau_M$ 's decrease in order of




FIGURE 2. Plots of the  $^{17}\text{O}$  transverse relaxation rate vs temperature for  $\text{Gd}^{3+}$  complexes of tetra-, bis-, and mono-amide derivatives of DOTA (unpublished data).




FIGURE 3. Relationship between coordinated water molecule lifetimes in Eu-tetraamide complexes vs the calculated solvent accessible surface around the bound water (adapted from ref 15). Although solvent accessible surface areas have not been reported for compounds 1, 8, and 9, their structures are included here because they are referred to in the text.

$\text{Eu}(\mathbf{1})^{3+}$  at  $1.3 \text{ ms}^{12} > \text{Eu}(\mathbf{4})^{3+}$  at  $780 \mu\text{s}^{11} \gg \text{Eu}(\mathbf{5})^{3+}$  at  $278 \mu\text{s}^9 > \text{Eu}(\mathbf{8})^{3+}$  at  $146 \mu\text{s}$  (unpublished)  $\sim \text{Eu}(\mathbf{9})^{3+}$  at  $156 \mu\text{s}^8 > \text{Eu}(\mathbf{10})^{3+}$  at  $120 \mu\text{s}^7$  (all for M isomers measured in wet acetonitrile). We have also observed that  $\text{Eu}^{3+}$ -bound water lifetimes for complexes that can be measured in more than one solvent are about 2-fold shorter in aqueous solution than in wet acetonitrile<sup>11</sup> and complexes with negatively charged side chains have shorter  $\tau_M$  values than their neutral counterparts. For example, the  $\tau_M$  found for  $\text{EuDOTA-4AmC}^-$  ( $300 \mu\text{s}$ , pH 7, aqueous) is somewhat shorter than the  $\tau_M$  for the corresponding ester,  $\text{EuDOTA-4AmCE}^{3+}$  ( $382 \mu\text{s}$ , pH 7.4, aqueous).<sup>11,31</sup> Both observations suggest that there is proton exchange contribution to  $\tau_M$  in aqueous media.

### Factors Affecting $\tau_M$ : Molecular Exchange versus Prototropic Exchange

Aime et al.<sup>32</sup> recently published a short review paper on the relaxivity contributions from the prototropic exchange

versus the water molecule exchange in  $\text{Gd}^{3+}$  complexes. It has been observed that prototropic exchange is accelerated in complexes such as these by acid and/or base and that an increase in prototropic exchange results in an increase in relaxivity below pH  $\sim 2$  and above pH  $\sim 9-10$ .<sup>28</sup> This early observation prompted us to examine more closely the water relaxivity of  $\text{GdDOTA-4AmP}$  as a function of pH. In this water-soluble complex,  $\tau_M$  as measured by  $^{17}\text{O}$  NMR was  $\sim 26 \mu\text{s}$  over the entire pH range, 6–9.5; yet the water  $^1\text{H}$  relaxivity steadily increases between pH 8.5 ( $r_1 = 4.5 \text{ mM}^{-1}\text{s}^{-1}$ ) and pH 6 ( $r_1 = 9.8 \text{ mM}^{-1}\text{s}^{-1}$ ).<sup>33</sup> Thus, upon protonation of the extended phosphonate groups (four  $\text{pK}_a$ 's ranging from 8.5 to 6), the protons on a single  $\text{Gd}^{3+}$ -bound water undergo prototropic exchange with bulk solvent. Such pH-dependent relaxivity is not a feature of current clinical  $\text{Gd}^{3+}$ -based CAs where water exchange is relatively fast ( $\tau_M < 240 \text{ ns}$ ), so the slow water exchange feature of  $\text{GdDOTA-4AmP}$  makes this an attractive agent for mapping extracellular pH in tissues by MRI.<sup>34</sup>




FIGURE 4. Plot of experimental  $\tau_M$  vs  $\text{Ln}^{3+}$  ionic radii for the LnDOTA-4AmCE complexes. The  $\tau_M$  values were obtained by fitting the temperature dependence of either the  $^{17}\text{O}$  or  $^1\text{H}$  NMR bound water line widths (from ref 14).

### Factors Affecting $\tau_M$ : Lanthanide Cation Size

Merbach et al.<sup>27</sup> found for series of LnDTPA-BMA complexes ( $\text{Ln}^{3+}$  = various lanthanide cations) that the  $\text{Ln}^{3+}$ -bound water lifetimes decreased along the series from 1.88  $\mu\text{s}$  ( $\text{Nd}^{3+}$ ) to 0.16  $\mu\text{s}$  ( $\text{Ho}^{3+}$ ) and suggested that this reflected a change in water exchange mechanism, from interchange ( $I_a$ ) for the larger cations to dissociative ( $D$ ) for the smaller cations. More recently, Parker et al.<sup>9</sup> reported that the bound water lifetime of a EuDOTA-tetra(amide)<sup>3+</sup> might be as much as 500 times that of the corresponding  $\text{Yb}^{3+}$  complex. We subsequently found that the bound water lifetimes of EuDOTA-4AmCE<sup>11</sup> and EuDOTA-4AmPE<sup>12</sup> are about 2-fold longer than those of corresponding  $\text{Gd}^{3+}$  complexes, even though  $\text{Eu}^{3+}$  and  $\text{Gd}^{3+}$  complexes are usually considered isostructural. To help clarify the relationship between  $\tau_M$  and  $\text{Ln}^{3+}$  ionic radius, we determined  $\tau_M$  values for the remaining complexes in this series and found that  $\tau_M$  is highly sensitive to ionic radii for those ions nearest  $\text{Eu}^{3+}$  (Figure 4).<sup>14</sup> Although the data appear to separate into three distinct subgroups, two crystal structures (the  $\text{Eu}^{3+}$  and  $\text{Tb}^{3+}$  complexes) and a lanthanide induced shift (LIS) analysis of paramagnetic  $^1\text{H}$  shift indicate that the entire series is isostructural (all largely the **M** isomer). Therefore, the sharp changes in  $\tau_M$  versus  $\text{Ln}^{3+}$  radii (Figure 4) must be ascribed to altered water exchange mechanisms along the series.

### Characterization of PARACEST Agents

**Two-Site Exchange Theory.** The basic physical process is that of chemical exchange between two pools of protons. Pool A is bulk water, and pool B is water bound to the chelated lanthanide ion. Pool B is chemically shifted by the paramagnetic lanthanide away from pool A. The basic NMR experiment is to apply RF irradiation at the Larmor frequency of pool B for a time sufficiently long so as to reach a steady-state in which the magnetizations in the rotating frame are time-independent.

The NMR behavior of the two exchanging pools of nuclei is described by the modified Bloch equations that include the effects of chemical exchange on the X, Y, and Z spin magnetizations of both pools of nuclei. These

equations are given below:<sup>35,36</sup>

$$\frac{dM_X^a}{dt} = -(\omega_a - \omega)M_Y^a - k_{2a}M_X^a + C_bM_X^b \quad (1)$$

$$\frac{dM_X^b}{dt} = -(\omega_b - \omega)M_Y^b - k_{2b}M_X^b + C_aM_X^a \quad (2)$$

$$\frac{dM_Y^a}{dt} = (\omega_a - \omega)M_X^a - k_{2a}M_Y^a + C_bM_Y^b - \omega_1M_Z^a \quad (3)$$

$$\frac{dM_Y^b}{dt} = (\omega_b - \omega)M_X^b - k_{2b}M_Y^b + C_aM_Y^a - \omega_1M_Z^b \quad (4)$$

$$\frac{dM_Z^a}{dt} = \frac{M_0^a}{T_{1a}} - k_{1a}M_Z^a + C_bM_Z^b + \omega_1M_Y^a \quad (5)$$

$$\frac{dM_Z^b}{dt} = \frac{M_0^b}{T_{1b}} - k_{1b}M_Z^b + C_aM_Z^a + \omega_1M_Y^b \quad (6)$$

where

$$k_{1a} = \frac{1}{T_{1a}} + C_a \quad (7)$$

$$k_{2a} = \frac{1}{T_{2a}} + C_a \quad (8)$$

and  $\omega_a$  is the Larmor frequency of nuclei in pool A. In these definitions,  $T_{1a}$  is the spin-lattice relaxation time of A,  $T_{2a}$  is the transverse relaxation time of A, and  $C_a$  is the transition rate of A nuclei in leaving A and is equal to  $1/\tau_a$ . Similar definitions apply to B nuclei;  $C_b = 1/\tau_M$ . Since exchange is a detailed-balance phenomenon,  $\tau_a = \tau_M/(M_0^b/M_0^a)$ . It is convenient to use the RF intensity  $B_1$  as expressed in Hertz:  $B_1 = \omega_1/2\pi$ .

A mathematical expression for  $M_Z^a/M_0^a$  was obtained from a solution of this set of equations under steady-state conditions in which the time derivatives of all the nuclear magnetizations are zero. The result is a complicated function of the seven NMR parameters ( $T_{1a}$ ,  $T_{2a}$ ,  $T_{1b}$ ,  $T_{2b}$ ,  $M_0^b$ ,  $M_0^a$ ,  $\tau_M$ , and the Larmor frequency difference  $\Delta\omega = \omega_a - \omega_b$ ) and the two experimental variables ( $B_1$  and irradiation frequency).

The maximum reduction in  $M_Z^a/M_0^a$  caused by chemical exchange occurs when the last two terms in eq 5 are zero. This situation occurs when (a) the B nuclei are completely saturated so that  $M_Z^b = 0$  and (b) there is no direct excitation of pool A nuclei by the  $B_1$  irradiation that excites B nuclei. Condition a is met when  $B_1$  is sufficiently large. Condition b is met when the relative Larmor frequency shift of A is very large so that  $\Delta\omega/\omega_1 \gg 1$ . Under these conditions, the steady-state solution of eq 5 gives

$$\frac{M_Z^a}{M_0^a} = \frac{\tau_a}{\tau_a + T_{1a}} = \frac{1}{1 + \frac{cq}{55.5} \frac{T_{1a}}{\tau_M}} \quad (9)$$




FIGURE 5. Z-spectra for EuDOTA-4AmCE (left) and NdDOTA-4AmCE (right) at 4.7 T obtained by using a 1 s presaturation pulse. The high-resolution  $^1\text{H}$  spectra (500 MHz) of the complexes show bound water resonances at +50 ppm ( $\text{Eu}^{3+}$ ) and -32 ppm ( $\text{Nd}^{3+}$ ) (from refs 16 and 38).

where  $c$  is the concentration of PARACEST agent and  $q$  is the number of bound water molecules per PARACEST complex.

A “useful” reduction in  $M_z^a/M_0^a$  requires that  $\tau_a$  be of the same order of magnitude or smaller than  $T_{1a}$ . Since  $\tau_a = \tau_M/(M_0^b/M_0^a)$ , short  $\tau_M$  values are needed when the lanthanide concentration ( $M_0^b$ ) is small. However, short  $\tau_M$  values also broaden the  $\text{Ln}^{3+}$ -bound water protons; hence, larger  $\Delta\omega$  values are required to avoid direct excitation of pool A. Such excitation obscures the actual CEST effect. Clearly, a paramagnetic complex with a large  $\Delta\omega$  value is advantageous. Note that eq 9 makes predictions only for limiting cases while the complete solution to the set of Bloch equations enables one to predict the  $M_z^a/M_0^a$  values for any general case.


**Z-Spectra (CEST Spectra).** One practical way to characterize chemical exchange in such systems is to selectively presaturate the sample in incremental steps over a range of frequencies and plot the remaining bulk water signal,  $M_s/M_0$ , versus saturation frequency. This was originally referred to as a Z-spectrum<sup>37</sup> and more recently as a CEST spectrum.<sup>19</sup> Figure 5 illustrates two such Z-spectra for samples of EuDOTA-4AmCE and NdDOTA-4AmCE.<sup>16,38</sup> The peaks at 0 ppm in each Z-spectrum represent direct saturation of bulk water, while the peaks centered near +50 and -32 ppm reflect chemical exchange between the respective  $\text{Eu}^{3+}$ -bound and  $\text{Nd}^{3+}$ -bound water molecules in these complexes. The peak at +50 ppm in the  $\text{Eu}^{3+}$  plot is noticeably sharper than the peak at -32 ppm in the  $\text{Nd}^{3+}$  plot, consistent with slower water exchange in the former complex. This is also evident in the width of the bound water resonances in the high-resolution  $^1\text{H}$  spectra shown plotted above the Z-spectra. Given that a peak due to chemical exchange should be detected in such Z-spectra for systems that meet the

Table 1.  $^1\text{H}$  Chemical Shifts and Lifetimes ( $\tau_M$ ) of  $\text{Ln}^{3+}$ -Bound Water in the LnDOTA-4AmCE Complexes<sup>38</sup>


complexes	observation of bound water	$\tau_M^{298}$ ( $\mu\text{s}$ )	$\delta(^1\text{H})$ (ppm)	$\Delta\omega \cdot \tau_M$		
				11.75 T	4.7 T	1.5 T
$\text{Pr}^{3+}$	Yes	20	-60	3.8	1.5	0.5
$\text{Nd}^{3+}$	Yes	80	-32	8.0	3.2	1.0
$\text{Sm}^{3+}$	Yes	320	-4	4.0	1.6	0.5
$\text{Eu}^{3+}$	Yes	382	50	60.0	24.0	7.7
$\text{Tb}^{3+}$	No	31	-600	58.5	23.4	7.5
$\text{Dy}^{3+}$	No	17	-720	38.5	15.4	4.9
$\text{Ho}^{3+}$	No	19	-360	21.5	8.6	2.8
$\text{Er}^{3+}$	No	9	200	5.7	2.3	0.7
$\text{Tm}^{3+}$	Yes	3	500	4.7	1.9	0.6
$\text{Yb}^{3+}$	Yes	3	200	1.9	0.5	0.2

intermediate to slow exchange criteria,  $\Delta\omega \cdot \tau_M \geq 1$ , one can then predict which complexes might serve as useful PARACEST agents at various  $B_0$  fields. These data, collected in Table 1, predict that most of the LnDOTA-4AmCE complexes should function as PARACEST agents at 11.75 T but only a few,  $\text{Eu}^{3+}$ ,  $\text{Tb}^{3+}$ ,  $\text{Dy}^{3+}$ , and  $\text{Ho}^{3+}$ , might be useable at the most common clinical MR imaging field strength, 1.5T. These data also illustrate that PARACEST agents will likely become more efficient at higher  $B_0$  fields.

One can further illustrate the impact of both  $\Delta\omega$  and  $\tau_M$  in determining the efficiency of a PARACEST agent by using predictions provided by the modified Bloch equations. Figure 6 shows 3D stacked plots of Z-spectra plotted as a function of bound water exchange rate ( $1/\tau_M$ ) for two PARACEST agents, one having a  $\text{Ln}^{3+}$ -bound water molecule at +50 ppm ( $\Delta\omega = 62\,832$  rad/s) and another having a  $\text{Ln}^{3+}$ -bound water at +800 ppm ( $\Delta\omega = 1 \times 10^6$  rad/s). The predictions assume presaturation of the bound water using a  $B_1$  field of 1500 Hz, a  $B_0$  of 4.7 T, and 20 mM agent. These stacked Z-spectra illustrate several important




**FIGURE 6.** Stacked plots of Z-spectra derived from the modified Bloch equations for two-site exchange. The plots show  $M_S/M_0$  (vertical axis) as a function of bound water exchanging rate ( $k_{ex} = 1/\tau_M$ ) for two paramagnetic complexes, one having a bound water signal at 50 ppm (left) and one at 800 ppm (right). The same set of parameters was used to generate these two plots;  $T_{1a} = 1.0$  s and  $T_{2a} = 0.3$  s for bulk water,  $T_{1b} = 0.1$  s and  $T_{2b} = 0.03$  s for bound water,  $B_0 = 4.7$  T,  $B_1 = 1500$  Hz, and a concentration of 20 mM.



**FIGURE 7.** Plot of the fractional decrease in bulk water signal intensity expected for a PARACEST agent having a bound water lifetime ( $\tau_M$ ) of 3  $\mu$ s and a bound water ( $^1$ H) paramagnetic shift of 500 ppm (based upon eq 9 by assuming  $T_{1a} = 1.0$  s).

points: (1)  $M_S/M_0$  falls off more rapidly with increasing exchange rate (shorter lifetime) for the 50 ppm agent than for the 800 ppm agent, (2) complexes with highly shifted bound water resonances are more efficient over a wider range of water exchange rates, and (3) the largest decrease in  $M_S/M_0$  will be achieved by using an agent with optimal water exchange rate.

What is the lower detection limit for an optimized PARACEST agent? Figure 7 shows one result predicted by the Bloch equations for a PARACEST agent having a bound water resonance at 500 ppm and a  $\tau_M$  of 3  $\mu$ s. Assuming that a sufficiently strong  $B_1$  field can be used to fully saturate the bound water signal, the Bloch equations predict that  $\sim 37\%$  of the bulk water signal can be eliminated by using  $\sim 100$   $\mu$ M agent, and even 5% of the bulk water signal can be eliminated by using  $\sim 10$   $\mu$ M agent. The latter concentration is well below the detection limit of a low molecular weight  $Gd^{3+}$ -based contrast agent with a typical relaxivity ( $r_1$ ) of 4–5  $mM^{-1} s^{-1}$ .<sup>39</sup> These data suggest that PARACEST agents have the potential of being as sensitive as or even more sensitive than  $Gd^{3+}$ -based  $T_1$  agents, assuming that all practical aspects can be solved. Just as one can increase the relaxivity of  $Gd^{3+}$ -based agents



**FIGURE 8.** Ratiometric plots for 30 mM LnDOTA–4AmC complexes at 312 K and 7.05 T. These data were generated by applying a train of frequency-selective e-burp pulses (at train of 20 ms shaped pulses for the NH protons and 1 ms shaped pulses for the OH protons) (adapted from ref 46).

substantially by conjugation to a polymer or formation of an aggregate,<sup>40–42</sup> similar modifications of PARACEST agents could be done and this would easily extend the lower detection limit into the submicromolar range. This was recently demonstrated for a series of polymers containing large numbers of exchangeable  $-NH$  groups.<sup>43</sup>

**pH Responsive PARACEST Agents.** Balaban et al.<sup>44</sup> first demonstrated that diamagnetic CEST agents having more than one type of proton exchangeable site may be used to measure pH using a “ratiometric method”. Aime et al.<sup>45,46</sup> used this same concept for PARACEST agents having both a slowly exchanging  $Ln^{3+}$ -bound water molecule and exchangeable  $-NH$  sites (Figure 8). For such complexes, exchange between the  $-NH$  sites and bulk water is quite sensitive to pH in the range 6–8, while proton and/or water molecule exchange is nearly independent over this same pH range. Hence, plots of the PARACEST effect expressed as a ratio for two different presaturation frequencies ( $Ln^{3+}-OH$  and  $Ln^{3+}-NH$ ) removes the effect of agent concentration and provides a direct measure of solution pH. One should note that the PARACEST ratio will be even more sensitive to pH when using an agent with larger  $\Delta\omega$  differences between the  $Ln^{3+}$ -bound water

and –NH resonances. The CEST efficiency at the two sites is field dependent (due to  $\Delta\omega$ ) so the pH responsiveness also becomes field dependent.<sup>31,47</sup> For example, the CEST effect for the Eu<sup>3+</sup>-bound water of EuDOTA–4AmC is independent of pH between pH 6 and 8 at 7.05 T<sup>46</sup> but not at 4.7 T.<sup>31</sup> Thus, one must establish ratiometric curves for each new compound and at each  $B_0$  field before placing this technique into practical use.

**Metabolite Responsive PARACEST Agents.** One example of this was given by a recent report showing that a PARACEST agent could be used to detect and quantify the glycolytic end product, lactate.<sup>48</sup> For this purpose, a Ln<sup>3+</sup>–tris-amide complex was used (a  $q = 2$  complex) so that lactate could bind into the inner coordination sphere of the Ln<sup>3+</sup> as a bidentate ligand. This resulted in a slow-exchanging ternary complex whose –NH resonances differed in chemical shift from the uncomplexed LnL –NH resonances by  $\sim 5$  ppm. Selective saturation of each –NH resonance (sequentially) resulted in a CEST effect at each site that was proportional to the concentration of each species, thereby allowing a quantitative comparison of their respective concentrations. It was shown the CEST ratio provides a direct readout of the lactate concentration in any equilibrium mixture that contains both LnL and LnL–lactate. Using this approach, one could envision the design of amide-based Ln<sup>3+</sup> chelates having a topological surface that allows selective binding of any metabolite within a complex biological mixture. This offers the possibility of in situ monitoring of metabolites by transmitting this binding information to bulk water via the CEST effect.

## Summary

Our general knowledge of lanthanide coordination chemistry was significantly advanced during the last two decades due to widespread scientific interest in Gd<sup>3+</sup>-based MRI contrast agents. Many important and surprising discoveries, especially concerning the dynamics of lanthanide complexes, that have come from these basic investigations will likely result in an even broader range of applications of lanthanide complexes to medical science. For example, we now know that complexes can be made with Ln<sup>3+</sup>-bound water lifetimes ( $\tau_M$ ) ranging from nanoseconds to milliseconds by simple modification of ligand side chain coordinating groups or by varying the central metal ion, solvent, or counterion. Complexes have been discovered wherein a single, bound water molecule is in such slow exchange with bulk water that a separate resonance appears in the high-resolution <sup>1</sup>H or <sup>17</sup>O NMR spectra of the complex, even at temperatures well above room temperature. This single feature alone already forms the basis of complexes that act as pH sensitive and metabolite specific MR agents. The recent demonstration that paramagnetic lanthanide complexes can act as efficient CEST agents offers the possibility of designing an array of new responsive agents for reporting biological events using conventional imaging technologies. Just as with Gd<sup>3+</sup>-based relaxation agents designed to respond

to enzyme activity,<sup>49</sup> pH,<sup>50</sup> Ca<sup>2+</sup>,<sup>51</sup> Zn<sup>2+</sup>,<sup>52</sup> and sugars,<sup>53</sup> we believe that PARACEST agents may offer a new avenue for research into biologically “responsive” agents capable of sensing molecular exchange phenomena in tissue.

*This work was supported in part by grants from the Robert A. Welch Foundation (AT-584), the National Institutes of Health (CA-84697), the Division of Research Resources, National Institutes of Health (RR-02584), and the Texas Advanced Technology Program.*

## References

- (1) Merbach, A. E.; Toth, E. *The Chemistry of Contrast Agents in Medical MRI*; Wiley: Chichester, 2001.
- (2) Lauffer, R. B. Paramagnetic metal complexes as water proton relaxation agents for NMR imaging: theory and design. *Chem. Rev.* **1987**, *87*, 901–927.
- (3) Caravan, P.; Ellison, J. J.; McMurray, T. J.; Lauffer, R. B. Gadolinium(III) Chelates as MRI Contrast Agents: Structure, Dynamics, and Applications. *Chem. Rev.* **1999**, *99*, 2293–2352.
- (4) Peters, J. A.; Huskens, J.; Raber, D. J. Lanthanide induced shifts and relaxation rate enhancements. *Prog. NMR Spectrosc.* **1996**, *28*, 283–350.
- (5) Springer, C. S., Jr. Measurement of metal cation compartmentalization in tissue by high-resolution metal cation NMR. *Annu. Rev. Biophys. Biophys. Chem.* **1987**, *16*, 375–399.
- (6) Bansal, N.; Germann, M. J.; Seshan, V.; Shires, G. T., III; Malloy, C. R.; Sherry, A. D. Thulium 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetrakis(methylene phosphonate) as a sodium-23 shift reagent for the in vivo rat liver. *Biochemistry* **1993**, *32*, 5638–5643.
- (7) Aime, S.; Barge, A.; Botta, M.; De Sousa, A. S.; Parker, D. Direct NMR spectroscopic observation of a lanthanide-coordinated water molecule whose exchange rate is dependent on the conformation of the complexes. *Angew. Chem., Int. Ed. Engl.* **1998**, *37*, 2673–2675.
- (8) Aime, S.; Barge, A.; Bruce, J. I.; Botta, M.; Howard, J. A. K.; Moloney, J. M.; Parker, D.; de Sousa, A. S.; Woods, M. NMR, Relaxometric, and Structural Studies of the Hydration and Exchange Dynamics of Cationic Lanthanide Complexes of Macrocyclic Tetraamide Ligands. *J. Am. Chem. Soc.* **1999**, *121*, 5762–5771.
- (9) Batsanov, A. S.; Beeby, A.; Bruce, J. I.; Howard, J. A. K.; Kenwright, A. M.; Parker, D. Direct NMR and luminescence observation of water exchange at cationic ytterbium and europium centres. *Chem. Commun.* **1999**, 1011–1012.
- (10) Dunand, F. A.; Aime, S.; Merbach, A. E. First <sup>17</sup>O NMR Observation of Coordinated Water on Both Isomers of [Eu(DOTAM)(H<sub>2</sub>O)]<sup>3+</sup>: A Direct Access to Water Exchange and its Role in the Isomerization. *J. Am. Chem. Soc.* **2000**, *122*, 1506–1512.
- (11) Zhang, S.; Wu, K.; Biewer, M. C.; Sherry, A. D. <sup>1</sup>H and <sup>17</sup>O NMR Detection of a Lanthanide-Bound Water Molecule at Ambient Temperatures in Pure Water as Solvent. *Inorg. Chem.* **2001**, *40*, 4284–4290.
- (12) Zhang, S.; Wu, K.; Sherry, A. D. Gd<sup>3+</sup> complexes with slowly exchanging bound-water molecules may offer advantages in the design of responsive MR agents. *Invest. Radiol.* **2001**, *36*, 82–86.
- (13) Dunand, F. A.; Dickins, R. S.; Parker, D.; Merbach, A. E. Towards rational design of fast water-exchanging Gd(dota-like) contrast agents? importance of the M/m ratio. *Chem. Eur. J.* **2001**, *7*, 5160–5167.
- (14) Zhang, S.; Wu, K.; Sherry, A. D. Unusually sharp dependence of water exchange rate versus lanthanide ionic radii for a series of tetra-amide complexes. *J. Am. Chem. Soc.* **2002**, *124*, 4226–4227.
- (15) Aime, S.; Barge, A.; Batsanov, A. S.; Botta, M.; Castelli, D. D.; Fedeli, F.; Mortillaro, A.; Parker, D.; Puschnann, H. Controlling the variation of axial water exchange rates in macrocyclic lanthanide(III) complexes. *Chem. Commun.* **2002**, 1120–1121.
- (16) Zhang, S.; Winter, P.; Wu, K.; Sherry, A. D. A Novel Europium(III)-Based MRI Contrast Agent. *J. Am. Chem. Soc.* **2001**, *123*, 1517–1518.
- (17) Wolf, S. D.; Balaban, R. S. Magnetization transfer contrast (MTC) and tissue water proton relaxation in vivo. *Magn. Reson. Med.* **1989**, *10*, 135–144.
- (18) Henkelman, R. M.; Stanisz, G. J.; Graham, S. J. Magnetization transfer in MRI: A review. *NMR Biomed.* **2001**, *14*, 57–64.
- (19) Ward, K. M.; Aletras, A. H.; Balaban, R. S. A New Class of Contrast Agents for MRI Based on Proton Chemical Exchange Dependent Saturation Transfer (CEST). *J. Magn. Reson.* **2000**, *143*, 79–87.
- (20) Sherry, A. D.; Zhang, S.; Wu, K. Paramagnetic metal ion-based macrocyclic magnetization transfer contrast agents and method of use. PCT Int. Appl. WO 0243775, 2002.

- (21) Micskei, K.; Helm, L.; Brucher, E.; Merbach, A. E. Oxygen-17 NMR study of water exchange on gadolinium polyaminopolyacetates  $[\text{Gd}(\text{DTPA})(\text{H}_2\text{O})]^{2-}$  and  $[\text{Gd}(\text{DOTA})(\text{H}_2\text{O})]^-$  related to NMR imaging. *Inorg. Chem.* **1993**, *32*, 3844–3850.
- (22) Powell, D. H.; Gonzalez, G.; Tissieres, V.; Micskei, K.; Bruecher, E.; Helm, L.; Merbach, A. E.  $\text{Gd}^{3+}$  chelates of interest in magnetic resonance imaging (MRI): studies using  $^{17}\text{O}$  NMR and EPR at several magnetic fields. *J. Alloys Compd.* **1994**, *207–208*, 20–4.
- (23) Cossy, C.; Helm, L.; Merbach, A. E. Water exchange kinetics on lanthanide(III) ions: a variable temperature and pressure oxygen-17 NMR study. *Inorg. Chim. Acta* **1987**, *139*, 147–149.
- (24) Cossy, C.; Helm, L.; Merbach, A. E. Oxygen-17 nuclear magnetic resonance kinetic study of water exchange on the lanthanide(III) aqua ions. *Inorg. Chem.* **1988**, *27*, 1973–1979.
- (25) Micskei, K.; Powell, D. H.; Helm, L.; Brucher, E.; Merbach, A. E. Water exchange on gadolinium (aqua)(propylenediamine tetraacetate) complexes  $[\text{Gd}(\text{H}_2\text{O})_8]^{3+}$  and  $[\text{Gd}(\text{PDTA})(\text{H}_2\text{O})_2]^-$  in aqueous solution: a variable-pressure, -temperature and -magnetic field oxygen-17 NMR study. *Magn. Reson. Chem.* **1993**, *31*, 1011–1020.
- (26) Aime, S.; Botta, M.; Fasano, M.; Paoletti, S.; Anelli, P. L.; Uggeri, F.; Virtuani, M. NMR Evidence of a Long Exchange Lifetime for the Coordinated Water in Ln(III)-Bis(methyl amide)-DTPA Complexes (Ln = Gd, Dy). *Inorg. Chem.* **1994**, *33*, 4707–4711.
- (27) Pubanz, D.; Gonzalez, G.; Powell, D. H.; Merbach, A. E. Unexpectedly Large Change of Water Exchange Rate and Mechanism on  $[\text{Ln}(\text{DTPA-BMA})(\text{H}_2\text{O})]$  Complexes along the Lanthanide(III) Series. *Inorg. Chem.* **1995**, *34*, 4447–4453.
- (28) Aime, S.; Barge, A.; Botta, M.; Parker, D.; De Sousa, A. S. Prototropic vs. whole water exchange contributions to the solvent relaxation enhancement in the aqueous solution of a cationic  $\text{Gd}^{3+}$  macrocyclic complex. *J. Am. Chem. Soc.* **1997**, *119*, 4767–4768.
- (29) Aime, S.; Botta, M.; Fasano, M.; Marques, M. P. M.; Geraldese, C. F. G. C.; Pubanz, D.; Merbach, A. E. Conformational and Coordination Equilibria on DOTA Complexes of Lanthanide Metal Ions in Aqueous Solution Studied by  $^1\text{H}$ -NMR Spectroscopy. *Inorg. Chem.* **1997**, *36*, 2059–2068.
- (30) Zhang, S.; Kovacs, Z.; Burgess, S.; Aime, S.; Terreno, E.; Sherry, A. D. {DOTA-bis(amide)} Lanthanide Complexes: NMR Evidence for Differences in Water-molecule Exchange Rates for Coordination Isomers. *Chem. Eur. J.* **2001**, *7*, 288–296.
- (31) Zhang, S.; Sherry, A. D. Paramagnetic CEST Agents: NH versus OH. *Int. Soc. Magn. Reson. Med.* **2002**, *10*, 2590.
- (32) Aime, S.; Botta, M.; Fasano, M.; Terreno, E. Prototropic and Water-Exchange Processes in Aqueous Solutions of Gd(III) Chelates. *Acc. Chem. Res.* **1999**, *32*, 941–949.
- (33) Zhang, S.; Wu, K.; Sherry, A. D. A novel pH-sensitive MRI contrast agent. *Angew. Chem., Int. Ed. Engl.* **1999**, *38*, 3192–3194.
- (34) Raghunand, N.; Howison, C.; Sherry, A. D.; Zhang, S.; Gillies, R. J. Renal and Systemic pH imaging by Contrast-enhanced MRI. *Magn. Reson. Med.* **2003**, *49*, 249–257.
- (35) McConnell, H. M. Reaction Rates by Nuclear Magnetic Resonance. *J. Chem. Phys.* **1958**, *28*, 430–431.
- (36) Woessner, D. E. Nuclear Transfer Effects in Nuclear Magnetic Resonance Pulse Experiments. *J. Chem. Phys.* **1961**, *35*, 41–48.
- (37) Grad, J.; Bryant, R. G. Nuclear magnetic cross-relaxation spectroscopy. *J. Magn. Reson.* **1990**, *90*, 1–8.
- (38) Zhang, S.; Sherry, A. D. Physical characteristics of lanthanide complexes that act as magnetization transfer (MT) contrast agents. *J. Solid State Chem.* **2003**, *171*, 38–43.
- (39) Ahrens, E. T.; Rothbacher, U.; Jacobs, R. E.; Fraser, S. E. A model for MRI contrast enhancement using  $T_1$  agents. *Proc. Natl. Acad. Sci. U.S.A.* **1998**, *95*, 8443–8448.
- (40) Lauffer, R. B.; Betteridge, D. R.; Padmanabhan, S.; Brady, T. J. Albumin binding of paramagnetic hepatobiliary contrast agents: enhancement of outer sphere relaxivity. *Nucl. Med. Biol.* **1988**, *15*, 45–46.
- (41) Toth, E.; Van Uffelen, I.; Helm, L.; Merbach, A. E.; Ladd, D.; Briley-Saebo, K.; Kellar, K. E. Gadolinium-based linear polymer with temperature-independent proton relaxivities: a unique interplay between the water exchange and rotational contributions. *Magn. Reson. Chem.* **1998**, *36*, S125–S134.
- (42) Bogdanov, A.; Matuszewski, L.; Bremer, C.; Petrovsky, A.; Weissleder, R. Oligomerization of Paramagnetic Substrates Results in Signal Amplification and Can Be Used for MR Imaging of Molecular Targets. *Mol. Imaging* **2001**, *1*, 1–9.
- (43) Goffeney, N.; Bulte, J. W. M.; Duyn, J.; Bryant, L. H.; van Zijl, P. C. M. Sensitive NMR detection of cationic-polymer-based gene delivery systems using saturation transfer via proton exchange. *J. Am. Chem. Soc.* **2001**, *123*, 8628–8629.
- (44) Ward, K. M.; Balaban, R. S. Determination of pH using water protons and chemical exchange dependent saturation transfer (CEST). *Magn. Reson. Med.* **2000**, *44*, 799–802.
- (45) Aime, S.; Barge, A.; Castelli, D. D.; Fedeli, F.; Mortrillaro, A.; Nielsen, F. U.; Terreno, E. Paramagnetic Lanthanide(III) complexes as pH-sensitive chemical exchange saturation transfer (CEST) contrast agents for MRI application. *Magn. Reson. Med.* **2002**, *47*, 639–648.
- (46) Aime, S.; Castelli, D. D.; Terreno, E. Novel pH-reporter MRI contrast agents. *Angew. Chem., Int. Ed.* **2002**, *41*, 4334–4336.
- (47) Zhang, S.; Michaudet, L.; Burgess, S.; Sherry, A. D. The amide protons of an ytterbium(III) dota tetraamide complex act as efficient antennae for transfer of magnetization to bulk water. *Angew. Chem., Int. Ed.* **2002**, *41*, 1919–1921.
- (48) Aime, S.; Delli Castelli, D.; Fedeli, F.; Terreno, E. A Paramagnetic MRI-CEST Agent Responsive to Lactate Concentration. *J. Am. Chem. Soc.* **2002**, *124*, 9364–9365.
- (49) Louie, A. Y.; Huber, M. M.; Ahrens, E. T.; Rothbacher, U.; Moats, R.; Jacobs, R. E.; Fraser, S. E.; Meade, T. J. *In vivo* visualization of gene expression using magnetic resonance imaging. *Nat. Biotechnol.* **2000**, *18*, 321–325.
- (50) Lowe, M. P.; Parker, D.; Reany, O.; Aime, S.; Botta, M.; Castellano, G.; Gianolio, E.; Pagliarin, R. pH-Dependent Modulation of Relaxivity and Luminescence in Macrocyclic Gadolinium and Europium Complexes Based on Reversible Intramolecular Sulfonamide Ligation. *J. Am. Chem. Soc.* **2001**, *123*, 7601–7609.
- (51) Li, W.-h.; Fraser, S. E.; Meade, T. J. A Calcium-Sensitive Magnetic Resonance Imaging Contrast Agent. *J. Am. Chem. Soc.* **1999**, *121*, 1413–1414.
- (52) Hanaoka, K.; Kikuchi, K.; Urano, Y.; Nagano, T. Selective sensing of zinc ions with a novel magnetic resonance imaging contrast agent. *J. Chem. Soc., Perkin Trans. 2* **2001**, 1840–1843.
- (53) Peters, J. A.; Frullano, L.; Rohovec, J. Sugar-targeted contrast agents for MRI. *J. Inorg. Biochem.* **2001**, *86*, 85–85.

AR020228M