

Formation of Carbon–Carbon Bonds under Catalysis by Transition-Metal Nanoparticles

MARCIAL MORENO-MAÑAS* AND
ROSER PLEIXATS

Department of Chemistry, Universitat Autònoma de
Barcelona, Cerdanyola, 08193-Barcelona, Spain

Received February 13, 2003

ABSTRACT

This Account presents the fast-growing field of catalysis by transition metal nanoparticles, its present situation concerning scope and limitations, as well as future prospects. The Account is limited to reactions involving carbon–carbon bond formation.

Introduction

Transition-metal nanoparticles have attracted a great deal of attention in the last 10 years; their preparation, structure determination, and applications are topics of current interest.^{1–14} Nanoparticles are defined as having 1–50 nm diameter, a size range where metals can show size-dependent properties. The smaller the cluster of atoms, the higher the percentage of atoms are on the surface, rendering nanoparticles very interesting in catalysis.^{6,11,14} Thus, a nanoparticle of 10 nm diameter has about 10% of its atoms in the surface, but one of 1 nm has 100%.⁷

The metal atoms constituting nanoparticles can be generated by (i) chemical reduction of a metal salt, (ii) thermal, photochemical, or sonochemical decomposition of a metal(0) complex, (iii) hydrogenation of a coordinating olefinic moiety, and (iv) vapor phase deposition. To this list proposed by Bradley² should be added (v) electrochemical reduction of higher valent species of the metal.⁹ During generation of nanoparticles, the following steps have been identified: (i) generation of atoms as above; (ii) nucleation to form an initial cluster of atoms; (iii) growing of the cluster until a certain volume is reached; and (iv) surrounding the cluster by a protecting shell that prevents agglomeration. Therefore, nanoparticles should be formed in the presence of a protecting agent. These protectors can be broadly divided into two categories: those providing electrostatic and those pro-

viding steric stabilization. The electrostatic stabilization is based upon the double electric layer formed when ions of the same sign are adsorbed at the nanoparticle surface (the charges in Figure 1 could be inverted). The counterions form a second layer that repels the neighboring nanoparticle; for example, sodium citrate acts by this mechanism. In other cases, protecting molecules of considerable length interact attractively with the surface of the nanoparticle. The volume of these surrounding molecules prevents mutual approach of metal surfaces at bonding distance. Popular protecting agents are polymers (poly(vinylpyrrolidone)), cyclodextrins, dendrimers, and so forth. Particularly well-known is the mechanism of stabilization by large molecules featuring functional groups with high affinity for the metal, i.e., thiols,¹⁵ sulfides, amines, or phosphines. Other common stabilizers are ionic surfactants, i.e., sodium dodecyl sulfate and lauryl trimethylammonium chloride. These compounds protect nanoparticles by both electrostatic and steric mechanisms (Figure 1). Heavily fluorinated compounds have also been found to stabilize metal nanoparticles.¹⁶

Size and dispersity are important properties of nanoparticles that are mainly determined by transmission electron microscopy (TEM). The use of metal nanoparticles for catalysis in reactions generating carbon–carbon bonds is in its infancy; however, we anticipate an important growth of activity in this field soon. This is the reason of this account.

Palladium-Catalyzed Reactions¹⁷

The Mizoroki–Heck Reaction. The Mizoroki–Heck (M–H) reaction is the arylation of olefins as indicated in Scheme 1.¹⁸ Aryl iodides are more active than bromides and chlorides in concordance with the energies of the C–X bonds. Main concerns are at present (i) finding catalytic systems that activate C–Cl bonds or alternatively, (ii) finding alternative cheaper leaving groups (e.g., diazonium cation, nitro, etc.), (iii) recovery and reuse of the catalysts without loss of activity, and (iv) development of phosphine-free palladium catalysts.


Reetz and co-workers showed in 1996 that tetraalkylammonium-stabilized palladium nanoparticles of 3 nm catalyzed the reaction of butyl acrylate, **1a**, with iodobenzene to afford butyl cinnamate **3a**.¹⁹ Reetz also reported M–H reactions of aryl bromides **2b** (R = H, *p*-COCH₃, *p*-NO₂) with styrene **1b**.²⁰ The catalysts were palladium nanoparticles (8–10 nm) generated either electrochemically or by thermolysis of palladium acetate (Scheme 2). The stabilizing shell was made of propylene carbonate. Chlorobenzene, **2c**, was only moderately active. Later on, Reetz and co-workers reported more details on the in situ generation of nanoparticles (1.3–1.6 nm) by thermolysis of palladium acetate (or palladium chloride plus sodium acetate) mainly in the presence of tetraalkylammonium

Marcial Moreno-Mañas was born in Barcelona, Spain, in 1941. After postdoctoral studies with Prof. Alan R. Katritzky at the University of East Anglia (U.K.), he joined the National Research Council in Spain (CSIC) and finally the "Universitat Autònoma de Barcelona", where he became full Professor in 1977. His interests embrace transition metals in organic chemistry, recovery and reuse of metal catalysts, fluorine chemistry, and nanoparticles.


Roser Pleixats studied chemistry at the "Universitat Autònoma de Barcelona", where she obtained her PhD in 1984. After postdoctoral research with Prof. R. J.P. Corriu and E. Colomer at the University of Montpellier II, she became Associate Professor at the "Universitat Autònoma de Barcelona" in 1991. Her research interests include palladium and other transition metal catalysts, metathesis reactions, nanoparticles, and catalyst recovery (fluorous biphasic systems, anchoring to organic and inorganic polymers).

* To whom correspondence should be addressed: Phone: 34-935811254. Fax: 34-935811265. E-mail: marcial.moreno@uab.es. Web page: <http://einstein.uab.es/mmorenos>.

Scheme 3. Stabilizer Used and Reactions Performed in Ref 43



Scheme 4. Precursors of Nanoparticles in Refs 44, 45, and 53



Our group reported the stabilization of palladium nanoparticles by the heavily fluorinated ketone **5** (Scheme 3). These unusually stabilized nanoparticles catalyzed the formation of ethyl cinnamate, **3a** (R = H), from iodobenzene, as well as ester **8** and nitrile **9** (Scheme 3). Efficient recovery and reuse without loss of activity was achieved by extraction of the catalytic material with fluoruous solvents.⁴³

Finally, Gladysz and co-workers have used palladium(II) complexes of fluoruous palladacycles **10**⁴⁴ and **11**⁴⁵ as catalyst precursor in M–H reactions (Scheme 4). They gave evidence favoring nonfluorous nanoparticles as the actual catalytic species. Palladacycle **10** acts as steady-state source of very reactive, DMF-soluble nanoparticles, probably stabilized by ammonium ions during the M–H reaction. TON's are as high as 1.51 million, making this catalytic system exceptional.

The Suzuki Cross-Coupling. The Suzuki cross-coupling between arylboronic acids **12** and aryl halides produces biphenyls **13** (Scheme 1).⁴⁶ Possibly the first report on the use of palladium nanoparticles in Suzuki coupling is due to Reetz and co-workers.^{19,21} This pioneering work¹⁹ described simple biphenyl formation from aryl bromides and even from activated aryl chlorides possessing electron-attracting groups.

Some groups already mentioned in connection with the M–H reaction tested their catalytic materials also in Suzuki couplings.^{27,32,35} More examples of palladium nanoparticles as catalysts in the Suzuki cross-coupling have been described.^{47–52} El-Sayed et al. concluded that vertex and edge atoms on the palladium particle surfaces are the active sites.⁴⁹ The most salient features of these contributions are some examples of recycling of the catalytic material^{35,51} and the activation of aryl chlorides.^{19,32,35} We have reported that palladium particles stabilized by fluo-

rous ketone **5** are also active in the Suzuki-type coupling of arylboronic acids with cinnamyl bromide (sp²–sp³ coupling).⁴³

Gladysz et al. reported Suzuki couplings in the presence of fluoruous sulfide complexes **14** and **15** (Scheme 4).^{45,53} As for the case of M–H reactions (vide supra), they concluded that the complexes were precursors of nanoparticles stabilized by nonfluorous shells.

Miscellaneous Reactions Catalyzed by Palladium Nanoparticles. Some examples of Stille-type^{27,32,35} and Sonogashira couplings^{30,35} have been reported (Scheme 5). Curiously, these Sonogashira couplings do not require copper(I) species in the catalytic system, as it happens with more conventional palladium catalysts.

Hydrocarbonylation of olefins,⁵⁴ aromatic couplings,⁵⁵ and hydrodechlorination of tetrachloromethane⁵⁶ are reactions involving C–C bond formation that have been catalyzed by palladium nanoparticles (Scheme 5).

For two cases of Tsuji–Trost allylation of carbon nucleophiles with palladium nanoparticle catalysis, see the discussion in the following cobalt paragraph.^{61,62}

Cobalt-Catalyzed Reactions

The Pauson–Khand (P–K) reaction permits preparation of cyclopentenones by condensation of alkenes, alkynes, and carbon monoxide (see Scheme 6 for the intramolecular version). The classical P–K reaction uses stoichiometric amount of Co₂(CO)₈ as source of carbon monoxide, but attempts to use catalytic metal in an atmosphere of carbon monoxide have been reported.⁵⁷ Colloidal cobalt (reduction of Co(II) or thermal decomposition of Co₂(CO)₈) has been applied successfully as catalyst in the reactions of Scheme 6,^{58–62} and in some cases recovery has been possible.^{58–61} Catalytic material containing both cobalt and palladium nanoparticles catalyze sequentially Pauson–Khand and Tsuji–Trost allylation of nucleophile (Scheme 6).^{61,62}




Reactions Catalyzed by Other Metal Nanoparticles

Nanoparticles made from several transition metals catalyze biphenyl formation by Suzuki cross-coupling of **2a**, **b** with **12**. The particles were prepared by reduction of pure or mixed metal chlorides with tetraoctylammonium formate. The following combinations gave interesting results: Cu, Pd, Ru, Cu/Pd, Pd/Pt, Pd/Ru, Cu/Pd/Pt, Cu/Pd/Ru, Pd/Pt/Ru, and Cu/Pd/Pt/Ru.⁶³ This paper particularly is noteworthy since it is a step forward in finding alternatives to palladium catalysts.



Some other cases pertinent to this Account have been described: acetylene polymerization under copper nanoparticle catalysis,⁶⁴ ring-opening metathesis polymerization under catalysis by a ruthenium complex supported on gold colloids,⁶⁵ McMurry-type couplings induced by nanostructured titanium clusters,⁶⁶ [3+2] cycloaddition reactions under nickel clusters catalysis,⁶⁷ and carbonylation of methanol catalyzed by rhodium colloids.⁶⁸

Scheme 5. Other Reactions Catalyzed by Palladium Nanoparticles





Stille-type cross-couplings




Sonogashira couplings



Other reactions


Scheme 6. The Pauson–Khand Reaction

 A = (ROCO)₂C, CH₂, O

Catalytic version



Ref. 61 and 62

expected soon. Some difficulties have to be solved before the use of nanoparticles in C–C and C-heteroatom bonds will be generalized. In our opinion, reproducibility in the properties of nanoparticles (size, dispersity, and performance), and recovery and reuse without loss of activity are of paramount importance. Therefore, efforts to deal with more reactions and to use more metals will be significant, but efforts to solve those important issues of reproducibility and recovery will be particularly welcome.

It might be that the use of ionic solvents could help to solve the problem of reutilization of the catalytic material. Studies related to the Mizoroki–Heck reaction have opened the question of whether nanoparticles are the actual catalytic material in ionic solvents, as well as when using *N*-heterocyclic carbenes as palladium sources.^{30,69,70}

Financial support has been obtained from "Ministerio de Ciencia y Tecnología" of Spain (Project BQU2002-04002) and from Generalitat de Catalunya (Project 2001SGR00181).

Conclusions

The papers collected below dealing directly with original research on the topic of this account were published in 1996 (4), 1997 (3), 1998 (3), 1999 (4), 2000 (9), 2001 (8), 2002 (15), and 2003 (3 until January). An increase is

Note Added after ASAP Posting

This paper first appeared on the Web with an error in Scheme 1. The correct version was published 7/9/2003.

References

- Lewis, L. N. *Chemical Catalysis by Colloids and Clusters*. *Chem. Rev.* **1993**, *93*, 2693–2730.
- Bradley, J. S. *The Chemistry of Transition Metal Colloids. In Clusters and Colloids, From Theory to Applications*; Schmid, G., Ed.; VCH: Weinheim, Germany, 1994; pp 459–544.
- In *Metal Clusters in Chemistry*; Braunstein, P.; Oro, L.; Raithby, P. R., Eds.; Wiley-VCH: Weinheim, Germany, 1998.
- In *Nanoparticles and Nanostructured Films. Preparation, Characterization and Applications*; Fendler, J. H., Ed.; Wiley-VCH: Weinheim, Germany, 1998.
- Klabunde, K. J.; Mohs, C. *Nanoparticles and Nanostructural Materials*. In *Chemistry of Advanced Materials. An Overview*; Interrante, L. V., Hampden-Smith, M. J., Eds.; Wiley-VCH: New York, 1998; Chapter 7, pp 271–327.
- Aiken III, J. D.; Finke, R. G. A Review of Modern Transition-Metal Nanoclusters: Their Synthesis, Characterization, and Applications in Catalysis. *J. Mol. Catal. A* **1999**, *145*, 1–44.
- Rao, C. N. R.; Kulkarni, G. U.; Thomas, P. J.; Edwards, P. P. *Metal Nanoparticles and Their Assemblies*. *Chem. Soc. Rev.* **2000**, *29*, 27–35.
- Horn, D.; Rieger, J. *Organic Nanoparticles in the Aqueous Phase – Theory, Experiment, and Use*. *Angew. Chem., Int. Ed. Engl.* **2001**, *40*, 4330–4361.
- Reetz, M. T.; Winter, M.; Breinbauer, R.; Thurn-Albrecht, T.; Vogel, W. Size-Selective Electrochemical Preparation of Surfactant-Stabilized Pd-, Ni- and Pt/Pd Colloids. *Chem. Eur. J.* **2001**, *7*, 1084–1094.
- Caruso, F. *Nanoengineering of Particle Surfaces*. *Adv. Mater.* **2001**, *13*, 11–22.
- Bönnemann, H.; Richards, R. M. *Nanoscope Metal Particles – Synthetic Methods and Potential Applications*. *Eur. J. Inorg. Chem.* **2001**, 2455–2480.
- Rao, C. N. R.; Kulkarni, G. U.; Thomas, P. J.; Edwards, P. P. *Size-Dependent Chemistry: Properties of Nanocrystals*. *Chem. Eur. J.* **2002**, *8*, 28–35.
- In *Metal Nanoparticles. Synthesis, Characterization, and Applications*; Feldheim, D. L., Foss, C. A., Jr., Eds.; Marcel Dekker: New York, 2002.
- Roucoux, A.; Schulz, J.; Patin, H. *Reduced Transition Metal Colloids: A Novel Family of Reusable Catalysts?* *Chem. Rev.* **2002**, *102*, 3757–3778.
- Templeton, A. C.; Wuelfing, W. P.; Murray, R. W. *Monolayer-Protected Cluster Molecules*. *Acc. Chem. Res.* **2000**, *33*, 27–36.
- Moreno-Mañas, M.; Pleixats, R. *Fluorous Nanoparticles*. In *Handbook of Fluorous Chemistry*; Gladysz, J. A., Curran, D. P., Horváth, I. T., Eds.; Wiley-VCH: in press.
- For a comprehensive book on palladium in organic chemistry, see: In *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., Ed.; Wiley-Interscience: New York, 2002.
- For most recent reviews see: (a) Crisp, G. T. *Variations on a Theme – Recent Developments on the Mechanism of the Heck Reaction and Their Implications for the Synthesis*. *Chem. Soc. Rev.* **1998**, *27*, 427–436. (b) Beletskaya, I. P.; Chepurkov, A. V. *The Heck Reaction as a Sharpening Stone of Palladium Catalysis*. *Chem. Rev.* **2000**, *100*, 3009–3066. (c) Whitcombe, N. J.; Hii, K. K.; Gibson, S. E. *Advances in the Heck Chemistry of Aryl Bromides and Chlorides*. *Tetrahedron* **2001**, *57*, 7449–7476. (d) Bhanage, B. M.; Arai, M. *Catalyst Product Separation Techniques in Heck Reaction*. *Catal. Rev.* **2001**, *43*, 315–344.
- Reetz, M. T.; Breinbauer, R.; Wanninger, K. *Suzuki and Heck Reactions Catalyzed by Preformed Palladium Clusters and Palladium/Nickel Bimetallic Clusters*. *Tetrahedron Lett.* **1996**, *37*, 4499–4502.
- Reetz, M. T.; Lohmer, G. *Propylene Carbonate Stabilized Nanostructured Palladium Clusters as Catalysts in Heck Reactions*. *Chem. Commun.* **1996**, 1921–1922.
- Reetz, M. T.; Westermann, E. *Phosphane-Free Palladium-Catalyzed Coupling Reactions: The Decisive Role of Pd Nanoparticles*. *Angew. Chem., Int. Ed. Engl.* **2000**, *39*, 165–168.
- Jeffery, T. *On the Efficiency of Tetraalkylammonium Salts in Heck Type Reactions*. *Tetrahedron* **1996**, *52*, 10113–10130.
- Beller, M.; Fischer, H.; Kühlein, K.; Reisinger, C.-P.; Herrmann, W. A. *First Palladium-Catalyzed Heck Reactions with Efficient Colloidal Catalyst Systems*. *J. Organomet. Chem.* **1996**, *520*, 257–259.
- Klingelhöfer, S.; Heitz, W.; Greiner, A.; Oestreich, S.; Förster, S.; Antoniotti, M. *Preparation of Palladium Colloids in Block Copolymer Micelles and Their use for the Catalysis of the Heck Reaction*. *J. Am. Chem. Soc.* **1997**, *119*, 10116–10120.
- Dhas, N. A.; Gedanken, A. *Sonochemical Preparation and Properties of Nanostructured Palladium Metallic Clusters*. *J. Mater. Chem.* **1998**, *8*, 445–450.
- Walter, J.; Heiermann, J.; Dyker, G.; Hara, S.; Shioyama, H. *Hexagonal or Quasi Two-Dimensional Palladium Nanoparticles – Tested at the Heck Reaction*. *J. Catal.* **2000**, *189*, 449–455.
- Pathak, S.; Greci, M. T.; Kwong, R. C.; Mercado, K.; Prakash, G. K. S.; Olah, G. A.; Thompson, M. E. *Synthesis and Applications of Palladium-Coated Poly(vinylpyridine) Nanospheres*. *Chem. Mater.* **2000**, *12*, 1985–1989.
- Zhao, F.; Bhanage, B. M.; Shirai, M.; Arai, M. *Heck Reactions of Iodobenzene and Methyl Acrylate with Conventional Supported Palladium Catalysts in the Presence of Organic and/or Inorganic Bases without Ligands*. *Chem. Eur. J.* **2000**, *6*, 843–848.
- (a) Ding, J. A.; Gin, D. L. *Catalytically Active Pd(0) Nanocomposites Based on a Liquid Crystal Template*. *Polymer Preprints* **1999**, *40*, 738–739. (b) Ding, J. H.; Gin, D. L. *Catalytic Pd Nanoparticles Synthesized Using a Lyotropic Liquid Crystal Polymer Template*. *Chem. Mater.* **2000**, *12*, 22–24.
- Deshmukh, R. R.; Rajagopal, R.; Srinivasan, K. V. *Ultrasound Promoted C–C Bond Formation: Heck Reaction at Ambient Conditions in Room-Temperature Ionic Liquids*. *Chem. Commun.* **2001**, 1544–1545.
- Mori, K.; Yamaguchi, K.; Hara, T.; Mizugaki, T.; Ebitani, K.; Kaneda, K. *Controlled Synthesis of Hydroxyapatite-Supported Palladium Complexes as Highly Efficient Heterogeneous Catalysts*. *J. Am. Chem. Soc.* **2002**, *124*, 11572–11573.
- Kogan, V.; Aizenshtat, Z.; Popovitz-Biro, R.; Neumann, R. *Carbon–Carbon and Carbon–Nitrogen Coupling Reactions Catalyzed by Palladium Nanoparticles Derived from a Palladium Substituted Keggin-Type Polyoxometalate*. *Org. Lett.* **2002**, *4*, 3529–3532.
- Choudary, B. M.; Chowdari, N. S.; Jyothi, K.; Kumar, N. S.; Kantam, M. L. *A New Bifunctional Catalyst for Tandem Heck-Asymmetric Dihydroxylation of Olefins*. *Chem. Commun.* **2002**, 586–587.
- Galow, T. H.; Drechsler, U.; Hanson, J. A.; Rotello, V. M. *Highly Reactive Heterogeneous Heck and Hydrogenation Catalysts Constructed through “Bottom-up” Nanoparticle Self-Assembly*. *Chem. Commun.* **2002**, 1076–1077.
- Choudary, B. M.; Madhi, S.; Chowdari, N. S.; Kantam, M. L.; Sreedhar, B. *Layered Double Hydroxide Supported Nanopalladium Catalyst for Heck-, Suzuki-, Sonogashira-, and Stille-Type Coupling Reactions of Chloroarenes*. *J. Am. Chem. Soc.* **2002**, *124*, 14127–14136.
- Bosman, A. W.; Vestberg, R.; Heumann, A.; Fréchet, J. M. J.; Hawker, C. J. *A Modular Approach toward Functionalized Three-Dimensional Macromolecules: From Synthetic Concepts to Practical Applications*. *J. Am. Chem. Soc.* **2003**, *125*, 715–728.
- Dhas, N. A.; Cohen, H.; Gedanken, A. *In Situ Preparation of Amorphous Carbon-Activated Palladium Nanoparticles*. *J. Phys. Chem. B* **1997**, *101*, 6834–6838.
- Le Bars, J.; Specht, U.; Bradley, J. S.; Blackmond, D. G. *A Catalytic Probe of the Surface of Colloidal Palladium Particles Using Heck Coupling Reactions*. *Langmuir* **1999**, *15*, 7621–7625.
- Yeung, L. K.; Crooks, R. M. *Heck Heterocoupling within a Dendritic Nanoreactor*. *Nanoletters* **2001**, *1*, 14–17.
- For reviews on catalysis by nanoparticles by Crooks group see: (a) Crooks, R. M.; Zhao, M.; Sun, L.; Chechik, V.; Yeung, L. K. *Dendrimer-Encapsulated Metal Nanoparticles: Synthesis, Characterization, and Applications in Catalysis*. *Acc. Chem. Res.* **2001**, *34*, 181–190. (b) Crooks, R. M.; Lemon, B. I., III; Sun, L.; Yeung, L. K.; Zhao, M. *Dendrimer-Encapsulated Metals and Semiconductors: Synthesis, Characterization, and Applications*. *Top. Curr. Chem.* **2001**, *212*, 81–135.
- Yeung, L. K.; Lee, C. T., Jr.; Johnston, K. P.; Crooks, R. M. *Catalysis in Supercritical CO₂ Using Dendrimer-Encapsulated Palladium Nanoparticles*. *Chem. Commun.* **2001**, 2290–2291.
- Rahim, E. H.; Kamounah, F. S.; Frederiksen, J.; Christensen, J. B. *Heck Reactions Catalyzed by PAMAM-Dendrimer Encapsulated Pd(0) Nanoparticles*. *Nanoletters* **2001**, *1*, 499–501.
- Moreno-Mañas, M.; Pleixats, R.; Villarroja, S. *Fluorous Phase Soluble Palladium Nanoparticles as Recoverable Catalysts for Suzuki Cross-Coupling and Heck Reactions*. *Organometallics* **2001**, *20*, 4524–4528.
- Rocaboy, C.; Gladysz, J. A. *Highly Active Thermomorphic Fluorous Palladacycle Catalyst Precursors for the Heck Reaction: Evidence for a Palladium Nanoparticle Pathway*. *Org. Lett.* **2002**, *4*, 1993–1996.
- Rocaboy, C.; Gladysz, J. A. *Thermomorphic Fluorous Imine and Thioether Palladacycles as Precursors for Highly Active Heck and Suzuki Catalysts: Evidence for Palladium Nanoparticle Pathways*. *New J. Chem.* **2003**, *27*, 39–49.
- For recent reviews on the Suzuki coupling see: (a) Suzuki, A. *Recent Advances in the Cross-Coupling Reactions of Organoboron Derivatives with Organic Electrophiles, 1995–1998*. *J. Organomet. Chem.* **1999**, *576*, 147–168. (b) Koitha, S.; Lahiri, K.; Kashinath, D. *Tetrahedron* **2002**, *58*, 9633–9695.

- (47) Li, Y.; Hong, X. M.; Collard, D. M.; El-Sayed, M. A. Suzuki Cross-Coupling Reactions Catalyzed by Palladium Nanoparticles in Aqueous Solution. *Org. Lett.* **2000**, *2*, 2385–2388.
- (48) Li, Y.; El-Sayed, M. A. The Effect of Stabilizers on the Catalytic Activity of Pd Colloidal Nanoparticles in the Suzuki Reactions in Aqueous Solution. *J. Phys. Chem. B* **2001**, *105*, 8938–8943.
- (49) Li, Y.; Boone, E.; El-Sayed, M. A. Size Effects of PVP–Pd Nanoparticles on the Catalytic Suzuki Reactions in Aqueous Solution. *Langmuir* **2002**, *18*, 4921–4925.
- (50) Ramarao, C.; Ley, S. V.; Smith, S. C.; Shirley, I. M.; DeAlmeida, N. Encapsulation of Palladium in Polyurea Microcapsules. *Chem. Commun.* **2002**, 1132–1133.
- (51) Kim, S.-W.; Kim, M.; Lee, W. Y.; Hyeon, T. Fabrication of Hollow Palladium Spheres and Their Successful Applications to the Recyclable Heterogeneous Catalyst for Suzuki Coupling Reactions. *J. Am. Chem. Soc.* **2002**, *124*, 7642–7643.
- (52) Strimbu, L.; Liu, J.; Kaifer, A. E. Cyclodextrin-Capped Palladium Nanoparticles as Catalysts for the Suzuki Reaction. *Langmuir* **2003**, *19*, 483–485.
- (53) Rocaboy, C.; Gladysz, J. A. Synthesis, Oxidations, and Palladium Complexes of Fluorous Dialkyl Sulfides: New Precursors to Highly Active Catalysts for the Suzuki Coupling. *Tetrahedron* **2002**, *58*, 4007–4014.
- (54) Bertoux, F.; Monflier, E.; Castanet, Y.; Mortreux, A. Palladium Catalyzed Hydroxycarbonylation of Olefins in Biphasic system: Beneficial Effect of Alkali Metal Salt and Protective-Colloid Agents on the Stability of the Catalytic System. *J. Mol. Catal. A* **1999**, *143*, 23–30.
- (55) Mukhopadhyay, S.; Rothenberg, G.; Gitis, D.; Sasson, Y. Tandem One-Pot Palladium-Catalyzed Reductive and Oxidative Coupling of Benzene and Chlorobenzene. *J. Org. Chem.* **2000**, *65*, 3107–3110.
- (56) Lokteva, E. S.; Lunin, V. V.; Golubina, E. V.; Simagina, V. I.; Egorova, M.; Stoyanova, I. V. C–C Bond Formation during Hydrodechlorination of CCl₄ on Pd-Containing Catalysts. *Stud. Surf. Sci. Catal.* **2000**, *130C*, 1997–2002.
- (57) For recent reviews on the Pauson-Khand reaction see: (a) Chung, Y. K. Transition Metal Alkyne Complexes: the Pauson-Khand Reaction. *Coord. Chem. Rev.* **1999**, *188*, 297–341. (b) Fletcher, A. J.; Christie, S. D. R. *J. Chem. Soc., Perkin Trans. 1* **2000**, 1657–1668.
- (58) Kim, S.-W.; Son, S. U.; Lee, S. S.; Hyeon, T.; Chung, Y. K. Colloidal Cobalt Nanoparticles: a Highly Active and Reusable Pauson-Khand Catalyst. *Chem. Commun.* **2001**, 2212–2213.
- (59) Son, S. U.; Lee, S. I.; Chung, Y. K.; Kim, S.-W.; Hyeon, T. The First Intramolecular Pauson-Khand Reaction in Water Using Aqueous Colloidal Cobalt Nanoparticles as Catalysts. *Org. Lett.* **2002**, *4*, 277–279.
- (60) Son, S. U.; Park, K. H.; Chung, Y. K. Cobalt Nanoparticles on Charcoal: A Versatile Catalyst in the Pauson-Khand Reaction, Hydrogenation, and the Reductive Pauson-Khand Reaction. *Org. Lett.* **2002**, *4*, 3983–3986.
- (61) Park, K. H.; Son, S. U.; Chung, Y. K. Sequential Actions of Palladium and Cobalt Nanoparticles Immobilized on Silica: One-Pot Synthesis of Bicyclic Enones by Catalytic Allylic Alkylation and Pauson-Khand Reaction. *Org. Lett.* **2002**, *4*, 4361–4363.
- (62) Son, S. U.; Park, K. H.; Chung, Y. K. Sequential Actions of Cobalt Nanoparticles and Palladium(II) Catalysts: Three-Step One-Pot Synthesis of Fenestranes from an Enyne and an Alkyne Diester. *J. Am. Chem. Soc.* **2002**, *124*, 6838–6839.
- (63) Thathagar, M. B.; Beckers, J.; Rothenberg, G. Copper-Catalyzed Suzuki Cross-Coupling Using Mixed Nanocluster Catalysis. *J. Am. Chem. Soc.* **2002**, *124*, 11858–11859.
- (64) Chen, K.; Zhang, Z.; Cui, Z.; Yang, D. IR Characterization of Products of Acetylene Polymerization over Nano-Cu Particles. *Gaofenzi Xuebao* **2000**, *180*–183; *Chem. Abstr.* **133**, 59490.
- (65) Bartz, M.; Küther, J.; Seshadri, R.; Tremel, W. Colloid-Bound Catalyst for Ring-Opening Metathesis Polymerization: A Combination of Homogeneous and Heterogeneous Properties. *Angew. Chem., Int. Ed. Engl.* **1998**, *37*, 2466–2468.
- (66) Reetz, M. T.; Quaiser, S. A.; Merk, C. Electrochemical Preparation of Nanostructured Titanium Clusters. Characterization and Use in McMurry-Type Coupling Reactions. *Chem. Ber.* **1996**, *129*, 741–743.
- (67) Reetz, M. T.; Breinbauer, R.; Wedemann, P.; Binger, P. Nanostructured Nickel-Clusters as Catalysts in [3+2] Cycloaddition Reactions. *Tetrahedron* **1998**, *54*, 1233–1240.
- (68) Wang, Q.; Liu, H.; Han, M.; Li, X.; Jiang, D. Carbonylation of Methanol Catalyzed by Polymer-Protected Rhodium Colloid. *J. Mol. Catal. A* **1997**, *118*, 145–151.
- (69) Xu, L.; Chen, W.; Xiao, J. Heck Reaction in Ionic Liquids and the in Situ Identification of *N*-Heterocyclic Carbene Complexes of Palladium. *Organometallics* **2000**, *19*, 1123–1127.
- (70) Caló, V.; Nacci, A.; Monopoli, A.; Laera, S.; Cioffi, N. Pd Nanoparticles Catalyzed Stereospecific Synthesis of β -Aryl Cinnamic Esters in Ionic Liquids. *J. Org. Chem.* **2003**, *68*, 2929–2933.
- (71) For reviews on ionic liquids see: (a) Welton, T. Room-Temperature Ionic Liquids. Solvents for Synthesis and Catalysis. *Chem. Rev.* **1999**, *99*, 2071–2083. (b) Earle, M.; Seddon, K. R. Ionic Liquids. Green Solvents for the Future. *Pure Appl. Chem.* **2000**, *72*, 1391–1398.

AR020267Y