

# Cyclic Halonium Ions with Five-Membered Rings


PAUL E. PETERSON\*

Department of Chemistry, St. Louis University, St. Louis, Missouri 63156

Received October 12, 1970

The reaction of the primary *p*-nitrobenzenesulfonate **1** in trifluoroacetic acid may be considered to proceed via two competing paths having rate constants designated  $k_{\Delta}$  and  $k_s$ , respectively (Scheme I). Recent work in our laboratories<sup>1</sup> indicates that reaction via the path having rate constant  $k_{\Delta}$  strongly predominates over the

**Scheme I**  
Competing Paths for Trifluoroacetylation of 4-Chloro-1-pentyl Nosylate




alternative path, the ratio  $k_{\Delta}/k_s$  being approximately 760. The intermediacy of the five-membered ring halonium ion **2** in the " $k_{\Delta}$  reaction" is strongly implied by the structure, **4**, of the isolated reaction product, which is seen to be the result of a 1,4 halogen shift. This example of neighboring group participation by the somewhat remote neighbor chlorine is representative of the relatively recently discovered reactions proceeding via five-membered-ring halonium ions, to be examined in this Account. The preparation of the ions as stable species observable by nmr in  $\text{SO}_2\text{-SbF}_5$  solutions<sup>2</sup> also will be described.

The concept, employed in Scheme I, that competing paths, one involving neighboring group participation, occur in a substantial number of reactions has been a fruitful one.<sup>3,4</sup> An analysis comparable to that in Scheme I was used in 1948 by Winstein, Grunwald, and Ingraham in adducing evidence for three-membered-ring halonium ion intermediates in the acetolysis (reaction in acetic acid) of *trans*-2-halocyclohexyl

brosylates (Scheme II, X = I, Br, and possibly Cl).<sup>5</sup> The label  $k_{\Delta}$  which they applied to the rate constant for the path involving participation was coined to signify that this additional path could lead to an increased total rate constant,  $k_T$ , for disappearance of reactant.

It should be noted that in 1948 the nonparticipation reaction was assumed to involve the formation of a planar carbonium ion in the rate-determining step.<sup>5a</sup> The associated rate constant was designated by the symbol  $k_c$ . More recent studies<sup>6</sup> make it clear that transition states for the solvolysis of secondary as well as primary substrates exhibit considerable bond formation to solvent, which results, ultimately in the formation of the product of Walden inversion.<sup>7</sup> The designation  $k_s$  has been used for the rate constant of this solvent-assisted process (as in Scheme II). Both the  $k_s$

**Scheme II**  
Example of a Solvolysis Reaction Thought to Involve a Halonium Ion Intermediate



and the  $k_{\Delta}$  reactions may, in a broad sense, be considered in most instances to be "carbonium ion reactions," since in the respective transition states of the rate-determining steps a partial positive charge may appear on the carbon atom bearing the leaving group (OBs in Scheme II).

The earliest postulation of a halonium ion intermediate in a reaction seems to have been made by Roberts and Kimball, in their now familiar rationalization of the trans addition of bromine to alkenes.<sup>8</sup> The general form of their postulate is illustrated (Scheme III) by some recent examples of addition reactions in which

\* Address correspondence to Chemistry Department, University of South Carolina, Columbia, S. C. 29208.

(1) (a) P. E. Peterson and J. F. Coffey, *J. Amer. Chem. Soc.*, **93**, 5208 (1971); (b) P. E. Peterson and J. F. Coffey, *Tetrahedron Lett.*, 3131 (1968).

(2) For a review, see G. A. Olah, *Science*, **168**, 1298 (1970).

(3) For a review, see B. Capon, *Quart. Rev., Chem. Soc.*, **18**, 21 (1964).

(4) A striking example of recent progress is the apparently precise determination of the extent of phenyl participation in the previously controversial solvolysis of 3-phenyl-2-butyl tosylate: H. C. Brown, C. J. Kim, C. J. Lancelot, and P. v. R. Schleyer, *J. Amer. Chem. Soc.*, **92**, 5245 (1970).


(5) (a) S. Winstein, E. Grunwald, and L. L. Ingraham, *ibid.*, **70**, 821 (1948); (b) E. Grunwald, *ibid.*, **73**, 5458 (1951).

(6) P. v. R. Schleyer, J. L. Fry, L. K. M. Lam, and C. J. Lancelot, *ibid.*, **92**, 2542 (1970), and accompanying papers.

(7) A. Streitwieser, Jr., T. O. Walsh, and J. R. Wolfe, Jr., *ibid.*, **87**, 3682 (1965).

(8) I. Roberts and G. E. Kimball, *ibid.*, **59**, 947 (1937).


**Scheme III**  
Examples of Addition Reactions Thought to Involve Halonium Ion Intermediates



chloronium, bromonium, and iodonium ion intermediates are thought to open with Walden inversion to give overall trans additions to alkenes.<sup>9</sup>

**Additions to Alkenes.** In our research group 1,4 halogen participation and the concomitant five-membered-ring halonium ions were first invoked<sup>10</sup> to explain the rates of reaction of 5-chloro-, 5-bromo-, and 5-iodo-1-hexene with trifluoroacetic acid (Scheme IV). It is

**Scheme IV**  
Competing Paths in the Addition of Trifluoroacetic Acid to 5-Halo-1-hexenes



interesting to note that the inference of halogen participation was made on the basis of rate evidence which a number of chemists might have regarded as an overinterpretation of small rate differences. Allen's data<sup>10a</sup> are given in Table I.

We argued that the reaction of 5-halo-1-hexenes would have been retarded to a greater extent than was observed (3.35- to 4.13-fold) owing to the inductive effect of halogen if a second reaction, 1,4 halogen participation, had not contributed to the overall rate.

(9) For a review, see R. C. Fahey, *Top. Stereochem.*, **3**, 237 (1968).  
(10) (a) P. E. Peterson and G. Allen, *J. Amer. Chem. Soc.*, **85**, 3608 (1963). (b) The idea for such participation had, as we shall mention later, been considered in Winstein's research group and in a few scattered references.

**Table I**  
Relative Rates of Reaction of 5-Substituted-1-hexenes with Trifluoroacetic Acid

Substituent, X	$k_H/k_X$ , 60°
H	1
Cl	4.13
Br	3.35
I	3.50
O <sub>2</sub> CCH <sub>3</sub>	32.6
O <sub>2</sub> CCF <sub>3</sub>	167

However, solvent-induced modification of the electronegativity of some substituents, owing to hydrogen bonding of trifluoroacetic acid to the substituent, prevented a quantitative assessment of the extent of neighboring group participation using the Hammett-Taft substituted substituent effect equation,  $\log k_X/k_H = \rho\sigma_I$ . The closely similar behavior of the various halogens was puzzling, as we shall see later.

The symmetrical halonium ion **9** should, in the absence of complications, react to give a product, **10**, which is 50% halogen shifted. An alkene corresponding to **8**, X = Cl, but having a deuterium label on the carbon bearing chlorine, reacted to give a product showing the presence of 40% CH<sub>3</sub>CHCl groups, in addition to 60% CH<sub>3</sub>CDCl groups.<sup>11</sup> This result suggested that reaction occurred predominantly *via* ion **9**.

Determinations of the rates of reaction of a large number of substituted alkenes, CH<sub>2</sub>=CH(CH<sub>2</sub>)<sub>n-2</sub>X, permitted us to give a more firmly based interpretation of the rates of reaction of substituted alkenes, including a quantitative separation of overall rate constants into the presumed components,  $k_A$  and  $k_S$ , for those cases where the substituent, X, participates. We assumed<sup>12</sup> that the rate-retarding inductive free energy effect of a substituent upon the normal addition would fall off by a constant factor,  $\epsilon$ , per methylene group—a rule of thumb long used by chemists.<sup>13,14</sup> Alkenes having non-participating substituents (cyano, trifluoroacetoxy, and acetoxy) obeyed this rule extraordinarily well, as seen in the linear graphical representation of the appropriate function<sup>14</sup> of the data (Figure 1), which we have termed an attenuation plot.<sup>15</sup> Deviations from the rule shown by 5-chloro-1-pentene (Figure 2) and the similar methoxy compound permitted evaluation of the additional contribution to the rate constant,  $k_A$ , arising from 1,4-chlorine (or methoxy) participation. The  $k_A/k_S$  values

(11) P. E. Peterson and E. V. P. Tao, *ibid.*, **86**, 4503 (1964).

(12) P. E. Peterson, C. Casey, E. V. P. Tao, A. Agtarap, and G. W. Thompson, *ibid.*, **87**, 5163 (1965).

(13) Cf. G. E. Branch and M. Calvin, "The Theory of Organic Chemistry," Prentice-Hall, New York, N. Y., 1941, p 204.

(14) We assume  $\log k_H - \log k_X \equiv \Delta \log k$  falls off by a constant factor,  $\epsilon$ , per methylene group ( $k_H$  and  $k_X$  are rate constants for reactions of substituted and unsubstituted compounds, respectively). Then

$$\Delta \log k_n = \epsilon^{n-m}(\Delta \log k_m)$$

where  $n$ ,  $m$  = numbers of carbon atoms in the aliphatic chains being compared. This leads to

$$\log \Delta \log k_n = (\log \epsilon)(n - m) + \log \Delta \log k_m$$

$$\text{Cf. } y = mx + b$$

(15) For an earlier use of this approach see H. Kloosterziel, *Recl. Trav. Chim. Pays-Bas*, **82**, 508 (1963).





Figure 1. Plot of rate data for addition of trifluoroacetic acid to alkenes having cyano ( $\Delta$ ), trifluoroacetoxy (O), and acetoxy ( $\bullet$ ) substituents, based on the assumption that the substituent effect changes by a constant factor per methylene group. Slopes of the lines shown here and in Figure 6 correspond to an attenuation factor,  $\epsilon$ , of 0.65. (Cf. ref 14.)

obtained in this way for 1,4 chlorine and 1,4 methoxyl participation were 7.5 and 6.1, respectively.

That substituents up to 11 carbon atoms from the reaction site could exert effects discernible above the "noise level" of presumably random, unpredictable influences upon reaction rates was somewhat surprising, but the remarkable linearity of the plots of Figure 1 gave assurance that the effects observed were real and regular. At the time of our study it was a bit unsatisfying that 5-chloro-1-pentene,  $\text{CH}_2=\text{CH}(\text{CH}_2)_3\text{Cl}$ , did not react with concomitant halogen shift. That this result is a consequence of the direction of ring opening of the intermediate chloronium ion has now been confirmed by its independent generation (Scheme I). Alkynes, encouragingly, *did undergo halogen shifts*, as outlined below.

**Additions to Alkynes.** The observation<sup>16</sup> of 1,4 chlorine shift in the reaction of 5-chloro-1-pentyne with trifluoroacetic acid (Scheme V) predated the already

**Scheme V**  
Competing Paths in the Reaction of Trifluoroacetic Acid with 5-Chloro-1-pentyne



mentioned<sup>11</sup> evidence of a chlorine shift in a deuterium-labeled alkene. Subsequent work substantiated the





Figure 2. Plot of rate data for addition of trifluoroacetic acid to alkenes having methoxy ( $\square$ ) and chloro (O) substituents, based on the assumption that the substituent effect changes by a constant factor per methylene group.

**Table II**  
Per Cent Cis Isomer in the Predominantly Trans Vinyl Halides from Halogen Shift Reactions of Alkynes 15, Scheme VI

Reactant	Cis product, %
5-Chloro-1-pentyne	8
5-Bromo-1-pentyne	8
5-Iodo-1-pentyne	5
6-Chloro-2-hexyne	9.6
6-Iodo-2-hexyne	16.6

original inference made by Duddey, then a graduate student, that the major product of reaction was the vinyl halide **13**, and showed that both rates and products were reasonably accounted for according to Scheme V.<sup>16b,17</sup> Later in this review the amounts of halogen-shifted products for a number of alkynes are compared with those expected from analysis of the reaction rates. In the absence of sufficient data to make attenuation plots, the Hammett-Taft equation,  $\log k_{\text{H}}/k_{\text{X}} = \rho\sigma_{\text{I}}$ , was used to estimate  $k_{\text{s}}$  despite the previously mentioned lack of a reliable basis for establishing  $\sigma_{\text{I}}$  values.

The use of deuterated trifluoroacetic acid solvent or alkynes bearing a terminal alkyl group permitted the stereochemistry of the addition to the triple bond during the halogen shift reaction to be determined. Predominant anti addition of D (or H) and X was found (cf. Scheme VI and Table II). The reasons for the




(16) (a) P. E. Peterson and J. Duddey, *J. Amer. Chem. Soc.*, **85**, 2865 (1963); (b) P. E. Peterson and J. Duddey, *ibid.*, **88**, 4990 (1966).

(17) P. E. Peterson, R. J. Bopp, and M. M. Ajo, *ibid.*, **92**, 2834 (1970).

stereochemical results remain speculative since the stereochemistry of such additions is presently unpredictable.<sup>9</sup>

The reasonable success in predicting the per cent of halogen shift from reaction rates suggested that the halonium ions **16** and similar ones do *not* open at the vinyl carbon. A few years later we were able to provide solid support to previous somewhat vague notions of the unreactivity of vinyl compounds by synthesizing the first simple, secondary, aliphatic vinyl sulfonates and determining<sup>18</sup> the relative rates of reaction in 50% methanol, shown in Scheme VII. Almost simultane-


**Scheme VII**  
Reactivity of Vinyl and Saturated Secondary Tosylates  
(OTs = *p*-OSO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>3</sub>)



ously, a number of workers independently reported related work in the area of vinyl reactivity.<sup>19</sup>

**Sulfonate Solvolyses.** Sulfonate solvolyses have been employed much more extensively than the reactions of alkenes and alkynes in the demonstration of neighboring group participation, including the previously mentioned early studies of 1,2 halogen participation in the cyclohexane series (*cf.* Scheme II). Using tosylate (*p*-toluenesulfonate) substrates designed for 1,4 chlorine participation we were able to obtain<sup>20</sup> a third kind of evidence for such participation, supplementing the studies of rate effects and halogen shifts, described above. Specifically we determined the stereochemical course of solvolysis of *erythro*-5-chloro-2-hexyl tosylate (**20**, Scheme VIII) and of the *threo* isomer. The results for reaction in several solvents are given in Table III.

**Scheme VIII**  
Possible Solvolysis Paths for *erythro*-5-Chloro-2-hexyl Tosylate



(18) P. E. Peterson and J. M. Indelicato, *J. Amer. Chem. Soc.*, **91**, 6194 (1969).

(19) For a review see M. Hanack, *Accounts Chem. Res.*, **3**, 209 (1970).

(20) P. E. Peterson, R. J. Bopp, D. M. Chevli, E. L. Curran, D. E. Dillard, and R. J. Kamat, *J. Amer. Chem. Soc.*, **89**, 5902 (1967).

**Table III**  
Per Cent Retention of Configuration in the Solvolysis of 5-Chloro-2-hexyl Tosylates in the Solvents Listed

Diastereomer	CF <sub>3</sub> CO <sub>2</sub> H	HCO <sub>2</sub> H	CH <sub>3</sub> CO <sub>2</sub> H
Erythro	92	61	29
Threo	89	49	22


**Table IV**  
Estimates of  $k_{\Delta}/k_s$  for Tosylate Solvolyses, Based on Analysis of Rate Data

Tosylate	$k_{\Delta}/k_s$ for reaction in		
	CF <sub>3</sub> CO <sub>2</sub> H	HCO <sub>2</sub> H	CH <sub>3</sub> CO <sub>2</sub> H
(erythro)	99	6.7	0.8
(threo)	65	4.1	
	33	3.6	0.7

Based on the assumption<sup>7,8</sup> that inversion (not racemization) would occur in the solvolysis without participation, as shown for the  $k_s$  path of Scheme VIII, and on the well-supported<sup>3</sup> presumption that 1,4 halogen participation should give net retention of configuration resulting from two inversions (Scheme VIII), the values in Table III represent direct measures of the per cent of reaction proceeding *via* the  $k_{\Delta}$  path. That the assumptions may be imperfect is indicated by the considerably higher estimates for the extent of participation in the form of  $k_{\Delta}/k_s$  values derived from analysis of reaction rates, using the attenuation plot approach mentioned earlier (Table IV).

The characteristic solvent effect (Tables III and IV) presumably reflects the slowing of the  $k_s$  reaction in the solvents of lower nucleophilicity relative to  $k_{\Delta}$ , which is insensitive to solvent nucleophilicity. Presumably because of this effect trifluoroacetic acid as a solvent is dramatically effective in promoting halogen participation. The culmination of efforts to produce clear-cut instances of 1,4 halogen participation has come in recent studies of the trifluoroacetolysis of primary sulfonate-secondary halides. In this system 1,4 shifts were "quantitative," as mentioned in the introduction (Scheme I).<sup>1</sup> Notably, the 1,2 shift of Cl (Scheme IX) also was "quan-

**Scheme IX**  
Main Path for the Trifluoroacetolysis of 2-Chloro-1-propyl Nosylate (ONs = *p*-OSO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>)




tative," in contrast with the result for *trans*-2-chlorocyclohexyl brosylate solvolysis,<sup>5</sup> in which chlorine participation was not definitely discernible.


Six-membered rings were formed less readily, but halogen-shifted products **28** and **29** predominated

Scheme X


## Products of Trifluoroacetyloysis of 5-Chloro-1-hexyl Nosylate



(Scheme X) in the reaction of the appropriate nosylate **27**. We note that **29** has also undergone a hydrogen shift. Even 6-chloro-1-heptyl nosylate (**32**) formed 17% of primary chlorides, presumably *via* the seven-membered-ring chloronium ion **33** (eq 1).<sup>1</sup> We found



that 3-chloro-1-butyl nosylate, which might form a four-membered halonium ion ring, was too unreactive for convenient study. The corresponding trifluoromethanesulfonate reacted at a convenient rate but gave no more than 6.6% chlorine shift. On the other hand, the corresponding bromide **34** reacted to give 46% of **36**, the product of 1,3 halogen shift (eq 2),<sup>21a</sup> presumably indicating that the four-membered-ring ion **35** was an intermediate in the reaction.<sup>21b</sup>



The extent to which various primary nosylate-secondary halides solvolyzed in trifluoroacetic acid *via* intermediate halonium ions is indicated in Table V where the results of the separation of rate constants into  $k_{\Delta}$  and  $k_s$  (using the attenuation plot method described above for alkenes) are given and compared with the per cents of halogen shift. Other rate data for sulfonates are summarized in a subsequent section of this review.

**Cyclopropanes, Epoxides, and Oxonium Salts.** Other reactions which have now been shown to proceed at least to some extent *via* five-membered-ring halonium ion intermediates include the ring opening in trifluoroacetic acid of the cyclopropane **37**<sup>22</sup> and epoxides **38** and **39**<sup>23</sup> and the trifluoroacetyloysis of *in situ* generated oxonium salt, **40** (Scheme XI).<sup>24</sup>

(21) (a) P. E. Peterson and W. Boron, *J. Amer. Chem. Soc.*, **93**, 4076 (1971). (b) Another instance has been found by C. E. Reincke and J. R. McCarthy, *ibid.*, **92**, 6376 (1970).

(22) P. E. Peterson and G. W. Thompson, *J. Org. Chem.*, **33**, 968 (1968).

(23) P. E. Peterson, B. R. Bonazza, and J. M. Indelicato, *Tetrahedron Lett.*, **13** (1971).

(24) P. E. Peterson and F. J. Slama, *J. Amer. Chem. Soc.*, **90**, 6516 (1968).

Table V


## Measures of Halogen Participation in Primary Nosylate Trifluoroacetyloyses

Nosylate	$k_{\Delta}/k_s$	100 $k_{\Delta}/(k_{\Delta} + k_s)$	Halogen shift, %
2-Chloro-1-propyl	2000	99.95	99 ± 1
3-Chloro-1-butyl <sup>a</sup>	≤ 7.1 <sup>b</sup>		≤ 6.6
4-Chloro-1-pentyl	760	99.87	99.5
5-Chloro-1-hexyl	7.1	88	90
6-Chloro-1-heptyl	0.20		17
4-Chloro-1-butyl	170	99.6	~50 <sup>c</sup>

<sup>a</sup> Data for 3-chloro-1-butyl trifluoromethanesulfonate. <sup>b</sup> Distribution between  $k_s$  and  $k_{\Delta}$  calculation from product data rather than from attenuation plot. <sup>c</sup> Data for deuterium-labeled nosylate.

Scheme XI


## Reactants Which Give Halonium Ion Intermediates upon Trifluoroacetyloysis



**Variation of the Halogen.** Since there is good evidence from inductive effects that nucleophilic solvents (including weakly nucleophilic trifluoroacetic acid) only partially "neutralize" the carbonium ion charge in non-participation ( $k_s$ ) processes,<sup>20</sup> it is attractive to assume that nucleophilic participating groups, including halogen, likewise react *via* transition states having some charge on the potential cationic carbon, C-1, and some on the participating group (*cf.* Scheme XII). We hypothesize that this balance of charge distribution may change as the ring size varies, as shown in Scheme XII.

Scheme XII

## Presumed Charge Distribution in Transition States Leading to Three- and Five-Membered-Ring Halonium Ions



The three-membered ring is pictured having the charge shifted toward halogen, since the dipole of the nonreacting C-X  $\sigma$  bond (shown inside the oval) would strongly destabilize a charge on C-1. Accordingly we postulate that three-membered-ring transition states exhibit the stability order I  $\gg$  Br  $\gg$  Cl presumed to be typical of highly charged halogen. We presume that the weaker C...X bond in the five-membered ring (which may be viewed as more of an electrostatic interaction between C-1 and the C-X dipole) has the stability order I  $\geq$  Br  $\geq$  Cl (I somewhat better than or equal to Br, etc.). Although "weak," the C...X bond is strong enough, in conjunction with entropy factors, to allow the  $k_{\Delta}$  process to compete with nucleophilic solvation in the  $k_s$

**Table VI**  
Halogen Dependence of  $k_{\text{trans}}/k_{\text{cis}}$  Ratios for Solvolysis of 2-Halocyclohexyl Tosylates

Halogen	$k_{\text{trans}}/k_{\text{cis}}$		
	CH <sub>3</sub> CO <sub>2</sub> H	HCO <sub>2</sub> H	CF <sub>3</sub> CO <sub>2</sub> H
Cl	3.8 <sup>a</sup>	6.8 <sup>b</sup>	36 <sup>b</sup>
Br	810 <sup>a</sup>	4800 <sup>b</sup>	10,000 <sup>b</sup>
I	1,700,000 <sup>c,d</sup>		

<sup>a</sup> Reference 5b. <sup>b</sup> Reference 25a. <sup>c</sup> Reference 5a. <sup>d</sup> Estimated as the  $k_{\Delta}/k_{\epsilon}$  value of ref 5b.


process.<sup>6</sup> Whether valid or not, the picture outlined correlates the striking differences noted for 1,2 participation as the halogen is changed, summarized below.

In Table VI the classic work originating in Winstein's group and extended by Grunwald<sup>5a,b</sup> (Scheme II) is collected with unpublished data obtained by Duddey<sup>25a</sup> to show that the 1,2 participation order  $I \gg Br \gg Cl$  (as reflected in the  $k_{\text{trans}}/k_{\text{cis}}$  ratio) is not an artifact of the solvent choice. 1,2 chlorine participation, not considered to be clearly established from  $k_{\text{trans}}/k_{\text{cis}}$  in acetic acid,<sup>5b</sup> becomes increasingly apparent in solvents of lower nucleophilicity. (Cis halogen is presumed not to participate; accordingly  $k_{\text{cis}} \cong k_{\text{s}}$ ;  $k_{\text{trans}} \cong k_{\Delta} + k_{\text{s}}$ .)

For the newly found 1,3 participation we tentatively postulate the halogen reactivity order  $Br > Cl$  based on the 46 and 6.6% (or less) bromine and chlorine shift, mentioned above.<sup>21a</sup>

In the case of 1,4 participation it seems appropriate to cite some unpublished data from Glick's Ph.D. thesis<sup>26</sup> reporting work done in Winstein's group, along with other results, indicating a very modest extent of participation in acetic acid and showing a moderate variation among the halogens in participation tendency (Table VII). Similar comparisons of 1,4 participation in reactions of alkenes and alkynes are summarized in Table VIII.

Fluorine, surprisingly, appeared to be a weakly 1,4 participating substituent in a number of the studies of reactions of alkenes, alkynes, and tosylates which have been mentioned. However, the recently reported<sup>17</sup> failures of the nosylate **41** to solvolyze with rate enhancement and of the triflate **42** to react with fluorine shift raise the suspicion that previous indications of fluorine participation were mistakes! We hasten to add that further studies may yet resolve the apparently



contradictory results in favor of some kind of participatory role for fluorine.

**Stable Halonium Ions in SbF<sub>5</sub>-SO<sub>2</sub>.** In the last six years Olah and coworkers have made an enormously important contribution to the general area of carbonium

(25) (a) J. E. Duddey, Ph.D. Thesis, St. Louis University, 1967. (b) Cf. D. D. Roberts and W. Hendrickson, *J. Org. Chem.*, **34**, 2415 (1969).

(26) R. E. Glick, Ph.D. Thesis, University of California at Los Angeles, 1954. Professor Saul Winstein called this work to our attention a few years prior to his untimely death in 1969.

**Table VII**  
Comparison of Halogens as Participating Groups in Acetolysis of 4-Halo-1-butyl Tosylates or Nosylate

Substrate	10%k, sec <sup>-1</sup> , 99.6°		
	( $k_{\Delta} + k_{\text{s}}$ ) <sup>a</sup>	$k_{\text{s}}$ (estd) <sup>b</sup>	$k_{\Delta}/k_{\text{s}}$
4-Chloro tosylate	1.25	1.03	0.22
4-Bromo tosylate	1.73	1.03	0.70
4-Iodo tosylate	4.12	1.03	3.09
4-Chloro nosylate			0.37 <sup>c</sup>
4-Bromo tosylate			0.79 <sup>d</sup>


<sup>a</sup> Reference 26. <sup>b</sup> Incorporates (cf. ref 26) a small correction for inductive effects (for 1-butyl tosylate 10%k was 1.53). <sup>c</sup> Based on per cent Cl shift; ref 1b. <sup>d</sup> Based on per cent shift, ref 27.

**Table VIII**  
Comparison of Halogens as Participating Groups in Reactions of Alkenes and Alkynes with Trifluoroacetic Acid



Substrate	$k_{\Delta}/k_{\text{s}}$ for X =		
	Cl	Br	I
CH <sub>2</sub> =CH(CH <sub>2</sub> ) <sub>2</sub> CHXCH <sub>3</sub>	14	14	8.1
CH <sub>2</sub> =CH(CH <sub>2</sub> ) <sub>3</sub> X	7.5	7.0	
CH≡C(CH <sub>2</sub> ) <sub>3</sub> X	3.4, <sup>a</sup> 10 <sup>b</sup>	4.3, <sup>a</sup> 23 <sup>b</sup>	6.1, <sup>a</sup> 27 <sup>b</sup>
CH <sub>3</sub> C≡C(CH <sub>2</sub> ) <sub>3</sub> X	5.8, 6.1	14	

<sup>a</sup> Based on analysis of rates. <sup>b</sup> Based on per cent halogen shift (probably the better value).

ion chemistry by showing that a large number of carbonium, halonium, and oxonium ions, many of which were known only as hypothetical intermediates in reactions, may be prepared in SbF<sub>5</sub>-SO<sub>2</sub> solutions and observed by nuclear magnetic resonance.<sup>2</sup> Following the report of the observation in SbF<sub>5</sub>-SO<sub>2</sub> of three-membered-ring halonium ions, the author spent a highly rewarding few weeks in Olah's laboratories investigating the preparation of the analogous five-membered rings. This collaborative effort resulted in the observation by nmr at low temperature of the nine ions **43**, **44**, and **45** (X = Cl, Br, I).<sup>28</sup> The reactions used are typified by



eq 3 and 4. Other ions (**46**, **47**) have been reported



from Olah's group.<sup>29</sup>




X = Cl, Br, I

(27) W. S. Traynovsky, G. L. Smyser, and M. D. Doyle, *Tetrahedron Lett.*, 3127 (1968).

(28) G. A. Olah and P. E. Peterson, *J. Amer. Chem. Soc.*, **90**, 4675 (1968).

Recently we have prepared halonium ions which were halogen substituted in the side chain (48)<sup>30</sup> or in the ring (49 and 50).<sup>31</sup>




The nmr spectra of all ions mentioned are of interest because of the strong deshielding effect of positive halogen which leads to peaks at approximately  $\delta$  5 (methylene next to positive chlorine) and/or  $\delta$  6 (methine next to positive chlorine).

(29) G. A. Olah, J. M. Bollinger, and J. Brinich, *J. Amer. Chem. Soc.*, **90**, 6989 (1968).

(30) P. E. Peterson, P. R. Clifford, and F. J. Slama, *ibid.*, **92**, 2840 (1970).

(31) P. E. Peterson and B. R. Bonazza, unpublished work.

It should be mentioned that Olah and coworkers have now reported the preparation of many alkyl halonium ions, including  $\text{CH}_3\text{ClCH}_3^+\text{Sb}_2\text{F}_{11}^-$  and  $\text{CH}_3\text{ICH}_3^+\text{SbF}_6^-$ , as stable solids at room temperature!<sup>32</sup> The cyclic halonium ions reviewed in this Account, accordingly, are merely the heterocyclic members of a potentially enormous class of reactive compounds. We may anticipate that rapid developments will occur in the structural elaboration of these ions and in the exploitation of their chemical properties, of which their high reactivity as alkylating agents (exemplified in the reaction of eq 5; cf. ref 31 and 32) has been found to be notable.



(32) G. A. Olah and J. R. DeMember, *J. Amer. Chem. Soc.*, **92**, 718 (1970).

## Polycations of Group VI

RONALD J. GILLESPIE\*

*Department of Chemistry, McMaster University, Hamilton, Ontario, Canada*

JACK PASSMORE

*Department of Chemistry, University of New Brunswick, Fredericton, New Brunswick, Canada*

*Received February 8, 1971*

Cation formation has normally been associated with the metallic elements, but over the past few years evidence has accumulated which shows that it is a fairly common occurrence among nonmetals as well, at least insofar as polycations of the type  $\text{A}_m^{n+}$  are concerned. These polycations are important in terms of their stereochemistry and bonding and constitute a novel class of compounds of intrinsic interest. For the transition metals homonuclear groupings are well known in the "cluster" compounds which, however, have ligands attached to the metal cluster, e.g.,  $[\text{Mo}_6\text{Cl}_8]^{4+}$ . In the nonmetal cations  $\text{A}_m^{n+}$  lone pairs of electrons take the place of ligands.

It has long been known that the mercurous ion is not  $\text{Hg}^+$ , but  $\text{Hg}_2^{2+}$ ,<sup>1,2</sup> and recently it has been shown that other metallic elements form similar cations. Thus, evidence has been presented for the formation of  $\text{Cd}_2^{2+}$ ,<sup>3</sup>  $\text{Zn}_2^{2+}$ ,<sup>4,5</sup>  $\text{Pb}_2^{2+}$ ,<sup>5</sup>  $\text{Mg}_2^{2+}$ ,<sup>6</sup>  $\text{Ca}_2^{2+}$ ,<sup>7</sup>  $\text{Sr}_2^{2+}$ ,<sup>8</sup> and  $\text{Ba}_2^{2+}$ <sup>8</sup> on

(1) A. Ogg, *Z. Phys. Chem. (Leipzig)*, **27**, 285 (1898).

(2) See F. A. Cotton and G. Wilkinson, "Advanced Inorganic Chemistry," 2nd ed, Interscience, New York, N. Y., 1967, p 611.

(3) J. D. Corbett, W. J. Burkhard, and L. F. Druding, *J. Amer. Chem. Soc.*, **83**, 76 (1961).

(4) D. H. Kerridge and S. A. Tariq, *J. Chem. Soc. A*, 1122 (1967).

(5) J. D. Van Norman, J. S. Bookless, and J. J. Egan, *J. Phys. Chem.*, **70**, 1276 (1966).

(6) M. Krumpelt, J. Fischer, and I. Johnson, *ibid.*, **72**, 506 (1968).

(7) A. S. Dworkin, H. R. Bronstein, and M. A. Bredig, *ibid.*, **70**, 2384 (1966).

(8) A. S. Dworkin, H. R. Bronstein, and M. A. Bredig, *ibid.*, **72**, 1892 (1968).

addition of the respective element to the corresponding  $\text{MCl}_2$  melt. Various bismuth polycations have been identified, and in the process many confusing results have been rationalized. The material thought to be  $\text{BiCl}$  has been shown by an X-ray study to contain  $\text{Bi}_9^{5+}$  in addition to  $\text{BiCl}_3^{2-}$  and  $\text{Bi}_2\text{Cl}_8^{2-}$ .<sup>9</sup> The  $\text{Bi}^+$ ,<sup>10</sup>  $\text{Bi}_5^{3+}$ ,<sup>10</sup> and  $\text{Bi}_3^{2+}$ <sup>11</sup> ions have been identified in melts and the compounds  $\text{Bi}_5(\text{AlCl}_4)_3$ <sup>12</sup> and  $\text{Bi}_3(\text{AlCl}_4)_2$ <sup>12</sup> isolated. Recently evidence has been given for a cation containing antimony in the  $\text{I}^+$  oxidation state.<sup>13</sup> ESR evidence has been presented for the formation of  $\text{Ag}_4^+$  or  $\text{Ag}_4^{3+}$ ,<sup>14</sup>  $\text{Ag}_2^+$ ,<sup>15</sup> and  $\text{Cd}_2^{3+}$ .<sup>15</sup>

There has long been evidence for positive oxidation states of iodine<sup>16</sup> and the  $\text{I}_3^+$  and  $\text{I}_5^+$  ions are now well established, although "I<sup>+</sup>" has been shown to be  $\text{I}_2^+$ .<sup>17</sup>

(9) A. Hershaft and J. D. Corbett, *Inorg. Chem.*, **2**, 979 (1963).

(10) N. J. Bjerrum, C. R. Boston, and G. P. Smith, *ibid.*, **6**, 1162 (1967).

(11) N. J. Bjerrum and G. P. Smith, *ibid.*, **6**, 1968 (1967).

(12) J. D. Corbett, *ibid.*, **7**, 198 (1968).

(13) R. C. Paul, K. K. Paul, and K. C. Malhotra, *Chem. Commun.*, 453 (1970); P. A. W. Dean and R. J. Gillespie, *ibid.*, 853 (1970).

(14) R. S. Eachus and M. C. R. Symons, *J. Chem. Soc. A*, 1329 (1970).

(15) R. S. Eachus, I. Marov, and M. C. R. Symons, *Chem. Commun.*, 633 (1970).

(16) J. Arotzky and M. C. R. Symons, *Quart. Rev., Chem. Soc.*, **16**, 282 (1962).

(17) R. J. Gillespie, J. B. Milne, and M. J. Morton, *Inorg. Chem.*, **7**, 2221 (1968), and references therein.