

# Transition-Metal Radicals: Chameleon Structure and Catalytic Function

DIDIER ASTRUC\*

Laboratoire de Chimie Organique et Organométallique, URA CNRS No. 35, Université de Bordeaux I, 351, Cours de la Libération, 33405 Talence Cédex, France


Received July 19, 1990 (Revised Manuscript Received October 19, 1990)

Radicals are of fundamental relevance to areas as varied as organic synthesis, homogeneous catalysis, the chemistry of metabolisms, the degradation of aerobic cells, the generation of tumors, materials science, photoprocesses, and nuclear chemistry. Transition-metal radicals are central to a number of these subjects.<sup>1</sup> I will give here an account of the trends governing their structure, reactivity, and role in electron-transfer (ET) catalysis and finally show that, on these bases, useful functions can be rationally designed with control of reactivity pathways. For this purpose, examples will be taken essentially from new families of piano-stool-shaped organometallic radicals containing several P or S ligands, the coordination of which has been the object of our scrutiny. Considerable attention by pioneers<sup>1,2</sup> in the field of transition-metal radicals has been devoted to metal carbonyls, which behave much as organic radicals in atom-transfer and dimerization reactions. On the other hand, in the carbonyl-free or carbonyl-poor radical chemistry presented here, dimerization is avoided. This feature parallels the relative inhibition of dimerization obtained with steric bulk in organic radicals such as triarylmethyl radicals and brings us more versatility in modulating ET versus radical-type reactions. ET is an essential, more inorganic oriented property of transition-metal radicals that is useful in electrocatalysis and for which the work by Kochi has brought much mechanistic insight during the last decade.<sup>1c</sup> In discussing the behavior of transition-metal radicals, one can compare them to chameleons: their structures have different, interchangeable facets, the modulation of their reactivity is remarkably facile, and their ground state is often not apparent from a given reaction.

## Chameleon Structure: What Electron Count?

The classical organometallic electron count formalism, extremely useful for even-electron complexes, may become more tedious with radicals because of their lability. In order to ascertain a count of valence electrons (VE), it is necessary to show experimentally the number and modes of coordination of all the ligands. This can only be achieved with X-ray crystal structures or, occasionally, with the help of infrared, ESR, or Mössbauer spectroscopies. Yet, several structures can have close

Didier Astruc has been a Professor of Chemistry at the University of Bordeaux I since 1983. He received his State Thesis from the University of Rennes in 1975 and was a NATO postdoctoral Fellow at the Massachusetts Institute of Technology, where he worked with Professor Richard R. Schrock (1977). Back at Rennes, he was a CNRS Fellow in 1978 and became "Maître de Recherche" in 1980. He was awarded the Prize of the Coordination Chemistry Division of the French Chemical Society in 1981 and the "Alexander von Humboldt" Senior Prize for the Sciences in 1989. His research interests are the synthetic, structural, and mechanistic aspects of transition-metal chemistry and its applications to organic synthesis and molecular electronics.



energy levels and rapidly interconvert under standard reaction conditions. This property is related to the very low kinetic barriers of these interconversion processes.


The complexes  $\text{Fe}^{\text{I}}\text{Cp}(\text{arene})$  were shown to have 19 VE.<sup>3a-c</sup> However, the accuracy of the spectroscopic techniques does not exclude a small fraction of other species in equilibrium. Indeed, a study of the kinetics of the reactions of this family with phosphines and phosphites,<sup>3d</sup> carried out at low temperature by observing the change of optical density at 710 nm, shows that substitution proceeds by a second-order process with activation parameters  $\Delta H^\ddagger = 13.3 \pm 1 \text{ kcal mol}^{-1}$  (1 cal = 4.184 J) and  $\Delta S^\ddagger = -22 \pm 3 \text{ cal mol}^{-1} \text{ K}^{-1}$  for  $\text{P}(\text{OMe})_3$  and arene = PhMe. The bimolecular reaction, confirmed by the relatively large, negative entropy of activation, indicates that, as known for 17e radicals<sup>4,5</sup>

\*Address until September 30, 1991: Department of Chemistry (Vollhardt's group), University of California at Berkeley, Berkeley, CA 94720.

(1) (a) Kochi, J. K. *Organometallic Mechanisms and Catalysis*; Academic Press: New York, 1978. (b) Sheldon, R. A.; Kochi, J. K. *Metal-Catalyzed Oxidations of Organic Compounds*; Academic Press: New York, 1981. (c) Kochi, J. K. *J. Organomet. Chem.* **1986**, *300*, 139.

(2) (a) Meyer, T. J.; Caspar, J. V. *Chem. Rev.* **1985**, *85*, 187. (b) Connelly, N. G.; Geiger, W. E. *Adv. Organomet. Chem.* **1984**, *23*, 1. (c) Connelly, N. G. *Chem. Soc. Rev.* **1989**, *18*, 153. (d) Baird, M. *Chem. Rev.* **1988**, *88*, 1217. (e) Brown, T. L. *Ann. N.Y. Acad. Sci.* **1980**, *333*, 80. (f) Troglor, W. C. *Int. J. Chem. Kinet.* **1987**, *19*, 1025. (g) Lappert, M. F.; Lednor, P. W. *Adv. Organomet. Chem.* **1976**, *14*, 345. (h) Stiegman, A. E.; Tyler, D. R. *Comments Inorg. Chem.* **1986**, *5*, 215; *Prog. Inorg. Chem.* **1988**, *36*, 125.


(3) (a) Astruc, D. *Acc. Chem. Res.* **1986**, *19*, 377. (b) Lacoste, M.; Astruc, D. *J. Chem. Soc., Chem. Commun.* **1987**, 667. (c) Lacoste, M.; Rabaà, H.; Astruc, D.; Le Beuze, A.; Saillard, J.-Y.; Précigoux, G.; Courseille, C.; Ardoin, N.; Bowyer, W. *Organometallics* **1989**, *8*, 2233. (d) Ruiz, J.; Astruc, D. *J. Chem. Soc., Chem. Commun.* **1989**, 815; *J. Am. Chem. Soc.* **1990**, *112*, 5471.



since the work by Poë<sup>4a</sup> on  $\text{Re}(\text{CO})_5^+$ , the mechanism involves an associative process. The EHT and  $X_\alpha$  calculations for  $[\text{Fe}^{\text{I}}\text{Cp}(\text{C}_6\text{H}_6)]$  (1) by Saillard, Le Beuze, and Rabañá show that the 19e species 1 and the 17e species  $[\text{Fe}^{\text{I}}\text{Cp}(\eta^4\text{-C}_6\text{H}_6)]$  (2) have close energies.<sup>3c,d</sup> Thus, given the too high energy of a hypothetical, sterically demanding 21e species, we proposed that the rate-limiting step in which the P donor attacks the iron center involves 2 and is preceded by a fast preequilibrium  $1 (19e) \rightleftharpoons 2 (17e)$ . By analogy, it is also reasonable to expect a series of  $19e \rightleftharpoons 17e$  interchanges for the various successive steps along the overall arene substitution (Scheme I).

Another reaction, dimerization of the 19e radicals 1 through the arene ligand,<sup>6,7</sup> is not attributable to the 19e form but to the minor 18e structure 4, which bears the radical center on one arene carbon, this ligand being then represented as a  $\eta^5$ -cyclohexadienyl radical (the rate of dimerization is low<sup>3d</sup>). The dominant, most electron rich 19e form is responsible for the many ET reactions with acceptors A,<sup>8</sup> but two other forms are also responsible for the reactivity of 1 (Scheme II).

For the piano-stool radicals  $\text{Fe}^{\text{I}}\text{Cp}^*\text{L}_n$  ( $n = 2$  or  $3$ ), isolobal with 1, one may draw even more numerous, rapidly interconverting structures of close energy, given the larger choice of ligands and the possibilities of intermolecular ligand dissociation. Coupling of the  $\text{C}_5\text{Me}_5$  ligand of the 18e transient radical  $[\text{Fe}^{\text{II}}(\eta^4\text{-C}_5\text{Me}_5)(\text{CO})_2(\text{PPh}_3)]$  (5) with the benzyl radical at a  $\text{Cp}^*$  carbon has been observed by Wrighton,<sup>8g,9</sup> but the ground state of 5 may be  $\text{Fe}^{\text{I}}$ . The 17e radicals  $[\text{Fe}^{\text{I}}\text{Cp}(\text{PPh}_3)_2]$  (6) and  $[\text{Fe}^{\text{I}}\text{Cp}^*(\eta^1\text{-dppe})(\text{CO})]$  (7) have also been



characterized by ESR as iron-centered radicals, and these structures may be responsible for H atom abstraction reactions to give  $[\text{Fe}^{\text{II}}\text{Cp}(\text{PPh}_3)_2(\text{H})]$  (8) and  $[\text{Fe}^{\text{II}}\text{Cp}^*(\eta^1\text{-dppe})(\text{CO})(\text{H})]$  (9).<sup>3d,10</sup> However, in the second case, this reactivity may also be taken into account by the intermediacy of the 18e ligand-centered radical  $[\text{Fe}^{\text{II}}\text{Cp}^*(\eta^2\text{-dppe})(\text{C}=\text{O})]$  (10) in equilibrium with 7. Kochi has demonstrated that 19e metal carbonyl radicals  $\text{Cr}(\text{CO})_6^{\cdot-}$  and  $\text{Mn}(\text{CO})_4(\text{PPh}_3)_2^{\cdot-}$  generated electrochemically can abstract a H atom from  $\text{HSnBu}_3$  to give formyl complexes.<sup>11</sup> This underlines the equilibrium between the metal-centered 19e radical and the 18e carbon-centered isomeric radical, as also observed by Symons<sup>12</sup> for  $\text{CoCp}(\text{CO})_2^{\cdot-}$ . Since transition-metal formyl complexes more or less rapidly decompose to metal hydrides,<sup>11</sup> one should not systematically conclude that hydride formation by H atom abstraction from a transition-metal carbonyl radical is a direct reaction of the 17e metal center.

### Chameleon Behavior: Influence of Bulk. ET Induced by a Salt

If, on the other hand, the 17e radical is too sterically crowded, the 19e form cannot be obtained by ligand addition as its energy becomes too high. In the piano-stool series, this is the case for the iron-methyl complexes 11<sup>13</sup> (Chart I), but many stable 17e radicals are known in other series,<sup>2b</sup> the pioneering example being ferricinium,<sup>14</sup> a useful one-electron oxidant<sup>15</sup> (vide infra).

The other extreme is the 17e radical 12, although not so structurally different from 10, for which the 19e structure is very easily accessible due to the entropy factor of the hanging dithiocarbamate (dtc) ligand.<sup>16</sup> Despite very fast scan cyclic voltammetry ( $5000 \text{ V s}^{-1}$ ), it was not possible to observe any reversibility in the anodic oxidation of the 18e precursor 10;<sup>16b,c</sup> oxidation

(4) (a) Poë, A. *Trans. Met. Chem.* **1982**, *7*, 65. (b) Herrinton, T. R.; Brown, T. L. *J. Am. Chem. Soc.* **1985**, *107*, 5700. (c) Shi, Q.; Richmond, T. G.; Troglor, W. C.; Basolo, F. *J. Am. Chem. Soc.* **1982**, *104*, 4032; **1984**, *106*, 71.

(5) Hershberger, J. W.; Klinger, R. J.; Kochi, J. K. *J. Am. Chem. Soc.* **1982**, *104*, 3034.

(6) Nesmeyanov, N. A.; Sinitsyna, N. S.; Kotova, L. S.; Kolesov, V. S.; Sivoi, V. F.; Vol'kenau, N. A. *Dokl. Akad. Nauk SSSR* **1978**, *242*, 1356.

(7) Hamon, J.-R.; Astruc, D. *Organometallics* **1988**, *7*, 1036.

(8) (a) Lacoste, M.; Varret, F.; Toupet, L.; Astruc, D. *J. Am. Chem. Soc.* **1987**, *109*, 6504. (b) Desbois, M.-H.; Astruc, D.; Guillin, J.; Varret, F.; Trautwein, A. X.; Villeneuve, G. *J. Am. Chem. Soc.* **1989**, *111*, 5800. (c) Desbois, M.-H.; Astruc, D. *Organometallics* **1989**, *8*, 1841. (d) Desbois, M.-H.; Astruc, D.; Guillin, J.; Varret, F. *Organometallics* **1989**, *8*, 1848. (e) Guerschais, V.; Román, E.; Astruc, D. *Organometallics* **1986**, *5*, 2505. (f) Román, E.; Astruc, D. *Bol. Soc. Chil. Quim.* **1989**, *34*, 39. (g) Astruc, D. *Chem. Rev.* **1988**, *88*, 1189. (h) Astruc, D.; Lacoste, M.; Toupet, L. *J. Chem. Soc., Chem. Commun.* **1990**, 558.

(9) Blaha, J. P.; Wrighton, M. S. *J. Am. Chem. Soc.* **1985**, *107*, 2694.

(10) (a) Lapinte, C.; Catheline, D.; Astruc, D. *Organometallics* **1984**, *3*, 817. (b) Lapinte, C.; Catheline, D.; Astruc, D. *Organometallics* **1988**, *7*, 1683. (c) Manriquez, J. M.; Astruc, D. *Bol. Soc. Chil. Quim.* **1986**, *31*, 161. (d) Ruiz, J.; Román, E.; Astruc, D. *J. Organomet. Chem.* **1987**, *322*, C13.

(11) (a) Narayanan, B. A.; Kochi, J. K. *J. Organomet. Chem.* **1984**, *272*.

(b) Gladysz, J. A. *Adv. Organomet. Chem.* **1982**, *20*, 1.

(12) Symons, M. C. R.; Bratt, S. W. *J. Chem. Soc., Dalton Trans.* **1979**, 1739.


(13) (a) Morrow, J.; Catheline, D.; Desbois, M.-H.; Manriquez, J.-M.; Ruiz, J.; Astruc, D. *Organometallics* **1987**, *6*, 2605. (b) Lacoste, M.; Astruc, D.; Garland, M.-T.; Varret, F. *Organometallics* **1988**, *7*, 2253.

(14) Wilkinson, G.; Rosenblum, M.; Whitting, M. C.; Woodward, R. B. *J. Am. Chem. Soc.* **1952**, *74*, 2125.


(15) (a) Catheline, D.; Astruc, D. *J. Organomet. Chem.* **1984**, *266*, C11. (b) Desbois, M.-H.; Astruc, D. *New J. Chem.* **1989**, *13*, 595.

(16) (a) Catheline, D.; Román, E.; Astruc, D. *Inorg. Chem.* **1984**, *23*, 4508. (b) Amatore, C.; Verpeaux, J.-N.; Madonik, A.; Desbois, M.-H.; Astruc, D. *J. Chem. Soc., Chem. Commun.* **1988**, 200. (c) Verpeaux, J.-N.; Desbois, M.-H.; Madonik, A. M.; Amatore, C.; Astruc, D. *Organometallics* **1990**, *9*, 630. (d) Desbois, M.-H.; Nunn, C.; Cowley, A.; Astruc, D. *Organometallics* **1990**, *9*, 640.

Scheme III



of 13 using ferricinium cleanly gives the 17e, stable complex 14 in which the dtc ligand is chelated<sup>17</sup> (eq 1).




In 14, CO exchange for a variety of ligands proceeds at low temperature, giving a whole new family of 17e complexes 15, by an associative process. The latter involves, as known in other cases,<sup>1,2</sup> 19e intermediates or transition states, the bimolecular mechanism being shown by kinetic studies using visible spectroscopy (isosbestic point at 600 nm). Even very weak ligands such as CH<sub>2</sub>Cl<sub>2</sub> can replace CO in 14 at low temperature,<sup>17,18</sup> and anionic ligands (CN<sup>-</sup>, SCN<sup>-</sup>) produce neutral 17e complexes that do not dimerize<sup>18</sup> (Scheme III).

In the case of the reaction of 14 or 15 with the 3e dtc ligand, the ligand-exchange reaction gives a stable 19e intermediate 16, which can be isolated or reduce H<sub>2</sub>O in situ to OH<sup>-</sup> and H<sub>2</sub>, and 17, the first Fe<sup>IV</sup> complex with a Cp-type ligand, is isolated in excellent yields from these reactions.<sup>17,18</sup> The neutral 19e Fe<sup>III</sup> radical 16 could also be cleanly synthesized by mono-electronic reduction of 17 using 1 equiv of the 19e synthon [Fe<sup>I</sup>Cp(C<sub>6</sub>Me<sub>6</sub>)]<sup>19</sup> (Scheme IV). This dichotomy of synthetic routes to 19e complexes is noteworthy. The

(17) Desbois, M.-H.; Astruc, D. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 460.


(18) Desbois, M.-H.; Astruc, D.; Varret, F., to be submitted for publication.

(19) Astruc, D.; Hamon, J.-R.; Lacoste, M.; Desbois, M.-H.; Madonik, A.; Román, E. *Organometallic Synthesis*; King, R. B., Ed.; Elsevier: Amsterdam, 1988; Vol. IV, p 172.

Scheme IV<sup>a</sup>

<sup>a</sup>The 17e (16B) and 19e (16A) forms equilibrate very fast in solution so that the more electron rich 19e form 16A must be responsible for the reduction of H<sub>2</sub>O.

Scheme V




ability of 19e complexes to behave as reducing agents is well-known.<sup>1c,2h,3,20,21</sup>

Carbonyl-free organoiron radicals with phosphine or phosphite ligands have been generated by mild exchange of the arene ligand in the 19e complex 18.<sup>3d</sup> This gives rise to a rapid equilibrium between 17e species Fe<sup>I</sup>Cp(PR<sub>3</sub>)<sub>2</sub> (3) and 19e species Fe<sup>I</sup>Cp(PR<sub>3</sub>)<sub>3</sub> (19) that depends on the size of the P donor. With P = PPh<sub>3</sub>, the sterically stabilized 17e radical can be characterized by the triplet found in the ESR spectrum<sup>3d</sup> but the too bulky 19e radical cannot form. On the contrary, with PMe<sub>3</sub>, ligand exchange from 18 in the presence of an aromatic acceptor A gives the salts

(20) Goldman, A. S.; Tyler, D. R. *Inorg. Chem.* **1987**, *26*, 253.

(21) (a) Fischer, E. O.; Jira, R. *Z. Naturforsch.* **1952**, *8B*, 327. (b) Kwiatek, J. *Catal. Rev.* **1967**, *1*, 37. (c) Miyake, A.; Kondo, A.; Aoyama, M. *Angew. Chem., Int. Ed. Engl.* **1969**, *81*, 498. (d) Van den Akker, M.; Jellinek, F. *Recl. Trav. Chim. Pays-Bas* **1971**, *90*, 1101.



<sup>a</sup> Radical reactions of the 17e species  $\text{Fe}^{\text{I}}\text{CpP}_2$  generated by reaction of  $\text{Fe}^{\text{I}}\text{CpC}_6\text{H}_5\text{R}$ ,  $\text{R} = \text{H}$  or  $\text{Me}$  (**1**, **18**), and P donors in toluene or THF (Scheme V) in the rigorous absence of  $\text{Na}^+\text{PF}_6^-$  (these reactions are inhibited by  $\text{Na}^+\text{PF}_6^-$ ). For the coupling to Hg, the radical  $\text{FeCp}(\text{PMe}_3)_2$  (**18**) was generated by Na/Hg reduction of  $[\text{FeCp}(\text{PMe}_3)_3]^+\text{PF}_6^-$  (**20d**).


$[\text{Fe}^{\text{II}}\text{Cp}(\text{PMe}_3)_3]^+\text{A}^-$  (**20**) when the thermodynamic potential of  $\text{A}^{0/-}$  is less negative than that of  $[\text{FeCp}(\text{PMe}_3)_3]^{+/0}$  ( $E^\circ \approx -2$  V vs SCE). This results from the formation of the extremely electron rich 19e species **19** subsequently giving monoelectronic reduction of A. In the absence of an acceptor, the radical reactions of the 17e form **3** are observed. In particular with phosphines, a series of carbonyl-free iron hydride complexes **8** have been made easily accessible in this way from **18**. Thus both the 17e and 19e forms are characterized alternatively depending on whether the reaction medium contains an acceptor A (Scheme V).

It is possible to reduce substrates A that have a thermodynamic potential more negative than that of **19/20** if the reduction of a is not reversible. This is the case of  $\text{CO}_2$ , rapidly reduced by **19** to CO and  $\text{CO}_3^{2-}$  at  $-20$  °C in THF +  $\text{Na}^+\text{PF}_6^-$ , whereas **20** and  $[\text{Fe}^{\text{II}}\text{Cp}(\text{PMe}_3)_2(\text{CO})]^+$  (**21**) are formed as  $\text{PF}_6^-$  salts.<sup>3d,22a</sup>

The radical reactions of the 17e iron-phosphine species are also varied,<sup>3d</sup> as shown in Scheme VI. The chameleon behavior of the radicals  $\text{Fe}^{\text{I}}\text{CpP}_n$  is illustrated by the complete switch from radical reactions to quantitative ET reactions observed by adding the simple sodium salt  $\text{Na}^+\text{PF}_6^-$ , even in a catalytic amount.<sup>3d</sup>

Indeed, in the course of the ligand-exchange reaction between **1** and P donors, ET between the two 19e species **1** and **19** giving the large ion pair **24** (eq 2) is not observed in the absence of salt because the decomposition of the 20e anion  $[\text{Fe}^0\text{Cp}(\text{toluene})]^-$  is slowed down by the stabilizing ion pairing with the large cation  $\text{FeCpP}_3^+$ ; the radical reactions represented in Scheme VI are faster.

(22) (a) Ruiz, J.; Guerchais, V.; Astruc, D. *J. Chem. Soc., Chem. Commun.* 1989, 812. (b) Ruiz, J.; Lacoste, M.; Astruc, D. *J. Chem. Soc., Chem. Commun.* 1989, 813.



This is no longer the case in the presence of the other ion pair  $[\text{Na}^+\text{PF}_6^-]$  (**23**) in THF because double ion exchange between the two ion pairs **22** and **23** is extremely fast and gives the very unstable ion pair **24** in which the 20e anion rapidly decomposes. In THF, a stoichiometric amount of **23** is needed because **20d** quantitatively precipitates, but in 2:1 THF/ $\text{CH}_3\text{CN}$ , the medium is homogeneous and **23** is recycled so that 0.07 equiv of **23** is enough to catalyze the ET reaction of eq 2 and totally inhibits the radical-type reactions (Scheme VII).<sup>3d</sup>

Similarly, in the absence of a P donor, the ion pair  $[\text{FeCp}(\text{THF})_3^+\text{Cp}^-]$  collapses to ferrocene so that, in THF, the catalytic Scheme VII applies. The dimerization of **1**, dominant in THF in the absence of a P donor and of **23**, is totally suppressed by the addition of 0.1 equiv of **23**.

This special salt effect<sup>23</sup> shows again the rapid interconversion between the 17e species  $\text{Fe}^{\text{I}}\text{CpP}_2$  and the 19e species  $\text{Fe}^{\text{I}}\text{CpP}_3$  (Scheme V, acceptor = **1**) and their reducing power. The ergonicity of the ET (eq 2) depends on the nature of P (in **19**) and of the arene (in **1**). The ET can be endergonic by up to 0.5 V using  $\text{P} = \text{P}(\text{OMe})_3$  and arene = mesitylene and  $\text{Na}^+\text{PF}_6^-$  still totally suppresses the radical reaction (Scheme VI) to induce ET.

### Catalytic Function of Transition-Metal Radicals: ET Chain

Ligand exchange is the most commonly encountered inorganic chain reaction<sup>25-33</sup> and can proceed either by H atom transfer chain<sup>24,25</sup> or by ET chain ( $\text{E}^{\cdot}\text{C}\text{E}^{\cdot}$ ),<sup>26-33</sup> but the latter reaction became most popular in the early 80s and was carefully studied by Kochi in mononuclear<sup>1c</sup> as well as in cluster chemistry.<sup>28</sup> Other ET chain inorganic reactions are known: isomerization,<sup>29</sup> insertion<sup>30</sup> and extrusion of  $\text{CO}$ ,<sup>1c</sup> and decomplexation<sup>31</sup> and

(23) (a) Winstein, S.; Clippinger, E.; Fainberg, A.; Robinson, G. C. *J. Am. Chem. Soc.* 1954, 76, 2597. (b) Loupy, A.; Tchoubar, B. *Effets de sel en Chimie Organique*; Bordas: Paris, 1988.

(24) (a) Chanon, M. *Acc. Chem. Res.* 1987, 20, 214. (b) Ebersson, L. *Electron-Transfer Reactions in Organic Chemistry*, Springer-Verlag: Berlin, 1987; pp 172-188.

(25) (a) Rich, R. L.; Taube, H. *J. Am. Chem. Soc.* 1954, 76, 2608. (b) Osborn, J. A. In *Organo-Transition Metal Chemistry*; Ishii, Y., Tsutsui, M., Eds.; Plenum: New York, 1975; p 65.

(26) (a) Kornblum, N. *Angew. Chem., Int. Ed. Engl.* 1975, 14, 734. (b) Russell, G. A. *Chem. Soc. Spec. Publ.* 1970, 24, 271. (c) Bunnett, J. F. *Acc. Chem. Res.* 1978, 11, 413. (d) Savéant, J.-M. *Ibid.* 1980, 13, 323.

(27) Feldberg, S. W. *J. Phys. Chem.* 1971, 75, 2377.

(28) Hershberger, J. W.; Kochi, J. K. *J. Chem. Soc., Chem. Commun.* 1982, 212.

(29) (a) Rieke, R. D.; Kojima, H.; Ofele, K. *J. Am. Chem. Soc.* 1976, 98, 6735. (b) Bond, A. M.; Carr, S. W.; Colton, R. *Inorg. Chem.* 1984, 23, 2343.


(30) Magnuson, R. H.; Meirowitz, R.; Zulu, S. J.; Giering, W. P. *Organometallics* 1983, 2, 460.

(31) Moinet, C.; Román, E.; Astruc, D. *J. Electroanal. Chem. Interfacial Electrochem.* 1981, 121, 241.


(32) (a) Astruc, D. *Angew. Chem., Int. Ed. Engl.* 1988, 27, 643. (b) Astruc, D. *Comments Inorg. Chem.* 1987, 6, 61. (c) Astruc, D. In *The Chemistry of the Metal-Carbon Bond*; Hartley, F. R., Patai, S., Eds.; Wiley: New York, 1987; Vol. 4, pp 625-731.

(33) Desbois, M.-H.; Astruc, D. *J. Chem. Soc., Chem. Commun.* 1988, 472.

Scheme VII



Scheme VIII



chelation.<sup>16b,c,32</sup> We have noted<sup>32</sup> from most reports that the ET chain reactions that work efficiently have an exergonic (thus fast<sup>34</sup>) cross-redox propagation step. There is only one way, oxidation or reduction, to insure the initiation providing this condition.

The rule (based on  $E^\circ$  values) works well for the substitution of electron-donating ligands such as MeCN initiated by oxidation in monometallic complexes<sup>1c,28</sup> and for CO substitution initiated by reduction in clusters.<sup>32a,c,35</sup> It is also useful to remove or replace a hydrocarbon ligand by more electron rich ligands in mononuclear complexes.<sup>3d,31,32a,36</sup> For instance,  $[\text{Fe}^{\text{II}}\text{Cp}(\text{PMe}_3)_3]^+\text{PF}_6^-$  and  $[\text{Fe}^{\text{II}}\text{Cp}(\text{diphos})\text{NCMe}]^+\text{PF}_6^-$  can be made on a large scale in quantitative yields in a few minutes at 20 °C by introduction of a few milligrams of  $[\text{Fe}^{\text{I}}\text{Cp}(\eta^6\text{-toluene})]$  to a THF suspension or an acetonitrile solution of  $[\text{Fe}^{\text{II}}\text{Cp}(\eta^6\text{-toluene})]^+\text{PF}_6^-$  containing the appropriate ligand.<sup>3d</sup> With  $\text{PMe}_3$  (Scheme VIII), the cross-redox step is indeed exergonic by 0.8 V ( $\Delta G^\circ = -18.4 \text{ kcal mol}^{-1}$ ), which provides a fast ET, avoiding the radical reactions of Scheme VIII. The mechanism of the other propagation step, ligand substitution, is associative (vide supra). The reaction works with all the P donors that fulfill the condition of an exergonic cross-redox step. This is not the case for

$\text{P}(\text{OPh})_3$ ,<sup>3d</sup> and the reaction indeed fails, although  $[\text{Fe}^{\text{II}}\text{Cp}\{\text{P}(\text{OPh})_3\}_3]^+$  is known.<sup>37</sup> The cross-redox step is endergonic by 0.1 V, and the radical reaction of Scheme VI is then observed.

This rule, stating that electrocatalytic reactions do not work when the cross-redox step is expected to be endergonic, ignores the other propagation step: the ligand exchange.<sup>1c</sup> It was thus desirable to try to catalyze the same reaction using either an oxidant or a reductant in the initiation. We have chosen the chelation of the dithiocarbamate ligand, for which the entropy drives the "chemical" propagation step.<sup>38</sup> For the first time, both electrocatalytic cycles were shown to work (Scheme IX), but with limited Coulombic efficiency. In the case of the oxidative initiation,<sup>16b,c</sup> the endergonicity of the cross-redox step slows it down and the isolable 17e chelated cation gives competing side reactions (precipitation in THF or CO exchange with the solvent shown in Scheme IX). With the reductive initiation,<sup>38</sup> the yields are somewhat better but the Coulombic efficiency is now limited by the other propagation step: the loss of the dithiocarbamate ligand competes with its chelation with CO loss. We believe that with suitably designed electron-reservoir ligands this duality of initiation could also be successfully applied.<sup>39</sup>

### Coupling Electrocatalysis with Organometallic Catalysis

It is possible, using the same rule, to design the rational synthesis of catalysts by ET chain catalyzed ligand exchange from 18e precatalysts. The oxidatively induced exchange of MeCN ligands by P donors in  $[\text{W}(\text{CO})_3(\text{NCMe})_3]$  (**25**) was known to proceed with high Coulombic efficiency.<sup>5</sup> Thus, we envisaged replacement of MeCN ligands by alkynes in this W complex to bring at least two alkyne ligands onto the W center. This ligand exchange involves an exergonic cross-redox propagation step. In this way, we meet the conditions of the well-known Chauvin-type<sup>40a</sup> metathesis mecha-

(37) Green, M. L. H.; Whiteley, R. N. *J. Chem. Soc. A* 1971, 1943.

(38) Desbois, M.-H.; Astruc, D. *J. Chem. Soc., Chem. Commun.* 1990, 943.

(39) (a) Summers, D. P.; Luong, J. C.; Wrighton, M. S. *J. Am. Chem. Soc.* 1981, 103, 5238. (b) Lahuerta, P.; Latorre, J.; Sanau, M. *J. Organomet. Chem.* 1985, 286.

(40) (a) Hérisson, J.-L.; Chauvin, Y. *Makromol. Chem.* 1970, 141, 161. (b) Katz, T. J.; Lee, S. J. *J. Am. Chem. Soc.* 1980, 102, 422.

(34) (a) Marcus, R. A. *J. Chem. Phys.* 1956, 24, 966. (b) Review: Marcus, R. A.; Sutin, N. *Biochim. Biophys. Acta* 1985, 811, 265.

(35) Bezems, G. J.; Rieger, P. H.; Visco, S. *J. Chem. Soc., Chem. Commun.* 1981, 265.

(36) Darchen, A. *J. Chem. Soc., Chem. Commun.* 1983, 768.



On the other hand, the first example of an ET chain reaction that can be initiated in either way also illustrates the problem of side reactions. Electron-reservoir ligands should be helpful in switching the mode of initiation of the ET chains.

An application of the suitably initiated electrocatalytic ligand exchange reaction is the coupling between  $\text{E}\ddot{\text{C}}\ddot{\text{E}}$  and organometallic catalysis with the aim of improving the reaction conditions, yields, and selectivities of transition-metal catalysis. Exemplified here by the facile induction of the polymerization of terminal alkynes, this principle should find considerable extension in the near future to other types of catalytic reactions. In more general terms, the coupling between several types of catalysis, some of which involve ET, is one of

the keys for future achievement and improvement of technological processes.<sup>41</sup>

*I am greatly indebted to the students, postdocs, and scientists whose names appear in the references for their ideas and efforts and especially to Drs. M.-H. Desbois, M. Lacoste, A. M. Madonik, and J. Ruiz. I also thank Dr. N. Ardoin and J. Moncada for their kind and efficient help in the preparation of the manuscript and the CNRS, the University of Bordeaux I, the Région Aquitaine, and the Humboldt Foundation for financial assistance.*

(41) (a) Lehn, J.-M. *Ann. N.Y. Acad. Sci.* **1986**, *471*, 41. (b) Rodriguez-Ubis, J. C.; Alpha, B.; Plancherel, D.; Lehn, J.-M. *Helv. Chim. Acta* **1984**, *67*, 2264. (c) Juris, A.; Barigeletti, F.; Campagna, S.; Balzani, V.; Belser, P.; von Zelewsky, A. *Coord. Chem. Rev.* **1988**, *84*, 85. (d) Meyer, T. J. *Acc. Chem. Res.* **1989**, *22*, 163. (e) Wrighton, M. S. *Comments Inorg. Chem.* **1985**, *4*, 269. (f) Fendler, J. H. *J. Phys. Chem.* **1985**, *89*, 2730.

## Identification of Molecular Adsorbates by LITD/FTMS: A Breakthrough for Surface Chemistry

DONALD P. LAND, CLAIRE L. PETTIETTE-HALL, JOHN C. HEMMINGER,\* and ROBERT T. MCIVER, JR.\*

*Department of Chemistry and Institute for Surface and Interface Science, University of California, Irvine, Irvine, California 92717*

*Received April 3, 1990 (Revised Manuscript Received October 1, 1990)*

### Introduction

During the last 20 years, enormous progress has been made in the development of new and more precise probes of the solid-gas interface. Techniques such as low-energy electron diffraction (LEED) and Auger electron spectroscopy (AES) provide information about the elemental composition and structure of the surface layer, and X-ray photoelectron spectroscopy (XPS) is a sensitive method for determining the elemental composition and oxidation state of surface adsorbates. These methods are of only limited usefulness to a chemist, however, because they reveal little about the *molecular* composition of the surface species. Electron energy loss spectroscopy (EELS) and infrared spectroscopy can be used to identify simple molecular adsorbates, but identifying complex species, such as are encountered in problems related to catalysis, lubrication, adhesion, corrosion, and microelectronics fabrication, remains a difficult problem. Part of the difficulty results because very high detection sensitivity is required. For example, one monolayer of adsorbed

material corresponds to about 3 pmol distributed over a surface area of 1 mm<sup>2</sup>. Another difficulty is that small signals from the adsorbate can be overwhelmed by those arising from the bulk substrate.

In this Account we describe an exciting new surface analysis method called LITD/FTMS.<sup>1-4</sup> Laser-induced thermal desorption (LITD) utilizes a pulsed laser beam to rapidly heat the surface and remove intact neutral molecular species. These are then ionized by an electron beam, trapped in a magnetic field, and detected by Fourier transform mass spectrometry (FTMS).<sup>5-8</sup> FTMS has two distinct advantages: high mass resolution and the ability to record all masses simultaneously from a *single* laser shot. LITD/FTMS is a highly sensitive and rapid method for identifying complex molecular species adsorbed on surfaces. Intermediates in catalytic reactions can be observed, and rate constants for surface-catalyzed reactions can be measured.

### Principles of LITD/FTMS

The sequence of events for a LITD/FTMS experiment is shown in Figure 1.<sup>9-12</sup> Figure 1a shows a pulsed

Donald Land received a B.A. in chemistry from Lawrence University in Wisconsin in 1984 and a Ph.D. in 1989 from UC Irvine. He is currently in Juelich, West Germany, on an Alexander von Humboldt Fellowship.

Claire Pettiette-Hall received a B.S. in chemistry from Yale University in 1983 and a Ph.D. in 1988 from Rice University.

John Hemminger received a B.S. in chemistry from UC Irvine in 1971 and a Ph.D. in chemical physics from Harvard University in 1976. After serving as a NSF Postdoctoral Fellow at UC Berkeley, he joined the faculty at UC Irvine in 1978. He has received an Alfred P. Sloan Fellowship and is the Director of the Institute for Surface and Interface Science at UC Irvine. His other areas of interest are interadsorbate interactions, vibrational spectroscopy at surfaces, and conducting polymer films.

Robert McIver received a B.S. in chemistry from the University of Kansas in 1967 and a Ph.D. from Stanford University in 1971. He joined the faculty at UC Irvine in 1971 and has made several key contributions to the development of Fourier transform mass spectrometry. He has received an Alfred P. Sloan Fellowship and several awards for outstanding teaching.

(1) Sherman, M. G.; Kingsley, J. R.; Dahlgren, D. A.; Hemminger, J. C.; McIver, R. T., Jr. *Surf. Sci.* **1985**, *148*, L25.

(2) Sherman, M. G.; Kingsley, J. R.; Hemminger, J. C.; McIver, R. T., Jr. *Anal. Chim. Acta* **1985**, *178*, 79.

(3) Land, D. P.; Wang, D. T.-S.; Tai, T.-L.; Sherman, M. G.; Hemminger, J. C.; McIver, R. T., Jr. In *Lasers and Mass Spectrometry*; Lubman, D. M., Ed.; Oxford University Press: New York, 1990.

(4) Land, D. P.; Pettiette-Hall, C. L.; Sander, D.; McIver, R. T., Jr.; Hemminger, J. C. *Rev. Sci. Instrum.* **1990**, *61*, 1674.

(5) Gross, M. L.; Rempel, D. L. *Science* **1984**, *226*, 261.

(6) Wilkins, C. L.; Chowdhury, A. K.; Nuwaysir, L. M.; Coates, M. L. *Mass Spectrom. Rev.* **1989**, *8*, 67.

(7) Russell, D. H. *Mass Spectrom. Rev.* **1986**, *5*, 167.

(8) Marshall, A. G. *Acc. Chem. Res.* **1985**, *18*, 167.

(9) Sherman, M. G.; Kingsley, J. R.; Dahlgren, D. A.; Hemminger, J. C.; McIver, R. T., Jr. *Surf. Sci.* **1985**, *148*, L25.