

The Unexpected Role of Vitamin E (α -Tocopherol) in the Peroxidation of Human Low-Density Lipoprotein¹

VINCENT W. BOWRY*[†] AND K. U. INGOLD[‡]

Department of Chemistry, University of New England, Armidale NSW 2351, Australia, and Steacie Institute for Molecular Sciences, National Research Council of Canada, Ottawa, Ontario, Canada K1A 0R6

Received February 3, 1998

Introduction

Atherosclerosis is the single most important cause of death in Western Society. "Fatty streaks" formed on the interior walls of arteries are the first visible signs of atherosclerosis. These fatty streaks contain peroxidized low-density lipoprotein (LDL)² particles, and oxidized LDL is strongly implicated in the initiation and development of atherosclerosis.³ Antioxidants would therefore be expected to attenuate atherosclerosis, and there are a number of human⁴ and animal studies⁵ indicating that this is the case. Naturally, there has been a veritable explosion of research into free-radical-induced peroxidation of LDL and its prevention by antioxidants.⁶


By far the major free-radical antioxidant in LDL lipid is α -tocopherol, α -TOH, biologically and chemically the most active form of vitamin E.⁷ The very high reactivity of α -TOH toward peroxy radicals⁷ and the relatively large amount of α -TOH present in LDL (vide infra) leads one to expect that LDL peroxidation would be very strongly inhibited. Yet despite this "expectation" and despite scores of experimental studies, there is still no consensus that enrichment of the lipoprotein with α -TOH will actually retard the oxidative modification of LDL in vitro.⁸

Our investigations into the kinetics of LDL peroxidation suggest that this lack of agreement arises from an "unrecognized consequence of lipid particle size"⁹ endowing α -TOH with a *prooxidant* activity that manifests itself in various ways according to experimental conditions (e.g., the type and amount of radical initiator).¹⁰ In this

Vincent W. Bowry was born in Australia, and educated at the University of Sydney (B.Sc. Hons) and the Australian National University (Ph.D.). After postdoctoral positions at the NRC-Canada and the Heart Research Institute-Sydney, he took up a lectureship at the University of New England. His research interests focus on free radical kinetics in organic and biological chemical systems.

K. U. Ingold received his B.Sc. degree from University College London and his D.Phil. from Oxford. He is a Distinguished Scientist in the Steacie Institute for Molecular Science, and his research has concentrated on free radicals in solution.

Scheme 1. α -Tocopherol-Inhibited Peroxidation of Esters of Polyunsaturated Fatty Acids (PUFA = LH)



Account we describe (i) how and when α -TOH becomes a prooxidant,¹¹ (ii) the inhibition of α -TOH's prooxidant activity by co-antioxidants,¹² (iii) use of a simple deuterium kinetic isotope effect to probe the peroxidation mechanism,¹³ and (iv) the implications of our findings for atherosclerosis research.

Background

Radical Scavenging in Homogeneous Systems. The principal elementary reactions involved in α -TOH-inhibited lipid peroxidation in a homogeneous system are shown in Scheme 1.¹⁴ Each molecule of α -TOH terminates two peroxidation chains (reactions 5 and 6), and the overall kinetic rate law is


$$d[\text{LOOH}]/dt = R_g k_p [\text{LH}] / 2k_{\text{inh}} [\alpha\text{-TOH}] \quad (I)$$

where R_g is the rate of generation of the radicals which initiate peroxidation chains. In homogeneous, lipid-like solvents, k_{inh} for α -TOH is greater¹⁵ than for any other lipid-soluble biological antioxidant and hence α -TOH is the most powerful natural inhibitor of lipid peroxidation.

The trapping of peroxy radicals by α -TOH is much slower in polar solvents; e.g., k_{inh} is 50-fold lower in ethanol than in cyclohexane solvent.¹⁶ Our recent investigations into the role of the solvent in hydrogen atom abstraction reactions^{17–20} show that slower trapping in polar solvents is due to hydrogen bonding of the substrate to hydrogen bond acceptor (HBA) solvents. Thus, the reactivity of α -TOH in an HBA solvent, **S**, depends on the fraction of the α -TOH that is *not* bound to the solvent (Scheme 2).²⁰ The solvent effect depends solely on the substrate–solvent interaction and is therefore *independent*

[†] University of New England.

[‡] National Research Council of Canada.



of the reactivity of the attacking radical, a remarkable property that has been confirmed over a very wide (10^{10} -fold!) kinetic range.¹⁸ The importance of this new principle is that if we know a single rate constant for H-abstraction from α -TOH by some radical, e.g., a peroxy radical, we can accurately predict the magnitude of this rate constant (e.g., k_{inh}) in many other solvents.

Radical Scavenging in Nonhomogeneous Systems.

Reported rate constants for peroxy radical trapping by α -TOH dispersed in SDS micelles²¹ and phospholipid liposomes²² are lower and very much lower, respectively, than the values reported for any homogeneous system, cf. $k_{\text{inh}}/\text{M}^{-1} \text{s}^{-1} = (5 \pm 1) \times 10^4$ (micelles) and 3×10^3 (liposomes) versus 5.1×10^5 (*tert*-butyl alcohol,¹⁶ a good HBA solvent). These results cannot be explained via Scheme 2 with $\text{S} = \text{H}_2\text{O}$ for the simple reason that water is not as good an HBA solvent as, say, *tert*-butyl alcohol.²³

Since the low reactivity of α -TOH in aqueous dispersions is not due to hydrogen bonding, it must be a consequence of a physical separation between at least some of the lipid peroxy radicals and some of the α -TOH molecules. Both inter- and intraparticle separations are possibilities. In the inter- case, α -TOH in particle A obviously cannot capture an $\text{LOO}\cdot$ in particle B until, e.g., $\text{LOO}\cdot$ diffuses from A to B. If B were completely depleted in α -TOH and $\text{LOO}\cdot$ exit from the particle was slow, then B could undergo extensive peroxidation despite the presence of abundant α -TOH in A (and hence in the overall system).

Thus, the *true* rate constant for the α -TOH/ $\text{LOO}\cdot$ reaction 5 in aqueous dispersions of lipid is unknown, and cannot be reliably derived from conventional measurements suitable for a homogeneous system.^{24,25} This is true not only for SDS micelles and phospholipid liposomes but also for LDL particles.

Low-Density Lipoprotein (LDL). The "peroxidation profile" of an LDL particle is shown in Figure 1.^{26,27} Of obvious importance to LDL peroxidation are (a) *substrate*, ~ 1450 bisallylic CH_2 groups (LHs) per particle (see Scheme 1), and (b) *antioxidant*, ~ 6 – 12 molecules of α -TOH per particle and lesser amounts of other real or putative²⁸ antioxidants (see Figure 1 and below). Of less obvious importance to peroxidation is

(c) *particle size*, Stokes radius ~ 10 nm, volume $3.8 \times 10^{-24} \text{ m}^3$ with a free lipid volume $V_{\text{lip}} \approx 3.2 \times 10^{-24} \text{ m}^3$ and hence a free molar volume, $N_A V_{\text{lip}} \approx 2 \times 10^3 \text{ dm}^3 \text{ mol}^{-1}$. What this means is that an LDL particle is simply too small to hold more than one radical for more than a very brief moment, e.g., ~ 1 s for a pair of $\alpha\text{-TO}\cdot$ radicals, ~ 1 ms for a pair of $\text{LOO}\cdot$ radicals, and $\sim 1 \mu\text{s}$ for an $\text{LOO}\cdot$ ($\text{ROO}\cdot$) and $\alpha\text{-TO}\cdot$ combination.²⁹




FIGURE 1. Schematic of an LDL particle. Each particle is a molecular assembly containing a protein moiety (apoprotein B-100, which intercalates and stabilizes the structure), a *core* of neutral lipids (cholesteryl and triglyceryl (TG) esters and free cholesterol), and a *coat* of polar lipids (mainly phosphatidylcholine (PC) diesters). Ch18:2 and Ch20:4 refer to the cholesteryl esters of linoleic and arachidonic acids, respectively.


In our studies (unlike many others) the LDL was very rapidly isolated. It contained ≤ 1 LOOH molecule per 500 particles³⁰ and retained most of its associated coenzyme Q₁₀ in its reduced form, CoQH₂, at the start of reaction (vide infra).

LDL Peroxidation

Theoretical Expectations. It can be seen from Scheme 1 (or eq 1) that lipid peroxy radicals partition between chain-terminating attack on α -TOH (reaction 5) and chain-propagating attack on LH (reaction 4) in the ratio $k_{\text{inh}}[\alpha\text{-TOH}]/k_p[\text{LH}]$. For example, when $k_{\text{inh}}[\alpha\text{-TOH}] = k_p[\text{LH}]$, 50% of the $\text{LOO}\cdot$ will be regenerated (via reactions 4 and 3) so that any peroxidation that does occur does not involve a radical chain reaction, or, in other words, in this situation the chain length, ν , would be 1.0. While there is some uncertainty regarding the effective magnitude of k_{inh} in LDL (vide supra), it probably lies between the value found in a saturated hydrocarbon solvent and the value estimated in water, viz. $k_{\text{inh}} \approx (68\text{--}8) \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$. Therefore, since $k_p \approx 100 \text{ M}^{-1} \text{ s}^{-1}$ per LH moiety,³¹ one molecule of α -TOH should be able to protect between 68 000 and 8000 LH moieties. The $[\text{LH}]/[\alpha\text{-TOH}]$ ratio in LDL is roughly $1450/9 = 160$ (Figure 1), and so it would appear that LDL is massively overprotected against peroxidation.

Experiments in which LDL was "oxidatively modified" using air and copper ions (usually $> 5 \text{ Cu}^{\text{II}}$ ions per LDL particle) lent support to the above analysis in that relatively little LOOH was formed until the α -TOH (and other endogenous antioxidants) had been consumed.³² Naturally this type of experiment led to the view that α -TOH is a powerful antioxidant for LDL, just as it is for bulk lipids. Unfortunately, there was a very poor correlation between the α -TOH content of LDL and the duration of the induction (or lag) period preceding lipid peroxidation.⁶ This little inconsistency led to the invention of a new variable, "unknown antioxidants" in the LDL. Naturally, these have remained unknown because they simply do not exist!

Scheme 3



Practical Reality. In 1990 two independent studies reported on the peroxidation of LDL exposed to organic peroxy radicals ($\text{ROO}\cdot$) generated at a steady rate (R_g) by water-soluble and lipid-soluble azo compounds (2,2'-azobis(2-amidinopropane hydrochloride), AAPH, and 2,2'-azobis(2,4-dimethylvaleronitrile), AMVN, respectively)^{33,34} (Scheme 3).


Both studies showed that LDL peroxidation occurred at much the same rate in the formally α -TOH inhibited period of peroxidation as in the subsequent uninhibited (α -TOH consumed) period. Both studies also noted that the minor quinolic antioxidant, ubiquinol-10 (CoQH_2), was consumed *before* α -TOH.

A more detailed investigation of the early period of peroxidation³⁵ revealed that the disappearance of CoQH_2 corresponded closely with the onset of a rapid oxidation phase.³⁶ In the presence of vitamin C (ascorbate, AscH^-), the CoQH_2 was still consumed first but now peroxidation was inhibited until the ascorbate was also depleted.


Even more intriguing was that peroxidation in the formally inhibited phase of LDL peroxidation was being propagated in a radical *chain*. Up to 100 LOOH molecules were formed per initiating radical, $\text{ROO}\cdot$, regardless of whether the peroxidation was induced by AAPH, AMVN, or (the oxidants generated by) stimulated neutrophils (scavenger cells).³⁵

These results were contrary to expectations for a peroxy-mediated, α -TOH-inhibited peroxidation (Scheme 1). That is, being 10-fold *less* reactive than α -TOH toward peroxy radicals in solution,³⁷ CoQH_2 was *expected* to be consumed after α -TOH and to have little effect on the peroxidation rate. Water-soluble vitamin C was also *expected* to have little or no effect on the peroxidation rate in AMVN-initiated LDL since it is known to be a very poor inhibitor of peroxidation induced by lipid-soluble $\text{ROO}\cdot$ radicals in micelles and liposomes.³⁸ And we certainly did *not expect* a radical chain in tocopherol-containing LDL.³⁹


Tocopherol-Mediated Peroxidation (TMP). These inconsistencies forced us to reconsider the fate of α -TO \cdot radicals, in particular, to consider whether the known, but very slow, chain-transfer reaction of α -TO \cdot with lipid



was contributing to peroxidation by regenerating $\text{LOO}\cdot$ radicals. This would help explain the high effectiveness of vitamin C since it was known that vitamin C (AscH^-) could reduce α -TO \cdot in PC liposome dispersions.³⁸



Scheme 4. Tocopherol-Mediated Peroxidation (TMP)



One can readily see that reaction 8 followed by coupling of the $\text{L}\cdot$ radical with oxygen and reaction of the resulting $\text{LOO}\cdot$ with α -TOH constitutes a *potential* radical chain—a tocopherol-mediated peroxidation (TMP) chain (Scheme 4)—but is it a plausible one? The problem here is that reaction 8 (chain transfer) is so slow that an α -TO \cdot radical would need to “wait around” for ~ 30 s between propagation steps ($k_{\text{TMP}}[\text{LH}] \approx 0.03 \text{ s}^{-1}$). At first sight this seemed highly unlikely in the face of chain-terminating reactions with rate constants of up to $3 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ (reaction 6) in a *homogeneous system*.⁴⁰

However, LDL is not homogeneous and diffusion of lipid-soluble compounds *between* LDL particles is known to be slow.^{41,42} It occurred to us that once an α -TO \cdot radical was formed in an LDL particle (reaction 5') it might remain trapped within that particle and thereby be forced to “wait around” for a relatively long time for a *second* $\text{ROO}\cdot$ radical to strike the particle (reaction 5'). Simple calculation revealed a “waiting period” on the order of several minutes for our reaction conditions; e.g., for a typical LDL + AAPH experiment, $[\text{LDL}]/R_g \approx 1 \mu\text{M}/1 \text{ nM s}^{-1} = 1000 \text{ s}$ or 17 min. This is, indeed, ample time for several LOOH molecules to be generated via the TMP chain in Scheme 4.

Models of TMP Kinetics. To prove the existence of TMP in LDL, we first needed to know what to look for in our experimental data. Kinetic models were developed from three basic premises: (i) no transfer of radicals *between* particles, (ii) a one-radical-per-particle limit (vide supra, particle size), and (iii) α -TOH retains its “normal” high radical scavenging activity in LDL.

(a) Simple Emulsion Polymerization Model. The third premise immediately told us that the vast majority of LOOH formed in the presence of α -TOH *must be* formed via reaction 5 (because a $\text{LOO}\cdot$ radical is ~ 200 -fold more likely to react with α -TOH than with LH in the LDL particle). Kinetic analysis of the resulting TMP-only system⁴³ yielded what we thought was a very unusual prediction: at steady state exactly half of the particles would contain one radical each and the other half none




FIGURE 2. Consumption of α -TOH and formation of cholesteryl ester hydroperoxide (CEOOH) in LDL initiated with various concentrations of AAPH. R_p^{\max} and the fractional rates of peroxidation ($\Phi = R_p^{\max}/[\text{LH}]$) were calculated from maximum slopes of the CEOOH vs time plots. Fractional peroxidation rates were nominally uniform between lipid classes and were proportional to the number of bisallylic methylene groups per lipid molecule (so, e.g., cholesteryl arachidonate, Ch24:4, peroxidized 3 times faster than cholesteryl linoleate, Ch18:2, $\Phi^{24:4} \approx 3\Phi^{18:2}$).

(i.e., $[\alpha\text{-TO}\cdot] = [\text{LDL}]/2$), and consequently the *peroxidation rate would be independent of the rate of radical generation*

$$R_p = -d[\text{LH}]/dt (\approx d[\text{LOOH}]/dt) = \frac{[\alpha\text{-TO}\cdot][\text{LH}]k_{\text{TMP}}}{([\text{LDL}]/2)[\text{LH}]k_{\text{TMP}}} \quad (\text{III})$$

In the event, the experimental rate of α -TOH-inhibited peroxidation was remarkably insensitive to the concentration of initiator (see Figure 2C). We had, in fact, uncovered the first example of *emulsion polymerization kinetics*^{44,45} in a nonpolymer system.

(b) Modified Emulsion Polymerization Model. Figure 2A demonstrates that eq III is not consistent with experiment since the oxidation rate (R_p) clearly becomes slower as the α -TOH is consumed. Further experiments showed that R_p was faster in α -TOH-enriched LDL.¹¹ This *prooxidant* activity of α -TOH could only be due to a high *phase-transfer* activity of α -TOH. That is, instead of particles being uniformly reactive to initiating radicals (as is probably true for emulsion polymerizations), the likelihood of a particle reacting with an incoming $\text{ROO}\cdot$ radical must (by premise iii) be dominated by α -TOH in nonperoxidizing particles ($[\alpha\text{-TOH}]k_{\text{ROO}\cdot/\alpha\text{-TOH}} \approx 6 \times 10^3 \text{ s}^{-1}$) and by $\alpha\text{-TO}\cdot$ in peroxidizing particles ($[\alpha\text{-TO}\cdot]k_{\text{ROO}\cdot/\alpha\text{-TO}\cdot} \approx 2 \times 10^5 \text{ s}^{-1}$).⁴⁶ Thus, assuming (i) *all* initiation occurred via reaction 5' (i.e., none via reaction 2) and (ii) *all* termination occurs via reaction 6', we found that a steady-state radical population required⁴⁷

$$k_5[\alpha\text{-TOH}][\text{ROO}\cdot] = 2k_6[\alpha\text{-TO}\cdot][\text{ROO}\cdot] \quad (\text{IV})$$


(rate of initiation) (rate of termination)

That is, $[\alpha\text{-TO}\cdot] = (k_5/2k_6)[\alpha\text{-TOH}]$, and therefore

$$R_p^{\text{inh}} = k_{\text{TMP}}[\text{LH}][\alpha\text{-TOH}](k_5/2k_6) \quad (\text{V})$$

(see Scheme 4). Equation V is consistent with the “sigmoid” shapes of the LOOH (CEOOH) vs *time* plots (Figure 2). After an early maximum in R_p , the rate falls as α -TOH is consumed, reaching a minimum just as the last few percent of α -TOH disappears, before accelerating again in the post-tocopherol period. Experimental data for AAPH- and AMVN-initiated native LDL yielded values of

Scheme 5. Radical Anti-TMP Mechanisms




$k_5/k_6 = 0.03$ and 0.006 , respectively, which may be compared with $k_{\text{ROO}\cdot/\alpha\text{-TOH}}/k_{\text{ROO}\cdot/\alpha\text{-TO}\cdot} = 0.005$ obtained using AMVN in an organic solution of α -TOH.⁴⁰

Co-antioxidants for LDL. *Plasma Antioxidants.* We tested the effect of some of the conditions and compounds found in blood plasma on azo-initiated peroxidation. It became clear from these experiments that TMP in LDL was strongly inhibited in plasma by CoQH₂, ascorbate, and bilirubin,⁴⁸ whereas other components such as free fatty acids, proteins, thiols, amino acids, and carotenoids, individually or in combination, had little or no effect on the peroxidation rate during the formal α -TOH-inhibited period. Lowering the oxygen tension from 150 mmHg (air) to a more physiological 15 mmHg also had negligible effect on the lipid peroxidation rate in tocopherol-containing LDL (cf. ref 49).

The putative plasma antioxidant, urate, proved interesting since it had a marked *prooxidant* effect on AAPH-initiated LDL. That is, addition of urate at its blood plasma concentration ($\sim 0.3 \text{ mM}$) *increased* the rate of LOOH formation by $\sim 50\%$ while slowing the rate of α -TOH consumption by $\sim 90\%$. Presumably, urate—as a good scavenger of aqueous peroxy but not $\alpha\text{-TO}\cdot$ radicals⁵⁰—was merely prolonging the effect of TMP by shielding the α -TOH from initiating radicals.

Anti-TMP Mechanisms. Since $\alpha\text{-TO}\cdot$ radicals drive peroxidation in tocopherol-containing LDL, *any* mechanism that reduces the rate of formation of $\alpha\text{-TO}\cdot$ radicals or increases their destruction once formed will retard lipid peroxidation. Three possible avenues for inhibiting TMP are open; they have been named radical export, interparticle radical transport, and radical import (Scheme 5).

Radical export is the irreversible transfer of a radical center from the LDL lipid to the aqueous phase. Ascorbate in the aqueous phase does this by rapidly reducing $\alpha\text{-TO}^\bullet$ (reaction 9) to yield $\alpha\text{-TOH}$ and the water-soluble, strongly reducing, ascorbyl radical. Even micromolar amounts of ascorbate prevented TMP from occurring in LDL. Ubiquinol-10 (CoQH_2) also reacts rapidly with $\alpha\text{-TO}^\bullet$ (reaction 12), but now radical export must occur indirectly



since the ubisemiquinone radical (CoQH^\bullet) will remain "tied" to its LDL particle by its long (C50) alkyl tail. We have proposed that radical export takes place via the reaction of CoQH^\bullet (or $\text{CoQ}^{\bullet-}$) with oxygen.









The resulting superoxide radical ($\text{HOO}^\bullet / \text{O}_2^{\bullet-}$) may prevent oxidation chains from starting by reacting with another radical in the aqueous phase, or it may terminate a chain (reaction 11, $\text{X} = \text{HOO}^\bullet / \text{O}_2^{\bullet-}$, which may help to explain why as little as one CoQH_2 molecule per three LDL particles reduces the peroxidation rate³⁵).

Interparticle radical transport, when enhanced, will inhibit TMP by shortening the lifetime of the LDL-trapped $\alpha\text{-TO}^\bullet$ radicals (Scheme 5). Millimolar amounts of many natural and synthetic compounds and, remarkably, even the "prooxidant" *tert*-butyl hydroperoxide slowed the rate of TMP in LDL. To be really effective at low (i.e., micromolar) concentrations, however, a radical "shuttling" anti-TMP agent (XH) needs to (a) *react rapidly* with $(\alpha\text{-TO}^\bullet)_{\text{LDL}}$, (b) yield an antioxidant radical, X^\bullet , that *diffuses rapidly* to other particles, and (c) yield an X^\bullet radical that *terminates rapidly* with $(\alpha\text{-TO}^\bullet)_{\text{LDL}}$ (i.e., reaction 11 rather than reacting with $\alpha\text{-TOH}$ to reinitiate TMP, reaction -10).⁵⁰ The role of each of these factors was examined in an extensive study of potential co-antioxidants (excerpted in Table 1).¹² In brief, all else being equal, the most reactive phenols in our study give the best inhibition (cf. factor a). For phenols with identical chemical reactivities, those with the greatest water solubility afforded the greatest inhibition of TMP (cf. factor b). Thus, in a series of $\alpha\text{-TOH}$ homologues, the co-antioxidant effectiveness decreased as the length of the alkyl tail (R) attached to the chromanol headgroup increased (see Table 2).

The third factor (selectivity for termination rather than reinitiation) is illustrated by the high co-antioxidant effectiveness of quinols compared to phenols (Table 1) since, in contrast to phenoxy radicals, semiquinone radicals ($\text{X}^\bullet = \text{HQ}^\bullet$) are strongly reducing species that react very rapidly with $\alpha\text{-TO}^\bullet$ (e.g., cf. $k_{\text{TO}^\bullet/\text{CoQH}^\bullet} = 10^9 \text{ M}^{-1} \text{ s}^{-1}$ and $k_{\text{TO}^\bullet/\text{BHT}^\bullet} = 10^5 \text{ M}^{-1} \text{ s}^{-1}$).


Radical import might be varied either to enhance chain termination (reaction 11) or to lower the reactivity of LDL particles toward initiating radicals. The most obvious way to achieve the latter goal is to remove $\alpha\text{-TOH}$ from the LDL, and indeed, a recent study by Stocker *et al.*⁵¹ shows

Table 1. Co-antioxidant Effectiveness of Some Compounds Added to AAPH- or AMVN-Initiated LDL at a Concentration Equal to That of the Endogenous $\alpha\text{-TOH}$ (10 μM)

Compound	$(1 - R_p^{\text{with}} / R_p^{\text{without}}) \times 100\%$ ^a	
	AAPH	AMVN
<i>Ascorbic Acid</i>		
AsH ⁺ 	99.8	99.9
<i>Phenols</i>		
BHT 	90	95.2
DBHA 	97	95.3
ED 	-3	-11
<i>Quinols and catechols</i>		
TBHQ 	99.9	99.9
2HED 	98.1	99.8

^a Peroxidation rates (R_p) at 37 °C calculated from the time point corresponding to 20% consumption of $\alpha\text{-TOH}$ in the unsupplemented (without) sample (see ref 12).

Table 2. Co-antioxidant Effectiveness of $\alpha\text{-TOH}$ Homologues Added to AAPH-Initiated LDL at a Concentration Equal to That of the Endogenous $\alpha\text{-TOH}$ (10 μM)

 , $\text{R} =$	Co antioxidant effectiveness $(1 - R_p^{\text{with}} / R_p^{\text{without}}) \times 100\%$ ^a
CO_2H (Trolox)	99.0
CH_2OH	99.2
CH_3	95.4
$n\text{-C}_6\text{H}_{11}$	80.9
$n\text{-C}_{11}\text{H}_{23}$	35.6
$n\text{-C}_{13}\text{H}_{27}$	0
$n\text{-C}_{15}\text{H}_{31}$	-7


^a See Table 1 footnote a.

that peroxide-free, tocopherol-depleted LDL is virtually inert to AAPH-initiated peroxidation!

Another approach to lowering LDL's reactivity toward initiating radicals is to convert the endogenous $\alpha\text{-TOH}$ into its deuterated form, $\alpha\text{-TOD}$.¹³ In a homogeneous system this will accelerate autoxidation by about a factor of 4 since $\alpha\text{-TOD}$ is about 4-fold less reactive than $\alpha\text{-TOH}$ toward LOO^\bullet ⁵³ (eq I). However, in LDL reaction 5' will be slower with $\alpha\text{-TOD}$ than with $\alpha\text{-TOH}$, and therefore the rate of peroxidation was predicted to be slower (eq V). In the event, the AAPH-initiated peroxidation of LDL was found to be 2.5 times *slower* in D_2O buffer than in H_2O buffer.^{54,55} The implied deuterium kinetic isotope

effect, $DKIE = k_{ROO\cdot/TOH}/k_{ROO\cdot/TOD} \approx 2.5$, is lower than the DKIE in a nonpolar solvent.⁵³

Transition Metal-Induced Peroxidation. Most in vitro studies of LDL oxidation have used copper ions and air as the oxidant. Despite extensive studies, the exact mechanism for this oxidation remains uncertain.⁵⁶ However, it now seems clear that α -TOH plays a crucial role in the early stages of oxidation since tocopherol-depleted, peroxide-free LDL is relatively inert to oxidation by copper ions (and, for that matter, iron ions, *lipoxygenase*, *horse-radish peroxidase*, AAPH, and even radiolytically generated hydroxyl radicals⁵¹). Thus, in fresh rapidly isolated LDL the initial radical-generating reaction is undoubtedly reduction of (lipoprotein associated⁵⁷) Cu^{2+} to Cu^{1+} by α -TOH.^{58,59}



The lack of any definite relationship between the (so-called) *lag phase* in copper ion-initiated oxidations and the tocopherol content of the lipoprotein⁶ supports the notion that reaction 14 plays a dominant role in early peroxidation. Thus, since the rate of initiation will be *proportional* to the concentration of the major *antioxidant*, α -TOH (per reaction 14), sample to sample variation in the lag period in copper ion-induced LDL oxidation is not expected to correlate with the α -TOH /LDL ratio.⁶⁰

The relative importance of the TMP ensuing from reaction 14 depends on how much copper is used. Large amounts of copper consume α -TOH so quickly that very little LOOH is produced in the inhibited period, whereas slower initiation (e.g., with less than two copper ions per LDL particle) leads to substantial peroxidation in the "inhibited period".⁶¹ Under very mild oxidizing conditions (such as those produced by a common cell culture medium, F-10) over 50% of the PUFA lipid in native LDL may be oxidized *before* the tocopherol is consumed,²⁶ a clear illustration of the importance of TMP in vitro if not in vivo.

TMP and Atherosclerosis

According to the oxidized lipoprotein hypothesis the initiating event for atherosclerosis is a modification of LDL by oxidants in the artery wall. This "oxidatively modified LDL" is recognized and scavenged by macrophages, transforming them into lipid-laden "foam" cells that deposit as fatty streaks and eventually plaque on the artery wall.⁶²

The role of α -TOH in this in vivo process is hard to define since the conditions and agents that modify LDL are unknown.⁵⁸ Our in vitro experiments have demonstrated that the lipid in LDL can be substantially oxidized *in the presence of* α -TOH. It is therefore tempting to ascribe an in vivo TMP process to the finding that in plaque containing normal amounts of vitamin E, "Approximately 30% of plaque Ch18:2 was oxidized..."⁶³ Unfortunately, however, this may be an oversimplification since the coexistence of oxidized lipid and antioxidant in plaque extracts may merely reflect the localized nature

of peroxidation; i.e., peroxidation may occur in sites that are sequestered from replenishment of endogenous antioxidants such as ascorbate. Indeed, lipid peroxidation may even be a *result* of scavenger-cell ingestion rather than its primary cause; i.e., the LDL modification leading to its uptake by macrophages may have been due to a more protein-specific reaction (e.g., with neutrophil-derived chlorine/hypochlorite⁶⁴), with lipid peroxidation occurring subsequently in the macrophage vacuole (possibly initiated by tyrosyl radicals⁶⁵ and sequestered from ascorbate).

Outlook

Careful in vitro kinetic studies revealed that in LDL the powerful antioxidant, α -TOH, can become a powerful prooxidant—particularly under mildly oxidizing conditions which consume α -TOH very slowly. It is reasonable to expect that the oxidizing conditions in vivo will be even milder than any applied in vitro. However, this does not necessarily mean that TMP must be important in vivo because vitamin C, ubiquinol, and many other natural and synthetic compounds can eliminate TMP—provided they have physical access to LDL/ α -TO \cdot particles. The prime research challenge today is to determine the importance of TMP in vivo and the magnitude of its contribution to atherosclerosis. Until answers to these problems are forthcoming, we simply recommend an adequate daily intake of both vitamins E and C as being a risk-free, cheap, and reasonable way to avert a possible heart attack or stroke.

References

- (1) Issued as NRCC No. 0000.
- (2) LDL is the main "cholesterol delivery vehicle" of human blood; LDL particles carry cholesterol—mainly in the form of cholesteryl esters—from the liver to tissues throughout the body. The name derives from the flotation density of LDL in aqueous suspension ($d = 1.00\text{--}1.1$ g/mL) which distinguishes it from the other lipid-bearing proteins in blood and lymph, notably the high-density lipoprotein (HDLs, $d > 1.15$ g/mL, the "cholesterol recovery vehicle" in blood), very low-density lipoproteins (VLDLs, $0.9\text{--}1.0$ g/mL), and chylomicrons.
- (3) Steinberg, D. P., S.; Carew, T. E.; Khoo, J. C.; Witztum, J. L. *New England J. Med.* **1989**, *320*, 915–924. Steinberg, D. *Lancet* **1995**, *346*, 36–38. Maggi, E.; Chiesa, R.; Melissano, G.; Castellano, R.; Astore, D.; Grossi, A.; Finardi, G.; Bellomo, G. *Arterioscler. Thromb.* **1994**, *14*, 1892–1899. Holvoet, P.; Collen, D. *FASEB J.* **1994**, *8*, 1279–1284. Palinski, W.; Ord, V. A.; Plump, A. S.; Breslow, J. L.; Steinberg, D.; Witztum, J. L. *Arterioscler. Thromb.* **1994**, *14*, 605–616.
- (4) See, e.g.: Gaziano, J. M. *Am. J. Med.* **1994**, *97*, 18S–21S. Hoffman, R. M.; Garewal, H. S. *Arch. Intern. Med.* **1995**, *155*, 241–246; Gey, K. F., Puska, P., Jordan, P.; Moser, U. K. *Am. J. Clin. Nutr.* **1991**, *53*, 326S–334S.
- (5) Sasahara, M.; Raines, E. W.; Chait, A.; Carew, T. E.; Steinberg, D.; Wahl, P. W.; Ross, R. *J. Clin. Invest.* **1994**, *94*, 155–164.
- (6) Review: Esterbauer, H.; Gebicki, J.; Puhl, H.; Jürgens, G. *Free Radical Biol. Med.* **1992**, *13*, 341–390.

- (7) Burton, G.; Ingold, K. U. *Acc. Chem. Res.* **1986**, *19*, 194–201.
- (8) Some reported correlations between α -TOH content and resistance of LDL to peroxidation: (Positive) Esterbauer, H.; Dieber Rotheneder, M.; Striegl, G.; Waeg, G. *Am. J. Clin. Nutr.* **1991**, *53*, 314s–321s. (Erratic or minor) McDowell, I. F. W.; Brennan, G. M.; McEneny, J.; Young, I. S.; Nicholls, D. P.; McVeigh, G. E.; Bruce, I.; Trimble, E. R.; Johnston, G. D. *Eur. J. Clin. Invest.* **1994**, *24*, 759–765. (Negative) Iwatsuki, M.; Niki, E.; Stone, D.; Darley-Usmar, V. M. *FEBS Lett.* **1995**, *360*, 271–276. Croft, K. D.; Williams, P.; Dimmitt, S.; Abu Amsha, R.; Beilin, L. *J. Biochim. Biophys. Acta* **1995**, *1254*, 250–256.
- (9) Ingold, K. U.; Bowry, V. W.; Stocker, R.; Walling, C. *Proc. Nat. Acad. Sci. U.S.A.* **1993**, *90*, 45–49.
- (10) Neuzil, J.; Thomas, S. R.; Stocker, R. *Free Radical Biol. Med.* **1997**, *22*, 57–71 and cited references.
- (11) Bowry, V. W.; Ingold, K. U.; Stocker, R. *Biochem. J.* **1992**, *288*, 341–344.
- (12) Bowry, V. W.; Mohr, D.; Cleary, J.; Stocker, R. *J. Biol. Chem.* **1995**, *270*, 5756–5763.
- (13) Witting, P. K.; Bowry, V. W.; Stocker, R. *FEBS Lett.* **1995**, *375*, 45–49.
- (14) Where LH refers to a bisallylic group in a polyunsaturated fatty acid ester (phospholipid, triglyceride, cholesterol ester, etc.), L^{\bullet} and LOO^{\bullet} are the derived carbon-centered radical and lipid peroxy radical, respectively, and ROO^{\bullet} is the peroxy radical which initiates lipid peroxidation.
- (15) Burton, G. W.; Doba, T.; Gabe, E.; Hughes, L.; Lee, F. L.; Prasad, L.; Ingold, K. U. *J. Am. Chem. Soc.* **1985**, *107*, 7053–7065.
- (16) Niki, E.; Saito, M.; Kawakami, A.; Kamiya, Y. *J. Biol. Chem.* **1984**, *259*, 4177–4182.
- (17) Avila, D. V.; Brown, C. E.; Ingold, K. U.; Luszytk, J. *J. Am. Chem. Soc.* **1993**, *115*, 466–470.
- (18) Valgimigli, L.; Banks, J. T.; Ingold, K. U.; Luszytk, J. *J. Am. Chem. Soc.* **1995**, *117*, 9966–9971. Valgimigli, L.; Ingold, K. U.; Luszytk, J. *J. Org. Chem.* **1996**, *61*, 7947–7950.
- (19) Avila, D. V.; Ingold, K. U.; Luszytk, J.; Green, W. H.; Procopio, D. R. *J. Am. Chem. Soc.* **1995**, *117*, 2929–2930.
- (20) MacFaul, P. A.; Ingold, K. U.; Luszytk, J. *J. Am. Chem. Soc.* **1996**, *118*, 1316–1321.
- (21) Castle, L.; Perkins, M. J. *J. Am. Chem. Soc.* **1986**, *108*, 6381–6382. Pryor, W. A.; Strickland, T.; Church, D. F. *J. Am. Chem. Soc.* **1988**, *110*, 2224–2229.
- (22) Barclay, L. R. C.; Baskin, K. A.; Locke, S. J.; Vindquist, M. R. *Can. J. Chem.* **1990**, *68*, 2258–2269.
- (23) Valgimigli, L.; Ingold, K. U.; Luszytk, J. *J. Am. Chem. Soc.* **1996**, *118*, 3545–3549.
- (24) Indeed, a recent study by Barclay²⁵ (using bisallylperoxy as a radical clock) yields a 22-fold higher k_{inh}/k_p ratio than was obtained from inhibited-rate measurements;²² cf. $k_{inh}/k_p = 170$ (inhibition) vs 3800 (peroxy clock).
- (25) Barclay, L. R. C.; Vinqvist, M. R.; Antunes, F.; Pinto, R. E. *J. Am. Chem. Soc.* **1997**, *119*, 5764–5765.
- (26) Bowry, V. W.; Stocker, R. *J. Am. Chem. Soc.* **1993**, *115*, 6029–6044.
- (27) Sattler, W.; Kostner, G. M.; Waeg, G.; Esterbauer, H. *Biochim. Biophys. Acta* **1991**, *1081*, 65–74.
- (28) Carotenoids—which give LDL its yellow-orange color—do not inhibit peroxidation in LDL. Gaziano, J. M.; Hatta, A.; Flynn, M.; Johnson, E. J.; Krinsky, N. I.; Ridker, P. M.; Hennekens, C. H.; Frei, B. *Atherosclerosis* **1995**, *112*, 187–195.
- (29) For two radicals cohabiting the same particle, $[\text{radical}] = 2/N_A V_{\text{lipid}}$ so that lifetime $\tau_{1/e} = N_A V_{\text{lipid}}/4k_t$, with $2k_t/M^{-1} s^{-1} \approx 10^3$ ($2 \alpha\text{-TO}^{\bullet}$), 10^6 (2LOO^{\bullet}), and 3×10^8 ($\text{LOO}^{\bullet} + \alpha\text{-TO}^{\bullet}$).
- (30) Bowry, V. W.; Stanley, K. K.; Stocker, R. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, *89*, 10316–10320.
- (31) Howard, J. A.; Ingold, K. U. *Can. J. Chem.* **1967**, *45*, 793–802.
- (32) See, e.g.: Esterbauer, H.; Striegl, G.; Puhl, H.; Rotheneder, M. *Free Radical Res. Commun.* **1989**, *6*, 67–75.
- (33) Stocker, R.; Barsamian, S.; Frei, B. *Free Radical Biol. Med.* **1990**, *9S*, 71.
- (34) Sato, K.; Niki, E.; Shimasaki, H. *Arch. Biochem. Biophys.* **1990**, *279*, 405–409.
- (35) Stocker, R.; Bowry, V. W.; Frei, B. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 1646–1660.
- (36) The rate of LOOH formation increased ~40-fold following the conversion of ~70% of the initial CoQH₂ into ubiquinone-10 (CoQ).
- (37) Barclay, L. R. C.; Vinqvist, M. R.; Mukai, K.; Itoh, S.; Morimoto, H. *J. Org. Chem.* **1993**, *58*, 7416–7420.
- (38) Doba, T.; Burton, G. W.; Ingold, K. U. *Biochim. Biophys. Acta* **1985**, *835*, 298–303.
- (39) That is, not until over 85% of the original α -TOH had been consumed, at which point many of the particles would be left unprotected.
- (40) Bowry, V. W.; Ingold, K. U. *J. Org. Chem.* **1995**, *60*, 5456–5467.
- (41) Massey, J. B. *Biochim. Biophys. Acta* **1984**, *793*, 387–392.
- (42) The diffusion half-life of α -TOH in LDL is on the order of 1000 s,⁴¹ and that of $\alpha\text{-TO}^{\bullet}$ may be even longer because it lacks the hydrophilic OH group.
- (43) With the simplest case assumption that particles are uniformly likely to react with the initiating ROO^{\bullet} .
- (44) Cf. for polymerization of a monomer (e.g., styrene) in an aqueous latex emulsion⁴⁵ $R_{\text{prop}} = -d[\text{monomer}]/dt = ((\text{no. of particles})/2)[\text{monomer}] \cdot k_{\text{polymer}/\text{monomer}}$.
- (45) Walling, C. W. *Free Radicals in Solution*; Wiley: New York, 1957.
- (46) Based on $k_{ROO^{\bullet}/\alpha\text{-TOH}} = 2 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$, $k_{ROO^{\bullet}/\alpha\text{-TO}^{\bullet}} = 3 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$, $[\alpha\text{-TOH}]_{\text{intraparticle}} = 3 \text{ mM}$, and $[\alpha\text{-TO}^{\bullet}]_{\text{intraparticle}} = 1 \text{ molecule/particle volume} \approx 0.5 \text{ mM}$.
- (47) Reference 26 gives a more exact analysis.
- (48) Neuzil, J.; Stocker, R. *J. Biol. Chem.* **1994**, *269*, 16712–16719.
- (49) Doba, T.; Burton, G. W.; Ingold, K. U.; Matsuo, M. *J. Chem. Soc., Chem. Commun.* **1984**, 461–462.
- (50) Simic, M. G.; Jovanovic, S. V. *J. Am. Chem. Soc.* **1989**, *111*, 5778–5782.
- (51) Kinetic analysis of Schemes 4 and 5 at the fast diffusion limit gives $R_p^{\text{inh}} = k_{\text{TMP}}[\text{LH}][\alpha\text{-TOH}](R_i/C_{\text{XH}}[\text{XH}])^{0.5}$ with $C_{\text{XH}} = k_{\text{TO}^{\bullet}/\text{XH}}(k_{\text{exit}}/k_{\text{X}^{\bullet}/\text{TOH}})(2k_{11}/k_{\text{exit}})$. The *co-antioxidant effectiveness factor*, C_{XH} , depends on the reactivity of XH to $\alpha\text{-TO}^{\bullet}$, the rate at which X^{\bullet} escapes the particle, and the relative rate at which an “escaped” X^{\bullet} radical reacts with nonoxidizing vs oxidizing particles. The predicted $R_p^{\text{inh}} \propto \{R_i/[\text{XH}]\}^{0.5}$ relationship has been confirmed for some compounds.
- (52) Neuzil, J.; Thomas, S. R.; Stocker, R. *Free Radical Biol. Med.* **1997**, *22*, 57–71.
- (53) Burton, G. W.; Ingold, K. U. *J. Am. Chem. Soc.* **1981**, *103*, 6472–6477.

- (54) Proton exchange between phenols and water is very facile, and thus, more generally, if an antioxidant or mixture of antioxidants is acting in chain-transfer (prooxidant) mode, peroxidation will be slower in a D₂O-washed reaction mixture than a H₂O-washed reaction mixture and vice versa.
- (55) Initiation efficiency was about 2-fold lower in D₂O-LDL, presumably because of the lower phase-transfer activity of α -TOD.
- (56) See, e.g.: (a) Abuja, P. M.; Albertini, R.; Esterbauer, H. *Chem. Res. Toxicol.* **1997**, *10*, 644–651. (b) Proudfoot, J. M.; Croft, K. D.; Puddey, I. B.; Beilin, L. J. *Free Radical Biol. Med.* **1997**, *23*, 720–728.
- (57) Gieseg, S. P.; Esterbauer, H. *FEBS Lett.* **1994**, *343*, 188–194.
- (58) See: Yoshida, Y.; Tsuchiya, J.; Niki, E. *Biochim. Biophys. Acta* **1994**, *1200*, 85–92. Lynch, S. M.; Frei, B. *J. Biol. Chem.* **1995**, *270*, 5158–5163. Kontush, A.; Meyer, S.; Finckh, B.; Kohlschutter, A.; Beisiegel, U. *J. Biol. Chem.* **1996**, *271*, 11106–11112.
- (59) Cu¹⁺ may then react with LOOH (generated from the α -TO[•]), i.e., $\text{Cu}^{1+} + \text{LOOH} \rightarrow \text{Cu}^{2+} + \text{LO}^{\bullet} + \text{OH}^-$. However, this is uncertain since (i) Cu¹⁺ autoxidizes in air and (ii) removal of Cu¹⁺ with neocuproin *accelerates* tocopherol oxidation.^{56b}
- (60) For example, linear regression analysis of data for LDL from 74 donors yielded lag time/min = 1.6 (mol of α -TOH/mol of LDL) + 59, $r^2 = 0.043$ (Figure 14 in ref 6).
- (61) A consequence of R_p being independent of the initiation rate.
- (62) Reviewed in Berliner, J. A.; Heinecke, J. W. *Free Radical Biol. Med.* **1996**, *20*, 707–727.
- (63) Suarna, C.; Dean, R. T.; May, J.; Stocker, R. *Arterioscler. Thromb.* **1995**, *15*, 1616–1624.
- (64) Leeuwenburgh, C.; Rasmussen, J. E.; Hsu, F. F.; Mueller, D. M.; Pennathur, S.; Heinecke, J. W. *J. Biol. Chem.* **1997**, *272*, 3520–3526. Hazell, L. J.; Van Den Berg, J. J. M.; Stocker, R. *Biochem. J.* **1994**, *302*, 297–304.
- (65) Savenkova, M. I.; Mueller, D. M.; Heinecke, J. W. *J. Biol. Chem.* **1994**, *269*, 20394–20400.

AR9500590