

Recovery of Carbonyl Compounds from *N,N*-Dialkylhydrazones

DIETER ENDERS,* LARS WORTMANN, AND RENÉ PETERS

Institut für Organische Chemie der Technischen Hochschule, Professor-Pirlet-Strasse 1, D-52074 Aachen, Germany

Received September 16, 1999

ABSTRACT

Deprotonation of enantiomerically pure hydrazones and subsequent trapping with suitable electrophiles generates new stereogenic centers with excellent stereoselectivity. To liberate the original carbonyl functionality in the final products, it is necessary to cleave the hydrazone moiety. In recent years, many reagents have been developed to regenerate carbonyl compounds from the corresponding dialkylhydrazones which are compatible with a wide range of functionalities. This has allowed the use of hydrazones in the total synthesis of complex natural products. This Account is meant to be an overview of methods which are classified as oxidative, hydrolytic, and reductive cleavage procedures.

Introduction



The development of efficient and highly selective methods for C–C and C–X bond formation has been, and continues to be, a challenging and exciting endeavor in organic chemistry. Procedures that allow the construction of C–C and C–X bonds α to the carbonyl group via electrophilic substitution are among the most important synthetic operations. Most of the problems in classical carbonyl chemistry, such as aldol-type self-condensation, di- and polyalkylation, control of regiochemistry, side reactions of products, and lack of reactivity of the corresponding enolates, have been solved by the use of *N,N*-dialkylhydrazones **2** as equivalents of the parent ketones and

Dieter Enders was born in 1946 in Butzbach, Germany. He studied chemistry at the Justus Liebig Universität Giessen and received his Dr. rer. nat. in 1974 under the supervision of Prof. Dieter Seebach. After postdoctoral studies at Harvard University with Prof. E. J. Corey, he went back to Giessen and obtained his habilitation in 1979. In 1980 he moved to the Universität Bonn as an associate professor and in 1985 to his present position as Professor of Organic Chemistry at the Rheinisch-Westfälische Technische Hochschule Aachen. His current research interests are asymmetric synthesis, new synthetic methods using organometallics, and the stereoselective synthesis of biologically active compounds.


Lars Wortmann was born in 1972 in Bad Soden am Taunus, Germany. He studied chemistry at the Johann-Wolfgang-Goethe Universität Frankfurt am Main and took his diploma with Prof. Johann Mulzer. In 1997 he moved to the Rheinisch-Westfälische Technische Hochschule Aachen, where he is currently working on his Ph.D. under the supervision of Prof. Dieter Enders. He is a member of the Graduiertenkolleg, and his main research interests are in natural product total synthesis as well as the development of new methodology for asymmetric synthesis.

René Peters was born in 1971 in Simmerath, Germany. He received his diploma of chemistry from the Rheinisch-Westfälische Technische Hochschule Aachen in 1997. In 1998 he was awarded the Springorum Award from the same university, where he is currently working on his Ph.D. under the supervision of Prof. Dieter Enders. His research interests are focused on organometallic chemistry, especially ligand design for asymmetric catalysis.

Scheme 1



Scheme 2




aldehydes **1**.¹ Deprotonation of enantiomerically pure hydrazones **2** and subsequent trapping with suitable electrophiles EX generates new stereogenic centers with excellent stereoselectivity. To liberate the original carbonyl functionality in the final products **4**, it is necessary to cleave the hydrazones **3**. Many new cleavage reagents have been developed over the past four decades which are compatible with a wide range of other functionalities. In general, cleavage procedures can be classified into oxidative, hydrolytic, and reductive methods. In the case of aldehyde dialkylhydrazones **5**, the removal of the hydrazone can lead to two different cleavage products (Scheme 2). Hydrolysis of **5** gives rise to the corresponding aldehydes **7** in a manner similar to the ketone cases. Another pathway is via oxidation or elimination, which leads to the corresponding nitriles **6**. As the nitrile functionality can be easily transformed into the corresponding aldehyde **7** by simple DIBAL reduction,² this Account will also mention reagents leading to the nitriles **6**.

Oxidative Cleavage Procedures

Ozone (O₃). *N,N*-Dimethylhydrazones (DMHs) and SAMP/RAMP hydrazones **8** [SAMP and RAMP = (*S*)- and (*R*)-1-

* To whom correspondence should be addressed. Fax: Int. +241/8888127. E-mail: Enders@RWTH-Aachen.de.





amino-2-methoxymethylpyrrolidine] of aldehydes and ketones may be oxidatively cleaved by ozonolysis in dichloromethane at -78°C .³ This reaction is very clean, and the only products formed are the desired carbonyl compounds **10** and 1 equiv of nitrosamine **9** in quantitative yield. The formation of the latter allows recycling of the chiral auxiliary RAMP or SAMP (**11**) after separation by distillation or chromatography and subsequent LAH reduction (Scheme 3).



Although the chemical yield of the recycling step is good ($\sim 80\%$), a partial racemization of up to 10% of the chiral auxiliary was observed in rare cases. Advantages of the oxidative cleavage by ozonolysis are extremely mild reaction conditions, low temperature (-78°C), neutral pH, short reaction times and the ease of detecting the end of the reaction (appearance of a green-colored mixture of yellow nitrosamine **9** and blue ozone or by addition of an indicator such as Sudan red), the excellent, almost quantitative yields, and the lack of any racemization of the sensitive aldehydes ($\text{R}^1 = \text{H}$) and ketones **10** during the operation. Of course, the method cannot be applied in the presence of other functional groups that are not compatible with ozone under these reaction conditions. However, because the $\text{C}=\text{N}$ double bond is highly reactive toward ozone, it is possible to obtain the desired carbonyl compounds chemoselectively, even in the presence of functional groups that are sensitive to oxidation, such as thioethers,⁴ protected α -hydrazino and aminoketones,⁵ and borane-protected phosphines,⁶ by TLC control or by the addition of certain diazo dyes as indicators.

Although the mechanism of the ozonolysis is not exactly known, it is conceivable that the oxidative $\text{C}=\text{N}$ bond cleavage starts with intermediates of type **12a** or **12b**, which directly decompose to carbonyl compound **10** and diazene **13**, which is further oxidized under the reaction conditions to the nitrosamine **9**. Indeed, 2 equiv of ozone per hydrazone are consumed (Scheme 4).⁷

Out of the manifold applications of SAMP hydrazone cleavage by ozonolysis, we would like to focus on some recent highlights in total synthesis of bioactive compounds. Swinholide A⁸ was impressively synthesized by Nicolaou et al. Our hydrazone methodology followed by ozonolysis was used to build up stereogenic centers at C-24 and C-24' (Scheme 5).



Holmes et al. used the SAMP/RAMP hydrazone method in their synthesis of medium-ring ethers such as model-compound (+)-octahydrodeacetylde bromolaurencin





(Scheme 6).⁹ After asymmetric alkylation (*de* = 90%) of the cycloheptanone SAMP hydrazone **14** followed by ozonolysis, chiral ketone **15** underwent a Baeyer–Villiger oxidation by treatment with trifluoroacetic acid to yield key intermediate **16**.

At an early stage of the total synthesis of (–)-C₁₀-desmethyl arteannuin B, the SAMP/RAMP hydrazone method made it possible to install a key stereogenic center to yield **17** as a single diastereomer (Scheme 7).¹⁰ During the ozonolysis, the conjugated double bond and dimethyl-acetal were resistant to the reaction conditions.

An elegant example of an iterative alkylation was used in the total synthesis of zaragozic acid A (Scheme 8) by Nicolaou et al.¹¹


Singlet Oxygen (¹O₂). The dye-sensitized photooxygenation of a variety of *N,N*-dimethylhydrazones and SAMP hydrazones in MeOH, THF, or dichloromethane at –78 to 20 °C followed by reduction with Ph₃P or DMS and hydrolysis yielded the parent carbonyl compounds **22** in moderate to good yields.¹² This method appears to have broad applicability for the regeneration of aldehydes and



ketones, as well as polyfunctional and relatively sensitive carbonyl compounds such as aldol adducts. Mechanistically, there are two obvious possibilities for the course of the oxidative cleavage, an ene-type reaction via a hydroperoxide **19** or a cycloaddition reaction via the dioxazetidine **20** as an intermediate (Scheme 9). Since dimethylnitrosamine **21** is not formed and the hydrazones **18** do not react in the absence of a sensitizer, the cycloaddition route and an autoxidation process can be excluded. Further evidence for the proposed ene-type reaction via **19** comes from the fact adamantanone-*N,N*-dimethylhydrazone **23** remains unchanged under the reaction conditions (Bredt's rule).

NaIO₄ and H₃IO₆. Dimethylhydrazones can be cleaved to the corresponding ketones in high yields under mild

Scheme 10



conditions using aqueous sodium periodate at pH 7 and room temperature.^{1b,c} The neutral periodate hydrolysis procedure is not satisfactory for the cleavage of dimethylhydrazone derivatives of aromatic or α,β -unsaturated aldehydes, since these usually afford mixtures of aldehydes and nitriles. However, under modified conditions, by the use of periodic acid H₅IO₆, nonconjugated aldehydes can be obtained in high yield from the corresponding dimethylhydrazones. In general, the production of nitrile at the expense of aldehyde is favored at higher pH. This method tolerates a variety of sensitive functional groups within the molecule such as acetals, silyl ethers, esters, and PMB ethers, as demonstrated in the total synthesis of ulapualide A by Panek et al. (Scheme 10).¹³

NaBO₃. *N,N*-Dimethylhydrazones and SAMP hydrazones are efficiently cleaved back to the parent ketones **24–30** under mild conditions at pH 7.¹⁴ The oxidative hydrolysis is effected in good yields (Table 1) with the very cheap bulk chemical sodium perborate in buffer solution and *tert*-butanol as cosolvent. Whereas DMH and SAMP hydrazones underwent smooth oxidative hydrolysis in 4–24 h, the cleavage of aromatic ketone hydrazones took 2–3 d. The procedure is also chemoselective, leaving a C=C double bond intact.

Tetrabutylammonium Peroxydisulfate. Tetrabutylammonium peroxydisulfate ((TBA)₂S₂O₈) has been shown to be a good source of a sulfate anion radical, which has the potential to be widely applicable in organic reactions.¹⁵ This radical anion is considered to oxidize the C=N double bond. It has been found that treatment of DMHs with

Table 1. Yields of *N,N*-Dialkylhydrazone Cleavages with NaBO₃

No.	Product	Hydrazone	Yield [%]
24		DMH	70
25		DMH	91
25		SAMP	86
26		DMH	95
27		DMH	85
28		DMH	90
29		SAMP	87
30		DMH	94



(TBA)₂S₂O₈ gave the parent ketones in excellent yields (89–97%). Because the reaction medium is at pH close to neutral, cleavage reactions have been found to proceed for hydrazones which contain sensitive functional groups, such as acetal and olefin moieties. Although the reaction mechanism is not yet clear, the cleavage appears to be initiated via the addition of sulfate radical to the carbon of C=N, and the oxygen of the carbonyl product probably originates from the peroxydisulfate.

[Hydroxy(tosyloxy)iodo]benzene (HTIB or Koser's Reagent). The carbonyl groups of various hydrazone derivatives of α -keto esters **31** were readily regenerated in high yield through oxidative hydrolysis using hypervalent organoiodine(III) reagent [hydroxy(tosyloxy)iodo]benzene (HTIB, **33**; Scheme 11).¹⁶ In contrast to **33**, the α -ketoesterdimethylhydrazones **31** were not cleaved by treatment with [bis(trifluoroacetoxy)iodo]benzene (BTI, **34**).


Magnesium Monoperoxyphthalate (MMPP). MMPP, a reagent with high stability at ambient temperature, has been shown to oxidize a wide range of substrates under milder conditions than MCPBA: the substrates include alkenes, ketones, sulfides and sulfoxides, and pyridine.¹⁷

N,N-Dimethylhydrazones and SAMP hydrazones of aliphatic or aromatic ketones undergo oxidative cleavage

Scheme 11



Scheme 12




on reaction with MMPP hexahydrate under mild conditions.^{18,19} The reaction is high-yielding (76–94%) and chemoselective and proceeds without racemization (>97%). In contrast, aldehyde hydrazones **35**, bearing a stereogenic center α to the hydrazone moiety, give rise to the parent nitriles **37** in good yields and without racemization via an oxy-Cope-type elimination via **36** (Scheme 12).²⁰




The broad applicability of this method has been demonstrated in several natural product syntheses, such as the enantioselective synthesis of a library of epothilones by Nicolaou et al., as depicted in Scheme 13.²¹ Key intermediate **38**, containing an isolated and conjugated double bond as well as a silyl ether group and a highly oxidation-sensitive thiazole moiety, was smoothly converted to the corresponding nitrile **39**, which could be further transformed into aldehyde **40** in excellent yield (81–90%).

In the first total synthesis of stigmatellin A, a strong inhibitor of the electron transport in chloroplasts and mitochondria, intermediate **41** was cleaved with MMPP to **42** in the presence of the paramethoxybenzyl ether moiety as an alcohol protecting group (Scheme 14).²²

MCPBA and MeCO₃H. The 3-chloroperoxybenzoic acid (MCPBA) oxidation of DMHs of aromatic aldehydes gives the corresponding nitriles (**43**–**51**) in good yields (Table 2).²³ The oxidation of aliphatic DMHs, however, gave rise to the formation of mixtures of nitriles and aldehydes (see **51**) and is therefore of no synthetic value. A further disadvantage is the long reaction time (several days). MCPBA was successfully used to cleave DMH **52** on a large

Scheme 13





product mixtures were obtained. The corresponding nitriles could be isolated in moderate yields.²⁵

H₂O₂/SeO₂ and H₂O₂/2-Nitrobenzeneselenic Acid (2-NBSeA). In comparison to the above-mentioned MCPBA procedure, H₂O₂ activated by SeO₂ or 2-nitrobenzeneselenic acid, respectively, resulted in shorter reaction times and higher yields. Surprisingly, no oxidation of aromatic aldehyde dimethylhydrazones was observed when the mixture was stirred with 30% H₂O₂, even for 12 d. Except for the more easily hydrolyzed aliphatic derivatives, all hydrazones were converted into nitriles in good yields (Table 2). The reaction catalyzed by SeO₂ was faster than that carried out in the presence of 2-nitrobenzeneselenic acid (2-NBSeA), but the yields were higher in the latter case. Oxidation of the aliphatic derivative with H₂O₂/2-NBSeA gave only the parent aldehyde, while the faster reaction with H₂O₂/SeO₂ produced a mixture of aldehyde and nitrile **51**. These results are in accordance with the tendency of aliphatic DMHs to undergo hydrolysis under the aqueous oxidation conditions. The H₂O₂/SeO₂ method was successfully applied to the cleavage of RAMP hydrazones **54** to provide chiral nitriles **55** for monomers in material science (Scheme 16).²⁶

H₂O₂. The H₂O₂ oxidation of the DMHs of cinnamaldehyde, benzaldehyde, and alkoxy-substituted benzaldehydes gave the corresponding nitriles in good yields. Oxidation of DMHs of *p*-nitrobenzaldehyde, *p*-chlorobenzaldehyde, *n*-valeraldehyde, and *n*-octanal gave mixtures of nitrile-containing products.²⁷ In the synthesis of aldol products via SAMP hydrazones of ketones, this method was applied yielding the products in good yields.²⁸


Dimethyldioxirane. The use of dimethyldioxirane (DMDO) is feasible for the cleavage of *N,N*-dimethylhydrazones, SAMP hydrazones, and aryl- and tosylhydrazones in high yields.²⁹ The procedure proceeds very rapidly; e.g., α -methylcyclohexanone SAMP hydrazone was cleaved within 5 min in 85% yield. Only minimal

Table 2. Comparison of MCPBA, H₂O₂/SeO₂, and H₂O₂/2-NBSeA

Product nitrile		MCPBA	H ₂ O ₂ /SeO ₂	H ₂ O ₂ /2-NBSeA
		Yield [%]	Yield [%]	Yield [%]
43		78	95	97
44		71	88	92
45		76	82	91
46		77	93	97
47		75	78	84
48		55	58	65
49		50	64	73
50		83	87	95
51		65	45	83 (aldehyde)

scale to ketone **53** without any epimerization of the axial methyl group (Scheme 15).²⁴

In a similar manner, peracetic acid has been used for aromatic aldehyde dimethylhydrazone cleavage at low temperatures, whereas at high temperatures undefined




racemization could be detected (less than 2%, ee = 94%). Under the reaction conditions, C=C double bonds were not attacked and acetoxy groups not hydrolyzed. Mechanistic studies revealed the reaction pathway shown in Scheme 17.

EPR studies suggested that the radical cation species in radical pair **56** was formed as an intermediate. Subsequent trapping of this with the bisoxyl radical anion in **56** (or direct electrophilic *O*-transfer from DMDO to the substrate) should provide **57**; the latter reacts with another equivalent of DMDO to give rise to the desired ketones **59** and nitrosamine **60** (pathway A). Pathway B explains the formation of small amounts of oxime **61** as side product.

MoF₆, **CoF₃**, **WF₆**, **UF₆**, and **MoOCl₃**. The covalent nature of transition metal fluorides in high oxidation states such as MoF₆,³⁰ CoF₃,³¹ WF₆,³² and UF₆³³ allows the use of nonaqueous solvents such as CHCl₃, CH₂Cl₂, and Freon 113 during the *N,N*-dimethylhydrazone cleavage. Although it is known that some of these compounds are fluorinating agents for carbonyl groups, no fluorinated products were observed. The reagents are easy to handle, and glass is not attacked. The desired carbonyl compounds may be obtained in moderate to excellent yields (Table 3). In the UF₆ case, it was demonstrated that ketone, ester, amide, nitrile, and nitro moieties were tolerated under the reaction conditions. The proposed mechanism of the MoF₆ cleavage is depicted in Scheme 18 and was shown to work similarly for the other cases. MoF₆ attacks in an electrophilic manner at the dialkylated nitrogen of hydrazone **62**. The treatment of the activated hydrazone salt

Table 3. Yields of Oxidative Transition Metal Halide-Mediated DMH Cleavages

Carbonyl Compound	MoF ₆ Yield [%]	CoF ₃ Yield [%]	WF ₆ Yield [%]	UF ₆ Yield [%]	MoOCl ₃ Yield [%]
	78	-	73	46	85
	96	46	83	-	76
	85	74	93	41	80
	66	89	91	-	88
	95	-	85	50	96
	94	94	94	-	-
	-	67	-	56	78
	-	76	-	69	-
	50	-	-	-	-




63 with water furnishes the carbonyl compound **65** and diazene **64** as byproduct.

In contrast to the ionic mechanism of the fluorides, the MoOCl₃ cleavage is proposed to proceed by a single electron transfer with intermediate radical cation **66**, which is trapped by the resulting Mo(IV) species (Scheme 19).³⁰

NOBF₄, **NO₂BF₄**, and **Clay-Supported Ferric Nitrate**. Nitronium and nitrosonium salts are capable of mediating

Scheme 19


Table 4. Yields of DMH Cleavage Using NOBF₄, NO₂BF₄, and Clay-Supported Ferric Nitrate 4


Carbonyl Compound	NOBF ₄	NO ₂ BF ₄	Clayfen
	Yield [%]	Yield [%]	Yield [%]
	75	73	87
	59	70	85
	82	50	-
	66	53	-
	86	80	-
	-	-	78
	-	-	80

hydrolysis either by electron transfer or by the nitrosylation route.³⁴ Accordingly, dimethylhydrazones were subjected to reaction with equimolar quantities of nitronium or nitrosonium tetrafluoroborate, yielding the corresponding ketones in moderate to good yields (Table 4).

Clayfen (ferric nitrate impregnated on K-10 bentonite clay) is a nitrosonium ion source comparable to NOBF₄ and is capable of effecting cleavage of the dimethylhydrazone group. The reaction is fast and exothermic, giving the corresponding carbonyl compounds in very good yields (Table 4).³⁵

MeReO₃/H₂O₂. The rhenium(VII)-catalyzed oxidative cleavage of aliphatic or aromatic DMH's with H₂O₂ gives the corresponding nitriles in good yields (83–89%).³⁶ Double bonds are resistant to the reaction conditions.


Scheme 20


Table 5. Effects of Base and Temperature on Chemical Yield and Enantiomeric Excess

solvent	base (equiv)	T (°C)	reaction		ee (%)
			time (h)	yield (%) ^a	
CH ₂ Cl ₂	DABCO (5.0)	40	2	nr	
CH ₂ Cl ₂	DMAP (2.0)	40	12	nr	
	DBU (5.5)	20	0.2	83	28
CH ₂ Cl ₂	DBU (4.0)	0	5.5	68	46
THF	DBU (1.2)	0	3.0	77–85	75–80

^a nr, no reaction.

Scheme 21




Methyl Iodide and DBU. Stupp et al. reported a convenient procedure to transform methiodide derivatives to nitriles under mildly basic conditions.³⁷ Furthermore, the application of this procedure to the synthesis of nitriles with absolute stereocontrol was demonstrated. To convert diastereomerically enriched hydrazone **67** (de = 80–82%) to the corresponding chiral nitrile **68** without racemization (Scheme 20), the reaction conditions were optimized as depicted in Table 5.

Chloroformates. Aliphatic aldehyde dimethylhydrazones have been shown to yield the parent nitriles by oxidative elimination of the carbomethoxyhydrazone salts (cy = 64–90%).³⁸


Methyl Propiolate and Dialkyl Acetylenedicarboxylate. During attempts to carry out [4 + 2] cycloaddition reactions with indole-3-carboxaldehyde *N,N*-dimethylhydrazone **69** with methyl propiolate or dimethyl acetylenedicarboxylate **70**, unexpectedly indole-3-carbonitriles **72** were formed (Scheme 21).³⁹

The proposed mechanism proceeds via aza-Michael

Scheme 22



Scheme 23



addition of the hydrazone dimethylamino group at methyl propiolate to form hydrazone **71** with subsequent oxidative elimination to the nitrile **72**.⁴⁰

Hydrolytic Cleavage Procedures


CuCl_2 and $\text{Cu}(\text{OAc})_2$. The Cu(II)-mediated hydrolysis is applicable to ketone and aldehyde DMHs.⁴¹ In the latter case, the formation of nitriles is not a significant side reaction. The hydrolysis is irreversible, since the resulting hydrazine is oxidized to the diazene by Cu(II).⁴²

Treatment of α -branched ketone SAMP hydrazones **3** with an aqueous CuCl_2 solution affords the parent ketones **4** in highly enantiomerically pure form (*ee* = 89–99%) and in moderate to excellent yields (52–94%, Scheme 22).⁴³ The cleavage conditions do not affect functionalities sensitive to oxidation or strong acids. For example, amines, alkenes, and thioethers as well as acid-labile protecting groups such as acetals and silyl ethers are compatible with this procedure. Compounds with these functionalities usually give low yields when subjected to other typical procedures, such as ozonolysis and the salt method. The alkylation/cleavage protocol can be carried out in one pot and affords the products without racemization. In addition, the Cu(II) method avoids the formation of nitrosamine byproducts. In contrast to the case with aldehyde DMHs, this method is not applicable to the cleavage of aldehyde SAMP hydrazones.


The broad applicability of this method has been demonstrated in the total synthesis of several complex natural products, such as the sesquiterpene lactone (+)-tuberiferine **73**⁴⁴ by Fraser-Reid et al. (Scheme 23).⁴⁵

In the total synthesis of complex pentacyclic marine diterpenoid (–)-stypoldion (**74**), characterized by a ben-

Scheme 24




Scheme 25



zoquinone furan moiety, the quaternary stereogenic center was introduced with excellent diastereoselectivity by direction of the neighboring dimethylhydrazone moiety followed by deprotection of the DMH **75** with CuCl_2 (Scheme 24).⁴⁶

Oxalic Acid. Cleavage of ketone SAMP hydrazones **3** with saturated aqueous oxalic acid solution gives the corresponding ketones **4** in excellent yields (84–98%) and with high enantiomeric purity (*ee* = 90–99%, Scheme 25).⁴⁷

This racemization-free cleavage procedure is compatible with functionalities that are sensitive to oxidative cleavage conditions, such as $\text{C}=\text{C}$ double bonds, or to strong acids, such as ketal groups. SAMP can be easily recovered from the corresponding oxalate salt **76** (*cy* = 80–90%) without any detectable racemization of the auxiliary. The SAMP hydrazones are hydrolyzed by aqueous oxalic acid in a short time. The acid-sensitive acetonide group was not hydrolyzed, and racemization was avoided by performing the reaction in a two-phase system



(water/ether or pentane). Usually, the ketone products **4** do not require chromatographic purification. This method is therefore suitable for large-scale reactions. No carcinogenic nitrosamine byproduct is formed as is the case in the ozonolysis method, and no toxic methylating agents are needed as in the salt method (*vide infra*). Oxalic acid has turned out to be the most versatile reagent of those our group has studied for the cleavage of ketone SAMP hydrazones **3**. Unfortunately, this method is not compatible for the hydrolysis of aldehyde SAMP hydrazones.

(NH₄)H₂PO₄. For the cleavage of dioxanone DMH **77** containing the acid-labile acetal moiety, conditions had to be carefully selected. Treatment with a saturated aqueous ammonium dihydrogen phosphate buffer at ambient temperature smoothly converted **77** into the ketone **78** in good yield (Scheme 26).⁴⁸





Methyl Iodide/HCl (Salt Method). According to Levisalles et al., DMHs are hydrolyzed under much milder conditions if they are first transformed to their corresponding trimethylhydrazonium iodides (MeI, EtOH, H₂O).⁴⁹ For the conversion of SAMP/RAMP hydrazones **3** of aldehydes and ketones to optically active carbonyl compounds **4**, a two-phase variant was developed.

Treatment of **3** with excess methyl iodide at 60 °C leads quantitatively to a mixture of the methiodides **79A** and **79B** (Scheme 27), which are hydrolyzed without further purification in a two-phase system (3–4 N HCl, *n*-pentane) to afford the parent carbonyl compounds **4** in good to excellent yields and in short reaction times (15–60 min).

This procedure also occurs without any racemization. Obviously the carbonyl compounds **4** are rapidly transferred with vigorous stirring into the pentane phase, which is free of acid. Chiral aldehydes and ketones bearing a stereogenic center in the position α to the carbonyl moiety are unexpectedly resistant against racemization in an acidic environment. Traces of base, however, lead to rapid and complete racemization.




When the aqueous layer, expected to contain the auxiliary in the form of salts **80**, is neutralized and extracted, SAMP hydrazone **81**, the hydrazine **82**, and SAMP were isolated in a mixture of 1:7:2 (yield = 40–50%). While the ammonium salts **80A** remain in the aqueous phase, trisubstituted hydrazines such as **82** are known to easily undergo air oxidation to formaldehyde hydrazones such as **81**, which itself is hydrolyzed to SAMP. In this way, at least a partial recycling of SAMP or RAMP is possible.

The procedure was employed in the synthesis of several natural products. For example, serricornin **83**, the sex pheromone of the female cigarette beetle, was synthesized in almost enantiomerically pure form (Scheme 28).⁵⁰




Nicolaou et al. have demonstrated the usefulness of the salt method by the total synthesis of epothilones A and B (Scheme 29).⁵¹ Compared to ozonolysis, the MeI/HCl method resulted in higher yields (O₃, 77%; MeI/HCl, 86%).

HCl. In some cases, the dialkylhydrazones are hydrolyzed efficiently even without the formation of methiodides. In the synthesis of RP 66471, a potent potassium channel opener, the SAMP/RAMP methodology was applied to build up the quaternary center α to the carbonyl



1. LDA, THF, TMEDA
2. **87**, $\text{BF}_3 \cdot \text{OEt}_2$, THF, -78°C
3. 6N HCl, Et_2O , rt
4. L-selectride, THF, -78°C




group. By acidic hydrolysis, the key intermediate **84** was obtained (Scheme 30).⁵²

Our group used acidic hydrolysis to synthesize ramulosin **85** and its analogues (Scheme 31).⁵³ After Michael addition of acetone SAMP hydrazones **86** to the unsaturated ester **87**, the auxiliary was removed by stirring in a two-phase system (Et_2O , 6 N HCl).

Silica Gel. The cleavage of hydrazones using acidic reagents suffers from the lack of selectivity in the presence of acid-sensitive groups. Using silica gel for the cleavage of DMHs gives the corresponding ketones containing other acid-sensitive functionalities such as THP groups, benzyl ether moieties, and acetal groups, which remain intact.⁵⁴ However, this method seems to be less versatile for the cleavage of α -branched ketones.

BiCl_3 and Microwave Irradiation. Hydrolytic cleavage of the C=N bond of DMHs by the use of catalytic quantities of BiCl_3 in wet THF under microwave irradiation yields the corresponding ketones or aromatic aldehydes.⁵⁵ The reaction proceeds efficiently in high yields (75–98%) at ambient pressure within a few minutes. For example, benzophenone DMH was cleaved in 2 min in 98% yield.



The corresponding control reaction, performed by refluxing for 8 h, produced only a 50% yield.

$\text{Pd}(\text{OAc})_2$ and SnCl_2 . Ketone DMHs undergo cleavage to the parent ketones by a catalytic amount of $\text{Pd}(\text{OAc})_2/\text{SnCl}_2$ in moderate to good yields.⁵⁶ The reaction proceeds without using acidic or oxidative reagents. Even α,β -unsaturated ketone DMHs can be hydrolyzed without rearrangement using this catalyst system. Functionalities such as halogen or nitro groups are also inert.


$\text{BF}_3 \cdot \text{OEt}_2$. $\text{BF}_3 \cdot \text{OEt}_2$ is a suitable Lewis acid to promote the hydrolysis of DMHs,⁵⁷ which is also applicable to the racemization-free cleavage of SAMP hydrazones of α -branched 2-oxoesters by stirring in acetone/water under addition of paraformaldehyde in an exchange reaction. Compared to the ozonolytic cleavage (yields = 80–90%), $\text{BF}_3 \cdot \text{OEt}_2$ gave higher yields (81–97%), but with longer reaction times.⁵⁸ Direct dithioketalization of *N,N*-dimethylhydrazones or SAMP hydrazones **88** of aliphatic, aromatic, and α,β -unsaturated aldehydes promoted by $\text{BF}_3 \cdot \text{OEt}_2$ or *p*-TsOH and 1,2-ethanedithiol in dry medium affords the corresponding dithiolanes **89** in nearly quantitative yields (Scheme 32).⁵⁹

Baker's Yeast. Ketones and aromatic aldehydes have been regenerated biocatalytically from their corresponding phenyl- and *N,N*-dimethylhydrazones in quantitative yield on incubating with baker's yeast.⁶⁰ In the case of the DMHs, no cleavage of α -branched carbonyl compounds has been reported.

Reductive Cleavage Procedures

Reductive cleavage procedures have been used in several cases to regenerate carbonyl compounds from their derivatives, such as oximes, 2,4-dinitrophenylhydrazones, tosylhydrazones, etc. However, in the case of dialkylhydrazone cleavage, very few studies have been carried out employing reductive methods. We were interested in recovering ferrocenyl ketones from their corresponding SAMP hydrazones by a reductive manner, as these compounds are often sensitive to oxidative and acidic reaction conditions. It could be demonstrated that some of these methods, known for related compounds, are also applicable to the cleavage of SAMP hydrazones.

TiCl_3 . Titanium(III) is well known for its ability to cleave the N–O bond of oximes and nitro compounds as well as the S–O bond of sulfoxides. Therefore, the cleavage of the N–N hydrazone bond was considered to be analogous. Trivalent titanium has been used frequently for deoxygenation procedures⁶¹ and for the cleavage of tosyl-⁶² and 2,4-dinitrophenylhydrazones.⁶³ The resulting imines generated from reduction are easily hydrolyzed to the carbonyl compounds. This method has also been applied



to the cleavage of ferrocenyl ketone SAMP hydrazones **90** (Scheme 33).⁶⁴

In some cases, the cleavage was accompanied by a small degree of racemization. Cp–Si bonds are cleaved under the reaction conditions, and hydroxy groups α to the ferrocenyl backbone are removed to form the corresponding alkyl group. Compounds containing Ph₂P in the *ortho* position do not yield the ketone **91**, but the corresponding methylene compound is formed by overreduction. This means that, for this special case, its applicability is limited to only a few functional groups.

SnCl₂. SnCl₂ has been used to recover carbonyl compounds from their 2,4-dinitrophenylhydrazones.⁶⁵ This method has been applied to cleave ferrocenyl ketone SAMP hydrazones **90** in good yields (70–85%).⁶⁴ The procedure tolerates many functionalities in the *ortho* position, such as alkyl, hydroxyalkyl, or halogen groups; however, the Cp–Si bonds are cleaved.

Cr(OAc)₂ and VCl₂. Cr(OAc)₂ effects reductive deoxygenation reactions which are highly efficient, even in the presence of acid- and base-sensitive functional groups such as ketals, hemiketals, esters, and epoxides.⁶⁶ The selection of the reagent was based on the premise that it would cause reductive fission of the oxime N–O linkage to give an imine which would suffer rapid hydrolysis to ketone at pH 5. Cr(OAc)₂ is also able to cleave ferrocenyl ketone SAMP hydrazones in good to excellent yields (65–99%), tolerating a thiophene moiety as R. An advantage is that the reaction products are obtained very pure by simple filtration.

VCl₂ is also an efficient deoxygenation reagent.⁶⁷ In comparison to Cr(II), for the cleavage of ferrocenyl ketone SAMP hydrazones **90**, V(II) gives lower yields (ca. 70%) but also very pure products.^{64b}

Conclusion

Many reagents have been developed to regenerate carbonyl compounds from the corresponding dialkylhydrazones which are compatible with a wide range of functionalities. This has allowed the use of hydrazones in the total synthesis of complex natural products. However, owing to the importance of hydrazone methodology, there is still an interest in designing new and mild reagents for hydrazone cleavage.

References

- (1) (a) Enders, D. Alkylation of Chiral Hydrazones. In *Asymmetric Synthesis*; Morrison, J. D., Ed.; Academic Press: Orlando, FL, 1984; Vol. 3, pp 275–339.
- (2) Zakharkin, L. I.; Khorlina, I. M. Preparation of Aldehydes by Reduction of Nitriles with Diisobutylaluminumhydride. *Dokl. Akad. Nauk. S.S.S.R.* **1957**, *116*, 422–424; *Chem. Abstr.* **1958**, *52*, 8040f.
- (3) (a) Enders, D.; Eichenauer, H. Asymmetrische Synthesen via metallierte chirale Hydrazone. Enantioselective Alkylierung von cyclischen Ketonen und Aldehyden. *Chem. Ber.* **1979**, *112*, 2933–2960.
- (4) Enders, D.; Schäfer, T.; Mies, W. Enantioselective Synthesis of α -Sulfonylated Ketones and Aldehydes via α -Thiolation of Metalated SAMP/RAMP-Hydrazones. *Tetrahedron* **1998**, *54*, 10239–10252.
- (5) Enders, D.; Joseph, R.; Poesz, C. Electrophilic Amination of Hydrazones: A New Synthetic Route to Protected α -Hydrazino and α -Aminoketones. *Tetrahedron* **1998**, *54*, 10069–10078.
- (6) Enders, D.; Berg, T.; Raabe, G.; Runsink, J. Enantioselective Synthesis of α -Phosphino Ketones and 2-Phosphino-Alcohols. *Liebigs Ann./Recl.* **1997**, 345–363.
- (7) (a) Erickson, R. E.; Andrulis, P. J., Jr.; Collins, J. C.; Lungle, M. L.; Merger, G. D. Mechanism of Ozonation Reaction. Carbon–Nitrogen Double Bonds. *J. Org. Chem.* **1969**, *34*, 2961–2966.
- (8) Nicolaou, K. C.; Patron, A. P.; Ajito, K.; Richter, P. K.; Khatuya, H.; Bertinato, P.; Miller, R. A.; Tomaszewski, M. J. Total Synthesis of Swinholide A, Preswinholide A, and Hemiswinholide A. *Chem. Eur. J.* **1996**, *2*, 847–868.
- (9) Burton, J. W.; Clark, J. S.; Derrer, S.; Stork, T. C.; Bendall, J. G.; Holmes, A. B. Synthesis of Medium Ring Ethers. The Synthesis of (+)-Laurencin. *J. Am. Chem. Soc.* **1997**, *119*, 7483–7498.
- (10) Schwaebé, M.; Little, R. D. Asymmetric Reductive Cyclization. Total Synthesis of (–)-C₁₀-Desmethyl Arteannuin B. *J. Org. Chem.* **1996**, *61*, 3240–3244.
- (11) Nicolaou, K. C.; Yue, E. W.; La Greca, S.; Nadin, A.; Yang, Z.; Leresche, J. E.; Tsurii, T.; Naniwa, Y.; De Riccardis, F. Synthesis of Zaragozaic Acid A/Squalestatin S1. *Chem. Eur. J.* **1995**, *1*, 467–494.
- (12) (a) Friedrich, E.; Lutz, W.; Eichenauer, H.; Enders, D. Mild Cleavage of *N,N*-Dimethylhydrazones to Carbonyl Compounds with Singlet Oxygen. *Synthesis* **1977**, 893–894. (b) Eichenauer, H.; Friedrich, E.; Lutz, W.; Enders, D. Regiospezifische und Enantioselective Aldol-Reaktionen. *Angew. Chem.* **1978**, *90*, 219–220; *Angew. Chem., Int. Ed. Engl.* **1978**, *17*, 206–207.
- (13) Panek, J. S.; Beresis, R. T.; Celatka, C. A. Studies Directed toward the Synthesis of Ulapualide A. Asymmetric Synthesis of the C26–C42 Fragment. *J. Org. Chem.* **1996**, *61*, 6494–6495.
- (14) (a) Enders, D.; Bhushan, V. Mild and Efficient Oxidative Cleavage of *N,N*-Dialkylhydrazones to Ketones with Perborate. *Z. Naturforsch.* **1987**, *42b*, 1595–1596. (b) McKillop, A.; Tarbin, J. A. Functional Group Oxidation Using Sodium Perborate. *Tetrahedron* **1987**, *43*, 1753–1758.
- (15) Choi, H. C.; Kim, Y. A. Facile Cleavage of *N,N*-Dimethylhydrazones to Ketones using Tetrabutylammonium Peroxydisulfate. *Synth. Commun.* **1994**, *24*, 2307–2311.
- (16) Barton, D. H. R.; Jaszberenyi, J. C.; Liu, W.; Shinada, T. Oxidation of Hydrazones by Hypervalent Organoiodine Reagents: Regeneration of the Carbonyl Group and Facile Syntheses of α -Acetoxy and α -Alkoxy Azo Compounds. *Tetrahedron* **1996**, *52*, 14673–14688.
- (17) Brougham, P.; Cooper, M. S.; Cummerson, D. A.; Heaney, H.; Thompson, N. Oxidation Reactions Using Magnesium Monoperoxyphthalate: A Comparison with *m*-Chloroperoxybenzoic Acid. *Synthesis* **1987**, 1015–1017.
- (18) Fernández, R.; Gasch, C.; Lassaletta, J. M.; Llera, J. M.; Vázquez, J. Simple and Efficient Conversion of *N,N*-Dimethylhydrazones and Aldehydes to Nitriles. *Tetrahedron Lett.* **1993**, *34*, 141–144.
- (19) Enders, D.; Plant, A. Oxidative Cleavage of *N,N*-Dialkylhydrazones to Ketones with Magnesium Monoperoxyphthalate. *Synlett* **1990**, 725–726.
- (20) Enders, D.; Plant, A. Enantioselective Synthesis of α -Substituted Nitriles by Oxidative Cleavage of Aldehyde SAMP-Hydrazones with Magnesium Monoperoxyphthalate. *Synlett* **1994**, 1054–1056.
- (21) Nicolaou, K. C.; Sarabia, F.; Ninkovic, S.; Ray, M.; Finlay, V.; Boddy, C. N. C. Variation der Ringgröße von Epithilonen—Totalsynthese von [14]-, [15]-, [17]-, und [18] Epithilon A. *Angew. Chem.* **1998**, *110*, 85–89; *Angew. Chem., Int. Ed.* **1998**, *37*, 81–84.
- (22) Enders, D.; Geibel, G.; Osborne, S. Diastereo- and Enantioselective Total Synthesis of Stigmatellin A. *Chem. Eur. J.* **2000**, in press.
- (23) Said, S. B.; Skarzewski, J.; Mlochowski, J. Conversion of Aldehydes into Nitriles via Oxidation of their Dimethylhydrazones. *Synthesis* **1989**, 223–224.
- (24) Duraisamy, M.; Walborsky, H. M. A Convenient Method for Hydrazone Hydrolysis. *J. Org. Chem.* **1984**, *49*, 3410–3411.
- (25) Horner, L.; Fernekess, H. Präparative Ergebnisse der Umsetzung von Hydrazinderivaten und Hydrazonen mit Peressigsäure. *Chem. Ber.* **1961**, *94*, 712–724.
- (26) Stupp, S. I.; Son, S.; Li, L. S.; Lin, H. C.; Keser, M. Bulk Synthesis of Two-Dimensional Polymers: The Molecular Recognition Approach. *J. Am. Chem. Soc.* **1995**, *117*, 5212–5227.

- (27) Smith, R. F.; Albright, J. A.; Waring, A. M. Oxidation of Aldehyde *N,N*-Dialkylhydrazones with Hydrogen Peroxide. A New Nitrile Synthesis. *J. Org. Chem.* **1966**, *31*, 4100–4102.
- (28) Eichenauer, H.; Friedrich, E.; Lutz, W.; Enders, D. Regiospezifische und Enantioselektive Aldol-Reaktionen. *Angew. Chem.* **1978**, *90* 219–220; *Angew. Chem., Int. Ed. Engl.* **1978**, *17*, 206–207.
- (29) (a) Altamura, A.; Curci, R.; Edwards, J. O. On the Facile Regeneration of Carbonyl Compounds by Oxidative Cleavage of Hydrazones using Dioxiranes. *J. Org. Chem.* **1993**, *58*, 7289–7293. (b) Altamura, A.; D'Accolti, L.; Detomaso, A.; Dinoi, A.; Fiorentino, M.; Fusco, C.; Curci, R. High-yield Synthesis of Nitriles by Oxidation of Aldehyde *N,N*-Dimethylhydrazones with Dimethyldioxirane. *Tetrahedron Lett.* **1998**, *39*, 2009–2012.
- (30) Olah, G. A.; Welch, J.; Prakash, G. K. S.; Ho, T. L. Oxidative Cleavage of Hydrazones with Molybdenyl Chloride (MoOCl_3) or Molybdenum Hexafluoride and Reductive Cleavage of Oximes with Molybdenyl Chloride/Zinc. *Synthesis* **1976**, 808–809.
- (31) Olah, G. A.; Welch, J.; Henninger, M. The Oxidative Cleavage of Hydrazones and Oximes with Cobalt(III) Trifluoride. *Synthesis* **1977**, 308–309.
- (32) Olah, G. A.; Prakash, G. K. S.; Ho, T. L. Tungsten Hexafluoride, a Mild Oxidizing Agent for the Cleavage of Hydrazones. *Synthesis* **1976**, 809–810.
- (33) Olah, G. A.; Welch, J.; Ho, T. L. Uranium Hexafluoride, a Convenient New Oxidizing Agent for Organic Synthesis. *J. Am. Chem. Soc.* **1976**, *98*, 6717–6718.
- (34) Olah, G. A.; Ho, T. L. Oxidative Cleavage of Oximes and Dimethylhydrazones with Nitronium and Nitrosonium Salts. *Synthesis* **1976**, 610–611.
- (35) Laszlo, P.; Polla, E. Conversion of *N,N*-Dimethylhydrazones to Carbonyl Compounds by Clay-Supported Ferric Nitrate. *Tetrahedron Lett.* **1984**, *25*, 3309–3312.
- (36) Rudler, H.; Denise, B. MTO Catalyzed Oxidation of Aldehyde *N,N*-Dimethylhydrazones with Hydrogenperoxide: High Yield Formation of Nitriles and *N*-Methylene-*N*-Methyl *N*-Oxide. *J. Chem. Soc., Chem. Commun.* **1998**, 2145–2146.
- (37) Moore, J. S.; Stupp, S. I. Cleavage of Aldehyde Hydrazone Iodides under Mild Conditions. A Convenient Route to Chiral Nitriles of High Enantiomeric Purity. *J. Org. Chem.* **1990**, *55*, 3377–3381.
- (38) Xiao, Z.; Timberlake, J. W. An Elimination Reaction of *N*-Carbomethoxy-*N,N*-dimethylhydrazone Salts to Alkyl Nitriles. *Tetrahedron* **1998**, *54*, 12715–12720.
- (39) (a) Moscardó, S. E.; Sepúlveda-Arques, J.; Abarca Gonzalez, B.; Ballesteros Campos, R.; Soriano Soto, C.; García Granda, S.; Gomez Beltran, F. Reactions of 3-Formylindole *N,N*-Dimethylhydrazones with Dienophiles. A Stereospecific Addition. *Heterocycles* **1992**, *34*, 61–66. (b) Biswas, G. K.; Nath, A. C.; Mukherjee, B.; Patra, A.; Chakrabarty, M. On Attempted *Diels-Alder* Reaction of 1-Ethoxycarbonylindole-3-carboxaldehyde *N,N*-Dimethylhydrazone. *Tetrahedron Lett.* **1992**, *33*, 117–118.
- (40) Gonzales, M. E.; Sancho, P.; Soriano, C.; Ballesteros, R.; Abarca, B.; Sepulveda, J. The Reactivity of 2-Formylthiophene, 1-Methyl-2-formylpyrrole, Benzaldehyde and 2-Formylfuran *N,N*-Dimethylhydrazones with Dimethyl Acetylenedicarboxylate. *Heterocycles* **1988**, *27*, 1227–1231.
- (41) Corey, E. J.; Knapp, S. Facile Conversion of *N,N*-Dimethylhydrazones to Carbonyl Compounds by Cupric Ion-Catalyzed Hydrolysis. *Tetrahedron Lett.* **1976**, 3667–3668.
- (42) Petredis, D.; Burke, A.; Balch, A. L. Cuprous Complexes of Phenyl diazene. *J. Am. Chem. Soc.* **1970**, *92*, 428–429.
- (43) Enders, D.; Hundertmark, T.; Lazny, R. Copper(II) Chloride Mediated Racemization-Free Hydrolysis of α -Alkylated Ketone SAMP-Hydrazones. *Synth. Commun.* **1999**, *28*, 27–33.
- (44) Okamura, H.; Yamauchi, K.; Miyawaki, K.; Iwagawa, T.; Nakatani, M. Synthesis and Biological Activities of Degraded Limonoids, (\pm)-Fraxinellonone and Its Related Compounds. *Tetrahedron Lett.* **1997**, *38*, 263–266.
- (45) Georges, M.; Fraser-Reid, B. Controlled Access to Furanose Precursors Related to Sesquiterpene Lactones. *J. Org. Chem.* **1985**, *50*, 5754–5758.
- (46) Falck, J. R.; Chandrasekhar, S.; Manna, S.; Chiu, C. C. S.; Mioskowski, C.; Wetzel, I. Total Synthesis of the Spiro-*o*-benzoquinonefuranone (–)-Stypoldione. *J. Am. Chem. Soc.* **1993**, *115*, 11606–11607.
- (47) Enders, D.; Hundertmark, T.; Lazny, R. Mild Racemization Free Cleavage of Ketone SAMP-Hydrazones with Oxalic Acid—Recycling of the Chiral Auxiliary. *Synlett* **1998**, 721–722.
- (48) Ulven, T.; Sørbye, K. A.; Carlsen, P. H. J. Synthesis of Racemic 3,5-*O*-Benzylidene-2-deoxypentoses from 2-Phenyl-1,3-dioxan-5-one Hydrate. *Acta Chem. Scand.* **1997**, *51*, 1041–1044.
- (49) Avaro, M.; Levisalles, J.; Rudler, H. *N,N*-Dimethylhydrazones as Protecting Groups in Synthesis. *J. Chem. Soc., Chem. Commun.* **1969**, 445–446.
- (50) Mori, K.; Niomi, H. Synthesis and Absolute Stereochemistry of Serricornin [(4*S*, 6*S*, 7*S*)-4,6-Dimethyl-7-hydroxy-3-nonanone]. *Tetrahedron* **1982**, *38*, 3705–3711.
- (51) Nicolaou, K. C.; Ninkovic, S.; Sarabia, F.; Vourloumis, D.; He, Y.; Vallberg, H.; Finlay, R. M. V.; Yang, Z. Total Synthesis of Epithilones A and B via a Macrolactonization-Based Strategy. *J. Am. Chem. Soc.* **1997**, *119*, 7974–7991.
- (52) Hart, T. W.; Guillochon, D.; Perrier, G.; Sharp, B. W.; Toft, M. P.; Vacher, B.; Walsh, R. J. A. The Synthesis of RP 66471. A Potent Potassium Channel Opener. *Tetrahedron Lett.* **1992**, *33*, 7211–7214.
- (53) Enders, D.; Kaiser, A. A Short Asymmetric Synthesis of Both Enantiomers of Ramulosin and Its Analogues. *Synthesis* **1996**, 209–214.
- (54) (a) Mitra, R. B.; Bhaskar Reddy, G. Selective Cleavage of Dimethylhydrazones to the Carbonyl Compounds Using Silica Gel and its Application in the Synthesis of (Z)-9-Tetradecenyl Acetate. *Synthesis* **1989**, 694–698. (b) Kotsuki, H.; Miyazaki, A.; Kadota, I.; Ochi, M. A Simple and Efficient Heterogeneous Hydrolysis of *N,N*-Dimethylhydrazone Derivatives. *J. Chem. Soc., Perkin Trans. 1* **1990**, 420–430.
- (55) Bi^{3+} -Catalyzed Regeneration of Carbonyl Compounds from Hydrazones under Microwave Irradiation. *Synlett* **1997**, 1251–1252.
- (56) Miino, T.; Hirota, T.; Yamashita, M. Palladium Catalyzed Hydrolysis of Dimethylhydrazones to Carbonyl Compounds. *Synlett* **1996**, 999–1000.
- (57) Gawley, R. E.; Termine, E. J. Mild Nonoxidative Procedure for the Hydrolysis of Dimethylhydrazones. *Synth. Commun.* **1982**, *12*, 15–18.
- (58) (a) Enders, D.; Dyker, H.; Raabe, G. Enantioselektive Synthese 3-substituierter 2-Ketoester. *Angew. Chem.* **1992**, *104*, 649–651; *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 618–620. (b) Enders, D.; Dyker, H.; Raabe, G.; Runsink, J. Enantio- and Diastereoselective Synthesis of 3-Substituted Cyclic Hemiketals of ω -Hydroxy-2-oxoesters. *Synlett* **1992**, 901–903.
- (59) Diéz, E.; López, A. M.; Pareja, C.; Martín, E.; Fernández, R.; Lassaletta, J. M. Direct Synthesis of Dithioketals from *N,N*-Dialkylhydrazones. *Tetrahedron Lett.* **1998**, *39*, 7955–7958.
- (60) Kamal, A.; Rao, M. V.; Meshram, H. M. Efficient Enzymatic Hydrazone Hydrolysis by Baker's Yeast. *Tetrahedron Lett.* **1991**, *32*, 2657–2658.
- (61) Timms, G. H.; Wildsmith, E. The Reduction of Oximes with Tervalent Titanium. A Mild Deoxygenation Procedure and the Partial Synthesis of Erythromyclamine. *Tetrahedron Lett.* **1971**, 195–198.
- (62) Chandrasekhar, B. P.; Sunthankar, S. V.; Telang, S. G. Cleavage of Steroidal Tosylhydrazones with Titanous Chloride under Mild Conditions. *Chem. Ind.* **1975**, 87–88.
- (63) McMurry, J. E.; Silvestri, M. A Method for Cleaving 2,4-Dinitrophenylhydrazones to Ketones. *J. Org. Chem.* **1975**, *40*, 1502–1504.
- (64) (a) Enders, D.; Peters, R.; Lochtman, R.; Runsink, J. Enantioselektive Synthesis of Planar Chiral *ortho*-Functionalised Ferrocenylketones. *Synlett* **1997**, 1462–1464. (b) Peters, R. Dissertation, RWTH Aachen, 2000.
- (65) Casnati, G. Recovery of Carbonyl Compounds from their dinitrophenylhydrazones. *Chem. Abstr.* **1958**, *52*, 17159.
- (66) Corey, E. J.; Richman, J. E. Reaction of Oxime *O*-Acetates with Chromous Acetate. A Method for the Conversion of Ketoximes to Ketones under Mild Conditions. *J. Am. Chem. Soc.* **1970**, *92*, 5276–5277.
- (67) Olah, G. A.; Arvanaghi, M.; Prakash, G. K. S. Reductive Cleavage of Oximes with Vanadium(II) Chloride. *Synthesis* **1980**, 220.

AR990062Y