

ACCOUNTS of CHEMICAL RESEARCH[®]

AUGUST 2000

Registered in U.S. Patent and Trademark Office; Copyright 2000 by the American Chemical Society

Nickel-Catalyzed Intermolecular Domino Reactions

SHIN-ICHI IKEDA

Faculty of Pharmaceutical Sciences, Nagoya City University,
Tanabe-dori, Mizuho-ku, Nagoya 467-8603, Japan

Received July 12, 1999

ABSTRACT

Nickel-catalyzed multiple-component reactions are a promising new type of domino reaction. For instance, a variety of starting materials such as alkenes, alkynes, and unsaturated carbonyl compounds can be connected with high selectivity under reasonably mild conditions. These reactions provide a new route to synthetically important key compounds including dienes, enynes, enol silyl ethers, and carbocycles. Furthermore, an asymmetric version of the reaction is successful. Thus, intermolecular domino reactions should be important reactions for years to come.


Introduction

Domino reactions are organic reactions where multiple carbon-carbon bond formation is achieved in a single operation without any troublesome experimental procedures. The procedures are advantageous environmentally and economically and should have wide practical application.¹⁻³ In particular, intramolecular domino reactions are powerful methods for constructing complex organic molecules. On the other hand, intermolecular domino reactions with high selectivity are a challenging issue in synthetic organic reactions. A range of strategies involving the sequential generation of radical and anionic species are useful for such intermolecular transformations.^{4,5} We have envisaged a new approach based on a transition-metal-catalyzed reaction.⁶ This Account outlines nickel-


Shin-ichi Ikeda was born in Nagoya, Japan, on Oct 26, 1965. He received his B.S. (1988) and M.S. (1990) degrees from Gifu University, Japan. In 1993 he received his Ph.D. from Osaka University. After obtaining his Ph.D., he was an Assistant Professor at the Nagoya City University, Faculty of Pharmaceutical Sciences and was promoted to an Associate Professor in 1997. His research interests include the use of transition metal complexes as catalytic reagents in organic synthesis.

Scheme 1


inter-intra or intra-intramolecular process



intra-inter or intra-intramolecular process



inter-intermolecular process



M = Pd, Ni, etc.

/ : newly formed C-C bond

catalyzed intermolecular domino reactions developed by our research group.

Regio- and Stereoselective Intermolecular Domino Reaction

The use of a transition-metal-catalyzed reaction as part of a domino process has gained increasing interest. In particular, processes including the Mizoroki-Heck reaction, i.e., palladium-catalyzed arylation and alkenylation of unsaturated carbon molecules, are well-known ex-

Scheme 2






Chart 1



amples (Scheme 1).^{3,6} These transformations proceed via the activation of a carbon unit (A) by a metal catalyst (M), e.g., oxidative addition to a Pd(0) complex, to generate an A–M species. The A–M species then adds to a carbon–carbon double or triple bond to lead to intermediate I or I', which is transformed into II or II' upon capturing another unit (B), e.g., coupling with carbon nucleophiles. In contrast to the domino process involving an intramolecular reaction, it would be difficult to develop a regio- and stereoselective intermolecular domino reaction of an unactivated alkene or alkyne with two independent carbon chains A and B.^{7–9}

As an approach to intermolecular multiple-component domino coupling, we noted that a nickel(0) complex catalyzed the carbonylative coupling of allyl chloride 1 with an unactivated alkyne, carbon monoxide, and an alcohol (Scheme 2).¹⁰ The key step is the preparation of alkenylnickel species 3 via the insertion of an alkyne into (π -allyl)nickel intermediate 2.¹¹ If 3 sequentially reacted with organometallics as carbon nucleophiles, a three-component coupling product would be obtained. Actually, the nickel(0)-catalyzed reaction of 1 with terminal alkynes 4 and alkenyltins 5 gave 6 with high regio- and stereoselectivities (Chart 1).^{12,13} Dimethylzinc (Me₂Zn) and trimethylaluminum (Me₃Al) also reacted with 1 and 4 to give 7 (Chart 2).¹⁴

Chart 2



Scheme 3






Chart 3




When an organophosphorus compound such as triphenylphosphine (PPh₃) was added to the reaction system (1–2 equiv vs Ni(0) catalyst), cross-coupling of 1 and organometallics, rather than three-component coupling, took place preferentially.¹⁵ This result indicates that the coordination of PPh₃ to 2 inhibits the production of 3' (Scheme 3).

Further investigations using enones 8 resulted in the development of a novel domino reaction. Alkynes 4 reacted with 5, 8, and chlorotrimethylsilane (Me₃SiCl) in




nickel(0)-catalyzed cyclization of alkynyl enones with organozincs (eq 4).²⁵





Diastereodifferentiating Domino Reaction with Norbornenes

C_5 -symmetric molecules contain at least two prochiral centers, which can be converted to chiral carbons by appropriate enantiotopic functionalization.²⁶ Torii and co-workers reported that a Pd(0) complex catalyzed an asymmetric desymmetrization of norbornene (eq 5).²⁷ Use




of a chiral-center-containing alkenyl iodide resulted in the differentiation of prochiral positions of the norbornene to create two chiral centers at C2 and C3. The combination of norbornene and a cyclic enone such as **8d**, which contains a prochiral face, should create a new domino process which is diastereodifferentiating. When **8d** was reacted with norbornene, an alkynyltin **5b**, and Me_3SiCl in the presence of nickel(0) species and a ligand **16**,²⁸ **18a** was obtained as a sole product after hydrolysis (Scheme 5).²⁹




The diastereodifferentiation can be explained by the pathway outlined in Scheme 6. Initially, a mixture of both intermediates **19a** and **19b** is formed under kinetic control. The subsequent thermodynamic equilibrium

Scheme 7



Scheme 8



between **19a** and **19b** would control the coupling of one enantiotopic face of **8d** with the one enantiotopic position in norbornene. The reaction using **5b** occurred in THF under refluxing conditions. At that point, the equilibrium lies so far toward thermodynamically favored **19a** that it leads to the formation of **17a**.³⁰

Enantioselective Intermolecular Domino Reaction

The introduction of a catalytic reaction with asymmetric induction to a domino process is also an important theme in recent synthetic studies. Overman et al. reported the sequential cyclization of a polyene by an asymmetric Heck reaction (Scheme 7).^{31,32} Enantioselection occurs in the first addition of the alkenylpalladium species to the alkenyl unit. The total asymmetric synthesis of (+)-xestoquinone by Key et al. has demonstrated the utility of this reaction.³³ Shibasaki et al. have provided an example of the process involving Suzuki coupling, i.e., palladium-catalyzed cross-coupling of organic electrophiles with organoboranes,³⁴ and the asymmetric Heck reaction (Scheme 8).³⁵ They also demonstrated the utility of an asymmetric Heck reaction–nucleophile (acetate, amine, and carbanion) capture process using an alkenyl triflate (Scheme 9).^{36,37}

We have found a nickel-catalyzed asymmetric multiple-component coupling using Me_2Zn (see Chart 5).³⁸ Inter-

Scheme 9







Chart 6





estingly, catalytic systems involving simple monodentate chiral oxazolines such as **20** (Chart 6),³⁹ which have been previously used as valuable chiral auxiliaries,⁴⁰ were very effective in giving optically active **12**. The results are summarized in eqs 6 and 7. In the reaction with an



internal alkyne such as 3-hexyne (eq 7), when a solution of **8g** and Me_3SiCl was slowly added to the reaction mixture including the catalyst, the corresponding **12e** was obtained in 81% ee (enantiomeric excess). In contrast, some optically active organophosphorus ligands such as BINAP⁴¹ and MOP,⁴² and bidentate oxazolines such as **21**–

Scheme 10

intra- or partially intermolecular cyclotrimerization*completely intermolecular cyclotrimerization*

three different starting materials

chemo- and regioselective cycloadduct

M = Co, Ni, Ru, Rh, Pd, etc. / : newly formed C–C bond

23,^{40,43,44} which have been extensively used as ligands for transition metals in asymmetric catalysis, did not give enantioselection.




Regioselective Three-Component Cyclotrimerization

Transition-metal-catalyzed cyclotrimerization (or [2 + 2] cycloaddition) leads to six-membered cyclic compounds with the formation of three new carbon–carbon bonds. Since the first example was reported by Reppe,⁴⁵ these reactions have been extensively studied by various research groups.^{46,47} Whereas the intramolecular and partially intermolecular modes of the reaction have been used as efficient synthetic methods,⁴⁸ there is no known example of the completely intermolecular mode. The catalytic reaction usually leads to a complex mixture of cycloadducts, which severely limits its utility (Scheme 10).⁴⁹

During our survey for domino coupling with alkynes **4** and enones **8**, we found that a binary metal system of nickel(0) and aluminum catalyzed a regioselective cyclotrimerization of one molecule of **8** and two molecules of **4** (Chart 7).^{50,51} In this reaction, the $\text{Me}_n\text{Al}(\text{OPh})_{3-n}$ ($n = 0-2$) cocatalyst, which was generated from the reaction with Me_3Al and phenol in situ,⁵² functions as a Lewis acid and activates enone **8**. The regiochemistries of the resulting cycloadducts **24** were determined by spectral analyses of


the corresponding aromatic compounds **25** and **26** produced upon subsequent treatment with 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) in air. Interestingly, the regioselectivities in the formation of the products are dependent on the alkyne **4** used in the cycloaddition. Thus, the reaction with **4a** or **4e** tends to selectively give isomer **25**, while another isomer **26** is derived from the reaction with (trimethylsilyl)acetylene (**4g**) or *tert*-butylacetylene (**4i**).

Finally, we have developed the first catalytic cyclotrimerization of three different unsaturated molecules.⁵³ A three-component cycloadduct **27a**, in which the Me_3Si group is adjacent to the carbonyl group, was predominantly obtained when **4e** was added dropwise to a mixture of **8d** and **4g** in the presence of the nickel and aluminum catalytic system (eq 8). In this procedure, the production of undesired cycloadducts **25a**, **26c**, and **28** was suppressed. A biaryl product **27b** was synthesized from the reaction with an aryl-substituted internal alkyne (eq 9).




As a related example of a nickel-catalyzed and Lewis acid-assisted reaction, it has been shown that the cycloaddition of diynes and enones **8** proceeded in the presence of a combination of nickel and zinc.⁵⁴ In this reaction, trimethylsilyl-substituted diynes reacted with an equimolar amount of **8b** to give completely regiochemical-controlled cycloadducts **29**, in which the tri-

Chart 7



methylsilyl group is adjacent to the carbonyl group (eqs 10 and 11).⁵⁵



Conclusion

Intramolecular domino transformations are useful for synthesizing complex compounds. On the other hand, the development of synthetically useful intermolecular multiple-component domino reactions is in its infancy. The nickel-catalyzed reactions summarized in this Account are among the practical intermolecular multiple-component reactions that have been developed to date and should find wide application in synthesis.⁵⁶

The author is deeply grateful to Professor Emeritus Yoshiro Sato (Faculty of Pharmaceutical Sciences, Nagoya City University) for

his kind discussion and to co-workers for their efforts. The author was supported by Grants-in Aid for Scientific Research on Priority Areas (A) (No. 283, "Innovative Synthetic Reactions") and Encouragement of Young Scientists from the Ministry of Education, Science, Sports and Culture, Government of Japan (Monbu-shō) and the Japan Society for the Promotion of Science, and by a Sasakawa Scientific Research Grant from The Japan Science Society.

References

- Hall, N. Chemists Clean Up Synthesis with One-Pot Reactions. *Science* **1994**, *266*, 32.
- (a) Ho, T.-L. *Tandem Organic Reactions*; Wiley-Interscience: New York, 1992. (b) Ho, T.-L. *Tactics of Organic Synthesis*; Wiley-Interscience: New York, 1994; p 79.
- (a) Tietze, L. F.; Belfuss, U. Sequential Transformations in Organic Chemistry: A Synthetic Strategy with a Future. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 131. (b) Bunce, R. A. Recent Advances in the Use of Tandem Reactions for Organic Synthesis. *Tetrahedron* **1995**, *51*, 13103. (c) Tietze, L. F. Domino Reactions in Organic Synthesis. *Chem. Rev.* **1996**, *96*, 115.
- Ryu, I.; Yamazaki, H.; Ogawa, A.; Kambe, N.; Sonoda, N. Four Carbon Component Coupling Reaction Based on Free-Radical Carbonylation: An Easy Access to β -Functionalized δ,ϵ -Unsaturated Ketones. *J. Am. Chem. Soc.* **1993**, *115*, 1187.
- (a) Shono, T.; Nishiguchi, I.; Sasaki, M. Zinc-Promoted One-Step Joining Reaction of Alkyl Halides, Activated Olefins, and Carbonyl Compounds. *J. Am. Chem. Soc.* **1978**, *100*, 4314. (b) Takai, K.; Ueda, T.; Ikeda, N.; Moriwake, T. Sequential Generation and Utilization of Radical and Anionic Species with a Novel Manganese-Lead Reducing Agent. Three-Component Coupling Reactions of Alkyl Iodides, Electron-Deficient Olefins, and Carbonyl Compounds. *J. Org. Chem.* **1996**, *61*, 7990. (c) Takai, K.; Matsukawa, N.; Takahashi, A.; Fujii, T. Three-Component Coupling Reactions of Alkyl Iodides, 1,3-Dienes, and Carbonyl Compounds by Sequential Generation of Radical and Anionic Species with CrCl_2 . *Angew. Chem., Int. Ed. Engl.* **1998**, *37*, 152. (d) Terao, J.; Saito, K.; Nii, S.; Kambe, N.; Sonoda, N. Regioselective Double Alkylation of Styrenes with Alkyl Halides Using a Titanocene Catalyst. *J. Am. Chem. Soc.* **1998**, *120*, 11822.
- (a) Oppolzer, W. Intramolecular, Stoichiometric (Li, Mg, Zn) and Catalytic (Ni, Pd, Pt) Metallo-Ene Reactions in Organic Synthesis. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 38. (b) Negishi, E. Zipper-Mode Cascade Carbometalation for Construction of Polycyclic Structures. *Pure Appl. Chem.* **1992**, *64*, 323. (c) Overman, L. E.; Abelman, M. M.; Kucera, D. J.; Tran, V. D.; Ricca, D. J. Palladium-Catalyzed Polyene Cyclizations. *Pure Appl. Chem.* **1992**, *64*, 1813. (d) Grigg, R. Heterocycle Synthesis by Palladium Catalyzed Cyclisation-Anion Capture Processes. A Power New Strategy. *J. Heterocycl. Chem.* **1994**, *31*, 631. (e) de Meijere, A.; Meyer, F. E. Fine Feathers Make Fine Birds: The Heck Reaction in Modern Garb. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 2379. (f) Negishi, E.; Copéret, C.; Ma, S.; Liou, S.-Y.; Liu, F. Cyclic Carbopalladation. A Versatile Synthetic Methodology for the Construction of Cyclic Organic Compounds. *Chem. Rev.* **1996**, *96*, 365. (g) Ojima, I.; Tzamaroudaki, M.; Li, Z.; Donovan, R. J. Transition Metal-Catalyzed Carbocyclizations in Organic Synthesis. *Chem. Rev.* **1996**, *96*, 635. (h) Grigg, R.; Sridharan. Palladium catalyzed cascade cyclization-anion capture, relay switches and molecular queues. *J. Organomet. Chem.* **1999**, *576*, 65. (i) de Meijere, A.; Brase, S. Palladium in action: domino coupling and allylic substitution reactions for the efficient construction of complex organic molecules. *J. Organomet. Chem.* **1999**, *576*, 88.
- For vicinal difunctionalization of activated alkenes, see: Chapdelaine, M. J.; Hulce, M. Tandem Vicinal Difunctionalization: β -Addition to α,β -Unsaturated Carbonyl Substrates Followed by α -Functionalization. *Org. React.* **1990**, *38*, 225.
- For vicinal difunctionalization of norbornenes, see: (a) Catellani, M.; Chiusoli, G. P. One-Pot Palladium-Catalyzed Synthesis of 2,3-Disubstituted Bicyclo[2.2.1]heptanes and Bicyclo[2.2.1]hept-5-enes. *Tetrahedron Lett* **1982**, *23*, 4517. (b) Catellani, M.; Chiusoli, G. P.; Mari, A. Palladium- or Nickel-Catalyzed Sequential Reaction of Organic Bromides, Bicyclo[2.2.1]hept-2-ene or Bicyclo[2.2.1]hepta-2,5-diene and Alkynes. *J. Organomet. Chem.* **1984**, *275*, 129. (c) Catellani, M.; Chiusoli, G. P.; Concari, S. A New Palladium Catalyzed Synthesis of *cis,Exo*-2,3-Diarylsusbstituted Bicyclo[2.2.1]heptanes or Bicyclo[2.2.1]hept-2-enes. *Tetrahedron* **1989**, *45*, 5263. (d) Kosugi, M.; Tamura, H.; Sano, H.; Migita, T. Palladium-Catalyzed Reactions of Organic Halides with Organotin Compounds Involving Insertion of Norbornene. Synthesis of 2,3-Disubstituted Norbornane. *Chem. Lett.* **1987**, 193. (e) Kosugi, M.;

- Tamura, H.; Sano, H.; Migita, T. Palladium-Catalyzed Reaction of Organic Halides with Organotin Compounds Involving Olefin Insertion: Synthesis of 2,3-Disubstituted Norbornanes. *Tetrahedron* **1989**, *45*, 961. (f) Kosugi, M.; Kimura, T.; Oda, H.; Migita, T. Synthesis of *cis*-Olefins via Palladium-Catalyzed Coupling of Organic Halides, Norbornadiene, and Organotin Compounds. *Bull. Chem. Soc. Jpn.* **1993**, *66*, 3522. (g) Oda, H.; Kobayashi, T.; Kosugi, M.; Migita, T. Synthesis of 5,12-DIHETE Derivative by Palladium-Catalyzed Ternary Coupling between Vinylic Halides, a Vinylic Tin Compound, and Norbornadiene. *Tetrahedron* **1995**, *51*, 695. (h) Torii, S.; Okumoto, H.; Ozaki, H.; Nakayasu, S.; Kotani, T. Palladium-Catalyzed Tandem Assembly of Norbornene, Vinylic Halides, and Cyanide Nucleophile Leading to *cis-Exo*-2,3-Disubstituted Norbornanes. *Tetrahedron Lett.* **1990**, *31*, 5319. (i) Albrecht, K.; Reiser, O.; Weber, M.; Knierem, B.; de Meijere, A. Palladium-Catalyzed Domino Coupling Reactions of Aryl Halides with Norbornene and Norbornene Derivatives—A Simple Route to Polycyclic Aromatic Compounds. *Tetrahedron* **1994**, *50*, 383. (j) Albrecht, K.; de Meijere, A. Palladium-Catalyzed Multiple Coupling Reactions of β -Bromostyrene with Norbornene and Dicyclopentadiene. *Chem. Ber.* **1994**, *127*, 2539.
- (9) For vicinal difunctionalization of activated alkynes, see: van Belzen, R.; Hoffmann, H.; Elsevier: C. J. Catalytic Three-Component Synthesis of Conjugated Dienes from Alkynes via Pd⁰, Pd^{II}, and Pd^{IV} Intermediates Containing 1,2-Diimine. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 1743.
- (10) (a) Chiusoli, G. P. Catalysis of Olefin and Carbon Monoxide Insertion Reactions. *Acc. Chem. Res.* **1973**, *6*, 422. (b) Cassar, L.; Chiusoli, G. P.; Guerrieri, F. Synthesis of Carboxylic Acids and Esters by Carbonylation Reactions at Atmospheric Pressure Using Transition Metal Catalysts. *Synthesis* **1973**, 509. (c) Llebaria, A.; Moretó, J. M. The [Ni(CO)₄]-Promoted Carbonylative Cycloaddition of Alkynes and Allyl Halides for the Synthesis of 2-Cyclopentene Derivatives. *J. Organomet. Chem.* **1993**, *452*, 1.
- (11) Billington, D. C. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon Press: Oxford, 1991; Vol. 3, p 423.
- (12) Ikeda, S.; Cui, D.-M.; Sato, Y. Regio- and Stereoselective Synthesis of 3,6-Diene-1-ynes by Nickel-Catalyzed Coupling Reaction of Allyl Chlorides, 1-Alkynes, and Alkynyltins. *J. Org. Chem.* **1994**, *59*, 6877.
- (13) Cui, D.-M.; Tsuzuki, T.; Miyake, K.; Ikeda, S.; Sato, Y. Nickel-Catalyzed Tandem Coupling of Allyl Electrophiles, Alkynes, and Alkynyltins. *Tetrahedron* **1998**, *54*, 1063.
- (14) Ikeda, S.; Miyashita, H.; Sato, Y. Tandem Coupling of Allyl Electrophiles, Alkynes, and Me₂Al or Me₂Zn in the Presence of a Nickel Catalyst. *Organometallics* **1998**, *17*, 4316.
- (15) Cui, D.-M.; Hashimoto, N.; Ikeda, S.; Sato, Y. Nickel-Catalyzed Cross-Coupling Reaction of Allyl Halides with Alkynyltins. *J. Org. Chem.* **1995**, *60*, 5752.
- (16) Ikeda, S.; Sato, Y. Synthesis of Stereodefined Enynes by the Nickel-Catalyzed Coupling Reaction of Alkynyltins, Alkynes, and Enones. *J. Am. Chem. Soc.* **1994**, *116*, 5975.
- (17) Ikeda, S.; Kondo, K.; Sato, Y. Nickel-Catalyzed Tandem Coupling of α,β -Enones, Alkynes, and Alkynyltins for the Regio- and Stereoselective Synthesis of Conjugated Enynes. *J. Org. Chem.* **1996**, *61*, 8248.
- (18) Ikeda, S.; Kondo, K.; Sato, Y. Nickel-Catalyzed Tandem Coupling of Enones, Alkynes, Alkynylzinc, and Chlorotrimethylsilane. *Chem. Lett.* **1999**, 1227.
- (19) Ikeda, S.; Yamamoto, H.; Kondo, K.; Sato, Y. Nickel-Catalyzed Tandem Coupling of Chlorotrimethylsilane, α,β -Enones, Alkynes, and Dimethylzinc. *Organometallics* **1995**, *14*, 5015.
- (20) Johnson, J. R.; Tully, P. S.; Mackenzie, P. B.; Sabat, M. A Practical Reversed-Polarity Alternative to Organocuprate Conjugate Addition Chemistry. Halocarbon Coupling Reactions of Enal- and Enone-Derived Allylnickel Reagents. *J. Am. Chem. Soc.* **1991**, *113*, 6172.
- (21) Studemann, T.; Knochel, P. New Nickel-Catalyzed Carbozincation of Alkynes: A Short Synthesis of (*Z*)-Tamoxifen. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 93.
- (22) Shirakawa, E.; Yamasaki, K.; Yoshida, H.; Hiyama, T. Nickel-Catalyzed Carbostannylation of Alkynes with Allyl-, Acyl-, and Alkynylstannanes: Stereoselective Synthesis of Trisubstituted Vinylstannanes. *J. Am. Chem. Soc.* **1999**, *121*, 10221.
- (23) Erdic, E. Transition Metal Catalyzed Reactions of Organozinc Reagents. *Tetrahedron* **1992**, *48*, 9577.
- (24) Grisso, B. A.; Johnson, J. R.; Mackenzie, P. B. Nickel-Catalyzed, Chlorotrialkylsilane-Assisted Conjugate Addition of Alkynyltin Reagents to α,β -Unsaturated Aldehydes. Evidence for a [1-(Trialkylsilyloxy)allyl]nickel(II) Mechanism. *J. Am. Chem. Soc.* **1992**, *114*, 5160.
- (25) (a) Montgomery, J.; Savchenko, A. V. Nickel-Catalyzed Cyclizations of Alkynyl Enones with Concomitant Stereoselective Tri- or Tetrasubstituted Alkene Introduction. *J. Am. Chem. Soc.* **1996**, *118*, 2099. (b) Montgomery, J.; Seo, J.; Chui, H. M. P. Competition Between Insertion and Conjugate Addition in Nickel-Catalyzed Coupling of Enones with Unsaturated Functional Groups. *Tetrahedron Lett.* **1996**, *37*, 6839. (c) Montgomery, J.; Oblinger, E.; Savchenko, A. V. Nickel-Catalyzed Organozinc-Promoted Carbocyclizations of Electron-Deficient Alkenes with Tethered Unsaturation. *J. Am. Chem. Soc.* **1997**, *119*, 4911. (d) Montgomery, J.; Chevliakov, M. V.; Briemann, H. L. Nickel-Catalyzed Heterocycle Construction with Stereoselective Exocyclic Alkene Introduction. *Tetrahedron* **1997**, *53*, 16449. (e) Chevliakov, M. V.; Montgomery, J. Nickel and Palladium Catalysis in the Stereoselective Synthesis of Functionalized Pyrrolidines: Enantioselective Formal Synthesis of (+)- α -Allokainic Acid. *Angew. Chem., Int. Ed. Engl.* **1998**, *37*, 3144. (f) Seo, J.; Chui, H. M. P.; Heeg, M. J.; Montgomery, J. Novel Chemoselectivity and Stereochemical Aspects of Nickel-Catalyzed [2 + 2 + 2] Cycloadditions. *J. Am. Chem. Soc.* **1999**, *121*, 476. (g) Chevliakov, M. V.; Montgomery, J. A Stereodivergent Approach to (–)-Kainic Acid and (+)-Allokainic Acid Utilizing the Complementarity of Alkyne and Allene Cyclizations. *J. Am. Chem. Soc.* **1999**, *121*, 11139.
- (26) Ward, R. S. Non-Enzymatic Asymmetric Transformations Involving Symmetrical Bifunctional Compounds. *Chem. Soc. Rev.* **1990**, *19*, 1.
- (27) (a) Torii, S.; Okumoto, H.; Ozaki, H.; Nakayasu, S.; Tadokoro, T.; Kotani, T. Asymmetric Multipoint Control by Diastereo-Differentiative Assembly of Three Components with Palladium Catalyst. *Tetrahedron Lett.* **1992**, *33*, 3499. (b) Torii, S.; Okumoto, H.; Kotani, T.; Nakayasu, S.; Ozaki, H. Asymmetric Multipoint Control by Palladium-Catalyzed Diastereoselective Tandem Carbon–Carbon Bond Formation. *Tetrahedron Lett.* **1992**, *33*, 3503.
- (28) (a) Brunner, H.; Obermann, U. Enantioselective Hydrosilylierung von Ketonen mit [Rh(COD)Cl]₂/Pyridinyloxazolone-Katalysatoren. *Chem. Ber.* **1989**, *122*, 499. (b) Bolm, C.; Weickhardt, K.; Zehnder, M.; Ranff, T. Synthesis of Optically Active Bis(2-oxazolines): Crystal Structure of a 1,2-Bis(2-oxazolonyl)benzene-ZnCl₂ Complex. *Chem. Ber.* **1991**, *124*, 1173.
- (29) Cui, D.-M.; Yamamoto, H.; Ikeda, S.; Hatano, K.; Sato, Y. Nickel-Catalyzed Diastereodifferentiative Tandem Coupling of α,β -Enones, Norbornenes, and Organometallics. *J. Org. Chem.* **1998**, *63*, 2782.
- (30) Kablaoui, N. M.; Hicks, F. A.; Buchwald, S. L. Titanocene-Catalyzed Cyclocarbonylation of α -Allyl Aryl Ketones to β -Butyrolactones. *J. Am. Chem. Soc.* **1997**, *119*, 4424.
- (31) Carpenter, N. E.; Kucera, D. J.; Overman, L. E. Palladium-Catalyzed Polyene Cyclization of Trienyl Triflates. *J. Org. Chem.* **1989**, *54*, 5846.
- (32) Shibasaki, M.; Boden, C. D. J.; Kojima, A. The Asymmetric Heck Reaction. *Tetrahedron* **1997**, *53*, 7371.
- (33) Maddaford, S. P.; Andersen, N. G.; Cristofoli, W. A.; Keay, B. A. Total Synthesis of (+)-Xestoninone Using an Asymmetric Palladium-Catalyzed Polyene Cyclization. *J. Am. Chem. Soc.* **1996**, *118*, 10766.
- (34) Miyaura, N.; Suzuki, A. Palladium-Catalyzed Cross Coupling Reactions of Organoboron Compounds. *Chem. Rev.* **1995**, *95*, 2457.
- (35) Kojima, A.; Takemoto, T.; Sodeoka, M.; Shibasaki, M. Catalytic Asymmetric Synthesis of Halenaquinone and Halenaquinol. *J. Org. Chem.* **1996**, *61*, 4876.
- (36) (a) Ohshima, T.; Kagechika, K.; Adachi, M.; Sodeoka, M.; Shibasaki, M. Asymmetric Heck Reaction – Carbanion Capture Process. Catalytic Asymmetric Total Synthesis of (–)- $\Delta^9(12)$ -Capnellene. *J. Am. Chem. Soc.* **1996**, *118*, 7108. Also see, (b) Kagechika, K.; Shibasaki, M. Asymmetric Heck Reaction: A Catalytic Asymmetric Synthesis of the Key Intermediate for $\Delta^9(12)$ -Capnellene-3 β ,8 β ,10 α -triol and $\Delta^9(12)$ -Capnellene-3 β ,8 β ,10 α ,14-tetrol. *J. Org. Chem.* **1991**, *56*, 4093. (c) Kagechika, K.; Ohshima, T.; Shibasaki, M. Asymmetric Heck Reaction–Anion Capture Process. A Catalytic Asymmetric Synthesis of the Key Intermediates for the Capnellens. *Tetrahedron* **1993**, *49*, 1773.
- (37) For recent examples of domino catalytic asymmetric reactions, see: (a) Arai, T.; Sasai, H.; Aoe, K.; Okamura, K.; Date, T.; Shibasaki, M. A New Multifunctional Heterobimetallic Asymmetric Catalyst for Michael Additions and Tandem Michael–Aldol Reactions. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 4876. (b) Yamada, K.; Arai, T.; Sasai, H.; Shibasaki, M. A Catalytic Asymmetric Synthesis of 11-Deoxy-PGF1 Using ALB, a Heterobimetallic Multifunctional Asymmetric Complex. *J. Org. Chem.* **1998**, *63*, 3666. (c) Sasai, H.; Hiroi, M.; Yamada, Y. M. A.; Shibasaki, M. The First Tandem Inter-Intramolecular Catalytic Asymmetric Nitroaldol Reaction Utilizing a LnLi₃tris((*R*)-binaphthoxide) Complex ((*R*)-LnLB) (Ln: Lanthanoid). *Tetrahedron Lett.* **1997**, *38*, 6031. (d) Mikami, K.; Matsukawa, S.; Nagashima, M.; Funabashi, H.; Morishima, H. Tandem and Two Directional Asymmetric Catalysis of the Mukaiyama Aldol Reaction. *Tetrahedron Lett.* **1997**, *38*, 579.

- (38) Ikeda, S.; Cui, D.-M.; Sato, Y. Nickel-Catalyzed Asymmetric Multiple-Component Tandem Coupling. Effects of Simple Monodentate Oxazolines as Chiral Ligands. *J. Am. Chem. Soc.* **1999**, *121*, 4712.
- (39) (a) Kurth, M. J.; Decker, O. H. W. Enantioselective Preparation of 3-Substituted-4-pentenoic Acids via the Claisen Rearrangement. *J. Org. Chem.* **1985**, *50*, 5769. (b) Leonard, W. R.; Romine, J. L.; Meyers, A. I. A Rapid and Efficient Synthesis of Chiral 2-Hydroxy-2-Oxazolines. *J. Org. Chem.* **1991**, *56*, 1961. (c) Meyers, A. I.; Shipman, M. Asymmetric Michael Additions to α,β -Unsaturated Oxazolines. An Efficient Preparation of Chiral β,β -Disubstituted Propionaldehydes. *J. Org. Chem.* **1991**, *56*, 7098. (d) Kamata, K.; Agata, I.; Meyers, A. I. An Efficient and Versatile Method for the Synthesis of Optically Active 2-Oxazolines: An Acid-Catalyzed Condensation of Ortho Esters with Amino Alcohols. *J. Org. Chem.* **1998**, *63*, 3113.
- (40) Ager, D. J.; Prakash, I.; Schaad, D. R. 1,2-Amino Alcohols and Their Heterocyclic Derivatives as Chiral Auxiliaries in Asymmetric Synthesis. *Chem. Rev.* **1996**, *96*, 835.
- (41) Noyori, R.; Takaya, H. BINAP: An Efficient Chiral Element for Asymmetric Synthesis. *Acc. Chem. Res.* **1990**, *23*, 345.
- (42) Uozumi, Y.; Hayashi, T. Catalytic Asymmetric Synthesis of Optically Active 2-Alkanols via Hydrosilylation of 1-Alkenes with a Chiral Monophosphine-Palladium Catalyst. *J. Am. Chem. Soc.* **1991**, *113*, 9887.
- (43) (a) von Matt, P.; Pfaltz, A. Chiral Phosphinoaryldihydroxazoles as Ligands in Asymmetric Catalysis: Pd-Catalyzed Allylic Substitution. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 566. (b) Sprinz, J.; Helmchen, G. Phosphinoaryl- and Phosphinoalkyloxazolines as New Chiral Ligands for Enantioselective Catalysis: Very High Enantioselectivity in Palladium Catalyzed Allylic Substitutions. *Tetrahedron Lett.* **1993**, *34*, 1769. (c) Dawson, G. J.; Frost, C. G.; Williams, J. M. J.; Coote, S. J. Asymmetric Palladium Catalyzed Allylic Substitution Using Phosphorus Containing Oxazoline Ligands. *Tetrahedron Lett.* **1993**, *34*, 3149.
- (44) Pfaltz, A. Chiral Semicorrins and Related Nitrogen Heterocycles as Ligands in Asymmetric Catalysis. *Acc. Chem. Res.* **1993**, *26*, 339.
- (45) Reppe, W.; Schweckendiek, W. J. Cyclisierende Polymerisation von Acetylen. III Benzol, Benzolderivate und Hydroaromatische Verbindungen. *Justus Liebigs Ann. Chem.* **1948**, *560*, 104.
- (46) (a) Schore, N. E. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon: Oxford, 1991; Vol. 5, Chapter 9.4, p 1129. (b) Grotjahn, D. B. In *Comprehensive Organometallic Chemistry II*; Hegedus, L. S., Ed.; Pergamon Press: Oxford, 1995; Vol. 12, Chapter 7.4.
- (47) (a) Schore, N. E. Transition-Metal-Mediated Cycloaddition Reactions of Alkynes in Organic Synthesis. *Chem. Rev.* **1988**, *88*, 1081. (b) Lautens, M.; Klute, W.; Tam, W. Transition Metal-Mediated Cycloaddition Reactions. *Chem. Rev.* **1996**, *96*, 49.
- (48) Vollhardt, K. P. C. Cobalt-Mediated [2 + 2 + 2]-Cycloadditions: a Maturing Synthetic Strategy. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 539.
- (49) For examples of stoichiometric process, see: (a) Wakatsuki, Y.; Kuramitsu, T.; Yamazaki, H. Cobaltacyclopentadiene Complexes as Starting Materials in the Synthesis of Substituted Benzenes, Cyclohexadienes, Thiophenes, Selenophenes, and Pyrroles. *Tetrahedron Lett.* **1974**, 4549. (b) Takahashi, T.; Kitora, M.; Xi, Z. Cycloaddition of Zirconacyclopentadienes to Alkynes using Copper Salts: Formation of Benzene Derivatives. *J. Chem. Soc., Chem. Commun.* **1995**, 361. (c) Takahashi, T.; Tasi, F.-Y.; Li, Y.; Nakajima, K.; Kitora, M. Carbon-Carbon Bond Formation Reaction of Zirconacyclopentadienes with Alkynes in the Presence of Ni(II)-complexes. *J. Am. Chem. Soc.* **1999**, *121*, 11093. For a cyclotrimerization of alkynes using a stoichiometric amount of zirconium complex and a catalytic amount of copper salt, see: (d) Takahashi, T.; Xi, Z.; Yamazaki, A.; Liu, Y.; Nakajima, K.; Kitora, M. Cycloaddition Reaction of Zirconacyclopentadienes to Alkynes: Highly Selective Formation of Benzene Derivatives from Three Different Alkynes. *J. Am. Chem. Soc.* **1998**, *120*, 1672.
- (50) Ikeda, S.; Mori, N.; Sato, Y. Regioselective Cyclic Cotrimerization of α,β -Enones and Alkynes by a Nickel-Aluminum Catalyst System. *J. Am. Chem. Soc.* **1997**, *119*, 4779.
- (51) For other regioselective cyclotrimerization of alkenes and alkynes, see: (a) Heimbach, P.; Ploner, K.-J.; Thömel, F. Preparation of Substituted 5-Vinylcyclohexa-1,3-dienes and Tricyclo[2.2.2.0^{2,6}]-oct-7-enes. *Angew. Chem., Int. Ed. Engl.* **1971**, *10*, 276. (b) Chalk, A. J. Catalysis of the Cyclotrimerization of Acetylenes with N-Substituted Maleimides. *J. Am. Chem. Soc.* **1972**, *94*, 5928. (c) Balaich, G. J.; Rothwell, I. P. Regio- and Stereoselective Formation and Isomerization of 1,3-Cyclohexadienes Catalyzed by Titanium Aryloxy Compounds. *J. Am. Chem. Soc.* **1993**, *115*, 1581. (d) Johnson, E. S.; Balaich, G. J.; Rothwell, I. P. Regio- and Stereoselective Synthesis of the 1,3-Cyclohexadiene Nucleus by [2 + 2 + 2] Cycloaddition Reactions Catalyzed by Titanium Aryloxy Compounds. *J. Am. Chem. Soc.* **1997**, *119*, 7685.
- (52) Maruoka, K.; Yamamoto, H. Precise Design of Aluminum Reagents That Exhibit Steric and Electronic Effects. *J. Synth. Org. Chem. Jpn.* **1993**, *51*, 1074 and references therein.
- (53) Mori, N.; Ikeda, S.; Sato, Y. Selective Cyclotrimerization of Enones and Alkynes by a Nickel and Aluminum Catalytic System. *J. Am. Chem. Soc.* **1999**, *121*, 2722.
- (54) (a) Ikeda, S.; Watanabe, H.; Sato, Y. Nickel- and Zinc-Promoted [2 + 2 + 2] Cycloaddition of Dienes and α,β -Enones. *J. Org. Chem.* **1998**, *63*, 7026. Also see: (b) Sambaiah, T.; Li, L.-P.; Huang, D.-J.; Lin, C.-H.; Rayabarapu, D. K.; Cheng, C.-H. Highly Regio- and Stereoselective Cocyclotrimerization and Linear Cotrimerization of α,β -Unsaturated Carbonyl Compounds with Alkynes Catalyzed by Nickel Complexes. *J. Org. Chem.* **1999**, *64*, 3663.
- (55) For regioselective cycloaddition of diynes with alkynes, see: (a) McDonald, F. E.; Zhu, H. Y. H.; Holmquist, C. R. Rhodium-Catalyzed Alkyne Cyclotrimerization Strategies for C-Arylglycoside Synthesis. *J. Am. Chem. Soc.* **1995**, *117*, 6605. (b) Witulski, B.; Stengel, T. Rhodium(I)-Catalyzed [2+2+2] Cycloadditions with N-Functionalized 1-Alkynylamines: A Conceptually New Strategy for the Regiospecific Synthesis of Substituted Indolines. *Angew. Chem., Int. Ed. Engl.* **1999**, *38*, 2426.
- (56) Recently, nickel-catalyzed domino coupling reactions using aldehydes have been extensively studied. For example: (a) Oblinger, E.; Montgomery, J. A New Stereoselective Method for the Preparation of Allylic Alcohols. *J. Am. Chem. Soc.* **1997**, *119*, 9065. (b) Qi, X.; Montgomery, J. New Three-Component Synthesis of 1,3-Dienes Employing Nickel Catalysis. *J. Org. Chem.* **1999**, *64*, 9310. (c) Kimura, M.; Matsuo, S.; Shibata, K.; Tamaru, Y. Nickel(0)-Catalyzed Three-Component Connection Reaction of Dimethylzinc, 1,3-Dienes, and Carbonyl Compounds. *Angew. Chem., Int. Ed. Engl.* **1999**, *38*, 3386. (d) Sato, Y.; Takanashi, T.; Mori, M. Studies on Transmetalation of a Nickelacycle with Organometallic Reagents. *Organometallics* **1999**, *18*, 4891.

AR9901016