

# Humanization of Lepidopteran Insect-Cell-Produced Glycoproteins

NOBORU TOMIYA,<sup>\*,†</sup>MICHAEL J. BETENBAUGH,<sup>‡</sup> AND YUAN C. LEE<sup>†</sup>

Department of Biology and Department of Chemical and Biomolecular Engineering, Johns Hopkins University, 3400 North Charles Street, Baltimore, Maryland 21218

Received November 7, 2002

## ABSTRACT

The insect cell–baculovirus expression vector system, widely used for glycoprotein production, is not ideal for pharmaceutical glycoprotein production due to the characteristics of the *N*-glycans in the expressed products. Insect cells lack several enzymes required for mammalian-type *N*-glycan synthesis and contain a specific *N*-acetylglucosaminidase that stunts the growth of chains and a core  $\alpha$ -1,3-fucosyltransferase that yields potentially allergenic glycoforms. Current knowledge on *N*-glycan processing in lepidopteran insect cells is summarized, and strategies to develop better glycoprotein expression systems suitable for pharmaceutical glycoprotein production are discussed.

## 1. Introduction

Several insect cell lines from lepidopteran insects (see Figure 1) are used for glycoprotein expression with the baculovirus expression vector system because this expression vector system can produce glycoproteins more readily than mammalian cell expression systems. In addition, lepidopteran insect cells grow well in serum-free medium and in suspension culture. However, this system is not yet ideal for the production of pharmaceutical glycoproteins because the insect-produced glycoproteins have significantly different *N*-glycans from those by mammalian cells (Figure 2). They are (i) the inability to synthesize sialylated complex-type *N*-glycans in contrast to mammalian cells<sup>1–3</sup> and (ii) the presence of potentially allergenic structure, Fuc $\alpha$ (1,3)GlcNAc-Asn, not seen in glycoproteins from mammalian cells. Insect cells tend to generate paucimannosidic or high mannose structures

terminating in Man or occasionally GlcNAc. Paucimannosidic structures have been repeatedly found on glycoproteins produced by lepidopteran insect cells derived from *Spodoptera frugiperda*,<sup>4–17</sup> *Mamestra brassica*,<sup>5,8</sup> *Bombyx mori*,<sup>4,8</sup> *Trichoplusia ni*,<sup>17,18</sup> and *Estigmene acrea*.<sup>9,10</sup> These small glycans consist of one to three Man and one or two GlcNAc residues with or without Fuc attached to the Asn-linked GlcNAc. Differences in *N*-glycan structures may result in different biological activities, pharmacokinetic behavior, and allergenic reactions in human.<sup>19–22</sup>

In the past decade, much effort has been devoted to analysis of *N*-glycan structures, activity measurements and gene cloning of *N*-glycan processing enzymes, and measurements of sugar nucleotide for the baculovirus-insect cell system. It is clear that the inability of most lepidopteran insect cells to produce mammalian-type *N*-glycans are mainly attributable to (i) the lack of certain glycosyltransferases, (ii) the inability to produce CMP-neuraminic acid, and (iii) the presence of a  $\beta$ -*N*-acetylglucosaminidase which removes GlcNAc from the Man $\alpha$ (1,3) branch. Recently, attempts have been made to improve *N*-glycan structures of lepidopteran cell-produced glycoproteins to alleviate these problems. In this Account, we will dissect the *N*-glycan processing pathways in lepidopteran cells with reference to enzyme activities and the levels of donor sugar nucleotides involved in *N*-glycan processing. We also review the status of engineering *N*-glycan processing pathways in this protein expression system. Readers are also referred to other recently published reviews on *N*-glycosylation in insects and insect cells.<sup>2,3</sup>

## 2. Analytical Methods

**2.1. Assay of Glycosyltransferase and Glycosidase Activity.** Most common measurement of glycosyltransferase and glycosidase activities is by the use of radiolabeled sugar nucleotides to trace the radiolabeled sugar transferred to the acceptor. Although this method is sensitive, its obstacles are strictly controlled storage, usage, and disposal of radioisotopes. Glycans can be labeled with a fluorescent tag such as pyridylamino (PA) group for structural analysis, glycosyltransferase, and glycosidase assays. 2-D HPLC analysis of PA-glycans enables fine separation and reliable identification of product glycans with subtle structural differences.<sup>23,24</sup>

Many insect cells contain core  $\alpha$ -1,3- and core  $\alpha$ -1,6-fucosyltransferases, which add Fuc to the Asn-linked GlcNAc via  $\alpha$ -1,3- and  $\alpha$ -1,6-linkages, respectively. The activities of these enzymes were assayed by measuring the transfer of radiolabeled Fuc from GDP-Fuc to acceptor substrates. However, this assay is unsatisfactory when the sample has both enzyme activities. Since PA-derivatized glycans could not be acceptors for Fuc-T C3,<sup>25</sup> fluorescent

Noboru Tomiya, Research Associate of Biology at Johns Hopkins University since 1998, was born in Japan (1952) and obtained his PhD degree from Nagoya University (Japan) in 1988. His interests have been in the biochemistry and engineering of glycoproteins. He developed the multidimensional HPLC technique with Dr. Noriko Takahashi (Nagoya City University, Japan) for glycan structural analysis.






Michael J. Betenbaugh, Professor of Chemical Engineering at Johns Hopkins University since 1999, was born in Atlanta, Georgia (1959) and obtained his PhD degree in 1988 from University of Delaware. He has been collaborating with the coauthors on the production of glycoproteins with insect cells. His interests have been in biotechnology, glycobiology, insect and mammalian cell culture, and metabolic engineering of glycoproteins.

Yuan, C. Lee, Professor of Biology at Johns Hopkins University since 1974, was born in Taiwan and obtained his PhD in Biochemistry in 1962. He has been collaborating with the coauthors on production of glycoproteins with insect cells. His interests have been in the chemistry and biochemistry of glycoconjugates. He was the recipient of the 2001 Hudson Award from American Chemistry Society.


\* To whom correspondence should be addressed. Tel: +1-410-516-7322. Fax: +1-410-516-8716. E-mail: ntomiyal@jhu.edu.

<sup>†</sup> Department of Biology.

<sup>‡</sup> Department of Chemical and Biomolecular Engineering.

					
Latin name	<i>Spodoptera frugiperda</i>	<i>Trichoplusia ni</i>	<i>Estigmene acrea</i>	<i>Bombyx mori</i>	<i>Mamestra brassicae</i>
Common name	Fall Armyworm	Cabbage looper	Salt marsh moth	Silkworm	Cabbage moth
Family name (Subfamily name)	Noctuidae (Amphipyriinae)	Noctuidae (Plusiinae)	Arctiidae	Bombycidae	Noctuidae (Hadeninae)
Cell line	Sf9/Sf21	BTI-Tn-5B1-4	Ea4	Bm-N	Mb0503
Tissue derived	Pupal ovary	Pupal ovary	Larval hemocytes	Ovary	Larval hemocytes

**FIGURE 1.** Lepidopteran cell lines often used for glycoprotein production. Pictures of *Spodoptera frugiperda*, *Trichoplusia ni*, and *Estigmene acrea* were obtained from "Web Images of North American Moth Species: <http://alpha.furman.edu/~snyder/leplist/>". Pictures of *Bombyx mori* and *Mamestra brassicae* were obtained from "Moths and Butterflies of Europe: <http://www.leps.it/>".




**FIGURE 2.** Comparison of typical *N*-glycans found in glycoproteins produced by human or lepidopteran cells. (a) Sialylated complex-type *N*-glycan, (b) high mannose-type *N*-glycan, (c) paucimannosidic *N*-glycan. GN, *N*-acetylglucosamine; F, fucose; M, mannose; G, galactose.

derivatized glycopeptides must be used as acceptor substrates, and fucosylated products were analyzed by reversed-phase HPLC<sup>26</sup> and MALDI-TOF mass spectrometry.<sup>27</sup> A recently developed assay of galactosyltransferase (GalT) uses microtiterplate-coated GlcNAc-BSA as the acceptor and Eu-labeled RCA120 lectin as the probe. The newly formed Gal $\beta$ (1,4)GlcNAc structure is measured as  $\beta$ 4GalT activity in both native and engineered lepidopteran insect cells.<sup>28</sup>

**2.2. Sugar Nucleotide Assay.** Prior methods for measurement of sugar nucleotides in biological samples, using anion-exchange HPLC or reversed-phase HPLC, could not separate UDP-Glc and UDP-GlcNAc from UDP-Gal and UDP-GalNAc, respectively. Our recent method, high-performance anionic exchange chromatography under mildly alkaline condition, can separate all sugar nucleotides in the extracts from lepidopteran insect cells simultaneously.<sup>29</sup>

### 3. Glycosyltransferases and Glycosidases Involved in *N*-Glycan Processing in Insect Cells

The processing pathway of *N*-glycans in lepidopteran insect cells is shown in Figure 3. A number of studies have suggested that initial processing of *N*-glycans in insect



**FIGURE 3.** *N*-Glycan processing pathway in lepidopteran insect cells. Symbols: ■, GlcNAc; ○, Man; ●, Gal; ◇, Fuc; □, Glc; △, Sia. The dotted arrow indicates pathways which are insufficient in lepidopteran insect cells.

cells is similar or identical to that of mammalian cells. However, insect cells appear to lack some of the processing pathways of mammalian cells but contain additional glycosylation activities absent in mammalian cells.


**3.1.  $\alpha$ -Glucosidase I, II, and  $\alpha$ -Mannosidase I.** Glc<sub>3</sub>-Man<sub>9</sub>GlcNAc<sub>2</sub> is processed by  $\alpha$ -glucosidase I, II, and  $\alpha$ -mannosidase I to generate Man<sub>5</sub>GlcNAc<sub>2</sub> structure. Many glycoproteins produced by lepidopteran insect cells have high-mannose-type glycans. For example, *N*-glycans on human IgG and hTF produced by Tn-5B1-4 cells included various high mannose-type and paucimannosidic glycans, with some incomplete complex-type glycans.<sup>18,30</sup> Expression of  $\alpha$ -glucosidase I and II in several lepidopteran insect cells appears adequate.<sup>31</sup> In addition,  $\alpha$ -mannosidase I has been purified from Sf21 cells<sup>32</sup> and cloned from Sf9 cells,<sup>33</sup> and its substrate specificity has been characterized.<sup>34</sup> These results suggest that lepidopteran insect cells contain ample  $\alpha$ -glucosidase I, II, and  $\alpha$ -mannosidase I.

**3.2. *N*-Acetylglucosaminyltransferase I and  $\alpha$ -Mannosidase II.** First, GlcNAc is added to Man $\alpha$ (1,3) branch of Man<sub>5</sub>GlcNAc<sub>2</sub> by *N*-acetylglucosaminyltransferase I (GlcNAcT-I); thereafter, two Man residues are removed by  $\alpha$ -mannosidase II. Substantial levels of GlcNAcT-I activities were observed in several insect cell lines including Sf9, Sf21, Mb0503, and Bm-N.<sup>35</sup> Mannosidase-II was active in Sf21, Mb0503, and Bm-N cells<sup>36</sup> and was purified from Sf21 cells.<sup>37</sup> Like its counterpart from mammalian cells,  $\alpha$ -mannosidase II from insect cells requires GlcNAc on the Man $\alpha$ (1–3) branch for its activity.<sup>36</sup> These studies suggest that lepidopteran insect cells have high levels of  $\alpha$ -mannosidase II and GlcNAcT-I in order to generate the precursor glycan required for the formation of complex-type *N*-glycans.

**3.3. *N*-Acetylglucosaminyltransferase II.** In mammalian cells, the product *N*-glycan of  $\alpha$ -mannosidase II reaction serves as an acceptor for the next reaction catalyzed by *N*-acetylglucosaminyltransferase II (GlcNAcT-II), which adds another GlcNAc to the Man $\alpha$ (1,6) branch. However, lepidopteran insect cells, including Sf9, Sf21, Mb0503, and Bm-N cells, have been shown to have only 1% or less of the endogenous GlcNAcT-II activity present in mammalian cells.

**3.4.  $\beta$ -1,4-Galactosyltransferase.** A terminal GlcNAc on either Man-branch is usually galactosylated by  $\beta$ -1,4-galactosyltransferase (GalT) in mammalian cells. In contrast, galactosylated *N*-glycans are rarely found in glycoproteins from lepidopteran cells. In fact, negligible levels of GalT activity were detected in Sf9, Tn-5B1-4, and Mb0503 cells.<sup>38–40</sup> GalT activities in Sf9 and Tn-5B1-4 cells were reexamined using an Eu-fluorescence assay method.<sup>28</sup> Tn-5B1-4 cells were observed to contain about 10% of GalT activity detected in Chinese hamster ovary (CHO) cells, while Sf9 cells did not contain any detectable levels of GalT activity.<sup>28</sup>

**3.5. Core  $\alpha$ -1,3- and  $\alpha$ -1,6-Fucosyltransferases.** *N*-Glycans with one or two GlcNAc on Man<sub>3</sub>-core can be further modified by core fucosyltransferases. Both core Fuc-T's require the presence of GlcNAc $\beta$ (1,2) on the Man $\alpha$ (1,3) branch for its action.<sup>41</sup> *N*-Glycans containing either one or both of Fuca $\alpha$ (1,3) and Fuca $\alpha$ (1,6) attached to the Asn-linked GlcNAc were identified on the membrane glycoproteins from Mb0503, Sf21, and Bm-N cells, in which glycoproteins from Mb0503 cells contained highest levels of  $\alpha$ -1,3-fucosylated *N*-glycans.<sup>8</sup> Similar results were reported for *N*-glycans from recombinant IgG,<sup>42</sup> human serum transferrin,<sup>30</sup> neuropsin,<sup>43</sup> and CR3 domain of latent TGF-beta binding protein-1<sup>44</sup> produced by Tn-5B1-4 cells. However, human interferon  $\omega$ 1 expressed in Sf9 cells showed no  $\alpha$ -1,3-fucosylated *N*-glycan.<sup>6</sup> Activities of core  $\alpha$ -1,3-fucosyltransferase (Fuc-T C3) and core  $\alpha$ -1,6-fucosyltransferase (Fuc-T C6) were detected in Mb0503 cells, in which the former was much higher than the latter.<sup>41</sup> Fuc-T C6, but not Fuc-T C3, was easily detected in Sf9 cells. Independently, Fuc-T C6 activity was detected in Sf9, Mb0503, and Bm-N cells.<sup>45</sup> Interestingly, in the same study, Fuc-T C3 was detected only in Mb0503 cells and



**FIGURE 4.** Sugar nucleotide contents of Sf9, Tn-5B1-4, and CHO cells (ref 29).

not in Sf9 and Bm-N cells. A gene encoding Fuc-T C3 has been recently cloned from *Drosophila melanogaster*.<sup>27</sup>


**3.6.  $\beta$ -*N*-Acetylglucosaminidase.** A  $\beta$ -*N*-Acetylglucosaminidase specific for the terminal GlcNAc on the Man $\alpha$ (1,3) branch was found in Sf21, Bm-N, and Mb0503 cells,<sup>46</sup> and it was suggested that this enzyme was localized in the microsome-like membrane fraction in Sf21 cells.<sup>46</sup> Similar enzymatic activity was also detected in the cell lysates and cell culture supernatant of insect cells derived from *Spodoptera frugiperda*, *Trichoplusia ni*, *Bombyx mori*, or *Malacosoma disstria*.<sup>47</sup> Structural analysis of *N*-glycans from human IgG<sup>42</sup> and hTf<sup>30</sup> expressed in Tn-5B1-4 cells suggested the presence of such a  $\beta$ -*N*-acetylglucosaminidase in Tn-5B1-4 cells. The further removal of additional Man residues by  $\alpha$ -mannosidase(s) can lead to the generation of structures with fewer than three Man residues, as has been observed in several studies.

**3.7. Sialyltransferase.** Sialyltransferase (SiaT) adds *N*-acetylneuraminic acid to the terminal Gal residues on *N*-glycans in mammalian cells. However, SiaT activity has yet to be detected in Sf9,<sup>40,48,49</sup> Sf21,<sup>49</sup> Tn-5B1-4,<sup>48</sup> Mb0503,<sup>48</sup> and Ea4<sup>49</sup> cells, even using highly sensitive assays with radiolabeled CMP-NeuAc or fluorescent CMP-Neu5Ac derivatives as the donor substrate.

## 4. Sugar Nucleotides

**4.1. Endogenous Sugar Nucleotide Levels in Lepidopteran Insect Cells.** All glycosyltransferases in the synthetic pathway for complex-type *N*-glycans require respective sugar nucleotides as donor substrates. Examination of the sugar nucleotide concentrations in lepidopteran insect cells demonstrated the presence of substantial levels of UDP-hexose, UDP-*N*-acetylhexosamine, GDP-Fuc, and GDP-Man in Sf9, Mb0503, and Tn-5B1-4 cells.<sup>48</sup> However, no CMP-NeuAc was detected in the same study.<sup>48</sup> Similar results were obtained on the sugar nucleotide levels in Sf9 and Tn-5B1-4 cells<sup>29</sup> (Figure 4).

**4.2. Enzymes Involved in Sialic Acid and CMP-Sialic Acid Synthesis.** Of particular significance is the absence in lepidopteran insect cells of the CMP-NeuAc necessary for sialylation of *N*-glycans. In mammalian cells, sialic



**FIGURE 5.** CMP-Neuraminic acid synthesis pathway. The dotted arrow indicates pathways which are insufficient in lepidopteran insect cells. 1, bifunctional UDP-GlcNAc 2-epimerase/ManNAc kinase; 2, *N*-acetylneuraminic acid 9-phosphate synthase; 3, *N*-acetylneuraminic acid 9-phosphate phosphatase; 4, CMP-neuraminic acid synthase. Exogenous ManNAc can be converted to ManNAc-6-P by *N*-acetylmannosamine kinase (5).

acids are synthesized from UDP-GlcNAc through multiple enzymatic reactions (Figure 5). The bifunctional enzyme, UDP-*N*-acetylglucosamine 2-epimerase/*N*-acetylmannosamine kinase, is believed to be a key enzyme in the biosynthesis of Neu5Ac in rat liver.<sup>50</sup> This enzyme converts UDP-GlcNAc to ManNAc-6-P, which is further converted to *N*-acetylneuraminic acids by *N*-acetylneuraminic acid 9-phosphate synthase (SAS) and *N*-acetylneuraminic acid 9-phosphate phosphatase. *N*-Acetylneuraminic acid is then converted to CMP-Neu5Ac by CMP-Neu5Ac synthase (CMP-SAS).

Effertz et al.<sup>51</sup> reported that the UDP-*N*-acetylglucosamine-2-epimerase activity in Sf9 cells was about 30 times less (in terms of specific activity) than that in rat liver cytosol fraction. Interestingly, Sf9 cells had 50 times higher *N*-acetylmannosamine kinase activity compared with the 2-epimerase activity.<sup>51</sup> We found that Sf9 cells contained negligible levels of neuraminic acids, and no detectable *N*-acetylneuraminic acid 9-phosphate synthase activity was present in the lysate of Sf9 cells.<sup>52</sup> We also found that Sf9 cells do not have detectable CMP-sialic acid synthase activity.<sup>53</sup>

## 5. Engineering of *N*-Glycan Processing Pathway

**5.1. Targets.** The general strategy for humanizing glycoproteins produced by the insect cell–baculovirus expression system is shown in Figure 6. The goal of engineering *N*-glycan processing is to develop a new insect cell–baculovirus expression vector system(s) that can express human-like sialylated multi-antennary complex-type *N*-glycans. As described in the earlier sections, several lines of evidence suggest that the majority of lepidopteran insect cells currently used for protein expression apparently lack several enzymes for such a goal. Moreover, lepidopteran insect cells contain the undesirable  $\beta$ -*N*-acetylglucosaminidase and Fuc-T C3. The former diminishes the key glycans containing GlcNAc $\beta$ (1,2)Man $\alpha$ (1,3) to stunt the normal growth of complex-type *N*-glycans, and the latter generates potentially allergenic *N*-glycans. Therefore, the *N*-glycan processing pathways need to be altered in the insect cells by enhancing or suppressing respective processing pathways.

**5.2. Improvement of *N*-Acetylglucosaminylation of the Man $\alpha$ (1,3)-Branch.**  $\beta$ -*N*-Acetylglucosaminidase was implicated as a problem in *N*-glycan elongation by its

absence of *Estigmene acrea* cells, which produced *N*-glycans containing terminal *N*-acetylglucosamine residues.<sup>54</sup> Sf9 cells are known to contain high levels of  $\beta$ -*N*-acetylglucosaminidase.<sup>54</sup> Using Sf9 cells, Wagner et al. succeeded in *N*-glycan elongation by coexpression of human *N*-acetylglucosaminyltransferase I and fowl plague virus hemagglutinin.<sup>9</sup> Watanabe et al.<sup>55</sup> examined the effect of a  $\beta$ -*N*-acetylglucosaminidase inhibitor, 2-acetamide-1,2-dideoxynojirimycin (2-ADN) on bovine interferon- $\gamma$  (bIFN- $\gamma$ ) production in Tn-5B1-4 cells. Inhibition of  $\beta$ -*N*-glucosaminidase led to expression of sialylated bIFN- $\gamma$ . The authors speculated that the inhibitor enhanced accumulation of substrates possessing a  $\beta$ (1,2)-linked GlcNAc, thereby leading to further elongation by GalT and SiaT to form sialylated *N*-glycans. However, the overall increase of *N*-glycans containing  $\beta$ (1,2)-linked GlcNAc was not determined.

**5.3. Improvement of Galactosylation.** Expression of a mammalian GalT by a baculovirus vector increased galactosylation of glycoprotein,<sup>56</sup> indicating that the mammalian enzyme expressed by baculovirus infection could function in the infected lepidopteran cells and that it could compete with the  $\beta$ -*N*-acetylglucosaminidase activity in insect cells.<sup>56</sup> Similar results were obtained when human serum transferrin (hTf) was expressed by Tn-5B1-4 cells infected with two baculoviruses, one encoding a gene for hTf and the other encoding a gene for a mammalian GalT.<sup>30</sup> In this study, 13% of the total *N*-glycans were galactosylated, and protection of GlcNAc on Man $\alpha$ (1,3) branch against  $\beta$ -*N*-acetylglucosaminidase by galactosylation was confirmed (Figure 7).

In a different strategy, when Sf-9 or Tn-5B1-4 cells stably transformed to include GalT into the host cell genome (Sf $\beta$ 4GalT<sup>38</sup> from Sf9 or Tn5 $\beta$ 4GalT<sup>57</sup> from Tn-5B1-4) were infected with native baculovirus, galactosylated virion glycoprotein, gp64<sup>38,57</sup> was produced (shown by lectin blot). Similarly, the infection of Sf $\beta$ 4GalT cells with a recombinant baculovirus expression vector encoding human tissue plasminogen activator (t-PA) also produced galactosylated t-PA.<sup>38</sup>

**5.4. Production of Biantennary Complex-Type *N*-Glycans.** Production of biantennary complex-type *N*-glycans was achieved recently by expressing a mammalian *N*-acetylglucosaminyltransferase II (GlcNAcT-II) in lepidopteran cells using a transgenic insect cell line, SfSWT-1,<sup>58</sup> or using baculovirus expression vector system.<sup>59</sup>

**5.5. Formation of Sialylated *N*-Glycans.** Sialylation of *N*-glycans was reported in Tn-5B1-4 cells when the cells were cultured in the presence of an hexosaminidase inhibitor (2-ADN).<sup>55</sup> This result is particularly intriguing since Tn-TB1-4 cells lack GalT, SiaT, and CMP-NeuAc synthase (see above). Unfortunately, analysis was only by lectin blot and not by quantitative chemical analysis of the exact structures.

Tn-4h is a clonal isolate from Tn-5B1-4<sup>60</sup> and expressed sialylated glycoproteins when cultured in the High Aspect Ratio Vessel (HARV) bioreactor in serum-bearing medium. These findings are especially interesting since closely related cell line (Tn-5B1-4) lack this capability.




FIGURE 6. General strategy for humanization of glycoproteins produced by lepidopteran cell-baculovirus expression system.




FIGURE 7. Changes in *N*-glycan structures on human serum transferrin expressed in Tn-5B1-4 cells by co-infection of a recombinant baculovirus expression vector encoding the gene for a mammalian GalT. ■, *N*-glycans containing nongalactosylated GlcNAc on the Man $\alpha$ (1,3) branch; □, *N*-glycans containing galactosylated GlcNAc on the same branch; gray shading, Paucimannosidic glycans (ref 30).

Sialylation was also detected in virion glycoprotein, gp64, when Sf9 cells were infected with a recombinant baculovirus vector encoding mammalian GalT and  $\alpha$ -2,6-sialyltransferase (SiaT6), while no sialylation was detected in the absence of either GalT or SiaT6.<sup>61</sup> Similar results were observed when Sf $\beta$ 4GalT cells, which constitutively express heterologous GalT, were infected with a recombinant baculovirus encoding SiaT6.<sup>62</sup> Subsequently, two transgenic insect cells, Sf $\beta$ 4GalT/ST6<sup>40</sup> and Tn5 $\beta$ 4GalT/ST6<sup>57</sup> cells, each having foreign genes encoding both mammalian GalT and SiaT6 in their genomic DNA, were shown to express both enzymes constitutively. When these cells were infected (in serum-containing medium) with a wild-type baculovirus or a recombinant baculovirus encoding insect  $\alpha$ -mannosidase-I, both gp64<sup>40,57</sup> and  $\alpha$ -mannosidase-I<sup>40</sup> were found to be sialylated.

Recent study indicated that inclusion of sialylated molecules in serum-free culture medium could increase sialylation.<sup>63</sup> These results suggested the presence of a sialic acid salvaging pathway in lepidopteran cells. The mechanism(s) for increased sialylation is not understood. Sialo-glycoconjugates might have to be taken up by insect cells, and sialic acid might be released in insect cells by sialidase.<sup>64</sup> However, how the CMP-SAS deficient lepidopteran cells<sup>53</sup> can accomplish substantial sialylation is not clear. Alternatively, sialylation might occur by trans-sialylation, which had been known in *Trypanosoma cruzi*.<sup>65</sup>




FIGURE 8. Improvement of CMP-Neu5Ac production by expression of *N*-acetylneuraminic acid 9-phosphate synthase (SAS) and CMP-neuraminic acid synthase (CMP-SAS) in Sf9 cells cultured in the medium with or without 10 mM ManNAc supplementation (ref 53).

**5.6. Synthesis of CMP-Neu5Ac.** The processing steps catalyzed by UDP-*N*-acetylglucosamine 2-epimerase/*N*-acetylmannosamine kinase, *N*-acetylneuraminic acid-9-phosphate synthase, and CMP-Neu5Ac synthase represent bottlenecks in the CMP-Neu5Ac synthesis pathway of lepidopteran cells (Figure 5). To overcome this problem, we cloned mammalian *N*-acetylneuraminic acid-9-phosphate synthase<sup>52</sup> and CMP-Neu5Ac synthase,<sup>53</sup> and expressed these enzymes in Sf9 cells. When Sf9 cells were infected with a recombinant baculovirus expression vector encoding *N*-acetylneuraminic acid-9-phosphate synthase and were cultured in a medium supplemented with *N*-acetylmannosamine (ManNAc), Sf9 cells produced high levels of *N*-acetylneuraminic acid (Neu5Ac).<sup>52</sup> Furthermore, when Sf9 cells, co-infected with two different baculoviruses, each encoding *N*-acetylneuraminic acid-9-phosphate synthase, or CMP-Neu5Ac synthase, were cultured in serum-free medium supplemented with ManNAc, Sf9 cells produced high levels of CMP-Neu5Ac (Figure 8). However, without ManNAc supplement, only ca. 1/20 of that of CHO cells was produced (Figure 8), despite the presence of a substantial amount of UDP-GlcNAc (Figure 4). These results are consistent with the observation that Sf9 cells lack UDP-*N*-acetylglucosamine 2-epimerase activity but have high levels of ManNAc kinase activity.<sup>51</sup> This study demonstrated that the metabolic pathway for synthesis of CMP-Neu5Ac could be enhanced by a combination of overexpression of required enzymes and feeding of Man-

NAC, a precursor of Neu5Ac. Our laboratories have shown recently that the supplementation of ManNAc can be superseded by expressing the UDP-*N*-acetylglucosamine 2-epimerase/*N*-acetylmannosamine kinase gene in combination with other sialylation pathway genes (unpublished data).

## 6. Conclusion and Future Prospects

Many lines of evidence have indicated that the inability of the vast majority of lepidopteran cells to synthesize mammalian-type *N*-glycans. The inability to obtain such *N*-glycans in lepidopteran cells can be attributed to the insufficient levels of GalT, GlcNAcT-II, SiaT, UDP-*N*-acetylglucosamine 2-epimerase/*N*-acetylmannosamine kinase, *N*-acetylneuraminic acid-9-phosphate synthase, and CMP-Neu5Ac synthase activities.  $\beta$ -*N*-Acetylglucosaminidase, which removes GlcNAc on the Man $\alpha$ (1,3) branch, was detected in several lines of lepidopteran cells. This enzyme apparently prevents synthesis of complex-type *N*-glycans by removing the key intermediate glycan containing GlcNAc $\beta$ (1,2)-Man $\alpha$ (1,3). In addition, Fuc-T C3 generates the potentially allergenic glycan structure, Fuc $\alpha$ -(1,3)GlcNAc-Asn, on glycoproteins expressed in lepidopteran cells. Substantial improvement of galactosylation levels has been achieved by expressing mammalian GalT in lepidopteran cells. Galactosylation prevented degradation of *N*-glycans by  $\beta$ -*N*-acetylglucosaminidase and the generation of truncated paucimannosidic glycans. Synthesis of di-branched, galactosylated complex-type *N*-glycans has also been achieved by coexpression of GalT and GlcNAcT-II. Furthermore, the appearance of sialylation in insect cells grown in serum-containing medium has been observed by expressing SiaT in lepidopteran cells in combination with GalT expression. A strategy to engineer the CMP-neuraminic acid synthesis pathway for cells grown in serum-free medium has been established by a combination of genetic engineering and substrate feeding techniques. Expression of glycoproteins having completely sialylated, branched complex-type *N*-glycans in lepidopteran cells grown in serum-free medium was actually accomplished by using a new transgenic insect cell line, SfSWT-3 with ManNAc feeding.<sup>66</sup> SfSWT-3 is a stably transformed insect cell line derived from Sf9 cells, and these cells express mammalian SAS and CMP-SAS in addition to mammalian GlcNAcT-I, GlcNAcT-II, GalT, and SiaT. Structures that are more complex would require inclusion of additional glycosyltransferases (e.g., GlcNAcT-IV and GlcNAcT-V).

In parallel, several analytical methods have been developed to provide for sensitive, high-throughput, and detailed analysis of enzyme activities, glycan structures, and donor sugar nucleotides. Continuous efforts are also underway to establish better analytical techniques for detecting intracellular metabolites and extracellular structures. By combining cellular engineering with efficient analysis of detailed structures, we should be able to characterize and manipulate glycosylation processing pathways in the future for a wide spectrum of applications.

## Abbreviations

CMP-Neu5Ac, cytidine-5'-monopho-*N*-acetylneuraminic acid; Fuc, fucose; Gal, galactose; GalT, galactosyltransferase; GDP-Man, guanosine-5'-diphosphomannose; GlcNAc, *N*-acetylglucosamine; GlcNAc-1-P, *N*-acetylglucosamine-1-phosphate; GlcNAc-6-P, *N*-acetylglucosamine-6-phosphate; Man, mannose; ManNAc, *N*-acetylmannosamine; ManNAc-6-P, *N*-acetylmannosamine-6-phosphate; Neu5Ac, 5-acetamido-3,5-dideoxy-D-glycero-D-galacto-2-nonulosonic acid (*N*-acetylneuraminic acid); Neu5Ac-9-P, *N*-acetylneuraminic acid-9-phosphate; UDP-Gal, uridine-5'-diphogalactose; UDP-GlcNAc, uridine-5'-diphopho-*N*-acetylglucosamine; SiaT, sialyltransferase.

*Preparation of this Account as well as research from our laboratory cited in this review was supported by grants from the National Science Foundation Grant BES9814100 from the Metabolic Engineering Program.*

## References

- (1) Marz, L.; Altmann, F.; Staudacher, E.; Kubelka, V. In *Glycoproteins*; Montreuil, J., Vliegthart, J. F. G., Schachter, H., Eds.; Elsevier: Amsterdam, 1995; Vol. 29a, pp 543–563.
- (2) Altmann, F.; Staudacher, E.; Wilson, I. B.; Marz, L. Insect cells as hosts for the expression of recombinant glycoproteins. *Glycoconjugate J.* **1999**, *16*, 109–123.
- (3) Marchal, I.; Jarvis, D. L.; Cacan, R.; Verbert, A. Glycoproteins from insect cells: sialylated or not? *Biol. Chem.* **2001**, *382*, 151–159.
- (4) Kulakosky, P. C.; Hughes, P. R.; Wood, H. A. N-Linked glycosylation of a baculovirus-expressed recombinant glycoprotein in insect larvae and tissue culture cells. *Glycobiology* **1998**, *8*, 741–745.
- (5) Lopez, M.; Coddeville, B.; Langridge, J.; Plancke, Y.; Sautiere, P.; Chaabihi, H.; Chirat, F.; Harduin-Lepers, A.; Cerutti, M.; Verbert, A.; Delannoy, P. Microheterogeneity of the oligosaccharides carried by the recombinant bovine lactoferrin expressed in *Mamestra brassicae* cells. *Glycobiology* **1997**, *7*, 635–651.
- (6) Voss, T.; Ergulen, E.; Ahorn, H.; Kubelka, V.; Sugiyama, K.; Maurer-Fogy, I.; Gloszl, J. Expression of human interferon omega 1 in Sf9 cells. No evidence for complex-type N-linked glycosylation or sialylation. *Eur. J. Biochem.* **1993**, *217*, 913–919.
- (7) Grabenhorst, E.; Hofer, B.; Nimtz, M.; Jager, V.; Conradt, H. S. Biosynthesis and secretion of human interleukin 2 glycoprotein variants from baculovirus-infected Sf21 cells. Characterization of polypeptides and posttranslational modifications. *Eur. J. Biochem.* **1993**, *215*, 189–197.
- (8) Kubelka, V.; Altmann, F.; Kornfeld, G.; Marz, L. Structures of the N-linked oligosaccharides of the membrane glycoproteins from three lepidopteran cell lines (Sf-21, IZD-Mb-0503, Bm-N). *Arch. Biochem. Biophys.* **1994**, *308*, 148–157.
- (9) Wagner, R.; Liedtke, S.; Kretzschmar, E.; Geyer, H.; Geyer, R.; Klenk, H. D. Elongation of the N-glycans of fowl plague virus hemagglutinin expressed in *Spodoptera frugiperda* (Sf9) cells by coexpression of human  $\beta$ 1,2-N-acetylglucosaminyltransferase I. *Glycobiology* **1996**, *6*, 165–175.
- (10) Ogonah, O. W.; Freedman, R. B.; Jenkins, N.; Patel, K.; Rooney, B. Isolation and characterization of an insect cell line able to perform complex N-linked glycosylation on recombinant proteins. *Bio/Technol.* **1996**, *14*, 197–202.
- (11) Kretzschmar, E.; Geyer, R.; Klenk, H. D. Baculovirus infection does not alter N-glycosylation in *Spodoptera frugiperda* cells. *Biol. Chem. Hoppe-Seyler* **1994**, *375*, 23–27.
- (12) Aeed, P. A.; Elhammer, A. P. Glycosylation of recombinant prorenin in insect cells: the insect cell line Sf9 does not express the mannose 6-phosphate recognition signal. *Biochemistry* **1994**, *33*, 8793–8797.
- (13) Manneberg, M.; Friedlein, A.; Kurth, H.; Lahm, H. W.; Fountoulakis, M. Structural analysis and localization of the carbohydrate moieties of a soluble human interferon gamma receptor produced in baculovirus-infected insect cells. *Protein Sci.* **1994**, *3*, 30–38.
- (14) Hogeland, K. E., Jr.; Deinzer, M. L. Mass spectrometric studies on the N-linked oligosaccharides of baculovirus-expressed mouse interleukin-3. *Biol. Mass Spectrom.* **1994**, *23*, 218–224.

- (15) James, D. C.; Freedman, R. B.; Hoare, M.; Ogonah, O. W.; Rooney, B. C.; Lariouov, O. A.; Dobrovolsky, V. N.; Lagutin, O. V.; Jenkins, N. N-glycosylation of recombinant human interferon-gamma produced in different animal expression systems. *Biotechnology (N.Y.)* **1995**, *13*, 592–596.
- (16) James, D. C.; Goldman, M. H.; Hoare, M.; Jenkins, N.; Oliver, R. W.; Green, B. N.; Freedman, R. B. Posttranslational processing of recombinant human interferon-gamma in animal expression systems. *Protein Sci.* **1996**, *5*, 331–340.
- (17) Kulakosky, P. C.; Shuler, M. L.; Wood, H. A. N-glycosylation of a baculovirus-expressed recombinant glycoprotein in three insect cell lines. *In Vitro Cell Dev. Biol. Anim.* **1998**, *34*, 101–108.
- (18) Hsu, T.-A.; Takahashi, N.; Tsukamoto, Y.; Kato, K.; Shimada, I.; Masuda, K.; Whiteley, E. M.; Fan, J.-Q.; Lee, Y. C.; Betenbaugh, M. J. Differential N-glycan patterns of secreted and intracellular IgG produced in *Trichoplusia ni* cells. *J. Biol. Chem.* **1997**, *272*, 9062–9070.
- (19) Hemmer, W.; Focke, M.; Kolarich, D.; Wilson, I. B.; Altmann, F.; Wohrl, S.; Gotz, M.; Jarisch, R. Antibody binding to venom carbohydrates is a frequent cause for double positivity to honeybee and yellow jacket venom in patients with stinging-insect allergy. *J. Allergy Clin. Immunol.* **2001**, *108*, 1045–1052.
- (20) Wilson, I. B.; Harthill, J. E.; Mullin, N. P.; Ashford, D. A.; Altmann, F. Core alpha1,3-fucose is a key part of the epitope recognized by antibodies reacting against plant N-linked oligosaccharides and is present in a wide variety of plant extracts. *Glycobiology* **1998**, *8*, 651–661.
- (21) Wilson, I. B.; Zeleny, R.; Kolarich, D.; Staudacher, E.; Stroop, C. J.; Kamerling, J. P.; Altmann, F. Analysis of Asn-linked glycans from vegetable foodstuffs: widespread occurrence of Lewis a, core alpha1,3-linked fucose and xylose substitutions. *Glycobiology* **2001**, *11*, 261–274.
- (22) Prenner, C.; Mach, L.; Glossl, J.; Marz, L. The antigenicity of the carbohydrate moiety of an insect glycoprotein, honey-bee (*Apis mellifera*) venom phospholipase A2. The role of alpha 1,3-fucosylation of the asparagine-bound N-acetylglucosamine. *Biochem. J.* **1992**, *284* (Pt. 2), 377–380.
- (23) Tomiya, N.; Lee, Y. C.; Yoshida, T.; Wada, Y.; Awaya, J.; Kurono, M.; Takahashi, N. Calculated two-dimensional sugar map of pyridylaminated oligosaccharides: elucidation of the jack bean alpha-mannosidase digestion pathway of Man9GlcNAc2. *Anal. Biochem.* **1991**, *193*, 90–100.
- (24) Tomiya, N.; Awaya, J.; Kurono, M.; Endo, S.; Arata, Y.; Takahashi, N. Analyses of N-linked oligosaccharides using a two-dimensional mapping technique. *Anal. Biochem.* **1988**, *171*, 73–90.
- (25) Staudacher, E.; Dalik, T.; Wawra, P.; Altmann, F.; Marz, L. Functional purification and characterization of a GDP-fucose: beta-N-acetylglucosamine (Fuc to Asn linked GlcNAc) alpha 1,3-fucosyltransferase from mung beans. *Glycoconjugate J.* **1995**, *12*, 780–786.
- (26) Roitinger, A.; Leiter, H.; Staudacher, E.; Altmann, F. HPLC method for the determination of Fuc to Asn-linked GlcNAc fucosyltransferases. *Glycoconjugate J.* **1998**, *15*, 89–91.
- (27) Fabini, G.; Freilinger, A.; Altmann, F.; Wilson, I. B. Identification of core alpha 1,3-fucosylated glycans and cloning of the requisite fucosyltransferase cDNA from *Drosophila melanogaster*. Potential basis of the neural anti-horseradish peroxidase epitope. *J. Biol. Chem.* **2001**, *276*, 28058–28067.
- (28) Abdur-Rahman, B.; Ailor, E.; Jarvis, D. L.; Betenbaugh, M. J.; Lee, Y. C.  $\beta$ -1,4-Galactosyltransferase Activity in Native and Engineered Insect Cells Measured with Time-Resolved Eu-Fluorescence. *Carbohydr. Res.* **2002**, *337*, 2181–2186.
- (29) Tomiya, N.; Ailor, E.; Lawrence, S. M.; Betenbaugh, M. J.; Lee, Y. C. Determination of nucleotides and sugar nucleotides involved in protein glycosylation by high-performance anion-exchange chromatography: sugar nucleotide contents in cultured insect cells and mammalian cells. *Anal. Biochem.* **2001**, *293*, 129–137.
- (30) Ailor, E.; Takahashi, N.; Tsukamoto, Y.; Masuda, K.; Rahman, B. A.; Jarvis, D. L.; Lee, Y. C.; Betenbaugh, M. J. N-glycan patterns of human transferrin produced in *Trichoplusia ni* insect cells: effects of mammalian galactosyltransferase. *Glycobiology* **2000**, *10*, 837–847.
- (31) Davis, T. R.; Schuler, M. L.; Granados, R. R.; Wood, H. A. Comparison of oligosaccharide processing among various insect cell lines expressing a secreted glycoprotein. *In Vitro Cell. Dev. Biol. Anim.* **1993**, *29A*, 842–846.
- (32) Ren, J.; Bretthauer, R. K.; Castellino, F. J. Purification and properties of a Golgi-derived (alpha 1,2)-mannosidase-I from baculovirus-infected lepidopteran insect cells (IPLB-SF21AE) with preferential activity toward mannosyl-6-N-acetylglucosamine2. *Biochemistry* **1995**, *34*, 2489–2495.
- (33) Kwar, Z.; Herscovics, A.; Jarvis, D. L. Isolation and characterization of an alpha 1,2-mannosidase cDNA from the lepidopteran insect cell line Sf9. *Glycobiology* **1997**, *7*, 433–443.
- (34) Kwar, Z.; Romero, P. A.; Herscovics, A.; Jarvis, D. L. N-Glycan processing by a lepidopteran insect alpha1,2-mannosidase. *Glycobiology* **2000**, *10*, 347–355.
- (35) Velardo, M. A.; Bretthauer, R. K.; Boutaud, A.; Reinhold: B.; Reinhold: V. N.; Castellino, F. J. The presence of UDP-N-acetylglucosamine: Alpha-3-D-mannoside beta 1,2-N-acetylglucosaminyltransferase I activity in *Spodoptera frugiperda* cells (IPLB-SF-21AE) and its enhancement as a result of baculovirus infection. *J. Biol. Chem.* **1993**, *268*, 17902–17907.
- (36) Altmann, F.; Marz, L. Processing of asparagine-linked oligosaccharides in insect cells: evidence for  $\alpha$ -mannosidase II. *Glycoconjugate J.* **1995**, *12*, 150–155.
- (37) Ren, J.; Castellino, F. J.; Bretthauer, R. K. Purification and properties of alpha-mannosidase II from Golgi-like membranes of baculovirus-infected *Spodoptera frugiperda* (IPLB-SF-21 AE) cells. *Biochem. J.* **1997**, *15*, 951–956.
- (38) Hollister, J. R.; Shaper, J. H.; Jarvis, D. L. Stable expression of mammalian beta 1,4-galactosyltransferase extends the N-glycosylation pathway in insect cells. *Glycobiology* **1998**, *8*, 473–480.
- (39) van Die, I.; van Tetering, A.; Bakker, H.; van den Eijnden, D. H.; Joziassse, D. H. Glycosylation in lepidopteran insect cells: identification of a beta 1 $\rightarrow$ 4-N-acetylglucosaminyltransferase involved in the synthesis of complex-type oligosaccharide chains. *Glycobiology* **1996**, *6*, 157–164.
- (40) Hollister, J. R.; Jarvis, D. L. Engineering lepidopteran insect cells for sialoglycoprotein production by genetic transformation with mammalian beta 1,4-galactosyltransferase and alpha 2,6-sialyltransferase genes. *Glycobiology* **2001**, *11*, 1–9.
- (41) Staudacher, E.; Marz, L. Strict order of (Fuc to Asn-linked GlcNAc) fucosyltransferases forming core-difucosylated structures. *Glycoconjugate J.* **1998**, *15*, 355–360.
- (42) Hsu, T. A.; Takahashi, N.; Tsukamoto, Y.; Kato, K.; Shimada, I.; Masuda, K.; Whiteley, E. M.; Fan, J. Q.; Lee, Y. C.; Betenbaugh, M. J. Differential N-glycan patterns of secreted and intracellular IgG produced in *Trichoplusia ni* cells. *J. Biol. Chem.* **1997**, *272*, 9062–9070.
- (43) Takahashi, N.; Tsukamoto, Y.; Shiosaka, S.; Kishi, T.; Hakoshima, T.; Arata, Y.; Yamaguchi, Y.; Kato, K.; Shimada, I. N-glycan structures of murine hippocampus serine protease, neuropsin, produced in *Trichoplusia ni* cells. *Glycoconjugate J.* **1999**, *16*, 405–414.
- (44) Rudd, P. M.; Downing, A. K.; Cadene, M.; Harvey, D. J.; Wormald, M. R.; Weir, I.; Dwek, R. A.; Rifkin, D. B.; Gleizes, P. E. Hybrid and complex glycans are linked to the conserved N-glycosylation site of the third eight-cysteine domain of LTBP-1 in insect cells. *Biochemistry* **2000**, *39*, 1596–1603.
- (45) Staudacher, E.; Kubelka, V.; Marz, L. Distinct N-glycan fucosylation potentials of three lepidopteran cell lines. *Eur. J. Biochem.* **1992**, *207*, 987–993.
- (46) Altmann, F.; Schwihla, H.; Staudacher, E.; Glossl, J.; Marz, L. Insect cells contain an unusual, membrane-bound beta-N-acetylglucosaminidase probably involved in the processing of protein N-glycans. *J. Biol. Chem.* **1995**, *270*, 17344–17349.
- (47) Licari, P. J.; Jarvis, D. L.; Bailey, J. E. Insect cell hosts for baculovirus expression vectors contain endogenous exoglycosidase activity. *Biotechnol. Prog.* **1993**, *9*, 146–152.
- (48) Lopez, M.; Tetaert, D.; Juliant, S.; Gazon, M.; Cerutti, M.; Verbert, A.; Delannoy, P. O-glycosylation potential of lepidopteran insect cell lines. *Biochim. Biophys. Acta* **1999**, *1427*, 49–61.
- (49) Hooker, A. D.; Green, N. H.; Baines, A. J.; Bull, A. T.; Jenkins, N.; Strange, P. G.; James, D. C. Constraints on the transport and glycosylation of recombinant IFN-gamma in Chinese hamster ovary and insect cells. *Biotechnol. Bioeng.* **1999**, *63*, 559–572.
- (50) Hinderlich, S.; Stasche, R.; Zeidler, R.; Reutter, W. A bifunctional enzyme catalyzes the first two steps in N-acetylneuraminic acid biosynthesis of rat liver. Purification and characterization of UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase. *J. Biol. Chem.* **1997**, *272*, 24313–24318.
- (51) Effertz, K.; Hinderlich, S.; Reutter, W. Selective loss of either the epimerase or kinase activity of UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase due to site-directed mutagenesis based on sequence alignments. *J. Biol. Chem.* **1999**, *274*, 28771–28778.
- (52) Lawrence, S. M.; Huddleston, K. A.; Pitts, L. R.; Nguyen, N.; Lee, Y. C.; Vann, W. F.; Coleman, T. A.; Betenbaugh, M. J. Cloning and expression of the human N-acetylneuraminic acid phosphate synthase gene with 2-keto-3-deoxy-D-glycero-D-galacto-nononic acid biosynthetic ability. *J. Biol. Chem.* **2000**, *275*, 17869–17877.
- (53) Lawrence, S. M.; Huddleston, K. A.; Tomiya, N.; Nguyen, N.; Lee, Y. C.; Vann, W. F.; Coleman, T. A.; Betenbaugh, M. J. Cloning and expression of human sialic acid pathway genes to generate CMP-sialic acids in insect cells. *Glycoconj. J.* **2001**, *18*, 205–213.

- (54) Wagner, R.; Geyer, H.; Geyer, R.; Klenk, H.-D. N-acetyl-.beta.-glucosaminidase accounts for differences in glycosylation of influenza virus hemagglutinin expressed in insect cells from a baculovirus vector. *J. Virol.* **1996**, *70*, 4103–4109.
- (55) Watanabe, S.; Kokuho, T.; Takahashi, H.; Takahashi, M.; Kubota, T.; Inumaru, S. Sialylation of N-glycans on the recombinant proteins expressed by a baculovirus-insect cell system under beta-N-acetylglucosaminidase inhibition. *J. Biol. Chem.* **2002**, *277*, 5090–5093.
- (56) Jarvis, D. L.; Finn, E. E. Modifying the insect cell N-glycosylation pathway with immediate early baculovirus expression vectors. *Nat. Biotechnol.* **1996**, *14*, 1288–1292.
- (57) Breitbach, K.; Jarvis, D. L. Improved glycosylation of a foreign protein by Tn-5B1-4 cells engineered to express mammalian glycosyltransferases. *Biotechnol. Bioeng.* **2001**, *74*, 230–239.
- (58) Hollister, J.; Grabenhorst, E.; Nimtz, M.; Conradt, H.; Jarvis, D. L. Engineering the protein N-glycosylation pathway in insect cells for production of biantennary, complex N-glycans. *Biochemistry* **2002**, *41*, 15093–15104.
- (59) Tomiya, N.; Howe, D.; Aumiller, J. J.; Pathak, M.; Park, J.; Palter, K.; Jarvis, D. L.; Betenbaugh, M. J.; Lee, Y. C. Complex-type biantennary N-glycans of recombinant human transferrin from *Trichoplusia ni* insect cells expressing mammalian  $\beta$ -1,4-galactosyltransferase and  $\beta$ -1,2-N-acetylglucosaminyltransferase II. *Glycobiology* **2003**, *13*, 23–34.
- (60) Joshi, L.; Shuler, M. L.; Wood, H. A. Production of a sialylated N-linked glycoprotein in insect cells. *Biotechnol. Prog.* **2001**, *17*, 822–827.
- (61) Jarvis, D. L.; Howe, D.; Aumiller, J. J. Novel baculovirus expression vectors that provide sialylation of recombinant glycoproteins in lepidopteran insect cells. *J. Virol.* **2001**, *75*, 6223–6227.
- (62) Seo, N.-S.; Hollister, J. R.; Jarvis, D. L. Mammalian Glycosyltransferase Expression Allows Sialoglycoprotein Production by Baculovirus-Infected Insect Cells. *Protein Expr. Purif.* **2001**, *22*, 234–241.
- (63) Hollister, J.; Conradt, H.; Jarvis, D. L. Evidence for a Sialic Acid Salvaging Pathway in Lepidopteran Insect Cells. *Glycobiology* **2003**, *13*, (in press).
- (64) Licari, P. J.; Jarvis, D. L.; Bailey, J. E. Insect cell hosts for baculovirus expression vectors contain endogenous exoglycosidase activity. *Biotechnol. Prog.* **1993**, *9*, 146–152.
- (65) Zingales, B.; Carniol, C.; de Lederkremer, R. M.; Colli, W. Direct sialic acid transfer from a protein donor to glycolipids of trypanosome forms of *Trypanosoma cruzi*. *Mol. Biochem. Parasitol.* **1987**, *26*, 135–144.
- (66) Aumiller, J. J.; Hollister, J. R.; Jarvis, D. L. A Transgenic Insect Cell Line Engineered to Produce CMP-Sialic Acid and Sialylated Glycoproteins. *Glycobiology*, in press.

AR020202V