

# Ligand Effects on Dioxygen Activation by Copper and Nickel Complexes: Reactivity and Intermediates

MASATATSU SUZUKI\*

Department of Chemistry, Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan

Received December 21, 2006


## ABSTRACT

Copper and nickel complexes having various active-oxygen species  $M_n-O_2$  ( $n = 1$  or  $2$ ), such as *trans*-( $\mu$ -1,2-peroxo) $Cu^{II}_2$ , bis( $\mu$ -oxo) $M^{III}_2$ , bis( $\mu$ -superoxo) $Ni^{II}_2$ , and ligand-based alkylperoxo- $M^I_m$ , can be produced by a series of tetradentate tripodal ligands (TMPA analogues) containing sterically demanding 6-methyl substituent(s) on the pyridyl group(s), where TMPA = tris(2-pyridylmethyl)amine. Roles of the methyl substituent(s) for the formation of the active-oxygen species and their oxidation reactivities are reported.

## Introduction

Dioxygen-activating nonheme transition-metal centers in metalloenzymes have structural diversity. They use a variety of donor atoms provided by proteins together with exogenous ligands, such as  $H_2O$ ,  $OH^-$ ,  $O^{2-}$ , and need structural flexibility and/or change during the catalytic cycles. To mimic such structural flexibility and/or change, a variety of ligands have been developed. Recent advances in synthetic model chemistry have provided a chemical basis for structures and physicochemical properties of various active-oxygen species  $M_n-O_2$  ( $M = Fe$ ,<sup>1–3</sup>  $Co$ ,<sup>4</sup>  $Ni$ ,<sup>4</sup> and  $Cu$ ,<sup>5–11</sup> and  $n = 1$  or  $2$ ). Although metalloenzymes use mainly manganese, iron, and copper ions, the chemistry of other transition-metal complexes is also important not only to provide an additional chemical basis for understanding the reaction mechanisms of the metalloenzymes but also to develop the artificial oxidation catalysts. Among those metal complexes, nickel and copper ions are capable of producing similar  $M_n-O_2$  species in the oxidation states of II and III as shown in Figure 1. Systematic studies of the structures and reactivities of those nickel and copper complexes depending upon the change in the d electron configurations and the d orbital energies are of particular importance to gain the fundamental basis for dioxygen-activation chemistry mediated by metal complexes. In this Account, we focus on the formation of the nickel and copper complexes having

Masatatsu Suzuki was born in Kanazawa, Japan, in 1948. He received his Ph.D. degree under the supervision of Prof. S. Kida from Kyushu University in 1975. Then, he joined Kyushu University as an Assistant Professor (1975–1983). During that period, he spent a year (1980–1981) as a postdoctoral fellow in Prof. Kazuo Nakamoto's group at Marquette University, Milwaukee, WI. Then, he moved to Kanazawa University and became an Associate Professor in 1988 and a Professor in 1994. His research interest is centered on syntheses of functional models for various dioxygen-activating metalloenzymes.




**FIGURE 1.** Representative  $M_n-O_2$  ( $M = Cu$  and/or  $Ni$ , and  $n = 1$  or  $2$ ) species. The framed species are structurally characterized species derived from TMPA analogues.

active-oxygen species supported by the  $N_4$  tetradentate tripodal ligands, tris(2-pyridylmethyl)amine (TMPA) analogues, containing 6-methyl substituents on the pyridyl groups.

Since Karlin et al. synthesized the first structurally characterized copper–dioxygen complex, *trans*-( $\mu$ -1,2-peroxo)dicopper(II) (Figure 1e), using a tetradentate tripodal ligand TMPA,<sup>12,13</sup> a variety of  $Cu_n-O_2$  complexes have been developed. Kitajima et al. synthesized the first example of the peroxo-dicopper(II) models for the  $O_2$  transport protein (hemocyanin) and aromatic ring oxidation enzyme (tyrosinase), which have a ( $\mu$ - $\eta^2$ : $\eta^2$ -peroxo) coordination mode (Figure 1f), using sterically demanding tridentate Tp derivatives (Tp = hydrotris(pyrazolyl)borate).<sup>5,14,15</sup> Since then, many ( $\mu$ - $\eta^2$ : $\eta^2$ -peroxo)dicopper(II) complexes of bi- and tridentate ligands have been developed.<sup>6–11</sup> Furthermore, Tolman et al. discovered the making and breaking of the dioxygen O–O bond performed by copper complexes; they demonstrated that some ( $\mu$ - $\eta^2$ : $\eta^2$ -peroxo)-dicopper(II) complexes of sterically demanding tridentate  $N,N',N''$ -trisubstituted tacn (tacn = 1,4,7-triazacyclononane) undergo facile interconversion with bis( $\mu$ -oxo)dicopper(III) species (Figure 1g),<sup>16,17</sup> which involves a reversible two-electron redox process. The relative stability between ( $\mu$ - $\eta^2$ : $\eta^2$ -peroxo)dicopper(II) and bis( $\mu$ -oxo)dicopper(III) species depends upon various factors, such as steric and electronic effects as well as solvents, counteranion, temperature, and so on.<sup>18–22</sup> In addition to copper complexes, Hikichi et al. expanded the bis( $\mu$ -oxo)dimetal chemistry to the cobalt and nickel complexes.<sup>4,23,24</sup> They synthesized bis( $\mu$ -oxo) $M^{III}_2$  complexes ( $M = Co$  and  $Ni$ ) of the Tp derivatives in the reaction of bis( $\mu$ -hydroxo) $M^{II}_2$  complexes with  $H_2O_2$ , which is in marked contrast to the formation of the ( $\mu$ - $\eta^2$ : $\eta^2$ -peroxo)dicopper(II) complexes having the same ligand system.<sup>15</sup>

In addition to the bi- and tridentate ligands, a wide variety of  $N_4$  tetradentate tripodal ligands have been

\* To whom correspondence should be addressed. Telephone: +81-76-264-5701. Fax: +81-76-264-5742. E-mail: suzuki@cacheibm.s.kanazawa-u.ac.jp.




**FIGURE 2.** Stereochemical diversity of metal centers and tetradentate tripodal ligands.

developed,<sup>5–10,25,26</sup> which are capable of accommodating the stereochemical diversity of metal complexes as shown in Figure 2. Among them, TMPA is one of the most widely used tetradentate tripodal ligands in inorganic chemistry. As mentioned, Karlin et al. first demonstrated that copper(I) complexes of TMPA and its analogues,  $[\text{Cu}(\text{L})(\text{CH}_3\text{CN})]^+$ , produce *trans*-( $\mu$ -1,2-peroxy)dicopper(II) complexes in a trigonal bipyramidal structure via the formation of end-on superoxo-copper(II) complexes ( $[\text{Cu}(\text{L})(\text{O}_2)]^+$ ).<sup>12,13,27,28</sup> Such tetradentate tripodal ligands tend to form trigonal bipyramidal *trans*-( $\mu$ -1,2-peroxy)dicopper(II) complexes. However, the introduction of the methyl substituent(s) into the 6 position of the pyridyl group(s) can modulate the stereochemical and electronic effects, which influence the stereochemistry and reactivity of metal complexes having active-oxygen species. The 6-methyl groups can play various roles, such as (1) controlling the stereochemistry of metal centers and modulating the donor properties by steric requirement, (2) constructing the hydrophobic cavity around the active-oxygen center, which protects the active-oxygen center from unfavorable decomposition reactions, and (3) acting as an oxidation substrate to probe the oxidation ability of the active-oxygen center. In this Account, we report how stereochemical and electronic effects of a series of TMPA analogues,  $\text{Me}_n\text{-tpa}$  ( $n = 1\text{--}3$ ) and  $\text{Me}_2\text{-etpy}$ , shown in Figure 2, influence the formation and reactivity of the nickel and copper complexes having active-oxygen species ( $\text{M}_n\text{-O}_2$ ). Such systematic studies for the  $\text{Fe}_n\text{-O}_2$  species using  $\text{Me}_n\text{-tpa}$  ligands have also been reported by Que et al.<sup>1,2,29</sup>

## Formation and Characterization of Bis( $\mu$ -oxo)dicopper(III) Complexes

TMPA forms an acetonitrile adduct,  $[\text{Cu}(\text{TMPA})(\text{CH}_3\text{CN})]^+$  ( $\mathbf{1}^{\text{TMPA}}$ ), whereas,  $\text{Me}_2\text{-tpa}$ ,  $\text{Me}_3\text{-tpa}$ , and  $\text{Me}_2\text{-etpy}$  do not form the adducts but form a four-coordinate trigonal pyramidal  $[\text{Cu}(\text{L})]^+$  (Figure 2a).<sup>30–32</sup> It has been shown that, in the metal complexes having more sterically crowded five- and six-coordinate environments, the M–N(6-methylpyridyl) distances are longer than the M–N(pyridyl) distances, owing to the steric interaction between the



**FIGURE 3.** Formation, molecular structure, and decomposition of  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-tpa}}$ ). Atoms are colored by atom type (copper, green; oxygen, red; nitrogen, dark blue; and carbon, gray).

6-methyl group and adjacent donor atom.<sup>29,33,34</sup> However, no significant elongation of the Cu–N(6-methylpyridyl) distances relative to the Cu–N(pyridyl) distances is observed in the present trigonal pyramidal copper(I) complexes, indicating that the 6-methyl group(s) has no significant influence on the steric interaction in the four-coordinate trigonal pyramidal structure.


The reaction of  $[\text{Cu}(\text{Me-tpa})]^+$  ( $\mathbf{1}^{\text{Me-tpa}}$ ) with dioxygen generated a *trans*-( $\mu$ -1,2-peroxy)dicopper(II) complex  $[\text{Cu}_2(\text{O}_2)(\text{Me-tpa})_2]^{2+}$  ( $\mathbf{2}^{\text{Me-tpa}}$ ) similar to the TMPA complex.<sup>35</sup> This peroxy complex decayed rapidly compared to the TMPA complex. Although the  $\nu_{\text{O-O}}$  of  $\mathbf{2}^{\text{Me-tpa}}$  ( $833\text{ cm}^{-1}$ ) is comparable to that of the TMPA complex ( $832\text{ cm}^{-1}$ ), the  $\nu_{\text{Cu-O}}$  ( $550\text{ cm}^{-1}$ ) is significantly lower than that of the TMPA complex ( $561\text{ cm}^{-1}$ ),<sup>36</sup> suggesting some stereochemical change in the  $\text{Me-tpa}$  complex because of the steric interaction between a 6-methylpyridyl group and peroxide, which seem to be responsible in lowering the thermal stability of  $\mathbf{2}^{\text{Me-tpa}}$ . Such observations were also made for some *trans*-( $\mu$ -1,2-peroxy)dicopper(II) complexes of sterically demanding ligands, such as  $\text{Me}_3\text{Bz}_3\text{-tren}$  (tris(*N*-benzyl-*N*-methylaminoethyl)amine).<sup>37</sup>

Unlike  $\mathbf{1}^{\text{Me-tpa}}$ , the introduction of more 6-methyl-2-pyridylmethyl pendant(s) and/or a 2-pyridylethyl pendant significantly influences the reactivity with  $\text{O}_2$ . The copper(I) complex  $[\text{Cu}(\text{Me}_2\text{-tpa})]^+$  ( $\mathbf{1}^{\text{Me}_2\text{-tpa}}$ ) generates a bis( $\mu$ -oxo)dicopper(III) complex  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-tpa}}$ ) at  $\sim -80\text{ }^\circ\text{C}$ , as shown in Figure 3.<sup>30</sup> Although the copper(I) complexes of the  $\text{Me}_2\text{-etpy}$  and  $\text{Me}_3\text{-tpa}$  also generate bis( $\mu$ -oxo)dicopper(III) complexes,  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-etpy})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-etpy}}$ ) and  $[\text{Cu}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_3\text{-tpa}}$ ), the reactions are slow compared to that of  $\mathbf{1}^{\text{Me}_2\text{-tpa}}$  and, during the formation, significant oxidative N-dealkylation of  $\text{Me}_2\text{-etpy}$  occurs for  $\mathbf{3}^{\text{Me}_2\text{-etpy}}$ ,<sup>32</sup> and oxidative N-dealkylation and oxidation of the 6-methyl group of the 6-methyl-2-pyridylmethyl pendant of  $\text{Me}_3\text{-tpa}$  take place for  $\mathbf{3}^{\text{Me}_3\text{-tpa}}$ .<sup>31</sup> In addition, they did not fully oxygenate even at  $-80\text{ }^\circ\text{C}$  under 1 atm of  $\text{O}_2$  (vide infra).

However, the decomposition of  $\mathbf{3}^{\text{Me}_2\text{-etpy}}$  can be significantly suppressed (stable for days at  $-80^\circ\text{C}$ ) using a deuterated ligand  $\text{Me}_2\text{-etpy-}d_4$ , in which four hydrogen atoms of the methylene groups are deuterated, and in the case of the  $\text{Me}_3\text{-tpa}$  complex, a perdeuterated  $\text{Me}_3\text{-tpa-}d_{15}$  was needed for the formation of bis( $\mu$ -oxo) species ( $\mathbf{3}^{\text{Me}_3\text{-tpa-}d_{15}}$ ), which is stable for 1 day at  $-80^\circ\text{C}$ . The results indicate that the decompositions of these complexes involve the C–H bond cleavage of the methylene and/or 6-methyl groups and significant kinetic isotope effects for those C–H bond cleavages.

The crystal structure of the bis( $\mu$ -oxo)  $\text{Me}_2\text{-tpa}$  complex revealed that each copper ion has a square planar structure composed of a  $\text{N}_2\text{O}_2$  donor with two 6-methyl-2-pyridylmethyl pendant arms, which interact weakly with copper ion in the axial positions [2.48(1) and 2.55(1) Å].<sup>30</sup> The average Cu–O (1.803 Å) and  $\text{Cu}\cdots\text{Cu}$  [2.758(4) Å] distances are comparable to those of the bis( $\mu$ -oxo)dicopper(III) complexes of tri- and bidentate ligands, such as  $[\text{Cu}_2(\text{O})_2(\text{Bn}_3\text{-tacn})_2]^{2+}$  (tacn = 1,4,7-triazacyclononane; 1.806 and 2.794 Å)<sup>38</sup> and  $[\text{Cu}_2(\text{O})_2(\text{L}_{\text{ME}})_2]^{2+}$  (1.806 and 2.743 Å) ( $\text{L}_{\text{ME}}$  = *N*-methyl-*N*-ethyl-*N'*-methyl-*N'*-ethyl-1,2-cyclohexanediamine).<sup>39</sup> The crystal structure of  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-etpy-}d_4)_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-etpy-}d_4}$ ) is similar to that of  $\mathbf{3}^{\text{Me}_2\text{-tpa}}$ .<sup>32</sup> Thus, the presence of two 6-methylpyridyl groups prevents the formation of *trans*-( $\mu$ -1,2-peroxo)dicopper(II) species in a trigonal bipyramidal structure because of a steric interaction and generates bis( $\mu$ -oxo)dicopper(III) complexes.

The electronic spectrum of  $\mathbf{3}^{\text{Me}_2\text{-tpa}}$  in dichloromethane at  $-80^\circ\text{C}$  showed intense absorption bands at 258 nm ( $\epsilon = \sim 36\,000\text{ M}^{-1}\text{ cm}^{-1}$ ), with a shoulder at 300 nm, and at 378 nm ( $\epsilon = \sim 19\,000\text{ M}^{-1}\text{ cm}^{-1}$ ) and a weak band at  $\sim 490\text{ nm}$  ( $\epsilon = \sim 330\text{ M}^{-1}\text{ cm}^{-1}$ ).<sup>30</sup> The spectral features of  $[\text{Cu}_2(\text{O})_2(\text{Me}_3\text{-tpa-}d_{15})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_3\text{-tpa-}d_{15}}$ ) and  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-etpy-}d_4)_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-etpy-}d_4}$ ) are similar to that of  $\mathbf{3}^{\text{Me}_2\text{-tpa}}$ .<sup>31,32</sup> The spectral features of these complexes are somewhat different from those of the bis( $\mu$ -oxo)dicopper(III) complexes of the bi- and tridentate ligands, which exhibit two intense absorption bands at  $\sim 300$  and  $\sim 400\text{ nm}$  assigned to the ligand-to-metal charge transfer (LMCT) transitions from the bridging oxides to  $\text{Cu}^{\text{III}}$  centers ( $\pi_\sigma^* \rightarrow d_{xy}$  and  $\sigma^* \rightarrow d_{xy}$ , respectively) by Solomon et al.<sup>40</sup> The absorption bands at  $\sim 300$  and 378–390 nm of the present complexes can also be assigned to the  $\pi_\sigma^* \rightarrow d_{xy}$  and  $\sigma^* \rightarrow d_{xy}$  transitions, respectively. The energies of the latter  $\sigma^* \rightarrow d_{xy}$  transitions of the present complexes are higher than those of the bis( $\mu$ -oxo) $\text{Cu}^{\text{III}}$  complexes of the bi- and tridentate ligands (390–448 nm).<sup>7</sup> This high energy shift seems to be attributable to high denticity of the present complexes, which increases the d orbital energy, although the interaction from the apical positions is weak. The resonance Raman (rR) spectra of  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-tpa}}$ ),  $[\text{Cu}_2(\text{O})_2(\text{Me}_3\text{-tpa-}d_{15})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_3\text{-tpa-}d_{15}}$ ), and  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-etpy-}d_4)_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-etpy-}d_4}$ ) showed the bands at 590–579  $\text{cm}^{-1}$ , assignable to the symmetric breathing mode of the  $\text{Cu}_2\text{O}_2$  core,<sup>40,41</sup> which are almost at the lowest end of those observed for the bis( $\mu$ -oxo)dicopper(III) complexes (630–580  $\text{cm}^{-1}$ ).



**FIGURE 4.** Formation pathway of bis( $\mu$ -oxo) $\text{Ni}^{\text{III}}_2$  (**5**) and bis( $\mu$ -superoxo) $\text{Ni}^{\text{II}}_2$  (**6**) in the reaction of bis( $\mu$ -hydroxo) $\text{Ni}^{\text{II}}_2$  (**4**) with  $\text{H}_2\text{O}_2$  and molecular structures of  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  (**5**<sup>Me<sub>2</sub>-tpa</sup>),  $[\text{Ni}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  (**5**<sup>Me<sub>3</sub>-tpa</sup>), and  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  (**6**<sup>Me<sub>2</sub>-tpa</sup>). Atoms are colored by atom type (nickel, yellow; oxygen, red; nitrogen, dark blue; carbon, gray; and hydrogen, light blue).

## Formation and Characterization of Bis( $\mu$ -oxo)dinickel(III) Complexes

As mentioned already, Hikichi et al. first synthesized bis( $\mu$ -oxo)dinickel(III) complexes of Tp derivatives in the reaction of bis( $\mu$ -hydroxo)dinickel(II) complexes with  $\text{H}_2\text{O}_2$ .<sup>4,23,24</sup> Similar bis( $\mu$ -oxo)dinickel(III) complexes of tridentate ligands having  $\text{N}_3$  and  $\text{S}_3$  (thioether) donors have also been reported.<sup>42–44</sup>  $\text{Me}_2\text{-tpa}$  and  $\text{Me}_3\text{-tpa}$  also produce the bis( $\mu$ -oxo)dinickel(III) complexes,  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  (**5**<sup>Me<sub>2</sub>-tpa</sup>) and  $[\text{Ni}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  (**5**<sup>Me<sub>3</sub>-tpa</sup>), in the reactions of the bis( $\mu$ -hydroxo)dinickel(II) complexes (**4**) with  $\text{H}_2\text{O}_2$ .<sup>45,46</sup>

Unlike the bis( $\mu$ -oxo)dinickel(III) complexes of  $\text{N}_3$  ligands, **5**<sup>Me<sub>2</sub>-tpa</sup> and **5**<sup>Me<sub>3</sub>-tpa</sup> are very reactive with  $\text{H}_2\text{O}_2$  to oxidize  $\text{H}_2\text{O}_2$  to superoxide, which produce bis( $\mu$ -superoxo)dinickel(II) complexes,  $[\text{Ni}_2(\text{O}_2)_2(\text{Me}_2\text{-tpa})_2]^{2+}$  (**6**<sup>Me<sub>2</sub>-tpa</sup>) and  $[\text{Ni}_2(\text{O}_2)_2(\text{Me}_3\text{-tpa})_2]^{2+}$  (**6**<sup>Me<sub>3</sub>-tpa</sup>), as shown in Figure 4. In the case of the  $\text{Me}_2\text{-tpa}$  complex, no **5**<sup>Me<sub>2</sub>-tpa</sup> was observed in the reaction of  $[\text{Ni}_2(\text{OH})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  (**4**<sup>Me<sub>2</sub>-tpa</sup>) with even 0.2 equiv of  $\text{H}_2\text{O}_2$ , indicating that **5**<sup>Me<sub>2</sub>-tpa</sup> is extremely reactive with  $\text{H}_2\text{O}_2$  compared to **5**<sup>Me<sub>3</sub>-tpa</sup>. Such a high reactivity of **5**<sup>Me<sub>2</sub>-tpa</sup> toward  $\text{H}_2\text{O}_2$  seems to be partly attributable to the structural feature of **5**<sup>Me<sub>2</sub>-tpa</sup>, in which there is a space around the bis( $\mu$ -oxo) $\text{Ni}^{\text{III}}_2$  core to which  $\text{H}_2\text{O}_2$  can access, as shown in Figure 4. However, **5**<sup>Me<sub>2</sub>-tpa</sup> can be isolated from the thermal decomposition of **6**<sup>Me<sub>2</sub>-tpa</sup> through disproportionation of the superoxo ligands (Figure 4). It is noted that such high reactivity toward  $\text{H}_2\text{O}_2$  has not been reported for the bis( $\mu$ -oxo)dinickel(III) complexes of  $\text{N}_3$  ligands. The origin of the high reactivity of the present complexes toward  $\text{H}_2\text{O}_2$  compared to those of  $\text{N}_3$  ligands is not known at present.

The crystal structures of **5**<sup>Me<sub>2</sub>-tpa</sup> and **5**<sup>Me<sub>3</sub>-tpa</sup> are similar to those of bis( $\mu$ -oxo)dicopper(III) complexes,<sup>45,46</sup> although the average Ni–N(axial) distance (2.27 Å) is significantly shorter than those in the copper complexes (2.51 Å), in the latter of which copper(III) ion having the  $d^8$  electron configuration tends to adopt a square planar structure. However, the average Ni–N(axial) bond distance is significantly longer than the average Ni–N(equatorial) bond distance (2.004 Å), which is attributable to the

Jahn–Teller effect in a low-spin  $d^7$  electron configuration and partly steric requirement of the 6-methyl substituents. The in-plane 6-methyl substituents of  $\mathbf{5}^{\text{Me}_3\text{-tpa}}$  form a hydrophobic cavity around a  $\text{Ni}^{\text{III}}(\mu\text{-O})_2\text{Ni}^{\text{III}}$  core, which may suppress the oxidation of  $\text{H}_2\text{O}_2$  compared to  $\mathbf{5}^{\text{Me}_2\text{-tpa}}$  as described above. Bis( $\mu$ -superoxo)dinickel(II) complexes of  $\text{Me}_2\text{-tpa}$  and  $\text{Me}_3\text{-tpa}$  have a distorted octahedral structure, in which two Ni centers are linked by two  $\mu$ -1,2-O–O bridges.

$[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{5}^{\text{Me}_2\text{-tpa}}$ ) shows an intense absorption band at 376 nm ( $\epsilon = \sim 6000 \text{ M}^{-1} \text{ cm}^{-1}$ ),<sup>46</sup> and  $[\text{Ni}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  ( $\mathbf{5}^{\text{Me}_3\text{-tpa}}$ ) shows an intense absorption band at 394 nm ( $\epsilon = \sim 4000 \text{ M}^{-1} \text{ cm}^{-1}$ ),<sup>45</sup> assignable to the  $\text{O}^{2-}$ -to- $\text{Ni}^{\text{III}}$  charge-transfer transition [LMCT: ( $\sigma_g + d_{x^2-y^2}(+)$ )  $\rightarrow$  ( $d_{xy}(-) + \sigma_u^*$ )] based on theoretical calculations by Riordan and Brunold et al. for  $[(\text{PhTt}^t\text{Bu})_2\text{Ni}_2(\mu\text{-O})_2]$  having a  $\text{S}_3$  (thioether) tridentate ligand [PhTt<sup>t</sup>Bu = phenyltris(*tert*-butylthio)methyl)borate].<sup>47</sup> Such LMCT transitions have also been observed for the five-coordinate bis( $\mu$ -oxo)dinickel(III) complexes containing tridentate  $\text{N}_3$  donor ligands ( $\lambda_{\text{max}} \sim 405\text{--}414 \text{ nm}$ ).<sup>23,24,42,43</sup> As in the case of the bis( $\mu$ -oxo)dinickel(III) complexes, the LMCT transition energies of the present six-coordinate complexes are higher than those of the five-coordinate complexes, such as  $[\text{Ni}_2(\text{O})_2(\text{Tp}^{\text{Me}_3})_2]$  ( $\lambda_{\text{max}} = 410 \text{ nm}$ ), where  $\text{Tp}^{\text{Me}_3} = \text{hydrotris}(3,4,5\text{-trimethylpyrazolyl})\text{borate}$ .<sup>24</sup> This may also be ascribed to high  $d$  orbital energy because of the increased coordination number from five to six. In addition, the LMCT transition energy of  $\mathbf{5}^{\text{Me}_2\text{-tpa}}$  is higher than that of  $\mathbf{5}^{\text{Me}_3\text{-tpa}}$ , which is also attributable to a stronger electron donation of  $\text{Me}_2\text{-tpa}$  compared to  $\text{Me}_3\text{-tpa}$ .

The rR spectra of  $\mathbf{5}^{\text{Me}_2\text{-tpa}}$  and  $\mathbf{5}^{\text{Me}_3\text{-tpa}}$  showed a characteristic intense band at 574 and 571  $\text{cm}^{-1}$ , respectively, assignable to a symmetric breathing mode of the  $\text{Ni}_2\text{O}_2$  core as observed for the  $\text{Cu}_2\text{O}_2$  cores. These values are lower than those of the bis( $\mu$ -oxo)dinickel(III) complexes of the tridentate nitrogen ligands (599–612  $\text{cm}^{-1}$ ),<sup>42,43</sup> suggesting that the Ni–O bonds for the present six-coordinate complexes  $\mathbf{5}^{\text{Me}_2\text{-tpa}}$  and  $\mathbf{5}^{\text{Me}_3\text{-tpa}}$  are weaker than those of the five-coordinate complexes.


The bis( $\mu$ -superoxo)dinickel(II) complexes  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{6}^{\text{Me}_2\text{-tpa}}$ ) and  $[\text{Ni}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  ( $\mathbf{6}^{\text{Me}_3\text{-tpa}}$ ) exhibit the  $\nu_{\text{O-O}}$  vibration at 1080–1100  $\text{cm}^{-1}$ .<sup>45,46</sup> Their electrospray ionization time of flight/mass spectrometry (ESI-TOF/MS) in acetone suggested the formation of a monomeric superoxo species in the solution state. However, ESR and X-ray absorption spectra of  $\mathbf{6}^{\text{Me}_2\text{-tpa}}$  clearly indicated that the  $\text{Ni}^{\text{II}}(\mu\text{-OO})_2\text{Ni}^{\text{II}}$  core structure remains intact in solution. This is in marked contrast to a monomeric superoxo nickel(II) complex of a sterically bulky tridentate thioether ligand,  $[\text{Ni}(\text{O})_2(\text{PhTt}^{\text{Ad}})]$ , where the superoxo ligand has been proposed to bind to the nickel center in a side-on fashion.<sup>48</sup>

## Reversible Four-Electron Redox Process of $\text{O}_2$ Performed by Copper Complexes

Interconversion between ( $\mu$ - $\eta^2$ : $\eta^2$ -peroxo)dinickel(II) and bis( $\mu$ -oxo)dinickel(III) cores has been well-demonstrated

for the complexes of various bi- and tridentate ligands.<sup>6,7</sup> The interconversion involves the reversible two-electron redox process. The most striking feature of the present complexes of the TMPA analogues having the 6-methyl substituents is the reversible cleavage and formation of the dioxygen O–O bond associated with the four-electron redox process.  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-tpa}}$ ) in dichloromethane at  $-80 \text{ }^\circ\text{C}$  exhibits reversible deoxygenation by bubbling  $\text{N}_2$  to regenerate the copper(I) complex  $\mathbf{1}^{\text{Me}_2\text{-tpa}}$ , as shown in Figure 3.<sup>30</sup> This is the first example of the reversible four-electron redox of dioxygen mediated by the metal complex.

The copper(I) complex  $[\text{Cu}(\text{Me}_2\text{-etpy-}d_4)]^+$  ( $\mathbf{1}^{\text{Me}_2\text{-etpy-}d_4}$ ) was not fully oxygenated in acetone even at  $-80 \text{ }^\circ\text{C}$  under 1 atm of  $\text{O}_2$ .  $^1\text{H}$  nuclear magnetic resonance (NMR) spectrum of  $\mathbf{1}^{\text{Me}_2\text{-etpy-}d_4}$  ( $[\text{Cu}] = \sim 19.0 \text{ mM}$ ) in the above conditions indicated that the formation ratio of  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-etpy-}d_4)_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-etpy-}d_4}$ ) is 70%.<sup>32</sup> A thermodynamic study for the oxygenation ( $2[\text{Cu}(\text{L})]^+ + \text{O}_2 \rightarrow [\text{Cu}_2(\text{O})_2(\text{L})_2]^{2+}$ ) showed that enthalpy and entropy changes are  $-53 + 2 \text{ kJ mol}^{-1}$  and  $-187 \pm 10 \text{ J mol}^{-1} \text{ K}^{-1}$ , respectively, which are significantly unfavorable compared to those of  $[\text{Cu}(\text{AN})]^+$  having a linear aliphatic triamine ligand (AN = 3,3'-iminobis(*N,N*-dimethylpropylamine)) reported by Karlin et al.<sup>21</sup> Further introduction of a 6-methyl group into  $\text{Me}_2\text{-tpa}$  significantly lowers the dioxygen affinity of the copper(I) complex  $[\text{Cu}(\text{Me}_3\text{-tpa-}d_{15})]^+$  ( $\mathbf{1}^{\text{Me}_3\text{-tpa-}d_{15}}$ ).<sup>31</sup> The oxygenation constant ( $K = \sim 3.8 \times 10^3 \text{ M}^{-2}$ ) of  $\mathbf{1}^{\text{Me}_3\text{-tpa-}d_{15}}$  is about 10 times lower than that of  $\mathbf{1}^{\text{Me}_2\text{-etpy-}d_4}$  ( $K = 4.0 \times 10^4 \text{ M}^{-2}$ ) at  $-80 \text{ }^\circ\text{C}$ . Thus, the relative dioxygen affinity of the copper(I) complex is  $\mathbf{1}^{\text{Me}_2\text{-tpa}} > \mathbf{1}^{\text{Me}_2\text{-etpy-}d_4} > \mathbf{1}^{\text{Me}_3\text{-tpa-}d_{15}}$ . This trend is well-correlated with the  $E_{1/2}(\text{Cu}^{\text{II}}/\text{Cu}^{\text{I}})$  values ( $\mathbf{1}^{\text{TMPA}} = -410$ ,  $\mathbf{1}^{\text{Me-tpa}} = -350$ ,  $\mathbf{1}^{\text{Me}_2\text{-tpa}} = -230$ ,  $\mathbf{1}^{\text{Me}_2\text{-etpy}} = -90$ , and  $\mathbf{1}^{\text{Me}_3\text{-tpa}} = -20 \text{ mV}$  versus  $\text{Fc}^+/\text{Fc}$ ).<sup>32</sup> Stepwise introduction of the 6-methyl-2-pyridylmethyl pendant and/or the 2-pyridylethyl pendant causes the positive shift of the  $E_{1/2}(\text{Cu}^{\text{II}}/\text{Cu}^{\text{I}})$  values. Although the  $E_{1/2}(\text{Cu}^{\text{II}}/\text{Cu}^{\text{I}})$  value is not a direct measure of the relative stability of copper(III) species, this seems to be also applicable to the copper(III) species. Thus, unlike bi- and tridentate ligands, the present type of the sterically demanding TMPA analogues preferentially form the bis( $\mu$ -oxo)dinickel(III) complexes and have a unique ability to stabilize both copper(I) and copper(III) oxidation states; they can take not only a square planar structure with weak ligation from the axial positions, which can fit to the copper(III) oxidation state with the  $d^8$  electron configuration, but also a trigonal pyramidal structure suitable for the copper(I) oxidation state. Consequently, they are capable of performing reversible conversion between copper(I) and bis( $\mu$ -oxo)dinickel(III) species involving a reversible four-electron redox process. The coexistence of copper(I) and bis( $\mu$ -oxo)dinickel(III) species implies that no appreciable redox reaction between these two species occurs in these systems. The hydrophobic cavities formed by the 6-methyl substituents of the pyridyl groups seem to be responsible to prevent such a redox reaction.



**FIGURE 5.** Formation pathway of the ligand-based coupling dimer  $[(\text{Me-tpa-CH}_2)_2]$  upon decomposition of  $5^{\text{Me}_2\text{-tpa}}$  under  $\text{N}_2$  and the oxidation products of the 6-methyl group of  $\text{Me}_2\text{-tpa}$  upon decomposition of  $5^{\text{Me}_2\text{-tpa}}$  under  $\text{O}_2$ .

### Oxidation Reactions by Bis( $\mu$ -oxo)dicopper(III) and Bis( $\mu$ -oxo)dinickel(III) Complexes

All of the bis( $\mu$ -oxo)dimetal complexes are reactive toward the supporting ligands as found for many bis( $\mu$ -oxo)dicopper(III) and dinickel(III) complexes. The reactivities highly depend upon the kind of metal ions and the supporting ligands. As mentioned already, the introduction of 6-methyl-2-pyridylmethyl pendant(s) and/or the 2-pyridylethyl pendant significantly suppresses dioxygen affinity, but the oxidation power of the bis( $\mu$ -oxo)dicopper(III) species toward the supporting ligands becomes stronger. As mentioned already,  $[\text{Cu}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  ( $3^{\text{Me}_3\text{-tpa}}$ ) and  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-etpy})_2]^{2+}$  ( $3^{\text{Me}_2\text{-etpy}}$ ) are very reactive toward the supporting ligands compared to  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $3^{\text{Me}_2\text{-tpa}}$ ). The order is  $3^{\text{Me}_3\text{-tpa}} > 3^{\text{Me}_2\text{-etpy}} > 3^{\text{Me}_2\text{-tpa}}$ . Decomposition of  $3^{\text{Me}_2\text{-tpa}}$  and  $3^{\text{Me}_2\text{-etpy}}$  in acetone at low temperatures under  $\text{N}_2$  causes the selective oxidative N-dealkylation of the 6-methyl-2-pyridylmethyl pendant.<sup>30,32</sup> However, decomposition of  $3^{\text{Me}_3\text{-tpa}}$  at  $-80^\circ\text{C}$  under  $\text{O}_2$  caused hydroxylation of one of the 6-methyl groups of  $\text{Me}_3\text{-tpa}$  ( $\text{Me}_2\text{-tpa-CH}_2\text{OH}$ ) and oxidative N-dealkylation of the 6-methyl-2-pyridylmethyl pendant.<sup>31</sup> In addition, a carboxylate complex  $[\text{Cu}(\text{Me}_2\text{-tpa-COO})]^+$ , in which one of the 6-methyl groups of the  $\text{Me}_3\text{-tpa}$  is oxidized to carboxylate, was also isolated.


In contrast to the selective oxidation of the methylene group in  $3^{\text{Me}_2\text{-tpa}}$ , the corresponding bis( $\mu$ -oxo)dinickel(III) complex  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $5^{\text{Me}_2\text{-tpa}}$ ) is capable of oxidizing both methylene and 6-methyl groups.<sup>46</sup> This seems to be attributable to the structural difference between copper and nickel complexes (vide infra). Thermal decomposition of  $5^{\text{Me}_2\text{-tpa}}$  in acetonitrile under  $\text{N}_2$  afforded a coupling dimer of the  $\text{Me}_2\text{-tpa}$  ligand ( $\text{Me-tpa-CH}_2$ )<sub>2</sub>, as shown in Figure 5, together with the oxidative N-dealkylation of the 6-methyl-2-pyridylmethyl pendant. The formation of  $(\text{Me-tpa-CH}_2)_2$  clearly indicates the presence of a ligand-based radical ( $\text{Me-}$

$\text{tpa-CH}_2\bullet$ ), which is stable enough for coupling with another  $\text{Me-tpa-CH}_2\bullet$ . Isotope-labeling experiments using  $5^{\text{Me}_2\text{-tpa}}$  and  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa-}d_1)_2]^{2+}$  ( $5^{\text{Me}_2\text{-tpa-}d_1}$ ) revealed that only an intramolecular ligand coupling occurs. The result strongly suggests the formation of a bis( $\text{Me-tpa-CH}_2\bullet$ )dinickel(II) species generated by stepwise hydrogen-atom abstractions from two methyl groups (Figure 5).


Thermal decomposition of  $5^{\text{Me}_2\text{-tpa}}$  under  $\text{O}_2$  gave no coupling dimer of  $\text{Me}_2\text{-tpa}$  but gave ligand-based alcohol ( $\text{Me-tpa-CH}_2\text{OH}$ ), carboxylate ( $\text{Me-tpa-COO}^-$ ), and a trace amount of aldehyde ( $\text{Me-tpa-CHO}$ ) together with a N-dealkylated ligand. Isotope-labeling experiments using  $^{18}\text{O}_2$  indicate that the oxygen source of the oxidized ligands is  $^{18}\text{O}_2$ , suggesting that the ligand-based radical,  $\text{Me-tpa-CH}_2\bullet$ , is very reactive with  $\text{O}_2$  to produce a peroxy radical ( $\text{Me-tpa-CH}_2\text{OO}\bullet$ ), and this reaction is much faster than the coupling of  $\text{Me-tpa-CH}_2\bullet$  radicals and the oxygen rebound from the dinickel center. Although this conversion seems to involve some complicated reactions, such as radical chain reaction(s) and/or autoxidation, one of the reactive intermediates having a ligand-based alkylperoxide,  $[\text{Ni}_2(\text{OH})(\text{Me}_2\text{-tpa})(\text{Me-tpa-CH}_2\text{OO})]^{2+}$  ( $7^{\text{Me}_2\text{-tpa}}$ ), which can generate the oxidation products ( $\text{Me-tpa-CHO}$ ,  $\text{Me-tpa-COO}^-$ , and  $\text{Me-tpa-CH}_2\text{OH}$ ), was isolated upon decomposition of the bis( $\mu$ -superoxo)dinickel(II) complex  $[\text{Ni}_2(\text{O}_2)_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $6^{\text{Me}_2\text{-tpa}}$ ) under  $\text{O}_2$  at  $-20^\circ\text{C}$ . A further oxidized complex  $[\text{Ni}_2(\text{Me-tpa-CH}_2\text{OO})_2]^{2+}$  ( $8^{\text{Me}_2\text{-tpa}}$ ) was also isolated in the reaction of  $[\text{Ni}_2(\text{OH})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $4^{\text{Me}_2\text{-tpa}}$ ) with a large excess of  $\text{H}_2\text{O}_2$  at  $-40^\circ\text{C}$ .<sup>49</sup>

Although the conversion from the ligand-based peroxy radical complex to the ligand-based alkylperoxo complex seems to involve some complicated reactions, such as radical chain reaction(s), a possible conversion pathway from  $6^{\text{Me}_2\text{-tpa}}$  is given in Figure 6: (1) disproportionation of the superoxo ligands in  $6^{\text{Me}_2\text{-tpa}}$  generates  $\text{O}_2$  and  $5^{\text{Me}_2\text{-tpa}}$  as mentioned already; (2)  $5^{\text{Me}_2\text{-tpa}}$  generates a ligand-based alkyl radical ( $\text{Me-tpa-CH}_2\bullet$ ) by hydrogen-atom abstraction; (3) the resulting  $\text{Me-tpa-CH}_2\bullet$  reacts with  $\text{O}_2$  to produce a ligand-based alkylperoxyl radical ( $\text{Me-tpa-CH}_2\text{OO}\bullet$ ); and (4 and 5) the reaction of  $\text{Me-tpa-CH}_2\text{OO}\bullet$  and a superoxo ligand of  $6^{\text{Me}_2\text{-tpa}}$  and/or hydrogen-atom abstraction of  $\text{Me-tpa-CH}_2\text{OO}\bullet$  afford  $[\text{Ni}_2(\text{OH})(\text{Me}_2\text{-tpa})(\text{Me-tpa-CH}_2\text{OO})]^{2+}$  ( $7^{\text{Me}_2\text{-tpa}}$ ) and  $\text{O}_2$ .

The bis( $\mu$ -alkylperoxo)dinickel(II) complex  $[\text{Ni}_2(\text{Me-tpa-CH}_2\text{OO})_2]^{2+}$  ( $8^{\text{Me}_2\text{-tpa}}$ ) is capable of producing  $\text{Me-tpa-COO}^-$  ( $\sim 62\%$ ) and  $\text{Me-tpa-CH}_2\text{OH}$  ( $\sim 37\%$ ).<sup>49</sup> Decomposition of  $8^{\text{Me}_2\text{-tpa}}$  in the presence of  $\text{H}_2^{18}\text{O}$  revealed that the oxygen atoms of both  $\text{Me-tpa-COO}^-$  and  $\text{Me-tpa-CH}_2\text{OH}$  are scrambled with  $^{18}\text{O}$ , suggesting the presence of some intermediate(s) that can react with water. A possible conversion pathway appears to involve the pathway shown in Figure 7: (1) the O–O bond cleavage of the peroxy ligand produces a ligand-based aldehyde; and (2) disproportionation of the aldehyde gives carboxylate and alkoxide via the Cannizzaro reaction, where the oxygen of aldehyde can be exchanged through acetal. Unusually facile disproportionation of aldehyde to car-



**FIGURE 6.** Possible sequential oxidation pathway of the 6-methyl group of  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $6^{\text{Me}_2\text{-tpa}}$ ) to  $[\text{Ni}_2(\text{OH})(\text{Me}_2\text{-tpa})(\text{Me-tpa-CH}_2\text{OO})]^{2+}$  ( $7^{\text{Me}_2\text{-tpa}}$ ). Atoms are colored by atom type (nickel, yellow; oxygen, red; nitrogen, dark blue; and carbon, gray).



**FIGURE 7.** Possible formation pathway from a ligand-based alkylperoxo complex  $[\text{Ni}_2(\text{OH})(\text{Me}_2\text{-tpa})(\text{Me-tpa-CH}_2\text{OO})]^{2+}$  ( $7^{\text{Me}_2\text{-tpa}}$ ) to alkoxy and carboxylato complexes.

boxylate and alkoxide appears to be attributable to the presence of the 6-methyl-2-pyridylmethyl pendant, which may act as a base to facilitate the Cannizzaro reaction. However, formation of a larger amount of the carboxylate ligand than the alkoxide ligand probably suggests that some other side reaction(s) takes place at the same time.

As in the case of the bis( $\mu$ -oxo)dicopper(III) complexes, the introduction of the 6-methyl-2-pyridylmethyl pendant destabilizes the nickel(III) oxidation state, leading to a stronger oxidant. This is in line with the successive positive shift of the  $E_{1/2}$  (II,III/II,II) values of  $[\text{Ni}_2(\text{OH})_2(\text{Me}_n\text{-tpa})_2]^{2+}$  (450, 550, 640, and 780 mV versus  $\text{Fc}/\text{Fc}^+$  for Tmpa,  $\text{Me-tpa}$ ,  $\text{Me}_2\text{-tpa}$ , and  $\text{Me}_3\text{-tpa}$ , respectively).<sup>46</sup>  $[\text{Ni}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  ( $5^{\text{Me}_3\text{-tpa}}$ ) decomposes within 1 h at  $-40^\circ\text{C}$ , whereas  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $5^{\text{Me}_2\text{-tpa}}$ ) is


stable over 1 day under the same conditions. The high reactivity of  $5^{\text{Me}_3\text{-tpa}}$  may be partly due to the proximity effect between the in-plane methyl groups and the oxo groups and partly due to the higher oxidation power of  $5^{\text{Me}_3\text{-tpa}}$ .

## Correlation between Reactivities and Structures of Bis( $\mu$ -oxo)dicopper(III) and Dinickel(III) Complexes


Both bis( $\mu$ -oxo)dicopper(III) complexes  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $3^{\text{Me}_2\text{-tpa}}$ ) and  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-etpy})_2]^{2+}$  ( $3^{\text{Me}_2\text{-etpy}}$ ) exhibited selective oxidation of the methylene group of the 6-methyl-2-pyridylmethyl pendant in the apical position,<sup>30,32</sup> whereas  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $5^{\text{Me}_2\text{-tpa}}$ ) oxidized both methylene and methyl groups of the 6-methyl-2-pyridylmethyl pendant.<sup>46</sup> The differential reactivities of copper and nickel complexes are closely related to the  $\text{O}_{\text{oxo}} \cdots \text{H}$  distances and their C–H bond energies. The molecular structure of  $3^{\text{Me}_2\text{-tpa}}$  suggests that the  $\text{O}_{\text{oxo}} \cdots \text{H}$  distances for the methyl groups ( $\sim 2.8\text{--}3.3 \text{ \AA}$ ) are slightly longer than those for the methylene groups ( $\sim 2.7 \text{ \AA}$ ), as shown in Figure 8. The short  $\text{O}_{\text{oxo}} \cdots \text{H}$  distances for the methylene groups together with a weaker C–H bond energy seem to be responsible for the selective oxidation of the methylene groups. In contrast, the  $\text{O}_{\text{oxo}} \cdots \text{H}$  distances for the methyl groups in the nickel complex  $5^{\text{Me}_2\text{-tpa}}$  ( $2.3\text{--}3.5 \text{ \AA}$ ) are shorter than those for the methylene groups ( $2.6\text{--}2.7 \text{ \AA}$ ). Such relative  $\text{O}_{\text{oxo}} \cdots \text{H}$  distances seem to be responsible for the parallel oxidation of the methyl and methylene groups in  $5^{\text{Me}_2\text{-tpa}}$ . Thus, the oxidation reactivities are highly dependent upon the d-electron configuration, which can modulate the stereochemistry of the complexes. The relative oxidation powers of the copper complexes toward the supporting ligands are stronger than those of the corresponding nickel complexes. This is in line with the general trend that the nickel complexes can access a higher oxidation state compared to the corresponding copper complex, owing to a higher d-orbital energy of the nickel complexes. Selective oxidation of the methyl group in  $[\text{Ni}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  ( $5^{\text{Me}_3\text{-tpa}}$ ) seems to be due to the proximity effect of the in-plane methyl group in  $5^{\text{Me}_3\text{-tpa}}$  ( $\text{O}_{\text{oxo}} \cdots \text{H} = 2.2\text{--}2.7 \text{ \AA}$ ).

## Formation of the Mononuclear Ligand-Based Alkylperoxo-Copper(II) Complex

Various metal complexes having active-oxygen species have been synthesized in the reaction of bis( $\mu$ -hydroxo)dimetal complexes with ROOH (R = H, alkyl, and acyl) similar to nickel complexes mentioned above. The reaction of  $[\text{Cu}_2(\text{OH})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  in acetonitrile with  $\text{H}_2\text{O}_2$  at  $-40^\circ\text{C}$  produced a ligand-based alkylperoxo complex,  $[\text{Cu}(\text{Me-tpa-CH}_2\text{OO})]^+$  ( $9^{\text{Me}_2\text{-tpa}}$ ), as shown in Figure 9.<sup>50</sup> The reaction of  $[\text{Cu}_2(\text{OH})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  with  $\text{H}_2\text{O}_2$  under  $\text{N}_2$  resulted in the reduction to the copper(I) complex  $[\text{Cu}(\text{Me}_2\text{-tpa})]^+$  ( $1^{\text{Me}_2\text{-tpa}}$ ), where no bis( $\mu$ -oxo)dicopper(III) species  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $3^{\text{Me}_2\text{-tpa}}$ ) was detected.  $1^{\text{Me}_2\text{-tpa}}$  further reacted with  $\text{H}_2\text{O}_2$  to produce a monomeric



**FIGURE 8.** Molecular structures near the bis( $\mu$ -oxo)dicopper cores of  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-tpa}}$ ),<sup>30</sup>  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-etpy-}d_4)_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-etpy-}d_4}$ ),<sup>32</sup>  $[\text{Ni}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{5}^{\text{Me}_2\text{-tpa}}$ ),<sup>46</sup> and  $[\text{Ni}_2(\text{O})_2(\text{Me}_3\text{-tpa})_2]^{2+}$  ( $\mathbf{5}^{\text{Me}_3\text{-tpa}}$ ).<sup>45</sup> Reaction sites are shown by arrows. The H atoms are placed at the calculated positions with the C–H bond = 0.95 Å. Atoms are colored by atom type (copper, green; nickel, yellow; oxygen, red; nitrogen, dark blue; carbon, gray; and hydrogen, light blue).



**FIGURE 9.** Formation of  $[\text{Cu}(\text{Me-tpa-CH}_2\text{OO})]^+$  ( $\mathbf{9}^{\text{Me}_2\text{-tpa}}$ ) and oxidation pathway. Atoms are colored by atom type (copper, green; oxygen, red; nitrogen, dark blue; and carbon, gray).

ligand-based alkylperoxo complex  $\mathbf{9}^{\text{Me}_2\text{-tpa}}$  by the selective oxidation of a methyl group of  $\text{Me}_2\text{-tpa}$ . Although a detailed formation mechanism of  $\mathbf{9}^{\text{Me}_2\text{-tpa}}$  is not known at present, a possible pathway involves a ligand-based radical  $\text{Me-tpa-CH}_2\bullet$  as found for the nickel complex. The reaction of  $\mathbf{1}^{\text{Me}_2\text{-tpa}}$  with  $\text{H}_2\text{O}_2$  may generate an active-oxygen species ( $\text{Cu}^{\text{II}}\text{-OH} + \bullet\text{OH}$  or  $\text{Cu}^{\text{III}}\text{=O} + \text{H}_2\text{O}$ ) by either O–O bond homo- or heterolysis, which is capable of performing the selective hydrogen-atom abstraction from the methyl group to generate the ligand-based radical  $\text{Me-tpa-CH}_2\bullet$ . This is in marked contrast to the reactivity of complex  $[\text{Cu}_2(\text{O})_2(\text{Me}_2\text{-tpa})_2]^{2+}$  ( $\mathbf{3}^{\text{Me}_2\text{-tpa}}$ ), which selectively oxidizes the methylene group of  $\text{Me}_2\text{-tpa}$ . Isotope-labeling experiments for the formation of  $[\text{Cu}(\text{Me-tpa-CH}_2\text{OO})]^+$  ( $\mathbf{9}^{\text{Me}_2\text{-tpa}}$ ) using  $\text{H}_2^{16}\text{O}_2$  under a large excess of  $^{18}\text{O}_2$  afforded both  $[\text{Cu}(\text{Me-tpa-CH}_2^{16}\text{O}^{16}\text{O})]^+$  and  $[\text{Cu}(\text{Me-tpa-CH}_2^{18}\text{O}^{18}\text{O})]^+$ , indicating that  $\text{Me-tpa-CH}_2\bullet$  reacts with not only  $\text{O}_2$  but also a reactive species derived from  $\text{H}_2\text{O}_2$ . Although a further pathway for the formation of  $\mathbf{9}^{\text{Me}_2\text{-tpa}}$  is not known at present, decomposition of  $\mathbf{9}^{\text{Me}_2\text{-tpa}}$  gave the alkoxo- and carboxylato-copper(II) complexes,  $[\text{Cu}(\text{Me-tpa-CO}_2)]^+$  ( $\mathbf{10}$ )

and  $[\text{Cu}(\text{Me-tpa-CH}_2\text{O})]^+$  ( $\mathbf{11}$ ), together with some copper(I) complexes. The formation pathways for  $\mathbf{10}$  and  $\mathbf{11}$  may be similar to those observed for the corresponding nickel complexes (Figure 7) through the Cannizzaro reaction, as shown in Figure 9.

## Concluding Remarks

A variety of nickel and copper complexes having active-oxygen species ( $\text{M}_2\text{-O}_2$ ) can be generated by TMPA analogues that have 6-methyl substituents on the pyridyl groups. In the case of copper complexes,  $\text{Me}_2\text{-tpa}$ ,  $\text{Me}_2\text{-etpy}$ , and  $\text{Me}_3\text{-tpa}$  destabilize the trigonal bipyramidal structure by the steric requirement of the methyl substituents and preferentially generate bis( $\mu$ -oxo)dicopper(III) complexes. They have a unique ability to stabilize copper(I) species in a trigonal pyramidal structure and copper(III) species in a square planar structure with weak ligation from the axial positions. Consequently, they are capable of performing reversible conversion between copper(I) and bis( $\mu$ -oxo)dicopper(III). Furthermore, dioxygen affinity of the copper(I) complexes can be modulated in the order  $\mathbf{1}^{\text{Me}_2\text{-tpa}} > \mathbf{1}^{\text{Me}_2\text{-etpy-}d_4} > \mathbf{1}^{\text{Me}_3\text{-tpa-}d_{15}}$  as successive introduction of the methyl group and/or pyridylethyl group. In contrast, the oxidizing powers of the bis( $\mu$ -oxo) species increase in the reverse order. These trends are well-correlated with the  $E_{1/2}(\text{Cu}^{\text{II}}/\text{Cu}^{\text{I}})$  values by successive introduction of the methyl group.

The hydrophobic cavity formed by the 6-methylpyridyl group(s) seems to play an important role in suppression of some unfavorable decomposition reactions and stabilization of  $\text{M}_n\text{-O}_2$  species. In the equilibrium between the copper(I) and bis( $\mu$ -oxo)dicopper(III) species, the hydrophobic cavity could be responsible for the suppression of the electron transfer between the copper(I) and bis( $\mu$ -oxo)dicopper(III) species. In addition, the hydrophobic cavity of the bis( $\mu$ -oxo)dinickel(III) complexes also plays an important role in the oxidation of  $\text{H}_2\text{O}_2$  by the bis( $\mu$ -oxo)Ni<sup>III</sup><sub>2</sub> center. The stabilization of the  $\text{M}_n\text{-O}_2$  species allows for isolation and structural characterization by X-ray crystallography at low temperatures. Structural information between the active center and substrate (the

methyl and methylene groups of the supporting ligand provides additional understanding of the oxidation reaction.

The methyl group of the present ligands also functions as an oxidation substrate. For the bis( $\mu$ -oxo)dinickel(III) complexes, oxidation of a methyl group affords ligand-based alcohol and carboxylate. The oxidation starts from the hydrogen-atom abstraction by the oxo group of the bis( $\mu$ -oxo)Ni<sup>III</sup><sub>2</sub> center. A series of reaction intermediates and products, such as a ligand-based coupling dimer and a ligand-based alkylperoxo complex, have been detected and/or isolated, which allow for the elucidation of the oxidation mechanism.

Although simple modifications of the supporting ligands presented in this Account provided useful information for dioxygen-activation chemistry mediated by metal complexes, development of further functional groups, such as the reaction cavity, which can bind exogenous oxidation substrates, is needed as a future challenge.

*I thank my co-workers and collaborators listed in the references. Their contributions to this work are greatly appreciated. This work was partly supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture, Japan.*

## References

- (1) Tshuva, E. Y.; Lippard, S. J. Synthetic Models for Non-heme Carboxylate-Bridged Diiron Metalloproteins: Strategies and Tactics. *Chem. Rev.* **2004**, *104*, 987–1012.
- (2) Costas, M.; Mehn, M. P.; Jensen, M. P.; Lawrence Que, L., Jr. Dioxygen Activation at Mononuclear Nonheme Iron Active Sites: Enzymes, Models, and Intermediates. *Chem. Rev.* **2004**, *104*, 939–986.
- (3) Solomon, E. I.; Brunold, T. C.; Davis, M. I.; Kemsley, J. N.; Lee, S.-K.; Lehnert, N.; Neese, F.; Skulan, A. J.; Yang, Y.-S.; Zhou, J. Geometric and Electronic Structure/Function Correlations in Non-heme Iron Enzymes. *Chem. Rev.* **2000**, *100*, 235–349.
- (4) Akita, M.; Hikichi, S. Inorganic Chemistry Based on Tp Ligands—From Dioxygen Complexes to Organometallic Systems. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 1657–1679.
- (5) Kitajima, N.; Moro-oka, Y. Copper–Dioxygen Complexes. Inorganic and Bioinorganic Perspectives. *Chem. Rev.* **1994**, *94*, 737–757.
- (6) Lewis, E. A.; Tolman, W. B. Reactivity of Dioxygen–Copper Systems. *Chem. Rev.* **2004**, *104*, 1047–1076.
- (7) Mirica, L. M.; Ottenwaelder, X.; Stack, T. D. Structure and Spectroscopy of Copper–Dioxygen Complexes. *Chem. Rev.* **2004**, *104*, 1013–1045.
- (8) Kopf, M. A.; Karlin, K. D. Models of Copper Enzymes and Heme–Copper Oxidases. In *Biomimetic Oxidations Catalyzed by Transition Metal Complexes*; Muenier, B., Ed; Imperial College Press: London, U.K., 2000; Chapter 7.
- (9) Hatcher, L. Q.; Karlin, K. D. Oxidant Types in Copper–Dioxygen Chemistry: The Ligand Coordination Defines the Cu<sub>2</sub>–O<sub>2</sub> Structure and Subsequent Reactivity. *J. Biol. Inorg. Chem.* **2004**, *9*, 669–683.
- (10) Solomon, E. I.; Chen, P.; Metz, M.; Lee, S.-K.; Palmer, A. E. Oxygen Binding, Activation, and Reduction to Water by Copper Proteins. *Angew. Chem., Int. Ed.* **2001**, *40*, 4570–4590.
- (11) Itoh, S.; Fukuzumi, S. Dioxygen Activation by Copper Complexes. Mechanistic Insights into Copper Monooxygenases and Copper Oxidases. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 2081–2095.
- (12) Jacobson, R. R.; Tyeklár, Z.; Farooq, A.; Karlin, K. D.; Liu, S.; Zubieta, J. A Cu<sub>2</sub>–O<sub>2</sub> Complex. Crystal Structure and Characterization of a Reversible Dioxygen Binding System. *J. Am. Chem. Soc.* **1988**, *110*, 3690–3692.
- (13) Tyeklár, Z.; Jacobson, R. R.; Wei, N.; Murthy, N. N.; Zubieta, J.; Karlin, K. D. Reversible Reaction of O<sub>2</sub> (and CO) with a Copper(I) Complex. X-ray Structures of Relevant Mononuclear Cu(I) Precursor Adducts and the *trans*-( $\mu$ -1,2-Peroxo)dicopper(II) Product. *J. Am. Chem. Soc.* **1993**, *115*, 2677–2689.
- (14) Kitajima, N.; Fujisawa, K.; Moro-oka, Y.  $\mu$ - $\eta^2$ : $\eta^2$ -Peroxo Binuclear Copper Complex, [Cu(HB(3,5-*i*-Pr<sub>2</sub>Pz<sub>3</sub>)]<sub>2</sub>(O<sub>2</sub>). *J. Am. Chem. Soc.* **1989**, *111*, 8975–8976.
- (15) Kitajima, N.; Fujisawa, K.; Fujimoto, C.; Moro-oka, Y.; Hashimoto, S.; Kitagawa, T.; Toriumi, K.; Tatsumi, K.; Nakamura, A. A New Model for Dioxygen Binding in Hemocyanin. Synthesis, Characterization, and Molecular Structure of the  $\mu$ - $\eta^2$ : $\eta^2$  Peroxo Dinuclear Copper(II) Complexes, [Cu(HB(3,5-*i*-Pr<sub>2</sub>Pz<sub>3</sub>)]<sub>2</sub>(O<sub>2</sub>) (R = *i*-Pr and Ph). *J. Am. Chem. Soc.* **1992**, *114*, 1277–1291.
- (16) Halfen, J. A.; Mahapatra, S.; Wilkinson, E. C.; Kaderli, S.; Young, V. G., Jr.; Que, L., Jr.; Zuberbühler, A. D.; Tolman, W. B. Reversible Cleavage and Formation of the Dioxygen O–O Bond within a Dicopper Complex. *Science* **1996**, *271*, 1397–1400.
- (17) Tolman, W. B. Making and Breaking the Dioxygen O–O Bond: New Insights from Studies of Synthetic Copper Complexes. *Acc. Chem. Res.* **1997**, *30*, 227–237.
- (18) Cahoy, J.; Holland, P. L.; Tolman, W. B. Experimental Studies of the Interconversion of  $\mu$ - $\eta^2$ : $\eta^2$ -Peroxo- and Bis( $\mu$ -oxo)dicopper Complexes. *Inorg. Chem.* **1999**, *38*, 2161–2168.
- (19) Lam, B. M. T.; Halfen, J. A.; Young, V. G., Jr.; Hagadorn, J. R.; Holland, P. L.; Lledós, A.; Cucurull-Sánchez, L.; Novoa, J. J.; Alvarez, S.; Tolman, W. B. Ligand Macrocyclic Structural Effects on Copper–Dioxygen Reactivity. *Inorg. Chem.* **2000**, *39*, 4059–4072.
- (20) Mahadevan, V.; Henson, M. J.; Solomon, E. I.; Stack, T. D. P. Differential Reactivity between Interconvertible Side-On Peroxo and Bis- $\mu$ -oxodicopper Isomers Using Peralkylated Diamine Ligands. *J. Am. Chem. Soc.* **2000**, *122*, 10249–10250.
- (21) Liang, H.-C.; Zhang, C. X.; Henson, M. J.; Sommer, R. D.; Hatwell, K. R.; Kaderli, S.; Zuberbühler, A. D.; Rheingold, A. L.; Solomon, E. I.; Karlin, K. D. Contrasting Copper–Dioxygen Chemistry Arising from Alike Tridentate Alkyltriamine Copper(I) Complexes. *J. Am. Chem. Soc.* **2002**, *124*, 4170–4171.
- (22) Taki, M.; Teramae, S.; Nagatomo, S.; Tachi, Y.; Kitagawa, T.; Itoh, S.; Fukuzumi, S. Fine-Tuning of Copper(I)-Dioxygen Reactivity by 2-(2-Pyridyl)ethylamine Bidentate Ligands. *J. Am. Chem. Soc.* **2002**, *124*, 6367–6377.
- (23) Hikichi, S.; Yoshizawa, M.; Sasakura, Y.; Akita, M.; Moro-oka, Y. First Synthesis and Structural Characterization of Dinuclear M(III) Bis( $\mu$ -oxo) Complexes of Nickel and Cobalt with Hydrotris(pyrazolyl)borate Ligand. *J. Am. Chem. Soc.* **1998**, *120*, 10567–10568.
- (24) Hikichi, S.; Yoshizawa, M.; Sasakura, Y.; Komatsuzaki, H.; Moro-oka, Y.; Akita, M. Structural Characterization and Intramolecular Aliphatic C–H Oxidation Ability of M<sup>III</sup>( $\mu$ -O)<sub>2</sub>M<sup>III</sup> Complexes of Ni and Co with the Hydrotris(3,5-dialkyl-4-X-pyrazolyl)borate Ligands Tp<sup>Me2,X</sup> (X = Me, H, Br) and Tp<sup>Pr2</sup>. *Eur. J. Chem.* **2001**, *23*, 5011–5028.
- (25) Wada, A.; Harata, M.; Hasegawa, K.; Jitsukawa, K.; Masuda, H.; Mukai, M.; Kitagawa, T.; Einaga, H. Structural and Spectroscopic Characterization of a Mononuclear Hydroperoxo-Copper(II) Complex with Tripodal Pyridylamine Ligands. *Angew. Chem., Int. Ed.* **1998**, *37*, 798–799.
- (26) Würtele, C.; Gaoutchenova, E.; Harms, K.; Holthausen, M. C.; Sundermeyer, J.; Schindler, S. Crystallographic Characterization of a Synthetic 1:1 End-On Copper Dioxygen Adduct Complex. *Angew. Chem., Int. Ed.* **2006**, *45*, 3867–3869.
- (27) Karlin, K. D.; Wei, N.; Jung, B.; Kaderli, S.; Niklaus, P.; Zuberbühler, A. D. Kinetics and Thermodynamics of Formation of Copper–Dioxygen Adducts: Oxygenation of Mononuclear Copper(I) Complexes Containing Tripodal Tetradentate Ligands. *J. Am. Chem. Soc.* **1993**, *115*, 9506–9514.
- (28) Karlin, K. D.; Kaderli, S.; Zuberbühler, A. D. Kinetics and Thermodynamics of Copper(I)/Dioxygen Interaction. *Acc. Chem. Res.* **1997**, *30*, 139–147.
- (29) Zang, Y.; Kim, J.; Dong, Y.; Wilkinson, E. C.; Appelman, E. H., Jr. Models for Nonheme Iron Intermediates: Structural Basis for Tuning the Spin States of Fe(TPA) Complexes. *J. Am. Chem. Soc.* **1997**, *119*, 4197–4205.
- (30) Hayashi, H.; Fujinami, S.; Nagatomo, S.; Ogo, S.; Suzuki, M.; Uehara, A.; Watanabe, Y.; Kitagawa, T. A Bis( $\mu$ -oxo)dicopper(III) Complex with Aromatic Nitrogen Donors: Structural Characterization and Reversible Conversion between Copper(I) and Bis( $\mu$ -oxo)dicopper(III) Species. *J. Am. Chem. Soc.* **2000**, *122*, 2124–2125.
- (31) Hayashi, H.; Uozumi, K.; Fujinami, S.; Nagatomo, S.; Shiren, K.; Furutachi, H.; Suzuki, M.; Uehara, A.; Kitagawa, T. Modulation of the Copper–Dioxygen Reactivity by Stereochemical Effect of Tetradentate Tripodal Ligands. *Chem. Lett.* **2002**, 416–417.
- (32) Mizuno, M.; Hayashi, H.; Fujinami, S.; Furutachi, H.; Nagatomo, S.; Otake, S.; Uozumi, K.; Suzuki, M.; Kitagawa, T. Ligand Effect on Reversible Conversion between Copper(I) and Bis( $\mu$ -oxo)dicopper(III) Complex with a Sterically Hindered Tetradentate Tripodal Ligand and Monooxygenase Activity of Bis( $\mu$ -oxo)dicopper(III) Complex. *Inorg. Chem.* **2003**, *42*, 8534–8544.
- (33) Goodson, P. A.; Oki, A. R.; Glerup, J.; Hodgson, D. J. Design, Synthesis, and Characterization of Bis( $\mu$ -oxo)dimanganese(III,III) Complexes. Steric and Electronic Control of Redox Potentials. *J. Am. Chem. Soc.* **1990**, *112*, 6248–6254.



- (34) Nagao, H.; Komeda, N.; Mukaida, M.; Suzuki, M.; Tanaka, K. Structural and Electrochemical Comparison of Copper(II) Complexes with Tripodal Ligands. *Inorg. Chem.* **1996**, *35*, 6809–6815.
- (35) Uozumi, K.; Hayashi, Y.; Suzuki, M.; Uehara, A. Reactivity of Copper(I) Complexes Containing Various Tetradentate Tripodal Ligands with Molecular Oxygen. *Chem. Lett.* **1993**, 963–966.
- (36) Baldwin, M. J.; Ross, P. K.; Pate, J. E.; Tyeklár, Z.; Karlin, K. D.; Solomon, E. I. Spectroscopic and Theoretical Studies of an End-On Peroxide-Bridged Coupled Binuclear Copper(II) Model Complex of Relevance to the Active Sites in Hemocyanin and Tyrosinase. *J. Am. Chem. Soc.* **1991**, *113*, 8671–8679.
- (37) Komiyama, K.; Furutachi, H.; Nagatomo, S.; Hashimoto, A.; Hayashi, H.; Fujinami, S.; Suzuki, M.; Kitagawa, T. Dioxygen Reactivity of Copper(I) Complexes with Tetradentate Tripodal Ligands Having Aliphatic Nitrogen Donors: Synthesis, Structures, and Properties of Peroxo and Superoxo Complexes. *Bull. Chem. Soc. Jpn.* **2004**, *77*, 59–72.
- (38) Mahapatra, S.; Halfen, J. A.; Wilkinson, E. C.; Pan, G.; Wang, X.; Young, V. G., Jr.; Cramer, C. J.; Que, L., Jr.; Tolman, W. B. Structural, Spectroscopic, and Theoretical Characterization of Bis( $\mu$ -oxo)dicopper Complexes, Novel Intermediates in Copper-Mediated Dioxygen Activation. *J. Am. Chem. Soc.* **1996**, *118*, 11555–11574.
- (39) Mahadevan, V.; Hou, Z.; Cole, A. P.; Root, D. E.; Lal, T. K.; Solomon, E. I.; Stack, T. D. P. Irreversible Reduction of Dioxygen by Simple Peralkylated Diamine–Copper(I) Complexes: Characterization and Thermal Stability of a  $[\text{Cu}_2(\mu\text{-O})_2]^{2+}$  Core. *J. Am. Chem. Soc.* **1997**, *119*, 11996–11997.
- (40) Henson, M. J.; Mukherjee, P.; Root, D. E.; Stack, T. D. P.; Solomon, E. I. Spectroscopic and Electronic Structural Studies of the  $\text{Cu(III)}_2$  Bis- $\mu$ -oxo Core and Its Relation to the Side-On Peroxo-Bridged Dimer. *J. Am. Chem. Soc.* **1999**, *121*, 10332–10345.
- (41) Holland, P. L.; Cramer, C. J.; Wilkinson, E. C.; Mahapatra, S.; Rodgers, K. R.; Itoh, S.; Taki, M.; Fukuzumi, S.; Que, L., Jr.; Tolman, W. B. Resonance Raman Spectroscopy as a Probe of the Bis( $\mu$ -oxo)dicopper Core. *J. Am. Chem. Soc.* **2000**, *122*, 792–802.
- (42) Itoh, S.; Bandoh, H.; Nagatomo, S.; Kitagawa, T.; Fukuzumi, S. Aliphatic Hydroxylation by a Bis( $\mu$ -oxo)dinickel(III) Complex. *J. Am. Chem. Soc.* **1999**, *121*, 8945–8946.
- (43) Itoh, S.; Bandoh, H.; Nakagawa, M.; Nagatomo, S.; Kitagawa, T.; Karlin, K. D.; Fukuzumi, S. Formation, Characterization, and Reactivity of Bis( $\mu$ -oxo)dinickel(III) Complexes Supported by A Series of Bis[2-(2-pyridyl)ethyl]amine Ligands. *J. Am. Chem. Soc.* **2001**, *123*, 11168–11178.
- (44) Mandimutsira, B. S.; Yamarik, J. L.; Brunold, T. C.; Gu, W.; Cramer, S. P.; Riordan, C. G. Dioxygen Activation by a Nickel Thioether Complex: Characterization of a  $\text{Ni}^{\text{III}}_2(\mu\text{-O})_2$  Core. *J. Am. Chem. Soc.* **2001**, *123*, 9194–9195.
- (45) Shiren, K.; Ogo, S.; Fujinami, S.; Hayashi, H.; Suzuki, M.; Uehara, A.; Watanabe, Y.; Moro-oka, Y. Synthesis, Structures, and Properties of Bis( $\mu$ -oxo)nickel(III) and Bis( $\mu$ -superoxo)nickel(II) Complexes: An Unusual Conversion of a  $\text{Ni}^{\text{III}}_2(\mu\text{-O})_2$  Core into a  $\text{Ni}^{\text{II}}_2(\mu\text{-OO})_2$  Core by  $\text{H}_2\text{O}_2$  and Oxygenation of Ligand. *J. Am. Chem. Soc.* **2000**, *122*, 254–262.
- (46) Cho, J.; Furutachi, H.; Fujinami, S.; Ohtsu, H.; Tosha, T.; Ikeda, O.; Suzuki, A.; Nomura, M.; Uruga, T.; Tanida, H.; Kawai, T.; Tanaka, K.; Kitagawa, T.; Suzuki, M. Sequential Reaction Intermediates in Aliphatic C–H Bond Functionalization Initiated by a Bis( $\mu$ -oxo)dinickel(III) Complex. *Inorg. Chem.* **2006**, *45*, 2873–2885.
- (47) Schenker, R.; Mandimutsira, B. S.; Riordan, C. G.; Brunold, T. C. Spectroscopic and Computational Studies on  $[(\text{PhTt}^{\text{tBu}})_2\text{Ni}_2(\mu\text{-O})_2]$ : Nature of the Bis- $\mu$ -oxo ( $\text{Ni}^{3+}$ )<sub>2</sub> “Diamond” Core. *J. Am. Chem. Soc.* **2002**, *124*, 13842–13855.
- (48) Fujita, K.; Schenker, R.; Gu, W.; Brunold, T. C.; Cramer, S. P.; Riordan, C. G. A Monomeric Nickel–Dioxygen Adduct Derived from a Nickel(I) Complex and  $\text{O}_2$ . *Inorg. Chem.* **2004**, *43*, 3324–3326.
- (49) Cho, J.; Furutachi, H.; Fujinami, S.; Suzuki, M. A Bis( $\mu$ -alkylperoxo)dinickel(II) Complex as a Reaction Intermediate for the Oxidation of the Methyl Groups of the  $\text{Me}_2\text{-tpa}$  Ligand to Carboxylate and Alkoxide Ligands. *Angew. Chem., Int. Ed.* **2004**, *43*, 3300–3303.
- (50) Mizuno, M.; Honda, K.; Cho, J.; Furutachi, H.; Tosha, T.; Matsumoto, T.; Fujinami, S.; Kitagawa, T.; Suzuki, M. A Mononuclear Alkylperoxocopper(II) Complex as a Reaction Intermediate in the Oxidation of the Methyl Group of the Supporting Ligand. *Angew. Chem., Int. Ed.* **2006**, *45*, 6911–6914.

AR600048G