

Understanding How the Thiolate Sulfur Contributes to the Function of the Non-Heme Iron Enzyme Superoxide Reductase

JULIE A. KOVACS* AND LISA M. BRINES

The Department of Chemistry, University of Washington, Box 351700, Seattle, Washington 98195-1700

Received December 21, 2006



ABSTRACT

Toxic superoxide radicals, generated via adventitious reduction of dioxygen, have been implicated in a number of disease states. The cysteine-ligated non-heme iron enzyme superoxide reductase (SOR) degrades superoxide via reduction. Biomimetic analogues which provide insight into why nature utilizes a *trans*-thiolate to promote SOR function are described. Spectroscopic and/or structural characterization of the first examples of thiolate-ligated Fe^{III}-peroxo complexes provides important benchmark parameters for the identification of biological intermediates. Oxidative addition of superoxide is favored by low redox potentials. The *trans* influence of the thiolate appears to significantly weaken the Fe–O peroxo bond, favoring proton-induced release of H₂O₂ from a high-spin Fe(III)–OOH complex.

Introduction

The bioinorganic chemistry of iron is rich and diverse,^{1–6} catalyzing reactions that range from the biosynthesis of neurotransmitters⁷ to the transport of O₂.⁸ A number of iron-promoted biosynthetic pathways involve dioxygen as the oxidant and/or oxygen atom source.^{1,5–7} Although O₂ oxidations are thermodynamically favored, they are kinetically slow because they are spin-forbidden, unless promoted by a transition metal, such as Fe²⁺.^{2,4} The intermediates formed in these reactions include iron–peroxo and oxo species,^{1,2,4,6} potent oxidants capable of functionalizing alkanes.⁶ Adventitious reduction of dioxygen can, on the other hand, result in the formation of toxic radicals, including superoxide (O₂^{•−}) which has been implicated in a number of disease states, including Alzheimer's, Parkinson's, and cancer.⁹ Due to its toxicity, organisms have evolved elaborate means for the degradation of O₂^{•−}. In aerobic organisms, manganese-, nickel-,

iron-, or copper- and zinc-containing enzymes known as superoxide dismutases (SODs) function to disproportionate (eq 2) adventitiously formed superoxide.^{10,11} In



anaerobic organisms, an iron-containing enzyme superoxide reductase (SOR) reduces (eq 1) superoxide;^{9,12–15} however, this requires an outside source of electrons. A cysteine sulfur bound to the iron site, as well as the positioning of the metal ion on the surface (vs the interior) of the protein,¹² alters the function of Fe-SOR relative to Fe-SOD.¹¹

Superoxide Reductase Enzyme Active Site Structure and Mechanism

Superoxide reductases (SORs) are intense blue, cysteine-ligated non-heme iron enzymes found in anaerobic microbes.^{3,12–14} The Fe^{II} active site is redox active, high-spin (*S* = 2), and ligated by four equatorial histidines and one apical cysteine *trans* to an open site.¹² The oxidized Fe^{III} resting state is high-spin (*S* = 5/2) and contains a glutamate (¹⁴Glu) coordinated to the sixth (axial) site.¹² This conserved ¹⁴Glu, as well as a highly conserved lysine (⁴⁷Lys), is proposed to be involved in the catalytic mechanism, possibly acting as a proton donor, and/or by attracting the anionic superoxide ion to the active site.^{12,16} Activity drops significantly in mutants lacking ⁴⁷Lys.¹⁶ The SOR-catalyzed conversion of superoxide (O₂^{•−}) to hydrogen peroxide (H₂O₂) requires one electron and two protons. Protons and solvent both play an important role in the SOR mechanism.¹⁴ Superoxide reduction is favored in the presence of a proton source [*E*_{1/2}(O₂^{•−}/H₂O₂) = 0.83 V at pH 7.5 vs −0.041 V at pH 14 (vs SCE)], whereas oxidation (to afford O₂) is favored in the absence of a proton source [*E*_{1/2}(O₂^{•−}/O₂) = −0.80 V at pH 14 vs −0.13 V at pH 0 (vs SCE)]. Our working hypothesis is that the unusual positioning of the metal ion on the surface of SOR provides the active site with a readily available source of protons which makes O₂ formation disfavored in anaerobic bacteria.

The mechanism by which SOR is proposed to reduce superoxide has been somewhat controversial, particularly with regard to the number of intermediates involved.^{14,15,17} The first step is generally agreed to involve the oxidative addition of O₂^{•−} to the open coordination site. Two transient intermediates (T₁ and T₂ in Figure 1) are observed in the reaction between SOR and O₂^{•−},¹⁵ and exogenous ligands (NO, N₃[−], and CN[−]) have been shown to bind to the SOR iron site,^{18,19} consistent with an inner-sphere mechanism. The first intermediate, T₁, forms at nearly diffusion controlled rates^{15,17} and is proposed to be an Fe^{III}-peroxo species.¹⁷ The second intermediate, T₂, forms more slowly and displays a pH-dependent *v*_{Fe–O} stretch that shifts with ¹⁸OH₂ and D₂O,²⁰ consistent with its assignment as an Fe^{III}–OH species.¹⁵ This Fe^{III}–OH

Julie Kovacs was born in Lansing, MI, on March 15, 1959, and received her B.S. degree from Michigan State University (where her father was on the faculty as a theoretical physicist) in 1981. She received her Ph.D. from Harvard University in 1986, where she synthesized the first analogue of the VFeS cluster site of nitrogenase with Richard Holm. She did her postdoctoral work at the University of California, Berkeley, with Bob Bergman and has been on the faculty at the University of Washington since 1988, where she is currently a full Professor. She currently serves as Chair-elect of the bioinorganic ACS subdivision and will Chair the 2008 Metals in Biology Gordon conference. Her research focuses on understanding how sulfur ligands influence function in metalloenzymes.

Lisa M. Brines was born in Spokane, WA, in 1980. She received a Bachelor of Science in Chemistry from Seattle University in 2002 and is currently a graduate student at the University of Washington. Her research in Julie Kovacs' lab involves studying the active sites of non-heme metalloenzymes using synthetic modeling techniques.

* Corresponding author.




FIGURE 1. Proposed mechanism for SOR-catalyzed reduction of superoxide via hydroperoxo (T₁) and solvent-bound (T₂) intermediates.

species presumably forms via H₂O-induced protonation of the proximal oxygen of the Fe^{III}-peroxo species, followed by H₂O₂ release and OH⁻ binding.^{15,20} Why protonation preferentially occurs at the proximal, as opposed to distal, oxygen is unclear at this point. Reduction of the Fe^{III}-OH intermediate in the presence of a proton regenerates the catalytically active five-coordinate Fe(II) form. The glutamate-bound resting state (R) shown in Figure 1 eventually forms in the absence of additional substrate (O₂⁻) or electrons. Although there is currently no evidence to suggest that this last step involves a proton, the difficulty with which Fe-OH bonds cleave relative to Fe-OH₂ bonds would imply that protonation occurs prior to ligand dissociation. The exogenous electron donor is currently unknown. Given that the addition of a proton and electron would be equivalent to the addition of a hydrogen atom, this introduces the interesting (albeit remote) possibility that the last step of the SOR mechanism may, perhaps, resemble the Fe(III)-OH intermediate-induced H-atom abstraction step of lipoxygenases.²¹⁻²³

Vibrational data to support the assignment of T₁ as a peroxo species have yet to be reported. However, when H₂O₂ is added to a mutant form of SOR,¹³ isotopically sensitive $\nu_{\text{O-O}}$ (850 cm⁻¹) and $\nu_{\text{Fe-O}}$ (438 cm⁻¹) stretches are observed by resonance Raman, consistent with the formation of a metal-peroxo species. This mutant peroxo species was originally proposed to be $\eta^2\text{-O}_2^{2-}$ side-on bound,¹³ on the basis of data reported for N-ligated synthetic non-heme iron-peroxo complexes,²⁴ but later was shown by X-ray crystallography to be an end-on Fe(III)-OOH species,²⁵ revealing a need for more thiolate-bound iron-peroxo complexes to provide benchmark vibrational parameters for interpreting biophysical data.

Relationship between SOR and P450 Active Sites and Mechanism

The primary coordination sphere of the iron active site of SOR is structurally¹² related to that of the heme enzyme cytochrome P450.^{26,27} Both enzymes have an apical cysteine *trans* to an open coordination site (Figure 2), and both react with oxygen-derived substrates to afford an Fe(III)-OOH intermediate. The mechanistic pathways taken following formation of the Fe(III)-peroxo intermedi-

ate are very different for these two enzymes, however. With P450, the hydroperoxo O-O bond is cleaved to afford a high-valent iron-oxo intermediate,^{6,27,28} whereas with SOR, the Fe-O(peroxo) bond is cleaved, releasing H₂O₂. The cysteine has been shown to play an important role in promoting P450-induced O-O bond cleavage,²⁹ yet O-O bond cleavage has not been observed with SOR. It is not clear why these two structurally similar systems follow divergent reaction pathways. Although a porphyrin was originally thought to be necessary for the stabilization of a high-valent iron-oxo complex, this was later shown not to be the case upon structural characterization of the first non-heme Fe(IV)=O complex.³⁰ Although theoretical calculations have shown that the site of protonation can influence the reaction pathway,²⁷ controlled, site-specific protonation with the metal ion sitting on the surface of the protein seems unlikely. Prior to the development of spectroscopic methods for probing non-heme iron,³¹ less was known about non-heme iron, relative to heme iron, mechanisms. Solomon has shown that Fe-peroxo spin states can influence the energetically preferred O-O versus Fe-O bond cleaving pathways.^{2,31} We have shown, in collaboration with Solomon, that the intense S → Fe(III) charge transfer bands characteristic of Fe(III)-SR compounds make it convenient to detect Cys-ligated non-heme iron sites and probe their reactivity.^{32,33}

Roles of Cysteinate Residues in Promoting Metalloenzyme Function

A comprehensive understanding of the influence of thiolate ligands on the properties of first-row transition metal ions is essential if we are to fully understand why nature utilizes cysteine residues to promote specific biological metalloenzyme functions.^{3,34} Cysteinate-ligated metalloenzymes promote a number of critical biological processes, including electron transfer,^{35,36} and strong bond activation.⁶ Cysteines form highly covalent bonds to transition metals, and this helps to facilitate redox changes.³⁷ Our work has shown that thiolate ligands significantly lower redox potentials,³⁸ make low-spin iron accessible in a non-heme environment,³⁹⁻⁴¹ stabilize iron in the +3 oxidation state,³⁹⁻⁴¹ and labilize sites *trans* to the thiolate, thereby promoting product release,³⁸ even with the typically inert low-spin Co(III) ion.^{42,43} Others have shown that the *trans* coordinated cysteine of P450 promotes O-O bond cleavage,^{6,29} and the subsequent Fe(IV)=O-promoted H-atom abstraction.⁴⁴

Biomimetic Models

A precise description of the correlation among the structure, key properties, and function of metalloenzyme active sites can be most readily obtained by building small molecular analogues.³⁴ Synthetic model complexes provide key parameters needed to fit spectroscopic data (e.g., EXAFS) and protein crystal structures.⁴⁵ Multidentate ligands are generally required for maintenance of a relatively rigid, well-defined synthetic active site model. The incorporation of thiolate ligands into synthetic models




FIGURE 2. Comparison of the thiolate-ligated non-heme, and heme, iron active sites of SOR (left) and P450 (right).




FIGURE 3. ORTEP diagram of thiolate-ligated, five-coordinate $[\text{Fe}^{\text{III}}(\text{S}_2^{\text{Me}_2}\text{N}_3(\text{Pr},\text{Pr}))]^+$ (**1**) synthesized by Kovacs group postdocs Steve Shoner and Jeff Ellison.

FIGURE 4. ORTEP diagram of thiolate-ligated, biomimetic superoxide reducing catalyst $[\text{Fe}^{\text{II}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))]^+$ (**2**) synthesized by Jason Shearer, a graduate student in the Kovacs lab.

can, however, be complicated by their propensity to oligomerize, as well as autoreduce and form disulfides,⁴⁶ especially when there is an open coordination site. Peroxides tend to react with thiolates to form sulfoxides and sulfones^{47,48} and with iron to form rust. Despite these synthetic challenges, much progress has been made in the biomimetic modeling of cysteine-ligated non-heme iron active sites in biology.^{32,38,40–42,48–57}

Reactive Five-Coordinate, Thiolate-Ligated Iron Complexes

A biomimetic system capable of reproducing the SOR reaction (Figure 1) would have two requirements. First, it would require an open coordination site for superoxide to bind to the metal. Second, it would require that the +3 oxidation state be reversibly accessible. Given the challenges associated with the synthesis of mononuclear thiolate-ligated transition metal complexes, especially those with higher oxidation states, we initially had to demonstrate that molecules meeting the criteria listed above could be synthesized. By incorporating gem-dimethyls adjacent to the thiolate, we were able to isolate a number of coordinatively unsaturated, mononuclear five-coordinate thiolate iron complexes, including $[\text{Fe}^{\text{III}}(\text{S}_2^{\text{Me}_2}\text{N}_3(\text{Pr},\text{Pr}))]^+$ (**1**) (Figure 3),⁴¹ $[\text{Fe}^{\text{III}}(\text{S}_2^{\text{Me}_2}\text{N}_3(\text{Et},\text{Pr}))]^+$,⁵⁸ and $[\text{Fe}^{\text{II}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))]^+$ (**2**) (Figure 4).⁵⁹ We found that we could isolate an $\text{Fe}^{\text{III}}\text{-SR}$ complex such as **1** via the in situ oxidation of an $\text{Fe}^{\text{II}}\text{-SR}$ precursor at low temperatures.^{41,58} Once oxidized, the Fe^{III} thiolates were

found to display rich spectroscopic features, including intense πS -to-metal charge transfer bands in the visible region,³² and low-spin EPR signals.^{33,40} The energy of these bands was found to be highly dependent on the local coordination environment and could therefore be used to monitor reactions.^{38,41,43,50,52–54} The $\text{Fe}^{\text{III}}\text{-SR}$ bonds were found to be highly covalent and favor a low-spin state, an observation that was surprising given the π -donor properties of thiolates.³²

A Functional SOR Model with a *cis*-Thiolate: $[\text{Fe}^{\text{II}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))]^+$

Using a tripodal amine ligand as a scaffold, we were able to assemble a functional SOR model, $[\text{Fe}^{\text{II}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))]^+$ (**2**),⁵⁹ that mimics each step of the proposed SOR mechanism (Figure 1). Like the SOR enzyme active site, complex **2** is high-spin ($S = 2$) and has a similar N_4S^1 ligand motif, but with the open site *cis* to the thiolate, rather than *trans* as in the enzyme. When superoxide is added to **2** in the presence of a proton donor, biomimetic activity is observed, resulting in the formation of H_2O_2 .^{50,53} At low temperatures (less than or equal to $-78\text{ }^\circ\text{C}$), a transient hydroperoxide intermediate, $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**), is observed.⁵³ In contrast to **2** which is colorless, intermediate **3** is tangerine orange and displays an intense charge transfer band at 452(2900) nm (Figure 5). When the solution is warmed, its color changes from orange to burgundy, and the band at 452 nm cleanly converts to a band at 511(1770) nm (Figure 5). The burgundy species was shown, via its independent synthesis, to be the




FIGURE 5. Low-temperature detection of the hydroperoxo intermediate $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) using electronic absorption spectroscopy, and its conversion to solvent-bound $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OMe})]^+$ (**4**) upon warming.




FIGURE 6. X-Band EPR spectrum of tangerine orange $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) at 7 K in an MeOH/EtOH (5:2) glass.

solvent-derived methoxide-bound species $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OMe})]^+$ (**4**). When the reaction between **2** and superoxide is monitored by stopped-flow at ambient temperature (in MeOH), intermediate **3** grows in at diffusion-controlled rates (too fast to measure) and then cleanly converts to **4** (releasing H_2O_2) at a rate $[65(1) \text{ s}^{-1}]^{53}$ similar to that of the enzyme (50 s^{-1}).^{60,61} The tangerine orange species displays a low-spin ($S = 1/2$) EPR signal (Figure 6), which converts to an intermediate-spin ($S = 3/2$) signal as the solution develops a burgundy color. These changes to the electronic absorption and EPR spectra indicated that a transient intermediate forms (at low temperatures) during the reduction of superoxide but did not reveal its identity. Vibrational data, obtained at $-78 \text{ }^\circ\text{C}$, unambiguously identified this intermediate as a hydroperoxo species. A Fermi doublet (at 786 and 784 cm^{-1}) is observed in the infrared spectrum of $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) in the $\nu_{\text{O-O}}$ stretching region (Figure 7). This doublet collapses to a sharpened singlet at 784 cm^{-1} upon addition of D_2O , indicating that a proton is associated with the moiety responsible for the vibration.⁵³ Addition of ^{18}O -labeled (23%) superoxide to **2** results in a new shifted $\nu_{\text{O-O}}$ stretch at 753 cm^{-1} , close to that predicted on the basis of Hooke's law for a diatomic oxygen species.




FIGURE 7. Low-temperature IR spectrum of ^{16}O -labeled (\cdots) and ^{18}O -labeled (23%; —) $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) showing the peroxy $\nu_{\text{O-O}}$ Fermi doublet that shifts upon incorporation of ^{18}O .




FIGURE 8. ORTEP diagram of solvent-bound $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{MeCN})]^{2+}$ showing that MeCN binds *trans* to the imine nitrogen and *cis* to the thiolate sulfur.

Although we have yet to crystallize our hydroperoxo intermediate, we have structurally characterized $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) using X-ray absorption spectroscopy, in collaboration with Rob Scarrow. Fits to the EXAFS data for **3** require a new short Fe–O bond 1.86(3) Å in length, which is not present in either the Fe^{II} precursor (**2**) or the methoxide-bound product **4**.⁵³ The XANES spectrum of **3** is consistent with an oxidized six-coordinate Fe^{III} intermediate. Addition of an outer-sphere oxygen at 2.79(6) Å improves the EXAFS fits slightly.⁵³ Together, these data are consistent with the oxidative addition of superoxide to $[\text{Fe}^{\text{II}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))]^+$ via an inner-sphere mechanism to form an end-on $\text{Fe}^{\text{III}}(\eta^1\text{-OOH})$ intermediate. Hydroperoxo $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) represented the first example of a thiolate-ligated peroxy species, and the first structurally characterized $\text{Fe}^{\text{III}}\text{-OOH}$ species in any ligand environment.⁵³ The coexistence of a thiolate (a reductant) and a peroxide (an oxidant) in the same molecule is quite remarkable. Evidence that the peroxide binds *cis* with respect to the thiolate comes from the isolation and characterization of a number of more stable derivatives, including $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{MeCN})]^{2+}$ (Figure 8), $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{CN})]^+$, $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{N}_3)]^+$ (Figure 9), and $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OAc})]^+$ (**5**; Figure 10),⁵⁴ where it was shown that all ligands bind in this mode.

Formation of hydroperoxo $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) is proton-dependent. No reaction occurs between $[\text{Fe}^{\text{II}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))]^+$ (**2**) and O_2^- in rigorously dried THF




FIGURE 9. ORTEP diagram of thiolate-ligated $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{N}_3)]^+$.




FIGURE 10. ORTEP diagram of acetate-bound $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OAc})]^+$ (**5**), a mimic for the glutamate-bound SOR resting state.

until an external proton donor is added. This rules out a mechanism involving abstraction of H^+ or a H-atom from the ligand. Addition of a variety of proton donors, including NH_4^+ , MeOH, and PhOH, rapidly induces the formation of $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**). The ammonium ion (NH_4^+) mimics the lysine residue proposed to be involved in the SOR mechanism.⁵⁰ The rate of formation of **3** is highly dependent on both the HA concentration and $\text{p}K_{\text{a}}$,⁶² indicating that a proton is transferred in the steps prior to or during the rate-determining step. The proton dependence of this reaction is consistent with three possible mechanisms involving initial protonation of O_2^- , the thiolate sulfur, or an Fe(II)–superoxo intermediate. These should be distinguishable on the basis of kinetics. Kinetic studies are currently underway in our laboratory, in an attempt to establish the most probable mechanism.⁶²

The Fe–O(peroxo) bond of $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) can be cleaved via the addition of a second, more acidic, proton donor.⁵⁰ This results in the release of H_2O_2 , via what appears to be a proton-dependent dissociative mechanism. Nucleophiles, such as OAc^- , do not react with **3**, and the rate of H_2O_2 release is dependent on the $\text{p}K_{\text{a}}$ of the proton donor.⁶² The addition of HOAc releases H_2O_2 from **3** six orders of magnitude faster than NH_4^+ does. Noncoordinating acids (HBF_4 and HClO_4) cleanly afford a common




FIGURE 11. Proton-induced conversion of $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OOH})]^+$ (**3**) to solvent-bound $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{MeOH})]^{2+}$ (**6**) as monitored by electronic absorption spectroscopy at low temperatures.




FIGURE 12. Catalytic cycle involving $[\text{Fe}^{\text{II}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))]^{2+}$ (**2**)–promoted reduction of superoxide to afford H_2O_2 , via the sequential protonation and Cp_2Co -promoted reduction of hydroperoxo- and solvent-bound intermediates.

eggplant purple intermediate, $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{MeOH})]^{2+}$ (**6**; $\lambda_{\text{max}} = 565 \text{ nm}$). Acetic acid also reacts with **3** to form solvent-bound **6** (Figure 11), which then converts to acetate-bound $[\text{Fe}^{\text{III}}(\text{S}^{\text{Me}_2}\text{N}_4(\text{tren}))(\text{OAc})]^+$ (**5**; Figure 10), a model for the Glu-bound oxidized SOR resting state, upon warming. A solvent-bound $[\text{Fe}^{\text{II}}\text{-i-OH(H)}]$ intermediate has also recently been identified in the mechanism of SOR,^{15,20} as a species distinct from its $\text{Fe}^{\text{III}}\text{-Glu}$ resting state. Following the release of H_2O_2 from our hydroperoxo intermediate **3**, we can regenerate the active catalyst, **2**, by adding an external reductant such as cobaltocene (Figure 12). Subsequent addition of superoxide results in regeneration of the hydroperoxo intermediate. Eight turnovers have been achieved in this stepwise manner.⁵⁰ The catalytic activity is most likely limited due to decomposition of the catalyst by H_2O_2 -promoted oxidation of the thiolate.




FIGURE 13. ORTEP diagram of *trans*-thiolate-ligated, biomimetic superoxide reducing catalyst $[\text{Fe}^{\text{II}}(\text{cyclam-PrS})]^+$ (**8**) synthesized by Terutaka Kitagawa, a graduate student in the Kovacs lab.

Influence of the *trans*-Thiolate Ligand in Promoting SOR Chemistry

A few examples of structural SOR models containing a *trans*-thiolate have been reported,^{28,55} and the thiolate ligand has been shown to lower $\text{Fe}^{3+/2+}$ redox potentials.⁵⁵ Although none of these structural models has been reported to react with superoxide, in one case, H_2O_2 addition was shown to afford a high-valent $\text{Fe}^{\text{IV}}=\text{O}$ species,²⁸ presumably via the cleavage of a transient Fe^{III} -peroxo O–O bond. A macrocyclic thiolate-ligated ferrous complex was recently reported,⁴⁹ which reacts with ROOH to afford the first example of a thiolate-ligated alkyl peroxy complex $[[[15]\text{aneN}_4]\text{Fe}^{\text{III}}(\text{SPh})(\text{OOR})]^+$ (**7**), with the thiolate presumably *trans* to the peroxy. Although **7** is a low-spin complex ($g = 2.20$ and 1.97), its $\nu_{\text{Fe}-\text{O}}$ stretch (612 cm^{-1}) is significantly lower than those of most low-spin Fe^{III} -OOR complexes.²⁴ Most likely, this is due to the *trans* influence of the thiolate sulfur, although one cannot conclusively say that ROO^- binds in this position. Macrocyclic cyclam ligands have been shown to fold to afford *cis*-ligated six-coordinate geometries.⁶³ Recently, we reported a rare example of a functional metalloenzyme active site model, $[\text{Fe}^{\text{II}}(\text{cyclam-PrS})(\text{BPh}_4)]^+$ (**8**; Figure 13), that like SOR reduces O_2^- , presumably via a *trans*-thiolate-ligated Fe^{III} -peroxo intermediate.³⁸ The thiolate ligand of **8** was shown to lower the redox potential by 565 mV, alter the spin state of the peroxo intermediate, and dramatically weaken the Fe–O(peroxo) bond, favoring O_2^- reduction and H_2O_2 release.³⁸ Consistent with previous observations regarding amine substituents,^{56,64} the secondary amines of **8** also appear to play an important role in the observed superoxide reduction chemistry. Upon addition of O_2^- (18-crown-6- K^+ salt) and a proton donor (i.e., MeOH) to **8** at $-78\text{ }^\circ\text{C}$ in CH_2Cl_2 , a metastable, high-spin ($g = 7.72$, 5.40 , and 4.15) burgundy [$\lambda_{\text{max}} = 530(1350)\text{ nm}$] intermediate (**9**) is observed (Figure 14). No reaction occurs in the absence of a proton donor, and O_2^- does not convert to H_2O_2 under the same conditions in the absence of **8**.³⁸ An $\nu_{\text{O}-\text{O}}$ stretch (Fermi doublet) is observed at 891 cm^{-1} in the vibrational (resonance Raman) spectrum of **9**, which shifts to 856 cm^{-1} in the ^{18}O -labeled (50% label) spectrum. Addition of D^+ (i.e., MeOD) causes the Fermi doublet to collapse. An intense $\nu_{\text{Fe}-\text{S}}$ stretch is also observed at 352 cm^{-1} , indicating that the thiolate remains coordinated to the metal, and an ^{18}O -sensitive $\nu_{\text{Fe}-\text{O}}$ stretch is observed




FIGURE 14. Low-temperature electronic absorption spectrum and DFT-calculated structure of the hydroperoxo intermediate $[\text{Fe}^{\text{III}}(\text{cyclam-PrS})(\text{OOH})]^+$ (**9**) formed upon addition of superoxide to reduced $[\text{Fe}^{\text{II}}(\text{cyclam-PrS})(\text{BPh}_4)]^+$ (**8**) in the presence of a proton source.




FIGURE 15. Catalytic cycle involving $[\text{Fe}^{\text{II}}(\text{cyclam-PrS})(\text{BPh}_4)]^+$ (**8**)-promoted reduction of superoxide to afford H_2O_2 , via the sequential protonation and Cp_2Co -promoted reduction of hydroperoxo- and acetate-bound intermediates.

at 419 cm^{-1} , which shifts to 400 cm^{-1} .³⁸ Altogether, these data are consistent with the proton-dependent oxidative addition of superoxide to **8** which forms a thiolate-ligated hydroperoxo intermediate $[\text{Fe}^{\text{III}}(\text{cyclam-PrS})(\text{OOH})]^+$ (**9**). The remarkably low $\nu_{\text{Fe}-\text{O}}$ stretch, and the high-spin state of this intermediate, strongly suggest that the hydroperoxide ligand binds *trans* to the thiolate, although one cannot conclusively assign its structure in the absence of crystallographic evidence.

Although clearly more examples are needed if one hopes to correlate spectroscopic parameters to peroxo and thiolate binding modes and ultimately SOR function, with three examples (**3**, **7**, and **9**) now in hand, we^{38,50,53} and others^{25,28,49} are closer to this goal. In comparison to other reported synthetic iron peroxides,²⁴ the $\nu_{\text{Fe}-\text{O}}$ stretch of **9** is significantly weakened (419 cm^{-1} for **9** vs reported range, $450\text{--}639\text{ cm}^{-1}$),²⁴ but it compares well with the only reported SOR peroxo stretch (438 cm^{-1}),¹³ which was recently shown by X-ray crystallography to contain a

trans-thiolate and end-on hydroperoxo.²⁵ The $\nu_{\text{O-O}}$ stretch of **9** is unusually high (reported range, 820–860 cm^{-1}). The DFT-optimized structure of **9** (Figure 14) contains an Fe–O distance of 1.95 Å and a force constant ($k_{\text{Fe-O}} = 1.20$ mdyn/cm) which is significantly longer (reported range, 1.76–1.86 Å) and weaker (reported force constant range, 2.2–2.1 mdyn/cm) than those of all other reported Fe–(η^1 -OOH) species.²⁴ These data strongly support a *trans* positioning of the thiolate in **9** and indicate that its influence is to significantly weaken the Fe–O(peroxo) bond, favoring Fe–O, as opposed to O–O, bond cleavage.

The Fe–O(peroxo) bond of $[\text{Fe}^{\text{III}}(\text{cyclam-PrS})(\text{OOH})]^+$ (**9**) can be cleaved via the addition of proton donors, resulting in the release of H_2O_2 . A blue acetate-bound derivative, $[\text{Fe}^{\text{III}}(\text{cyclam-PrS})(\text{OAc})]^+$ (**10**), modeling the glutamate-bound SOR resting state, forms following the low-temperature addition of acetic acid.³⁸ Once H_2O_2 is released, the active catalyst **8** can be regenerated via the addition of a sacrificial reductant (Cp_2Co). Subsequent addition of superoxide results in regeneration of hydroperoxo intermediate **9**. Five turnovers have been achieved in this stepwise manner. In comparison to our *cis*-thiolate-ligated hydroperoxo complex **3** which releases H_2O_2 extremely slowly in MeOH at -78 °C ($t_{1/2} = 63.9$ h), hydroperoxo complex **9** rapidly releases H_2O_2 under the same conditions (within seconds). The macrocyclic amine ligand in combination with the presumed *trans* positioning of the thiolate converts the spin state from an $S = 1/2$ state in *cis*-ligated **3**⁵³ to an $S = 5/2$ state, promoting faster Fe–O bond cleavage under similar conditions. If the thiolate is indeed *trans*, it would also increase the basicity of the proximal peroxo oxygen relative to that of the *cis*-thiolate-ligated system, and this would be expected to influence product release rates if Fe–OOH bond cleavage is proton-induced. The relative rates of proximal oxygen Fe–OOH protonation (to induce H_2O_2 release) versus O–O(peroxide) bond cleavage must play a critical role in governing whether a SOR or P450 reaction pathway is followed.

Concluding Remarks

The biological chemistry of iron is exquisitely controlled by its ligand environment. Supporting ligands play an important role in determining function by controlling metal ion redox potential and spin state, as well as the basicity of bound substrates. Spin states control reaction pathways by fine-tuning relative bond strengths. Although there are many parallels between heme and non-heme iron systems, porphyrin ligands alter reaction pathways by facilitating oxidation via the delocalization of charge onto the ligand. Thiolate ligands also have the potential of facilitating the delocalization of charge.⁶⁵ As described in this Account, we are beginning to understand the functional role of the SOR thiolate ligand in promoting superoxide reduction. The intense π -thiolate sulfur-to- Fe^{III} charge transfer band, and its sensitivity to local coordination environment, provide a convenient method for monitoring reactions involving thiolate-ligated non-heme

iron complexes.³² By successfully mimicking the individual proton and electron transfer steps of the SOR catalytic cycle,^{38,50} we have begun investigating the molecular level details of the SOR mechanism and its dependence on the positioning of the thiolate ligand relative to the substrate binding site. The thiolate ligand favors superoxide reduction, regardless of its position in the coordination sphere, by lowering the redox potential of the metal ion.⁵⁶ In a protein environment, the thiolate provides an efficient electron transfer pathway.^{35,37} Spectroscopic characterization of the first reported examples of SOR peroxo intermediate analogues has allowed us,^{38,53} and others,^{49,66} to determine how the thiolate influences key properties affecting reaction pathways. Despite the structural similarities between the heme iron enzyme P450 and non-heme iron enzyme SOR peroxo intermediates, the two systems follow very different reaction pathways.⁴ The spin state favored by the supporting amine ligands influences the relative strength of the Fe–O(peroxo) versus O–O bond, and presumably the basicity of the proximal peroxo oxygen. The combination of a high-spin state and *trans*-thiolate affords a labile Fe^{III} -OOH species susceptible to protonation by protic solvents (e.g., MeOH or H_2O), resulting in facile H_2O_2 release. The biomimetic analogues described herein have allowed us to establish many of the key properties that are critical for the identification and function of cysteinyl-ligated metalloenzymes.

J.A.K. is grateful to have had the privilege to work with a number of very talented graduate students and postdocs who have contributed in one way or another to the work described herein. The National Institutes of Health is also gratefully acknowledged for its support of the work carried out in the Kovacs laboratory (Grant GM 45881-16).

References

- (1) Costas, M.; Mehn, M. P.; Jensen, M. P.; Que, L. J. Dioxygen Activation at Mononuclear Nonheme Iron Active Sites: Enzymes, Models, and Intermediates. *Chem. Rev.* **2004**, *104*, 939–986.
- (2) Solomon, E. I.; Decker, A.; Lehnert, N. Non-heme iron enzymes: Contrasts to heme catalysis. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 3589–3594.
- (3) Kovacs, J. A. Synthetic Analogues of Cysteine-Ligated Non-Heme Iron, and Non-Corrinoid Cobalt Enzymes. *Chem. Rev.* **2004**, *104*, 825–848.
- (4) Kovacs, J. A. How Iron Activates O_2 . *Science* **2003**, *299*, 1024–1025.
- (5) Hausinger, R. P. Fe(II) α -Ketoglutarate-Dependent Hydroxylases and Related Enzymes. *Crit. Rev. Biochem. Mol. Biol.* **2004**, *39*, 1–47.
- (6) Denisov, I. G.; Makris, T. M.; Sliagar, S. G.; Schlichting, I. Structure and Chemistry of Cytochrome P450. *Chem. Rev.* **2005**, *105*, 2253–2277.
- (7) Kappock, T. J.; Caradonna, J. P. Pterin-Dependent Amino Acid Hydroxylases. *Chem. Rev.* **1996**, *96*, 2659–2756.
- (8) Momenteau, M.; Reed, C. A.; Guiset, F. Synthetic Heme Dioxygen Complexes. *Chem. Rev.* **1994**, *94*, 659–698.
- (9) Kurtz, D. M., Jr. Microbial Detoxification of Superoxide: The Non-Heme Iron Reductive Paradigm for Combating Oxidative Stress. *Acc. Chem. Res.* **2004**, *37*, 902–908.
- (10) Wuerges, J.; Lee, J.-W.; Yim, Y.-I.; Yim, H.-S.; Kang, S.-O.; Carugo, K. D. Crystal structure of nickel-containing superoxide dismutase reveals another type of active site. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 8569–8574.
- (11) Jackson, T. A.; Brunold, T. C. Combined Spectroscopic/Computational Studies on Fe and Mn-Dependent Superoxide Dismutases: Insights into Second Sphere Tuning of Active Site Properties. *Acc. Chem. Res.* **2004**, *37*, 461–470.
- (12) Yeh, A. P.; Hu, Y.; Jenney, F. E., Jr.; Adams, M. W. W.; Rees, D. C. Structures of the superoxide reductase from *Pyrococcus furiosus*

- in the oxidized and reduced states. *Biochemistry* **2000**, *39*, 2499–2508.
- (13) Mathe, C.; Mattioli, T. A.; Horner, O.; Lombard, M.; Latour, J.-M.; Fontecave, M.; Niviere, V. Identification of Iron(III) Peroxo Species in the Active Site of the Superoxide Reductase SOR from *Desulfoarculus baarsii*. *J. Am. Chem. Soc.* **2002**, *124*, 4966–4967.
- (14) Niviere, V.; Asso, M.; Weill, C. O.; Lombard, M.; Guigliarelli, B.; Favaudon, V.; and Houe'e-Levin, C. Superoxide Reductase from *Desulfoarculus baarsii*: Identification of Protonation Steps in the Enzymatic Mechanism. *Biochemistry* **2004**, *43*, 808–818.
- (15) Rodrigues, J. V.; Abreu, I. A.; Cabelli, D.; Teixeira, M. Superoxide Reduction Mechanism of *Archaeoglobus fulgidus* One-Iron Superoxide Reductase. *Biochemistry* **2006**, *45*, 9266–9278.
- (16) Lombard, M.; Houe'e-Levin, C.; Touati, D.; Fontecave, M.; Niviere, V. Superoxide Reductase from *Desulfoarculus baarsii*: Reaction Mechanism and Role of Glutamate 47 and Lysine 48 in Catalysis. *Biochemistry* **2001**, *40*, 5032–5040.
- (17) Kurtz, D. M., Jr. Avoiding high-valent iron intermediates: Superoxide reductase and rubrerythrin. *J. Inorg. Biochem.* **2006**, *100*, 679–693.
- (18) Clay, M. D.; Jenney, F. E., Jr.; Hagedoorn, P. L.; George, G. N.; Adams, M. W. W.; Johnson, M. K. Spectroscopic Studies of *Pyrococcus furiosus* Superoxide Reductase: Implications for Active-Site Structures and the Catalytic Mechanism. *J. Am. Chem. Soc.* **2002**, *124*, 788–805.
- (19) Clay, M.; Cosper, C. A.; Jenney, F. E., Jr.; Adams, M. W. W.; Johnson, M. K. Nitric oxide binding at the mononuclear active site of reduced *Pyrococcus furiosus* superoxide reductase. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 3796–3801.
- (20) Mathé, C.; Niviere, V.; Mattioli, T. A. Fe³⁺-Hydroxide Ligation in the Superoxide Reductase from *Desulfoarculus baarsii* Is Associated with pH Dependent Spectral Changes. *J. Am. Chem. Soc.* **2005**, *127*, 16436–16441.
- (21) Lehnert, N. I. S. E. Density-functional investigation on the mechanism of H-atom abstraction by lipoyxygenase. *J. Biol. Inorg. Chem.* **2003**, *8*, 294–305.
- (22) Goldsmith, C. R.; Stack, T. D. P. Hydrogen Atom Abstraction by a Mononuclear Ferric Hydroxide Complex: Insight into the Reactivity of Lipoyxygenase. *Inorg. Chem.* **2006**, *45*, 6048–6055.
- (23) Roth, J. P.; Mayer, J. M. Hydrogen Transfer Reactivity of a Ferric Bi-imidazoline Complex That Models the Activity of Lipoyxygenase Enzymes. *Inorg. Chem.* **1999**, *38*, 2760–2761.
- (24) Roelfes, G.; Vrajmasu, V.; Chen, K.; Ho, R. Y. N.; Rohde, J.-U.; Zondervan, C.; Crois, R. M.; Schudde, E. P.; Lutz, M.; Spek, A. L.; Hage, R.; Feringa, B. L.; Munck, E.; Que, L., Jr. End-on and Side-on Peroxo Derivatives of Non-Heme Iron Complexes with Pentadentate Ligands: Models for Putative Intermediates in Biological Iron/Dioxygen Chemistry. *Inorg. Chem.* **2003**, *42*, 2639–2653.
- (25) Katona, G.; Capentier, P.; Niviere, V.; Amara, P.; Adam, V.; Ohana, J.; Tsanov, N.; Bourgeois, D. Raman-Assisted Crystallography Reveals End-On Peroxide Intermediates in a Nonheme Iron Enzyme. *Science* **2007**, *316*, 449–453.
- (26) Schlichting, I.; Berendzen, J.; Chu, K.; Stock, A. M.; Maves, S. A.; Benson, D. E.; Sweet, R. M.; Ringe, D.; Petsko, G. A.; Sligar, S. G. The Catalytic Pathway of Cytochrome P450cam at Atomic Resolution. *Science* **2000**, *287*, 1615–1622.
- (27) Harris, D. L.; Loew, G. H. Theoretical Investigation of the Proton Assisted Pathway to Formation of Cytochrome P450 Compound I. *J. Am. Chem. Soc.* **1998**, *120*, 8941–8948.
- (28) Bukowski, M. R.; Koehntop, K. D.; Stubna, A.; Bominaar, E. L.; Halfen, J. A.; Münck, E.; Nam, W.; Que, L., Jr. A Thiolate-Ligated Nonheme Oxoiron(IV) Complex Relevant to Cytochrome P450. *Science* **2005**, *310*, 1000–1002.
- (29) Auclair, K.; Moenne-Loccoz, P.; Ortiz de Montellano, P. R. Roles of the Proximal Heme Thiolate Ligand in Cytochrome P450cam. *J. Am. Chem. Soc.* **2001**, *123*, 4877–4885.
- (30) Rohde, J.-U.; In, J. H.; Lim, M. H.; Brennessel, W. W.; Bukowski, M. R.; Stubna, A.; Münck, E.; Nam, W.; Que, L., Jr. Crystallographic and Spectroscopic Characterization of a Nonheme Fe(IV)=O complex. *Science* **2003**, *299*, 1037–1039.
- (31) Solomon, E. I. Geometric and Electronic Structure Contributions to Function in Bioinorganic Chemistry: Active Sites in Non-Heme Iron Enzymes. *Inorg. Chem.* **2001**, *40*, 3656–3669.
- (32) Kennepohl, P.; Neese, F.; Schweitzer, D.; Jackson, H. L.; Kovacs, J. A.; Solomon, E. I. Spectroscopy of Non-Heme Iron Thiolate Complexes: Insight into the Electronic Structure of the Low-Spin Active Site of Nitrile Hydratase. *Inorg. Chem.* **2005**, *44*, 1826–1836.
- (33) Lugo-Mas, P.; Dey, A.; Xu, L.; Davin, S. D.; Benedict, J.; Kaminsky, W.; Hodgson, K. O.; Hedman, B.; Solomon, E. I.; Kovacs, J. A. How Does Single Oxygen Atom Addition Affect the Properties of a Fe-Nitrile Hydratase Analogue? The Compensatory Role of the Unmodified Thiolate. *J. Am. Chem. Soc.* **2006**, *128*, 11211–11221.
- (34) Holm, R. H.; Kennepohl, P.; Solomon, E. I. Structural and Functional Aspects of Metal Sites in Biology. *Chem. Rev.* **1996**, *96*, 2239–2314.
- (35) Gray, H. B.; Malström, B.; Williams, R. J. P. Copper coordination in blue proteins. *J. Biol. Inorg. Chem.* **2000**, *5*, 551–559.
- (36) Rao, P. V.; Holm, R. H. Synthetic Analogues of the Active Sites of Iron-Sulfur Proteins. *Chem. Rev.* **2004**, *104*, 527–560.
- (37) Glaser, T.; Hedman, B.; Hodgson, K. O.; Solomon, E. I. Ligand K-Edge X-ray Absorption Spectroscopy: A Direct Probe of Ligand-Metal Covalency. *Acc. Chem. Res.* **2000**, *33*, 859–868.
- (38) Kitagawa, T.; Dey, A.; Lugo-Mas, P.; Benedict, J.; Kaminsky, W.; Solomon, E.; Kovacs, J. A. A Functional Model for the Metalloenzyme Superoxide Reductase. *J. Am. Chem. Soc.* **2006**, *128*, 14448–14449.
- (39) Shoner, S.; Barnhart, D.; Kovacs, J. A. A Model for the Low-Spin, Non-Heme, Thiolate-Ligated Iron Site of Nitrile Hydratase. *Inorg. Chem.* **1995**, *34*, 4517–4518.
- (40) Jackson, H. L.; Shoner, S. C.; Rittenberg, D.; Cowen, J. A.; Lovell, S.; Barnhart, D.; Kovacs, J. A. Probing the Influence of Local Coordination Environment on the Properties of Fe-Type Nitrile Hydratase Model Complexes. *Inorg. Chem.* **2001**, *40*, 1646–1653.
- (41) Ellison, J. J.; Nienstedt, A.; Shoner, S. C.; Barnhart, D.; Cowen, J. A.; Kovacs, J. A. Reactivity of Five-Coordinate Models for the Thiolate-Ligated Fe Site of Nitrile Hydratase. *J. Am. Chem. Soc.* **1998**, *120*, 5691–5700.
- (42) Shearer, J.; Kung, I. Y.; Lovell, S.; Kaminsky, W.; Kovacs, J. A. Why Is There an "Inert" Metal Center in the Active Site of Nitrile Hydratase? Reactivity and Ligand Dissociation from a Five-Coordinate Co(III) Nitrile Hydratase Model. *J. Am. Chem. Soc.* **2001**, *123*, 463–468.
- (43) Shearer, J.; Jackson, H. L.; Schweitzer, D.; Rittenberg, D. K.; Leavy, T. M.; Kaminsky, W.; Scarrow, R. C.; Kovacs, J. A. The first example of a nitrile hydratase model complex that reversibly binds nitriles. *J. Am. Chem. Soc.* **2002**, *124*, 11417–11428.
- (44) Green, M. T.; Dawson, J. H.; Gray, H. B. Oxoiron(IV) in Chloroperoxidase Compound II Is Basic: Implications for P450 Chemistry. *Science* **2004**, *304*, 1653–1656.
- (45) Sommerhalter, M.; Lieberman, R. L.; Rosenzweig, A. C. X-ray Crystallography and Biological Metal Centers: Is Seeing Believing. *Inorg. Chem.* **2005**, *44*, 770–778.
- (46) Corwin, D. T., Jr.; Gruff, E. S.; Koch, S. Zinc, cobalt, and cadmium thiolate complexes. *J. Chem. Soc., Chem. Commun* **1987**, *13*, 966–967.
- (47) Grapperhaus, C. A.; Darensbourg, M. Y. Oxygen Capture by Sulfur in Nickel Thiolates. *Acc. Chem. Res.* **1998**, *31*, 451–459.
- (48) Kung, I. Y.; Schweitzer, D.; Shearer, J.; Taylor, W. D.; Jackson, H. L.; Lovell, S.; Kovacs, J. A. How Do Oxidized Thiolate Ligands Affect the Electronic and Reactivity Properties of a Nitrile Hydratase Model Compound. *J. Am. Chem. Soc.* **2000**, *122*, 8299–8300.
- (49) Krishnamurthy, D.; Kasper, G. D.; Namuswe, F.; Kerber, W. D.; Sarjeant, A. A. N.; Moëne-Loccoz, P.; Goldberg, D. P. Fe(III)-OOR SOR model. *J. Am. Chem. Soc.* **2006**, *128*, 14222–14223.
- (50) Theisen, R. M.; Kovacs, J. A. The Role of Protons in Superoxide Reduction by a Superoxide Reductase Analogue. *Inorg. Chem.* **2005**, *44*, 1169–1171.
- (51) Tyler, L. A.; Noveron, J. C.; Olmstead, M. M.; Mascharak, P. K. Modulation of the pK_a of Metal-Bound Water via Oxidation of Thiolate Sulfur in Model Complexes of Co(III) Containing Nitrile Hydratase. *Inorg. Chem.* **2003**, *42*, 5751–5761.
- (52) Schweitzer, D.; Ellison, J. J.; Shoner, S. C.; Lovell, S.; Kovacs, J. A. A Synthetic Model for the NO-Inactivated Form of Nitrile Hydratase. *J. Am. Chem. Soc.* **1998**, *120*, 10996–10997.
- (53) Shearer, J.; Scarrow, R. C.; Kovacs, J. A. Synthetic models for the cysteinylated non-heme iron enzyme superoxide reductase: Observation and structural characterization by XAS of an Fe^{III}-OOH intermediate. *J. Am. Chem. Soc.* **2002**, *124*, 11709–11717.
- (54) Shearer, J.; Fitch, S. B.; Kaminsky, W.; Benedict, J.; Scarrow, R. C.; Kovacs, J. A. How does cyanide inhibit superoxide reductase? Insight from synthetic Fe^{III}N₄S model complexes. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 3671–3676.
- (55) Fiedler, A. T.; Halfen, H. L.; Halfen, J. A.; Brunold, T. C. Synthesis, Structure Determination, and Spectroscopic/Computational Characterization of a Series of Fe(II)-Thiolate Model Complexes: Implications for Fe-S Bonding in Superoxide Reductases. *J. Am. Chem. Soc.* **2005**, *127*, 1675–1689.
- (56) Theisen, R. M.; Shearer, J. W. K.; Kovacs, J. A. Steric and Electronic Control Over the Reactivity of a Thiolate-Ligated Fe(II) Complex with Dioxygen and Superoxide. Reversible mu-oxo Dimer Formation. *Inorg. Chem.* **2004**, *43*, 7682–7690.
- (57) Noveron, J. C.; Olmstead, M. M.; Mascharak, P. K. A Synthetic Analogue of the Active Site of Fe-Containing Nitrile Hydratase with Carboxamido N and Thiolate S as Donors: Synthesis, Structure, and Reactivities. *J. Am. Chem. Soc.* **2001**, *123*, 3247–3259.
- (58) Schweitzer, D.; Shearer, J.; Rittenberg, D. K.; Shoner, S. C.; Ellison, J. J.; Loloee, R.; Lovell, S. C.; Barnhart, D.; Kovacs, J. A. Enhancing

- Reactivity via Ligand Constraints. *Inorg. Chem.* **2002**, *41*, 3128–3136.
- (59) Shearer, J.; Nehring, J.; Kaminsky, W.; Kovacs, J. A. Modeling the Reactivity of Superoxide Reducing Metalloenzymes with a Nitrogen and Sulfur Coordinated Iron Complex. *Inorg. Chem.* **2001**, *40*, 5483–5484.
- (60) Kurtz, D. M.; Coulter, E. D. The mechanism(s) of superoxide reduction by superoxide reductases in vitro and in vivo. *J. Biol. Inorg. Chem.* **2002**, *7*, 653–658.
- (61) Emerson, J. P.; Coulter, E. D.; Cabelli, D. E.; Phillips, R. S.; Kurtz, D. M., Jr. Kinetics and Mechanism of Superoxide Reduction by Two-Iron Superoxide Reductase from *Desulfovibrio vulgaris*. *Biochemistry* **2002**, *41*, 4348–4357.
- (62) Nam, E.; Alokolaro, P.; Kovacs, J. A. Kinetics paper, 2007, manuscript in preparation .
- (63) Lemma, K.; Ellern, A.; Bakac, A. ¹H NMR Studies of Ligand and H/D Exchange Reactions of *cis*- and *trans*-([14]aneN₄)(H₂O)RhH²⁺ in Aqueous Solutions. *Inorg. Chem.* **2003**, *42*, 3662–3669.
- (64) Berry, J. F.; Bill, E.; Garcia-Serres, R.; Neese, F.; Weyhermuller, T.; Wieghardt, K. Effect of N-Methylation of Macrocyclic Amine Ligands on the Spin State of Fe(III): A Tale of Two Fluoro Complexes. *Inorg. Chem.* **2006**, *45*, 2027–2037.
- (65) Green, M. T. Evidence for Sulfur-Based Radicals in Thiolate Compound I Intermediates. *J. Am. Chem. Soc.* **1999**, *121*, 7939–7940.
- (66) Bukowski, M. R.; Halfen, H. L.; van der Berg, T. A.; Halfen, J. A.; Que, L., Jr. Spin-State Rationale for the Peroxo-Stabilizing Role of the Thiolate Ligand in Superoxide Reductase. *Angew. Chem., Int. Ed.* **2005**, *44*, 584–587.

AR600059H