

Non-Heme Fe(IV)–Oxo Intermediates

CARSTEN KREBS,^{*,†,‡}
 DANICA GALONIĆ FUJIMORI,[§]
 CHRISTOPHER T. WALSH,^{*,§} AND
 J. MARTIN BOLLINGER, JR.^{*,†,‡}

Department of Biochemistry and Molecular Biology and Department of Chemistry, The Pennsylvania State University, University Park, Pennsylvania 16802, and Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, Massachusetts 02115

Received March 15, 2007

ABSTRACT

High-valent non-heme iron–oxo intermediates have been proposed for decades as the key intermediates in numerous biological oxidation reactions. In the past three years, the first direct characterization of such intermediates has been provided by studies of several α KG-dependent oxygenases that catalyze either hydroxylation or halogenation of their substrates. In each case, the Fe(IV)–oxo intermediate is implicated in cleavage of the aliphatic C–H bond to initiate hydroxylation or halogenation. The observation of non-heme Fe(IV)–oxo intermediates and Fe(II)-containing product(s) complexes with almost identical spectroscopic parameters in the reactions of two distantly related α KG-dependent hydroxylases suggests that members of this subfamily follow a conserved mechanism for substrate hydroxylation. In contrast, for the α KG-dependent non-heme iron halogenase, CytC3, two distinct Fe(IV) complexes form and decay together, suggesting that they are in rapid equilibrium. The existence of two distinct conformers of the Fe site may be the key factor accounting for the divergence of the halogenase reaction from the more usual hydroxylation pathway after C–H bond cleavage. Distinct transformations catalyzed by other mononuclear non-heme enzymes are likely also to involve initial C–H bond cleavage by Fe(IV)–oxo complexes, followed by diverging reactivities of the resulting Fe(III)–hydroxo/substrate radical intermediates.

Introduction

A large, functionally and mechanistically diverse family of enzymes utilize similar, mononuclear non-heme Fe(II) centers to couple the activation of oxygen to the oxidation of their substrates.^{1–3} In most cases, oxygen is inserted into an unactivated C–H bond of the substrate (hydroxylation), but many other outcomes, including halogenation,

desaturation, cyclization, epoxidation, and decarboxylation, are known.^{3,4} Each of these reactions is a two-electron oxidation. The remaining two reducing equivalents required for the four-electron reduction of oxygen are often provided by a cosubstrate. The reducing cosubstrates used by various family members include α -keto-glutarate (α KG) (in the α KG-dependent enzymes³), tetrahydrobiopterin (in the pterin-dependent aromatic amino acid hydroxylases⁵), reduced nicotinamides [in the Rieske dioxygenases and (S)-2-hydroxypropylphosphonic acid epoxidase¹], and ascorbic acid (in 1-aminocyclopropane 1-carboxylic acid oxidase⁶). A few of the enzymes oxidize their substrates by four electrons and thus do not require a reducing cosubstrate. This subset includes the extradiol dioxygenases,¹ isopenicillin N-synthase (IPNS),⁷ and two enzymes, 4-hydroxymandelate synthase and (4-hydroxyphenyl)pyruvate dioxygenase (HPPD), which effect distinct four-electron oxidations of their common substrate.⁸ The latter two reactions are mechanistically similar to those catalyzed by the α KG-dependent enzymes, because both involve oxidative decarboxylation of an α -keto acid moiety to provide two electrons.

This remarkable array of oxidative transformations is made possible by the tuning of a largely conserved mononuclear non-heme iron cofactor unit, which is coordinated by as few as two protein ligands and thus has as many as four sites available to coordinate substrates. In the most common coordination sphere, three protein ligands of a (His)₂-(Asp/Glu) motif, known as the “facial triad” because they occupy one face of an octahedron, leave three remaining sites on the opposite face for substrate coordination.⁹ Reaction mechanisms proposed for these enzymes have invoked several intermediates following the addition of oxygen to the Fe(II) center.^{1,2} Two types of intermediates have been proposed: Fe-coordinated (su)peroxo complexes with an intact O–O bond, [Fe–O₂]^{2+/3+}, and high-valent Fe(IV)–oxo intermediates [or even Fe(V)–oxo for the Rieske dioxygenases], [Fe=O]^{4+/5+}. In particular, the high-valent Fe–oxo intermediates have been suggested to initiate substrate oxidation. In most cases, activation of the substrate involves abstraction of the H-atom of the target C–H bond by the Fe(IV)–oxo intermediate to yield a substrate radical and

Carsten Krebs obtained his Ph.D. in 1997 from the Ruhr-Universität Bochum and was a postdoctoral fellow at Emory University until 2001. He is Assistant Professor in the Department of Biochemistry and Molecular Biology and the Department of Chemistry at The Pennsylvania State University. His research program focuses on the use of kinetic and spectroscopic methods in studying non-heme enzymes.

Danica Galonić Fujimori obtained her Ph.D. in 2005 from the University of Illinois at Urbana-Champaign and joined Chris Walsh's group at Harvard Medical School in 2005. She received a postdoctoral fellowship from the Damon Runyon Cancer Research Foundation.

Christopher T. Walsh is the Hamilton Kuhn Professor of Biological Chemistry and Molecular Pharmacology at Harvard Medical School. His research focuses on enzymes and enzyme inhibitors, with recent specialization on natural products biosynthesis. He is a member of the National Academy of Sciences, the Institute of Medicine, and the American Philosophical Society.

* To whom correspondence should be addressed. C.T.W.: Department of Biological Chemistry and Molecular Pharmacology, 240 Longwood Ave., Boston, MA 02115; phone, (617) 432-1715; fax, (617) 432-0438; e-mail, christopher_walsh@hms.harvard.edu. J.M.B.: Department of Biochemistry and Molecular Biology, 208 Althouse Laboratory, University Park, PA 16802; phone, (814) 863-5707; fax, (814) 863-7024; e-mail, jmb21@psu.edu. C.K.: Department of Biochemistry and Molecular Biology, 306 S. Frear Laboratory, University Park, PA 16802; phone, (814) 865-6089; fax, (814) 863-7024; e-mail, ckrebs@psu.edu.


† Department of Biochemistry and Molecular Biology, The Pennsylvania State University.

‡ Department of Chemistry, The Pennsylvania State University.

§ Harvard Medical School.

J. Martin Bollinger, Jr., is Associate Professor in the Departments of Biochemistry and Molecular Biology and Chemistry at The Pennsylvania State University. His research program focuses on the use of kinetic and spectroscopic methods to dissect the mechanisms of enzymes that activate oxygen and cleave C–H bonds.

Scheme 1. Reactions Proposed To Be Mediated by High-Valent Fe–Oxo Intermediates



an Fe(III)–OH complex (Scheme 1). The so-called oxygen rebound, which was originally proposed for heme enzymes¹⁰ and formally involves recombination of a coordinated hydroxyl radical equivalent with the substrate radical, yields the hydroxylated product and a coordinatively unsaturated Fe(II) center. In addition to substrate hydroxylation, many other outcomes following H-atom abstraction by the Fe(IV)–oxo intermediate are documented. These include transfer (formally as the radical) of a ligand of the Fe center to the substrate radical. Examples include transfer of a halogen atom in the α KG-dependent halogenases⁴ and transfer of a thyl group in IPNS.⁷


Alternative reactivities that do not involve radical recombination with a ligand include desaturation and cyclization of the substrate. Formally, these reactions involve abstraction of a second H-atom by the Fe(III)–OH complex to yield the desaturated or cyclized product and an Fe(II)–OH₂ complex. However, other pathways are possible, making delineation of the mechanisms of these alternative outcomes a high priority for future studies.

Other substrate oxidations by high-valent Fe–oxo intermediates that do not involve H-atom abstraction include electrophilic attack on the aromatic ring of the substrate by the pterin-dependent hydroxylases⁵ and *cis* dihydroxylation of an aromatic substrate by the Rieske dioxygenases.^{1,2}

Significant insight into the geometric and electronic structures of high-valent non-heme Fe–oxo complexes and their reactivity was obtained in parallel from elegant studies of inorganic complexes^{11–15} (see ref 1 for a recent review), but these studies will not be reviewed here due to the brevity of this Account.

First Non-Heme Fe(IV)–Oxo Intermediate

A powerful approach to studying the mechanism of a metalloenzyme-catalyzed reaction is the direct detection of intermediates and their detailed characterization by a combination of kinetic and spectroscopic methods. Using this approach, one monitors changes in the geometric and/or electronic structure of the metal center during the reaction. This methodology had been used successfully in the 1990s to study O₂ activation by the non-heme diiron proteins methane monooxygenase and the R2 subunit of class I ribonucleotide reductase but was only recently applied to the mononuclear non-heme iron enzymes. The first direct detection of an intermediate in the reaction of a mononuclear non-heme iron enzyme with dioxygen was reported for HPPD.¹⁶ A transient absorption feature at 490 nm that forms with a second-order rate constant of 140 mM⁻¹ s⁻¹ and decays with a first-order rate constant of 7.8 s⁻¹ was noted. More detailed spectroscopic characterization of the associated intermediate has not yet been reported. Shortly after this work, we reported the detection and characterization of two transient states in the reaction of taurine: α KG dioxygenase (TauD). The α KG-dependent oxygenases are the largest and functionally most diverse subgroup of mononuclear non-heme iron enzymes.³ They catalyze many important reactions, including steps in the biosyntheses of antibiotics¹⁷ and collagen,¹⁸ the sensing of oxygen,^{19–23} the repair of alkylated DNA,^{24,25} and the regulation of transcription by demethylation of histones.^{26–28} A chemical mechanism was initially proposed more than 20 years ago by Hanauske-Abel and Günzler specifically for the enzyme prolyl-4-hydroxylase (P4H),²⁹ but its success in accom-

Scheme 2. General Mechanism of α KG-Dependent Dioxygenases

modating ensuing experimental data for many other α KG-dependent hydroxylases led to its becoming adopted as a consensus mechanism for the subfamily (Scheme 2).^{1–3,30}

Two iron-based intermediates were detected in TauD by stopped-flow (SF) absorption and freeze-quench (FQ) Mössbauer spectroscopies. The first intermediate, termed J, forms with second-order kinetics (first-order in O_2 and enzyme concentrations; $k = 130 \text{ mM}^{-1} \text{ s}^{-1}$). It is characterized by an absorption feature that is maximal at $\sim 318 \text{ nm}$ and a new Mössbauer quadrupole doublet with unusual parameters (isomer shift, δ , of 0.30 mm/s and quadrupole splitting, ΔE_Q , of -0.88 mm/s).³¹ It has a nearly axial $S = 2$ ground state with a positive zero-field splitting parameter, D , of 10.5 cm^{-1} .^{31,32} The large substrate deuterium kinetic isotope effect (^2H -KIE) on decay of J ($k_{\text{H}}/k_{\text{D}} \approx 50$) implied that it is the hydrogen-abstracting intermediate,^{33,34} which the Hanauske-Abel and Günzler mechanism predicted to be an Fe(IV)–oxo complex.^{1,2,29}

The presence of the Fe(IV)–oxo group in J was confirmed by resonance Raman spectroscopy, which revealed a band at 821 cm^{-1} that shifted to 787 cm^{-1} upon use of $^{18}\text{O}_2$,³⁵ and X-ray absorption spectroscopy, which demonstrated a short (1.62 \AA) interaction between the Fe and one of its ligands.³⁶ A recent comparison of experimentally determined spectroscopic parameters with those predicted by DFT calculations for several model structures suggested that J has either a trigonal bipyramidal or octahedral coordination environment.³⁷ The second accumulating state is an Fe(II)-containing TauD-product(s) complex (V in Scheme 2).³⁸

A Consensus Mechanism for the α KG-Dependent Dioxygenases

The TauD work proved that (1) a non-heme Fe(IV)–oxo complex could be trapped and characterized despite its





FIGURE 1. Comparison of the spectroscopic features of the Fe(IV)–oxo intermediates from TauD (top), P4H (middle), and CytC3 (bottom). (A) Comparison of the kinetics of the Fe(IV)–oxo intermediates monitored by SF absorption spectroscopy using unlabeled (red) and selectively deuterated substrates (blue). (B) Set of 4.2 K, 53 mT (left) and 4.2 K, 8 T (right) Mössbauer spectra of the Fe(IV)–oxo intermediates.

Scheme 3. Reactions Catalyzed by the α KG-Dependent Hydroxylase Thymine Hydroxylase (TH)

anticipated high reactivity and (2) the lifetime of the Fe(IV)-oxo intermediate could be extended to a remarkable degree by deuterium substitution of the target C-H bond, due to the large ²H-KIE. We next applied this insight to a prolyl-4-hydroxylase (P4H), because the hydroxylation of proline residues is biologically significant (e.g., in collagen biosynthesis and oxygen sensing). The monomeric P4H from *Paramecium bursaria Chlorella virus 1*, which was known to modify peptide substrates containing a (Pro-Ala-Pro-Lys)_n motif, was selected.³⁹ The combined stopped-flow absorption and freeze-quench Mössbauer data for the reaction of the P4H·Fe(II)· α KG·(Pro-Ala-Pro-Lys)₃ complex with O₂ demonstrated the accumulation of two kinetically competent intermediates.⁴⁰ The first intermediate is a high-spin Fe(IV) complex, which exhibits a large substrate ²H-KIE on its decay ($k_{\text{H}}/k_{\text{D}} \approx 210/3.4 \approx 60$), suggesting that it abstracts hydrogen from the substrate. The second accumulating state contains a high-spin Fe(II) center and is presumably an Fe(II)·product(s) complex. The spectroscopic properties of these two intermediates are strikingly similar to those of the TauD intermediates (Figure 1), suggesting that the distantly related α KG-dependent dioxygenases employ the same chemical mechanism and supporting the prevailing view of a conserved mechanism for the hydroxylase subfamily.



Alternative Reactivities of α KG-Dependent Dioxygenases

Studies by Stubbe and co-workers on thymine hydroxylase shed light on alternative reactivities that can occur in the Fe(II)/ α KG reaction manifold. This enzyme catalyzes three oxidations of the methyl group of thymine to the hydroxymethyl, aldehyde, and carboxylic acid in three separate O₂ activation events, each of which involves cleavage of a C-H bond (Scheme 3A). The authors demonstrated that the enzyme can catalyze epoxidation

of an olefin, conversion of an alkyne to a ketene, and successive S-oxidations of a thioether to sulfoxide and then sulfone (Scheme 3B).^{41–43} These substrates lack C-H bonds at the position that would normally be targeted by the Fe(IV)-oxo complex, and the alternative transformations that ensue parallel those previously seen for high-valent heme iron enzyme intermediates and inorganic model complexes. It is therefore tempting to speculate that the alternative oxidations are also effected by the Fe(IV)-oxo intermediate, but it is still conceivable that fundamentally different mechanisms might be operant [e.g., interception of an earlier intermediate, such as an Fe(III)-superoxide complex like **I** in Scheme 2].

α KG-Dependent Halogenases

The recent discovery of a new class of halogenating enzymes that carry out chlorination of unactivated carbon centers in the biosyntheses of several natural products of nonribosomal peptide origin established yet another reactivity for the Fe(II)/ α KG-dependent oxygenases.^{44–47} Aliphatic halogenases chlorinate the terminal methyl groups of amino acids tethered via a thioester linkage to the phosphopantetheine cofactor of peptidyl-carrier proteins (PCPs). They require iron, α KG, oxygen, and chloride for their activity. Initial insight into their catalytic mechanism was derived from the crystal structure of SyrB2, which chlorinates the γ -methyl group of L-threonine in syringomycin biosynthesis.⁴⁸ The Fe center is coordinated by two protein-derived histidines, bidentate α KG, water, and chloride. The carboxylate of the “facial triad” that normally coordinates the Fe(II) center is replaced with an alanine in the protein primary structure, presenting a coordination site for the chloride ligand. On the basis of this observation, a Cl-Fe(IV)-oxo intermediate was proposed as the C-H bond-cleaving intermediate, and chlorination was proposed to proceed via “chlorine rebound”

Scheme 4. Hydroxylation versus Halogenation Rebound Reactions**Scheme 5. Reaction Catalyzed by the Halogenase CytC3**

rather than “oxygen rebound” (Scheme 4). This hypothesis was tested experimentally for the non-heme iron halogenase, CytC3 from soil *Streptomyces*.⁴⁹ CytC3 chlorinates the γ -methyl group of L-2-aminobutyric acid (L-Aba) tethered to the PCP domain CytC2, L-Aba-S-CytC2 (Scheme 5).⁴⁷ Evidence of the accumulation of two transient states of the catalytic cycle was obtained in the reaction of the CytC3-Fe(II) $\cdot\alpha$ KG-Cl⁻-L-Aba-S-CytC2 complex with oxygen-saturated buffer. The first intermediate absorbs at 318 nm. Decay of A_{318} is markedly slowed by use of a deuterated substrate, L-4,4,4- d_3 -Aba-S-CytC2, demonstrating a ²H-KIE on decay of the intermediate and implicating it as the C–H bond-cleaving complex (see Figure 1). Mössbauer spectra revealed the presence of *two* high-spin Fe(IV) intermediates. Their proportions are constant with reaction time, suggesting that they are in rapid equilibrium. These results contrast with those of the α KG-dependent hydroxylases, for which the C–H bond-cleaving state comprises only *one* high-spin Fe(IV)–oxo complex. We speculated that the two intermediates are distinct conformers of the Cl-Fe(IV)–oxo complex and that the occurrence of two Fe(IV) complexes prior to C–H bond cleavage may also indicate the presence of two Fe(III)–OH complexes after H-atom abstraction. The presence of two conformers might be the key factor allowing halogenation in preference to the conventional hydroxylation outcome. Ongoing efforts are directed toward (i) elucidating the structures of the two conformers by employing other spectroscopic methods (e.g., X-ray absorption, resonance Raman, and magnetic circular dichroism) and (ii) defining the factors that affect the ratio of the two Fe(IV) complexes by perturbing the reaction conditions or by using modified substrates.

Exclusive halogenation (rather than hydroxylation) was also observed in an inorganic non-heme iron complex, in which the Fe center is coordinated to chloride and hydroxide.⁵⁰ The reaction is thought to proceed via hydrogen atom abstraction followed by rebound of the coordinated ligand. It was proposed that the preference for chlorination reflects the lower reduction potential of

chlorine radical ($\text{Cl}^\bullet + e^- \rightarrow \text{Cl}^-$, 1.36 V) relative to hydroxyl radical ($\text{HO}^\bullet + e^- \rightarrow \text{HO}^-$, 2.02 V).⁵⁰


The second intermediate observed during the reaction of the CytC3 halogenase is an Fe(II) complex that is spectroscopically distinct from the reactant. In analogy to the hydroxylases, it is presumably a product(s) complex. The accumulation of two intermediate states in the CytC3 system, the C–H bond-cleaving Fe(IV)–oxo intermediate and the Fe(II)–product(s) complex, underscores the mechanistic similarity of the α KG-dependent hydroxylases and halogenases and suggests that halogenation activity may have evolved from hydroxylation by iron ligand replacement (among other less apparent adaptations). Figure 1 further emphasizes these striking similarities by showing the kinetics of formation and decay of the Fe(IV)–oxo complexes in TauD, P4H, and CytC3 with both protium- and deuterium-containing substrates (left panel) and their Mössbauer spectra (center and right panels).

Mechanistic Diversity of Presumptive Fe(IV)–Oxo Intermediates in β -Lactam Biosyntheses


Nature has exploited the versatility of mononuclear non-heme iron enzymes in the biosynthetic pathways of a wide variety of β -lactam antibiotics.¹⁷ Isopenicillin *N*-synthase (IPNS) is one of the more well-understood mononuclear non-heme iron enzymes due to the extensive studies by Baldwin and co-workers. IPNS catalyzes the four-electron oxidation of δ -(L- α -aminoacyl)-L-cysteine-D-valine (ACV) tripeptide to isopenicillin N.⁵¹ Elegant biochemical and crystallographic studies provided convincing evidence that the reaction proceeds via two successive two-electron oxidations (Scheme 6).^{7,52} The first oxidation is thought to involve cleavage of the $C_{\text{Cys},\beta}$ -H bond by a formally Fe(III)–superoxo intermediate, followed by formation of the β -lactam ring, possibly by attack of the valine amidate on the thioaldehyde. This view is supported by several lines of evidence. First, the crystal structure of the IPNS-Fe(II)·ACV complex with NO as a surrogate for oxygen revealed the O-atom of the NO group to be in the proximity of the target $C_{\text{Cys},\beta}$ -H bond.⁵³ Second, with the substrate analogue δ -(L- α -aminoacyl)-L-cysteine-D- α -hydroxyvaleryl ester (Scheme 7A), $C_{\text{Cys},\beta}$ -H bond cleavage but not formation of the β -lactam ring was observed.⁵⁴ Third, with ACV containing C_{β} -deuterated cysteine as the substrate, a kinetic isotope effect on $k_{\text{cat}}/K_{\text{M}}$ (selection effect) of 1.4 was measured.⁵²

The second oxidation is proposed to involve $C_{\text{Val},\beta}$ -H bond cleavage by an Fe(IV)–oxo intermediate. Recombination of the resultant substrate radical with the coordinated sulfur (formally as the thiyl radical) results in formation of the thiazolidine ring of the substrate (Scheme 6). This sequence of events, i.e., cleavage of the $C_{\text{Val},\beta}$ -H bond by the Fe(IV)–oxo intermediate followed by transfer of a coordinated ligand other than the hydroxide, is analogous to that envisaged for the halogenases.

Scheme 6. Proposed Mechanism of IPNS



Scheme 7. Alternative Reactions of the Proposed Fe(IV)–Oxo Intermediate in IPNS




Although none of the proposed intermediates has been directly detected, there is convincing evidence of the intermediacy of a β -valinyl radical from studies of substrate analogues (Scheme 7). For example, formation of the [6.2.0] (Scheme 7B) ring system in the reaction of IPNS with the cyclopropane-containing substrate provides evidence that supports generation of a cyclopropylcarbinyl radical, lending credence to the proposed H-atom abstraction by the Fe(IV)–oxo intermediate.⁵⁵

Substrate analogues in which the D-valine residue is modified provided further insight into the reactivity of the presumptive Fe(IV)–oxo intermediate in IPNS. Of particular interest are conversion of the thioether of the D-S-methylcysteine-containing analogue to the sulfone⁵⁶ and oxidation of the olefinic D-allylglycine- and D-vinylglycine-containing substrates to yield desaturated and oxygenated bicyclic products (Scheme 7C–E).^{57,58} Isotopic labeling studies demonstrated that the O-atom incorporated into


the hydroxylated bicyclo[5.2.0] product of the D-allylglycine-containing substrate is derived from dioxygen (Scheme 7D).⁵⁹

Clavaminate synthase (CAS) from *Streptomyces clavuligerus* performs three two-electron oxidation reactions in the biosynthesis of clavulanic acid: hydroxylation, oxidative ring closure, and dehydrogenation (Scheme 8).^{60–65} The hydroxylation reaction is believed to proceed via the canonical mechanism involving H-atom abstraction by the Fe(IV)–oxo intermediate, followed by hydroxyl radical rebound. It has been proposed that a different mode of reactivity of the Fe(III)–OH/substrate radical state yields the second and third reactions.^{17,66} For the dehydrogenation reaction, the Fe(III)–OH intermediate could abstract a second H-atom from the substrate radical, yielding a hexacoordinate Fe(II)–OH₂ complex and the olefinic product.⁶⁷ Similarly, the cyclization reaction could proceed by successive H-atom abstractions by the Fe(IV)–oxo


Scheme 8. Reaction Catalyzed by CAS



Scheme 9. Reaction Catalyzed by DAOCS



Scheme 10. Reaction Catalyzed by CarC



and Fe(III)–OH intermediates (the latter from the hydroxyl group of the substrate) and a radical coupling step.¹⁷ It has been proposed that subtleties in the structures of the CAS-intermediate complexes favor the proposed second H-abstraction steps over other possible pathways for decay of the Fe(III)–OH/substrate radical states (e.g., oxygen rebound).

Deacetoxycephalosporin C synthase (DAOCS) from *S. clavuligerus* is an Fe(II)-containing α KG-dependant oxidase that catalyzes expansion of the five-membered thiazolidine ring of penicillin N to the six-membered dihydrothiazine ring of cephalosporins (Scheme 9).⁶⁸ It is believed that ring expansion is initiated by abstraction of hydrogen from the C2-methyl substituent by a ferryl species. This primary alkyl substrate radical then converts to the more stable tertiary radical in the ring expansion. The intermediacy of an episulfide species, with radical character localized on the sulfur, was proposed for this transformation.⁶⁹ Subsequent abstraction of the second hydrogen from C3 of the substrate, perhaps carried out by an Fe(III)–OH species, results in the formation of the endocyclic double bond and generation of the cephalosporin nucleus.^{70–73} The delicate balance of desaturation and hydroxylation pathways observed in CAS can also be observed in DAOCS upon use of the [3-²H]penicillin N substrate, in which the hydrogen target of the second abstraction is substituted with deuterium.⁷³ In this case, hydroxylation of the six-membered ring occurs, presumably by recombination of the more stable tertiary radical with the hydroxyl radical from the Fe(III)–OH intermediate.

Carbapenam synthase (CarC) from *Pectobacterium carotovorum* is a bifunctional α KG-dependent enzyme that epimerizes the unactivated C5 position of its (3*S*,5*S*)-carbapenam-3-carboxylate substrate and installs a double bond between C2 and C3, resulting in the formation of the carbapenam binucleus (Scheme 10).¹⁷ Isotopic labeling studies by Townsend and co-workers showed that the

C5-bound hydrogen is exchanged during the CarC-catalyzed epimerization.⁷⁴ The epimerization consumes α KG, despite the fact that the substrate is not oxidized in the transformation.⁷⁵ The resulting (3*S*,5*R*)-carbapenam-3-carboxylate is converted to (5*R*)-carbapenam-3-carboxylate by the same enzyme.⁷⁵ Computational studies suggested that the reaction is initiated by abstraction of the hydrogen atom of C5, presumably by the Fe(IV)–oxo intermediate generated by decarboxylation of α KG.⁷⁶ The Fe(III)–OH intermediate may serve as the source of the hydrogen atom in the formation of the (3*S*,5*R*)-carbapenam-3-carboxylate intermediate and regenerate the Fe(IV)–oxo complex.¹⁷ The Fe(IV)–oxo intermediate may then abstract the H-atom from C3, forming a stabilized, captodative C3 radical. Subsequent desaturation across the C2–C3 linkage via H-atom abstraction from C2 by the Fe(III)–OH species would be directly analogous to the mechanisms proposed for DAOCS and CAS.¹⁷

Summary and Outlook

The first examples of non-heme Fe(IV)–oxo enzyme intermediates have recently been detected in several α KG-dependent oxygenases. In each case, a large ²H-KIE on the decay of the intermediate has established that it is the C–H bond-cleaving complex. The spectroscopic parameters of the Fe(IV)–oxo complexes observed in hydroxylases TauD and P4H are almost identical, suggesting a conserved mechanism for substrate hydroxylation. The presence of two Fe(IV)–oxo complexes, which are apparently in rapid equilibrium and are presumably different

conformers, in CytC3 may be relevant to the divergent reactivity of the halogenases. Ongoing efforts are aimed at elucidating the molecular structure of the Fe(IV) complexes by a combination of spectroscopic and computational methods. We anticipate that further insight into the factors that determine the outcome of the oxidations catalyzed by high-valent Fe-oxo intermediates may be obtained from studies of other mononuclear non-heme enzymes (e.g., enzymes involved in the biosyntheses of β -lactam antibiotics) for which alternative reaction pathways have been proposed.

This work conducted in our laboratories and described herein was supported by the National Institutes of Health (Grant GM-69657 to J.M.B. and C.K. and Grants GM-20011 and GM-49338 to C.T.W.), the donors of the American Chemical Society Petroleum Research Fund (Grant ACS-PRF 41170-G3 to C.K.), the Arnold and Mabel Beckman Foundation (Young Investigator Award to C.K.), and the Dreyfus Foundation (Camille Dreyfus Teacher Scholar Award to C.K.). D.G.F. is supported by the Damon Runyon Cancer Research Foundation Postdoctoral Fellowship (DRG-1893-05).

References

- Costas, M.; Mehn, M. P.; Jensen, M. P.; Que, L., Jr. Dioxygen Activation at Mononuclear Nonheme Iron Active Sites: Enzymes, Models, and Intermediates. *Chem. Rev.* **2004**, *104*, 939–986.
- Solomon, E. I.; Brunold, T. C.; Davis, M. I.; Kemsley, J. N.; Lee, S.-K.; Lehnert, N.; Neese, F.; Skulan, A. J.; Yang, Y.-S.; Zhou, J. Geometric and Electronic Structure/Function Correlations in Non-Heme Iron Enzymes. *Chem. Rev.* **2000**, *100*, 235–349.
- Hausinger, R. P. Fe(II) α -ketoglutarate-dependent hydroxylases and related enzymes. *Crit. Rev. Biochem. Mol. Biol.* **2004**, *39*, 21–68.
- Vaillancourt, F. H.; Yeh, E.; Vosburg, D. A.; Garneau-Tsodikova, S.; Walsh, C. T. Nature's Inventory of Halogenation Catalysts: Oxidative Strategies Predominate. *Chem. Rev.* **2006**, *106*, 3364–3378.
- Fitzpatrick, P. F. Mechanism of Aromatic Amino Acid Hydroxylation. *Biochemistry* **2003**, *42*, 14083–14091.
- Rocklin, A. M.; Tierney, D. L.; Kofman, V.; Brunhuber, N. M. W.; Hoffman, B. M.; Christoffersen, R. E.; Reich, N. O.; Lipscomb, J. D.; Que, L., Jr. Role of the nonheme Fe(II) center in the biosynthesis of the plant hormone ethylene. *Proc. Natl. Acad. Sci. U.S.A.* **1999**, *96*, 7905–7909.
- Baldwin, J. E.; Bradley, M. Isopenicillin N synthase: Mechanistic studies. *Chem. Rev.* **1990**, *90*, 1079–1088.
- Neidig, M. L.; Decker, A.; Choroba, O. W.; Huang, F.; Kavana, M.; Moran, G. R.; Spencer, J. B.; Solomon, E. I. Spectroscopic and electronic structure studies of aromatic electrophilic attack and hydrogen-atom abstraction by non-heme iron enzymes. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 12966–12973.
- Koehntop, K. D.; Emerson, J. P.; Que, L., Jr. The 2-His-1-carboxylate facial triad: A versatile platform for dioxygen activation by mononuclear non-heme iron(II) enzymes. *J. Biol. Inorg. Chem.* **2005**, *10*, 87–93.
- Groves, J. T. Key elements of the chemistry of cytochrome P-450. The oxygen rebound mechanism. *J. Chem. Educ.* **1985**, *62*, 928–931.
- Rohde, J.-U.; In, J.-H.; Lim, M. H.; Brennessel, W. W.; Bukowski, M. R.; Stubna, A.; Münck, E.; Nam, W.; Que, L., Jr. Crystallographic and Spectroscopic Characterization of a Nonheme Fe(IV)=O complex. *Science* **2003**, *299*, 1037–1039.
- Lim, M. H.; Rohde, J.-U.; Stubna, A.; Bukowski, M. R.; Costas, M.; Ho, R. Y. N.; Münck, E.; Nam, W.; Que, L., Jr. An Fe^{IV}=O complex of a tetradentate tripodal nonheme ligand. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 3665–3670.
- Tiago de Oliveira, F.; Chanda, A.; Banerjee, D.; Shan, X.; Mondal, S.; Que, L., Jr.; Bominaar, E. L.; Münck, E.; Collins, T. J. Chemical and Spectroscopic Evidence for an Fe^V-Oxo Complex. *Science* **2007**, *315*, 835–838.
- Pestovsky, O.; Stoian, S.; Bominaar, E. L.; Shan, X.; Münck, E.; Que, L., Jr.; Bakac, A. Aqueous Fe^{IV}=O: Spectroscopic identification and oxo-group exchange. *Angew. Chem., Int. Ed.* **2005**, *44*, 6871–6874.
- Kaizer, J.; Klinker, E. J.; Oh, N. Y.; Rohde, J.-U.; Song, W. J.; Stubna, A.; Kim, J.; Münck, E.; Nam, W.; Que, L., Jr. Nonheme Fe^{IV}O Complexes That Can Oxidize the C-H Bonds of Cyclohexane at Room Temperature. *J. Am. Chem. Soc.* **2004**, *126*, 472–473.
- Johnson-Winters, K.; Purpero, V. M.; Kavana, M.; Nelson, T.; Moran, G. R. (4-Hydroxyphenyl)pyruvate Dioxygenase from *Streptomyces avermitilis*: The Basis for Ordered Substrate Addition. *Biochemistry* **2003**, *42*, 2072–2080.
- Kershaw, N. J.; Caines, M. E. C.; Sleeman, M. C.; Schofield, C. J. The enzymology of clavam and carbapenem biosynthesis. *Chem. Commun.* **2005**, 4251–4263.
- Kivirikko, K. I.; Myllyharju, J. Prolyl 4-hydroxylases and their protein disulfide isomerase subunit. *Matrix Biol.* **1998**, *16*, 357–368.
- Ivan, M.; Kondo, K.; Yang, H.; Kim, W.; Valiando, J.; Ohh, M.; Salic, A.; Asara, J. M.; Lane, W. S.; Kaelin, W. G., Jr. HIF α targeted for VHL-mediated destruction by proline hydroxylation: Implications for O₂ sensing. *Science* **2001**, *292*, 464–468.
- Jaakkola, P.; Mole, D. R.; Tian, Y.-M.; Wilson, M. I.; Gielbert, J.; Gaskell, S. J.; von Kriegsheim, A.; Hebestreit, H. F.; Mukherji, M.; Schofield, C. J.; Maxwell, P. H.; Pugh, C. W.; Ratcliffe, P. J. Targeting of HIF α to the von Hippel-Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation. *Science* **2001**, *292*, 468–472.
- Epstein, A. C. R.; Gleadle, J. M.; McNeill, L. A.; Hewitson, K. S.; O'Rourke, J.; Mole, D. R.; Mukherji, M.; Metzén, E.; Wilson, M. I.; Dhanda, A.; Tian, Y.-M.; Masson, N.; Hamilton, D. L.; Jaakkola, P.; Barstead, R.; Hodgkin, J.; Maxwell, P. H.; Pugh, C. W.; Schofield, C. J.; Ratcliffe, P. J. *C. elegans* EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation. *Cell* **2001**, *107*, 43–54.
- Bruick, R. K.; McKnight, S. L. A conserved family of prolyl-4-hydroxylases that modify HIF. *Science* **2001**, *294*, 1337–1340.
- Hirota, K.; Semenza, G. L. Regulation of hypoxia-inducible factor 1 by prolyl and asparaginyl hydroxylases. *Biochem. Biophys. Res. Commun.* **2005**, *338*, 610–616.
- Duncan, T.; Treweek, S. C.; Koivisto, P.; Bates, P. A.; Lindahl, T.; Sedgwick, B. Reversal of DNA alkylation damage by two human dioxygenases. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 16660–16665.
- Falnes, P. Ø.; Johansen, R. F.; Seeberg, E. AlkB-mediated oxidative demethylation reverses DNA damage in *Escherichia coli*. *Nature* **2002**, *419*, 178–182.
- Tsukada, Y.-i.; Fang, J.; Erdjument-Bromage, H.; Warren, M. E.; Borchers, C. H.; Tempst, P.; Zhang, Y. Histone demethylation by a family of JmjC domain-containing proteins. *Nature* **2006**, *439*, 811–816.
- Cloos, P. A. C.; Christensen, J.; Agger, K.; Maiolica, A.; Rappsilber, J.; Antal, T.; Hansen, K. H.; Helin, K. The putative oncogene GASC1 demethylates tri- and dimethylated lysine 9 on histone H3. *Nature* **2006**, *442*, 307–311.
- Klose, R. J.; Yamane, K.; Bae, Y.; Zhang, D.; Erdjument-Bromage, H.; Tempst, P.; Wong, J.; Zhang, Y. The transcriptional repressor JHD3A demethylates trimethyl histone H3 lysine 9 and lysine 36. *Nature* **2006**, *442*, 312–316.
- Hanuske-Abel, H. M.; Günzler, V. A stereochemical concept for the catalytic mechanism of prolylhydroxylase. Applicability to classification and design of inhibitors. *J. Theor. Biol.* **1982**, *94*, 421–455.
- Bollinger, J. M., Jr.; Price, J. C.; Hoffart, L. M.; Barr, E. W.; Krebs, C. Mechanism of Oxygen Activation by Taurine: α -Ketoglutarate Dioxygenase (TauD) from *Escherichia coli*. *Eur. J. Inorg. Chem.* **2005**, *2005*, 4245–4254.
- Price, J. C.; Barr, E. W.; Tirupati, B.; Bollinger, J. M., Jr.; Krebs, C. The First Direct Characterization of a High-Valent Iron Intermediate in the Reaction of an α -Ketoglutarate-Dependent Dioxygenase: A High-Spin Fe(IV) Complex in Taurine/ α -Ketoglutarate Dioxygenase (TauD) from *Escherichia coli*. *Biochemistry* **2003**, *42*, 7497–7508.
- Krebs, C.; Price, J. C.; Baldwin, J.; Saleh, L.; Green, M. T.; Bollinger, J. M., Jr. Rapid Freeze-Quench ⁵⁷Fe Mössbauer Spectroscopy: Monitoring Changes of an Iron-Containing Active Site during a Biochemical Reaction. *Inorg. Chem.* **2005**, *44*, 742–757.
- Price, J. C.; Barr, E. W.; Glass, T. E.; Krebs, C.; Bollinger, J. M., Jr. Evidence for Hydrogen Abstraction from C1 of Taurine by the High-Spin Fe(IV) Intermediate Detected during Oxygen Activation by Taurine: α -Ketoglutarate Dioxygenase (TauD). *J. Am. Chem. Soc.* **2003**, *125*, 13008–13009.
- Bollinger, J. M., Jr.; Krebs, C. Stalking intermediates in oxygen activation by iron enzymes: Motivation and method. *J. Inorg. Biochem.* **2006**, *100*, 586–605.
- Proshlyakov, D. A.; Henshaw, T. F.; Monterosso, G. R.; Ryle, M. J.; Hausinger, R. P. Direct Detection of Oxygen Intermediates in the Non-Heme Fe Enzyme Taurine/ α -Ketoglutarate Dioxygenase. *J. Am. Chem. Soc.* **2004**, *126*, 1022–1023.

- (36) Riggs-Gelasco, P. J.; Price, J. C.; Guyer, R. B.; Brehm, J. H.; Barr, E. W.; Bollinger, J. M., Jr.; Krebs, C. EXAFS Spectroscopic Evidence for an Fe=O Unit in the Fe(IV) Intermediate Observed during Oxygen Activation by Taurine: α -Ketoglutarate Dioxygenase. *J. Am. Chem. Soc.* **2004**, *126*, 8108–8109.
- (37) Sinnecker, S.; Svendsen, N.; Barr, E. W.; Ye, S.; Bollinger, J. M., Jr.; Neese, F.; Krebs, C. Spectroscopic and Computational Evaluation of the Structure of the High-Spin Fe(IV)-Oxo Intermediates in Taurine: α -Ketoglutarate Dioxygenase from *Escherichia coli* and its His99Ala Ligand Variant. *J. Am. Chem. Soc.* **2007**, *129*, 6168–6179.
- (38) Price, J. C.; Barr, E. W.; Hoffart, L. M.; Krebs, C.; Bollinger, J. M., Jr. Kinetic Dissection of the Catalytic Mechanism of Taurine: α -Ketoglutarate Dioxygenase (TauD) from *Escherichia coli*. *Biochemistry* **2005**, *44*, 8138–8147.
- (39) Eriksson, M.; Myllyharju, J.; Tu, H.; Hellman, M.; Kivirikko, K. I. Evidence for 4-hydroxyproline in viral proteins. Characterization of a viral prolyl 4-hydroxylase and its peptide substrates. *J. Biol. Chem.* **1999**, *274*, 22131–22134.
- (40) Hoffart, L. M.; Barr, E. W.; Guyer, R. B.; Bollinger, J. M., Jr.; Krebs, C. A Unified Mechanism for the α -Ketoglutarate-Dependent Dioxygenases: Evidence for a C-H-Cleaving Fe(IV) Complex in Prolyl-4-Hydroxylase. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 14738–14743.
- (41) Thornburg, L. D.; Lai, M. T.; Wishnok, J. S.; Stubbe, J. A non-heme iron protein with heme tendencies: An investigation of the substrate specificity of thymine hydroxylase. *Biochemistry* **1993**, *32*, 14023–14033.
- (42) Thornburg, L. D.; Stubbe, J. Mechanism-based inactivation of thymine hydroxylase, an α -ketoglutarate-dependent dioxygenase, by 5-ethynyluracil. *Biochemistry* **1993**, *32*, 14034–14042.
- (43) Lai, M.-t.; Wu, W.; Stubbe, J. Characterization of a Novel, Stable Norcaradiene Adduct Resulting from the Inactivation of Thymine Hydroxylase by 5-Ethynyluracil. *J. Am. Chem. Soc.* **1995**, *117*, 5023–5030.
- (44) Vaillancourt, F. H.; Yin, J.; Walsh, C. T. SyrB2 in syringomycin E biosynthesis is a nonheme FeII α -ketoglutarate- and O₂-dependent halogenase. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 10111–10116.
- (45) Vaillancourt, F. H.; Yeh, E.; Vosburg, D. A.; O'Connor, S. E.; Walsh, C. T. Cryptic chlorination by a non-haem iron enzyme during cyclopropyl amino acid biosynthesis. *Nature* **2005**, *436*, 1191–1194.
- (46) Galonić, D. P.; Vaillancourt, F. H.; Walsh, C. T. Halogenation of Unactivated Carbon Centers in Natural Product Biosynthesis: Trichlorination of Leucine during Barbamide Biosynthesis. *J. Am. Chem. Soc.* **2006**, *128*, 3900–3901.
- (47) Ueki, M.; Galonić, D. P.; Vaillancourt, F. H.; Garneau-Tsodikova, S.; Yeh, E.; Vosburg, D. A.; Schroeder, F. C.; Osada, H.; Walsh, C. T. Enzymatic Generation of the Antimetabolite γ,γ -Dichloroaminobutyrate by NRPS and Mononuclear Iron Halogenase Action in a Streptomyces. *Chem. Biol.* **2006**, *13*, 1183–1191.
- (48) Blasiak, L. C.; Vaillancourt, F. H.; Walsh, C. T.; Drennan, C. L. Crystal structure of the non-haem iron halogenase SyrB2 in syringomycin biosynthesis. *Nature* **2006**, *440*, 368–371.
- (49) Galonić, D. P.; Barr, E. W.; Walsh, C. T.; Bollinger, J. M., Jr.; Krebs, C. Two Interconverting Fe(IV) Intermediates in Aliphatic Chlorination by the Halogenase CytC3. *Nat. Chem. Biol.* **2007**, *3*, 113–116.
- (50) Kojima, T.; Leising, R. A.; Yan, S.; Que, L., Jr. Alkane functionalization at nonheme iron centers. Stoichiometric transfer of metal-bound ligands to alkane. *J. Am. Chem. Soc.* **1993**, *115*, 11328–11335.
- (51) Samson, S. M.; Belagaje, R.; Blankenship, D. T.; Chapman, J. L.; Perry, D.; Skatrud, P. L.; VanFrank, R. M.; Abraham, E. P.; Baldwin, J. E.; Queener, S. W.; Ingolia, T. D. Isolation, sequence determination and expression in *Escherichia coli* of the isopenicillin N synthetase gene from *Cephalosporium acremonium*. *Nature* **1985**, *318*, 191–194.
- (52) Baldwin, J. E.; Abraham, E. Biosynthesis of penicillins and cephalosporins. *Nat. Prod. Rep.* **1988**, *5*, 129–145.
- (53) Roach, P. L.; Clifton, I. J.; Hensgens, C. M. H.; Shibta, N.; Schofield, C. J.; Hajdu, J.; Baldwin, J. E. Structure of isopenicillin N synthase complexed with substrate and the mechanism of penicillin formation. *Nature* **1997**, *387*, 827–830.
- (54) Ogle, J. M.; Clifton, I. J.; Rutledge, P. J.; Elkins, J. M.; Burzlauff, N. I.; Adlington, R. M.; Roach, P. L.; Baldwin, J. E. Alternative oxidation by isopenicillin N synthase observed by X-ray diffraction. *Chem. Biol.* **2001**, *8*, 1231–1237.
- (55) Baldwin, J. E.; Adlington, R. M.; Domayne-Hayman, B. P.; Knight, G.; Ting, H.-H. Use of the cyclopropylcarbonyl test to detect a radical-like intermediate in penicillin biosynthesis. *J. Chem. Soc., Chem. Commun.* **1987**, 1661–1663.
- (56) Burzlauff, N. I.; Rutledge, P. J.; Clifton, I. J.; Hensgens, C. M. H.; Pickford, M.; Adlington, R. M.; Roach, P. L.; Baldwin, J. E. The reaction cycle of isopenicillin N synthase observed by X-ray diffraction. *Nature* **1999**, *401*, 721–724.
- (57) Baldwin, J. E.; Adlington, R. M.; Bradley, M.; Turner, N. J.; Pitt, A. R. Evidence for epoxide formation from isopenicillin N synthase. *J. Chem. Soc., Chem. Commun.* **1989**, 978–981.
- (58) Elkins, J. M.; Rutledge, P. J.; Burzlauff, N. I.; Clifton, I. J.; Adlington, R. M.; Roach, P. L.; Baldwin, J. E. Crystallographic studies on the reaction of isopenicillin N synthase with an unsaturated substrate analogue. *Org. Biomol. Chem.* **2003**, *1*, 1455–1460.
- (59) Baldwin, J. E.; Adlington, R. M.; Flitsch, S. L.; Ting, H. H.; Turner, N. J. Penicillin biosynthesis: The origin of hydroxy groups in β -lactams derived from unsaturated substrates. *J. Chem. Soc., Chem. Commun.* **1986**, 1305–1308.
- (60) Elson, S. W.; Baggaley, K. H.; Gillett, J.; Holland, S.; Nicholson, N. H.; Sime, J. T.; Woronicki, S. R. Isolation of two novel intracellular β -lactams and a novel dioxygenase cyclizing enzyme from *Streptomyces clavuligerus*. *J. Chem. Soc., Chem. Commun.* **1987**, 1736–1738.
- (61) Salowe, S. P.; Marsh, E. N.; Townsend, C. A. Purification and characterization of clavamate synthase from *Streptomyces clavuligerus*: An unusual oxidative enzyme in natural product biosynthesis. *Biochemistry* **1990**, *29*, 6499–6508.
- (62) Salowe, S. P.; Krol, W. J.; Iwata-Reuyl, D.; Townsend, C. A. Elucidation of the order of oxidations and identification of an intermediate in the multistep clavamate synthase reaction. *Biochemistry* **1991**, *30*, 2281–2292.
- (63) Baldwin, J. E.; Lloyd, M. D.; Wha-Son, B.; Schofield, C. J.; Elson, S. W.; Baggaley, K. H.; Nicholson, N. H. A substrate analogue study on clavaminic acid synthase: Possible clues to the biosynthetic origin of proclavaminic acid. *J. Chem. Soc., Chem. Commun.* **1993**, 500–502.
- (64) Busby, R. W.; Townsend, C. A. A single monomeric iron center in clavamate synthase catalyzes three nonsuccessive oxidative transformations. *Bioorg. Med. Chem.* **1996**, *4*, 1059–1064.
- (65) Lloyd, M. D.; Merritt, K. D.; Leeb, V.; Sewell, T. J.; Wha-Son, B.; Baldwin, J. E.; Schofield, C. J.; Elson, S. W.; Baggaley, K. H.; Nicholson, N. H. Product-substrate engineering by bacteria: Studies on clavamate synthase, a trifunctional dioxygenase. *Tetrahedron* **1999**, *55*, 10201–10220.
- (66) Zhou, J.; Kelly, W. L.; Bachmann, B. O.; Gunsior, M.; Townsend, C. A.; Solomon, E. I. Spectroscopic Studies of Substrate Interactions with Clavamate Synthase 2, a Multifunctional α -KG-Dependent Non-Heme Iron Enzyme: Correlation with Mechanisms and Reactivities. *J. Am. Chem. Soc.* **2001**, *123*, 7388–7398.
- (67) Townsend, C. A. New reactions in clavulanic acid biosynthesis. *Curr. Opin. Chem. Biol.* **2002**, *6*, 583–589.
- (68) Lloyd, M. D.; Lee, H.-J.; Harlos, K.; Zhang, Z.-H.; Baldwin, J. E.; Schofield, C. J.; Charnock, J. M.; Garner, C. D.; Hara, T.; Terwisscha van Scheltinga, A. C.; Valegård, K.; Viklund, J. A. C.; Hajdu, J.; Andersson, I.; Danielsson, A.; Bhikhabhai, R. Studies on the active site of deacetoxycephalosporin C synthase. *J. Mol. Biol.* **1999**, *287*, 943–960.
- (69) Valegård, K.; Terwisscha van Scheltinga, A. C.; Dubus, A.; Ranghino, G.; Öster, L. M.; Hajdu, J.; Andersson, I. The structural basis of cephalosporin formation in a mononuclear ferrous enzyme. *Nat. Struct. Mol. Biol.* **2004**, *11*, 95–101.
- (70) Pang, C. P.; White, R. L.; Abraham, E. P.; Crout, D. H.; Lutstorf, M.; Morgan, P. J.; Derome, A. E. Stereochemistry of the incorporation of valine methyl groups into methylene groups in cephalosporin C. *Biochem. J.* **1984**, *222*, 777–788.
- (71) Townsend, C. A.; Theis, A. B.; Neese, A. S.; Barrabee, E. B.; Poland, D. Stereochemical fate of chiral methyl of valine in the ring expansion of penicillin N to deacetoxycephalosporin C. *J. Am. Chem. Soc.* **1985**, *107*, 4760–4767.
- (72) Baldwin, J. E.; Adlington, R. M.; Kang, T. W.; Lee, E.; Schofield, C. J. The ring expansion of penams to cepams: A possible biomimetic process. *Tetrahedron* **1988**, *44*, 5953–5937.
- (73) Baldwin, J. E.; Adlington, R. M.; Crouch, N. P.; Schofield, C. J.; Turner, N. J.; Aplin, R. T. Cephalosporin biosynthesis: A branched pathway sensitive to an isotope effect. *Tetrahedron* **1991**, *47*, 9881–9900.
- (74) Stapon, A.; Li, R.; Townsend, C. A. Carbapenem Biosynthesis: Confirmation of Stereochemical Assignments and the Role of CarC in the Ring Stereoconversion Process from L-Proline. *J. Am. Chem. Soc.* **2003**, *125*, 8486–8493.
- (75) Stapon, A.; Li, R.; Townsend, C. A. Synthesis of (3S,5R)-carbapenam-3-carboxylic acid and its role in carbapenem biosynthesis and the stereoconversion problem. *J. Am. Chem. Soc.* **2003**, *125*, 15746–15747.
- (76) Topf, M.; Sandala, G. M.; Smith, D. M.; Schofield, C. J.; Easton, C. J.; Radom, L. The unusual bifunctional catalysis of epimerization and desaturation by carbapenem synthase. *J. Am. Chem. Soc.* **2004**, *126*, 9932–9933.

AR700066P