

Metal Unsaturation and Ligand Hemilability in Suzuki Coupling

ZHIQIANG WENG,* SHIHUI TEO, AND
T. S. ANDY HOR*

Department of Chemistry, National University of Singapore,
Kent Ridge, Singapore 117543

Received August 1, 2006

ABSTRACT

The combinative and complementary use of a hemilabile difunctional ligand on a metal, notably palladium, that is coordinatively and electronically unsaturated has led to the isolation of a string of unexpected low-valent complexes that are structurally intriguing. The ligands of interest are primarily ferrocenes functionalized by [P,N] and [P,O] donors. The characterization of these active Suzuki catalysts, which support sp^2 – sp^2 couplings, give valuable insights into the key Suzuki intermediates such as those arising from the reductive elimination, transmetalation, and oxidative addition steps. In this Account, we shall review and discuss our recent results in relation to selected developments in other laboratories.


1. Introduction

Transition metal-catalyzed C–C and C–X (X = heteroatom) bond formation is a powerful synthetic methodology in organic syntheses.^{1,2} A typical example is found in palladium-catalyzed (Miyaura) Suzuki cross-coupling³ (eq 1) of haloarenes with aryl boronic acids. Recent catalyst developments have expanded the application scope to natural product syntheses, materials design, and large-scale pharmaceutical production etc.⁴ The attractive features of this coupling include the flexible use of organic solvents with common inorganic bases, readily available substrates, air and moisture stability, functional group tolerance, coupling of sterically demanding groups, low toxicity of boronic acids, and facile removal of the boron-containing byproducts.

Zhiqiang Weng (born in 1971 in China) came to Singapore to join the graduate program of the National University of Singapore (NUS) after he graduated from Fuzhou University with B.Sc. (1994) and M.Sc. (1997) degrees. Upon graduation with a Ph.D. (2003) degree in the area of organochromium chemistry, he joined the Institute of Chemical and Engineering Sciences (ICES) of Singapore in 2003 as a Research Fellow, starting work in homogeneous catalysis. In October 2005, he returned to NUS as a Research Fellow to start his independent research, specializing in the synthesis of new and sensitive organotransition metal complexes and their catalytic applications in organic synthesis and olefin oligomerization.

Shihui Teo (born in 1981 in Singapore) graduated with a B.Sc. degree in 2003 from NUS. She then joined ICES as a Research Officer and started working in metallocene catalysis. In 2006, she returned to the graduate program of NUS under the advisory of Z. Weng and T. S. A. Hor. She is interested in chiral and functional catalysis.

T. S. Andy Hor (born in 1956 in Hong Kong) is Professor and Head of Chemistry of NUS and President of the Singapore National Institute of Chemistry. He joined NUS in 1984 after graduating from Imperial College [B.Sc.(Honors)], Oxford (D.Phil.) (with D. M. P. Mingos), and postdoctoral research at Yale University (with R. D. Adams). He has authored ~190 international papers on heteropolymetallic aggregates, supramolecular complexes, and organotransition metal catalysts. He was a fellow of Humboldt, Commonwealth, ASAIHL, Anthony Mason, Wilsomour, Frances Lion, etc. He currently serves the advisory board of *Inorganica Chimica Acta* and *Chemistry Asian Journal*.




The common Suzuki catalysts are phosphine complexes such as $Pd(PPh_3)_4$ with strong P-donors. The coupling pathway requires a sequential oxidative addition at an active Pd(0) by aryl halide, activation followed by transmetalation [this is not a typical transmetalation because boron is strictly not a metal; however, the idea of organo–ligand crossover between an organometalloid and an organopalladium to prepare the latter for reductive elimination is consistent with the concept of transmetalation; details of this step are described in the literature (e.g., Corbet, J.-P.; Mignani, G. *Chem. Rev.* 2006, 106, 2651–2710)] with the organoboronate substrate, and reductive elimination to yield the desirable product. A precatalyst such as $Pd(PPh_3)_4$ would need to enter the catalytic cycle through two successive ligand dissociations to give the 14-electron active catalytic complex $Pd^0(PPh_3)_2$. As such low-coordinate and low-valent palladium complexes are notoriously unstable, their formation is energetically unfavorable. Facile decomposition of such active catalysts also contributes to poor turnover. It is therefore desirable if one could design a “smart catalyst” that enters the catalytic cycle without going through a high energetic barrier and yet is both chemically stable and catalytically active. A strategy that we have developed recently is the complementary use of electronically and coordinatively tunable hemilabile ligands⁵ and reactive low-coordinate metals.⁶ These ligands are sensitive to the dynamic needs of the metal at different stages of the catalytic cycle. Ideally, such a ligand not only can protect the catalytic intermediates but also can activate all the key steps.


Accordingly, as an extension of our ongoing work in 1,1'-bis(diphenylphosphino)ferrocene (dppf) chemistry,⁷ we designed a series of ferrocene-based hemilabile ligands with hybrid and contrasting donors. The ferrocenyl core plays the important role of a spectator in giving the difunctional donors the desirable coordinative mobility and serving as an electronic reservoir needed for pendant/donating switches. The redox-active Fe(II) provides an additional electronic buffer that allows the catalytic metal to remain active in different redox stages. Such a smart catalyst helps in tackling challenging problems such as C–Cl and C–F bond activation, sp^3 – sp^3 coupling, and sterically hindered, room temperature, and aqueous reactions. Some of these challenges in Suzuki coupling can be achieved through the use of ferrocene-based [P,N], [P,O], [P,CN σ], [P,CN π], and [N,N] ligands in conjunction with active Pd compounds. Some unusual and highly active catalytic species that are both electronically and coordinatively unsaturated complexes have been isolated and crystallographically characterized. We shall summarize and review these from the perspective of other selected developments in this field.

* To whom correspondence should be addressed. E-mail: andyhor@nus.edu.sg. Fax: (65) 6873 1324.

Scheme 1



Scheme 2



2. Suzuki Coupling with Phosphine-Free Catalysts


Our earlier use of phosphine-free catalysts such as [N,N]-functionalized ferrocene-based ligands is targeted to provide a nontoxic system that is moisture- and air-stable. Other related approaches are found in N-heterocyclic carbene complexes,⁸ imidazole,⁹ palladium powder,¹⁰ acetate,^{11a} metal-free^{11b} and ligand-free^{11c} systems, and N-coordinated palladium systems, which include cyclometallated palladium complexes with imine,¹² oxime,^{13a} oxazoline,^{13b} diazabutadiene,^{14a} Dabco,^{14b} bis(oxazolinyl)pyrrole,¹⁵ and aminopyridyl.¹⁶ In sharp contrast to the rich coordination chemistry of its phosphorus analogue viz. dppf, the ligand chemistry of [N,N] ferrocene-based ligands is ill-developed.

The diimine functionalized ferrocenyl ligand $\text{Fe}(\eta\text{-C}_5\text{H}_4\text{N}=\text{CHPh})_2$ (**1**) coordinates easily at room temperature with $\text{PdCl}_2(\text{MeCN})_2$ to give a stable chelate complex $\text{PdCl}_2\text{Fe}(\eta\text{-C}_5\text{H}_4\text{N}=\text{CHPh})_2$ (**2**) (Scheme 1).¹⁷ Similar reaction using a bidentate diamine with a larger bite angle, $\text{Fe}[\eta\text{-C}_5\text{H}_4\text{C}_2\text{H}_4\text{N}(\text{CH}_3)_2]_2$ (**3**), leads to the isolation of a stable metallamacrocyclic complex $\text{Pd}_2\text{Cl}_4\{\text{Fe}[\eta\text{-C}_5\text{H}_4(\text{CH}_2)_2\text{N}(\text{CH}_3)_2]\}_2$ (**4**) (Scheme 2).¹⁸

Complex **2** is highly active toward Suzuki cross-coupling of aryl iodides and bromides with aryl boronic acids as a suspension in aqueous media. The products can be conveniently isolated, whereas the recovered catalyst remains active for up to five runs.

Complex **4** can be viewed as a dimeric version of **2**. Two Pd(II) centers are doubly bridged by two metallo-ligands to give effectively a $\{\text{Pd}_2\text{Fe}_2\}$ heterometallic “molecular rectangle”. Use of soluble metal polyions as catalysts is rare but could offer advantages such as chemical stability, bimetallic cooperation, and single catalysts with multiple sites. Complex **4**, which is air- and moisture-stable, promotes cross-coupling of aryl boronic acids with deactivated, electron-rich aryl bromides and

Chart 1. P,O Ligands for Suzuki Coupling





activated, electron-poor substrates under ambient conditions, giving generally high yields even under low catalytic loads. There is no evidence of any Fe → Pd dative bonding stabilization in **4** or its catalytic intermediates. Active and unsaturated catalysts that are supported by only weak N donors are generally too reactive to be isolated.

3. Suzuki Coupling with Catalysts Containing Hemilabile Ligands

Currently, the most common ligands used for Suzuki coupling reaction are electron-rich and sterically hindered monophosphine.¹⁹ The use of N-heterocyclic carbenes as phosphine alternatives is emerging.²⁰ These ligands have good donating properties, giving stable catalysts, but are handicapped by the lack of coordinative flexibility. Therefore, designing a new class of ligands that have hybrid and difunctional properties to support different catalytic intermediates formed within a single system is appealing.

The P,O and P,N ligands reported by Buchwald et al. exemplify the value of hybrid ligands. The palladium complexes of a new ligand, 2-(2',6'-dimethoxybiphenyl)-dicyclohexylphosphine (SPhos, **5**) (Chart 1), efficiently catalyze highly hindered aryl boronic acid and aryl halides.²¹ Milstein et al. prepared an electron-rich, bulky methoxy-benzyl phosphine (dmobp, **6**) to stabilize unsaturated Pd(0), Pd(I), and Pd(II). Coordination of the methoxy group in the proximity highlights the hemilabile effect.²²

Other P,O ligands used in Suzuki coupling include amide phosphine (**7**),^{23a,b} acetal phosphines (**8**),^{23c} ferrocenyl-supported alkoxy phosphines (**9**),^{23d} and **10**.^{23e} Other donor combinations such as P,N and P,S ligands (Chart 2) have also been used. Examples include 2-(dicyclohexylphosphino)-2'-(N,N-dimethylamino)biphenyl (**11**),^{24a} 2-(dimethylamino)-2'-diphenylphosphino-1,1'-binaphthyl (MAP) through P,C_σ coordination (**12**),^{24b} phosphino-substituted N-aryl pyrroles (PAP, **13**),^{24c} N-(2-diphenylphosphino)phenyl-2,6-diisopropylanilide (**14**),^{24d} axially dissymmetric P,S-heterodonor ligands (**15**),^{25a} (1-phenylphosphino)-1'-(methylthio)ferrocene (**16**),^{25b} iminophosphine,^{25c} aminophosphine,^{25d} 1-phosphabarrelene phosphine sulfide-substituted ligands,^{25e} and the P,S-chelating thioether aminophosphonites.^{25f}

Chart 2. P,N and P,S Ligands for Suzuki Coupling

Chart 3. Ferrocene-Based P,N and P,O Ligands for Suzuki Coupling


To fully exploit the hemilabile function, the desirable ligand should have the following features. (1) It must be a difunctional ligand with at least two basic sites that have significantly different donating abilities. (2) The electronic character of each donor atom can be tuned by chemical alteration of its attached or nearby substituents. (3) The two donating sites are separated by a metallocenyl moiety that is stereogeometrically flexible and redox active. (4) Its potential as a unidentate, chelating, and bridging ligand must be demonstrated. (5) It can support unsaturated metal through its electronic and spatial effects. On the basis of these considerations, the ferrocene-based [P,N] (**17**) and [P,O] ligands (**18**) were prepared (Chart 3).


A mixture of $\text{Pd}_2(\text{dba})_3$ and $[\eta\text{-C}_5\text{H}_4\text{CH}=\text{N}(\text{C}_6\text{H}_5)]\text{Fe}[\eta\text{-C}_5\text{H}_4\text{P}(t\text{-Bu})_2]$ (**17**) (dba = dibenzylideneacetone) efficiently catalyzes the Suzuki cross-couplings of a range of aryl boronic acids and aryl chlorides, affording the desired biaryl products in high isolated yields.²⁶ For example, 4-chlorobenzonitrile couples with phenyl boronic acid at a low catalyst load (0.01 mol % Pd) to give 4-biphenylcarbonitrile in a good TON of 10 000. Other catalysts that give exceedingly high TONs have been identified.²⁷

The advantage of using hybrid P,O ligands such as **18a-c** over monodentate monophosphine (**19**) or bidentate diphosphine (**20**) ligand is evident in the coupling between 4-chlorobenzonitrile and phenyl boronic acid (Table 1). Under similar catalytic conditions, the isolated yield of 4-acetylbiphenyl is quantitative for **18**, whereas the controls (**19** and **20**) gave insignificant products.²⁸ The co-presence of a strong coordinating group (phosphorus) and a much weaker and restrictive donor (oxygen) gives the ligands (**18**) the needed flexibility to produce high activities that are comparable to the activities of some other P,O ferrocenyl systems, such as aryl-MOPFs (**9**).^{23d}

Table 1. Effect of Ligand on the Suzuki Cross-Coupling^a

entry	ligand	isolated yield (%)
1	18a	84
2	18b	100
3	18c	100
4	19	trace
5	20	26


^a With a 0.2 mol % Pd load.

Scheme 3


The hemilability function meets the needs of the metal, which sequentially switches between saturated and unsaturated states. Use of these hybrid ligands helped us trap a number of saturated and unsaturated species under both stoichiometric and catalytic conditions. Reaction of $\text{Pd}_2(\text{dba})_3$ with **17** (1:1 L:Pd ratio) in a NMR tube gives two main products, $[\eta\text{-C}_5\text{H}_4\text{CH}=\text{N}(\text{C}_6\text{H}_5)]\text{Fe}[\eta\text{-C}_5\text{H}_4\text{P}(t\text{-Bu})_2]\text{Pd}(\text{dba})$ (**21**) and $\text{Pd}\{[\eta\text{-C}_5\text{H}_4\text{CH}=\text{N}(\text{C}_6\text{H}_5)]\text{Fe}[\eta\text{-C}_5\text{H}_4\text{P}(t\text{-Bu})_2]\}_2$ (**22**), formed in a ratio of ~4:1 (Scheme 3). When the concentration of **17** is doubled, **22** becomes the sole species. The latter can be synthesized (94%) independently from a reaction of **17** with $(\text{C}_5\text{H}_5)\text{Pd}(\text{C}_3\text{H}_5)$ (2:1).²⁹


Complex **21** is a 16-electron Pd(0) with a heterodifunctional chelate. It provides a crystallographic proof that an imine nitrogen, albeit weakly basic, can coordinate when the metal is unsaturated. The exposed metal is further stabilized by olefinic coordination from the adventitious dibenzylideneacetone (dba). dba-coordinated Pd(0) complexes have been identified in the literature,³⁰ including those in P,C (**23–25**),^{31,32} P,O (**26**),^{23c} P,N (**27**),³³ and P,P (**28**) complexes³⁴ (Chart 4). Although the role of dba in many catalytic cycles has been largely ignored, it could play a stabilizing function to keep the active Pd(0) in solution and suppress decomposition to Pd metal. The use of dba or hemilabile ligands to support molecular Pd(0) could be one of our best strategies for developing ligandless catalysts in truly homogeneous systems, as

opposed to those ligandless catalysts that are based on Pd/C.¹⁰ The benefits of coordinatively deficient [PdL] catalysts in Suzuki and other couplings have been described.³⁵



(21)

Some pertinent bond parameters of **21** are compared with those of other LPd⁰(η^2 -dba) complexes (Table 2). Their Pd–C_{olefin} bond distances are comparable and similar to that of (η^2 -dppf)Pt(η^2 -dba).³⁶ The Pd–P bond length [2.3385(10) Å] of **21** is notably longer and its chelate P–Pd–N angle [110.94(8)°] larger than the others. These indicate the steric and spatial demand of ligand **17**, which also helps to protect Pd(0) from overexposure. The weakness of the Pd–P bond in **21** may suggest a weak chelate. Indeed, in **22**, the ligand opens up to give a unidentate metalloligand with a dangling imine. This is an unusual complex not only because it is a 14-electron linear Pd(0), some of which²⁹ are given in Chart 5, but also because it is also rare to witness two potentially bidentate ligands on the same metal, each of which is unidentate thus resulting in a highly unsaturated metal. The ferrocenyl moiety stabilizes the active metal through a spatial shield.




[Pd(1)···H(23B) 2.931 Å]

(22)

A list of the Pd–P lengths and P–Pd–P angles for **22** and other Pd⁰(PR₃)₂ complexes is given in Table 3. Not all two-coordinated Pd(0) complexes are linear. The strict linearity is observed in **22** and **30–32**. Complexes **29** and **36** are slightly distorted, whereas **33–35** prefer a bent geometry. A supplementary intramolecular Pd–arene interaction in **33** could explain its angular distortion. In **22**, the closest Pd–X nonbonding contacts are found in Pd···H(23B)

Chart 4. A Sample of dba-Coordinated Complexes of Pd(0)



(2.931 Å), which is significantly longer than those in **29** and **31** (2.73 and 2.508 Å, respectively).²⁹ This rules out the possibility of agostic interaction and supports it being an authentic two-coordinate Pd(0).

The dynamic properties of **17** help to stabilize **21** and **22** under different stoichiometric conditions with or without supplementary support from the adventitious dba. These highly active complexes catalyze the formation of 4-biphenylcarbonitrile in near-quantitative yields at 90 °C under 0.25 mol % Pd (Table 4). At a lower catalyst load of 0.01 mol % (Pd), the activity of **22** is superior to that of **21** (100 and 52%, respectively).

An advantage of an unsaturated low-valent metal is that it could undergo oxidative addition with an aryl halide directly, without going through the barrier of reduction or ligand dissociation. In fact, this step could be the r.d.s. of palladium-catalyzed cross-coupling reactions.³⁷ Isolation of a catalytic active species like **21** or **22** enabled us to carry out the oxidative addition stoichiometrically. A mixture of **17** and Pd₂(dba)₃ (giving mainly **21** in situ) or a solution of **22** with C₆F₅I in THF at room temperature gives [η -C₅H₄CH=N(C₆H₅)]Fe[η -C₅H₄P(*t*-Bu)₂]Pd(II)(C₆F₅) (**37**) (41 or 31% yield, respectively) as the only identifiable product (Scheme 4).


Similar oxidative addition has been reported by Milstein et al. in the reaction between Pd(0) complex **34** (Scheme 5) and haloarenes which gives **38** with a P,O chelating Pd(II).²²

Complex **37** is a Pd(II) oxidative addition product with the expected P,N chelate and the more *trans*-labilizing aryl ligand opposite the weaker Pd–N bond. Preliminary experiments revealed that **37** reacts with phenylboronic acid in the presence of KF/Cs₂CO₃ in dioxane to give the hetero-coupling product 2,3,4,5,6-pentafluorobiphenyl in ca. 75% yield. This is consistent with **37** being a key intermediate in Suzuki coupling.

It is commonly assumed that metal-containing intermediates in Suzuki and related couplings are mononuclear complexes, such as that found in **37**. This is not necessarily correct especially in cases when phosphine is deficient

Table 2. Comparison of Selected Bond Parameters of [Pd⁰L(η²-dba)]

complex	Pd–P bond length (Å)	Pd–C _{olefin} bond length (Å) (mean)	P–Pd–E angle (E = N, O, or C) (deg)
21	2.3385(10)	2.129(3)	110.94(8)
23	2.2850(8)	2.133(3)	83.01(7)
24	2.3001(17)	2.143(6)	–
25	2.3269(18)	2.145(6)	96.25(18)
26	2.2924(7)	2.102(2)	87.38(5)
27	2.2655(12)	2.110(4)	97.96(6)
28	2.296(1)	2.150(4)	88.26(5)

Chart 5. Complexes of [Pd⁰(PR₃)₂]

Table 3. Comparison of Selected Bond Parameters of [Pd⁰(PR₃)₂]

complex	Pd–P bond length (Å) (mean)	P–Pd–P angle (deg)
22	2.2828(16)	180.0
29	2.278(4)	176.8(1)
30	2.285(3)	180.0
31	2.276(1)	180.0
32	2.2764(7)	180.0
33	2.2761(11)	154.82(4)
34	2.2913(19)	166.86(7)
35	2.2685(8)	164.48(3)
36	2.2861(4)	173.964(14)

Table 4. Catalytic Yield Comparison of **21 and **22****

catalyst	21	22
0.25 mol % (Pd)	98%	100% yield
0.01 mol % (Pd)	52%	100% yield



or when the hemilabile ligand carries a donor that is poorly basic. This is evident when the P,O ligand **18** is mixed with Pd₂(dba)₃ and C₆F₅I stoichiometrically. Several oxidative addition products (**39–42**) have been identified (Scheme 6); among them, **39** and **42** have been crystallographically established. Complex **39a** is a dinuclear Pd(II) with doubly bridging iodide and the ligand **18a** being a P-only donor with dangling cyclic acetal. It suggests that facile dimerization could provide a stabilizing mechanism for active oxidative addition intermediates. Although such dinuclear structure is saturated, it could return to its active form through bridge cleavage in a

coordinating solvent or through re-chelation of a hemilabile ligand. The latter is spectroscopically observed in **40**.

When the L:M ratio is ≥2:1, the mononuclear form regenerates with an active *trans*–*cis* isomerization between **41** and **42**. The latter formation allows productive *trans*-metallation to take place followed by *cis* coupling of the aryl residues. Isolation of these complexes pointed to two alternative catalytic pathways via the dinuclear (route A) or geometric isometric (route B) forms, both of which can be supported by a hemilabile ligand (Scheme 7).

Guram et al. isolated two similar oxidative addition products from the reaction of phenyl backbone-derived P,O ligands with Pd(dba)₂ and excess 4-*tert*-Bu-C₆H₄Br, viz., (P,O)₂Pd(4-*tert*-Bu-C₆H₄)Br (**43**) and (P,O)Pd(4-*tert*-Bu-C₆H₄)Br (**44**) (Scheme 8).³⁸ It is notable that introducing a methyl group into the dioxolane can influence the catalytic intermediates by favoring the monophosphine over the *trans*-bisphosphine species.

Grotjahn et al. also reported a new type of hemilabile imidazol-2-ylphosphine ligand system, which allows steric adjustment at the P-donor and basic imidazole N-3 (Scheme 9).^{29e} The two types of oxidative addition products are represented by the expected *trans*-bis(phosphine)Pd(R)(X) **45** and the mononuclear P,N chelating **46**. The nature of haloarenes and R on the phosphine also influence the product preference. Complexes **44** and **46** are analogous to **37**, whereas **43** and **45** are related to **42**.

Scheme 4

Scheme 5



Since the M:L ratios are different in the two routes, one could in principle push the catalytic path toward route B from route A by increasing the ligand dosage. A slight increase in TON (from ~ 1000 to 2000) is observed for **18a**, while there is an ~ 20 -fold TON increase [from ~ 2000 (max) to ~ 44600] for most of the electron-rich substrates when the ligand dosage of **18b** to Pd is doubled. There is little effect on the ortho-substituted electron-neutral substrates, which support the possibility that different types of substrates would favor different mechanistic pathways. It is likely that both pathways are operative whereby the dominance of either is influenced by the choice of ligands, substrates, and steric factors.²⁸






A promising future of hemilabile ligand research is to design intelligent ligands that have the dynamic qualities to cater to different types of substrates by promoting and supporting different catalytic pathways as a means of

achieving maximum yields. This is possible only if the hemilabile ligands can stabilize and activate different forms of reaction intermediates. A related prospect is to take advantage of the switchable coordination modes of the hybrid ligand to trigger a sequential activation and stabilization mechanism. This would provide a lead in designing stable catalysts without compromising their activities. Another approach can be demonstrated by Hartwig's recent work with a new class of structurally unique three-coordinated arylpalladium halide complexes with a hindered phosphine ligand (Scheme 10).³⁹ Instead of using an unsaturated Pd(0) to promote the oxidative addition, one could promote the subsequent transmetalation and reductive elimination through the generation of an unsaturated oxidative addition complex intermediate. It remains to be seen whether coordinatively unsaturated Pd(II) or Pd(I) intermediate could be stabilized by hemilabile ligands.

4. Suzuki Coupling by a Pd(I) Catalyst

In most Pd-mediated cross-coupling reactions, Pd(II) or Pd(0) sources such as Pd(OAc)₂ or Pd₂(dba)₃ with phosphine ligands are used by default. Recently, a rare use of Pd(I) has been reported (Chart 6).⁴⁰ Hartwig et al. and Prashad et al. have reported high activity of a Pd(I) dimer, [Pd₂(μ-Br)₂(P-*t*-Bu₃)₂] (**47**), in cross-coupling reactions.^{40a,b} Vilar et al. used the new Pd(I) dimer **48** as a precatalyst for amination of aryl chloride.^{40c} Barder used the isolated phosphine arene-ligated Pd(I) dimer **49** as a precatalyst to promote Suzuki reaction.^{40d} Milstein et al. reported that partial reduction of

Scheme 6


Scheme 7

Scheme 8

Chart 6. Pd(I) Dimer for Cross-Coupling Reactions

Scheme 9

Scheme 10


$[\text{Pd}(2\text{-methylallyl})\text{Cl}]_2$ in the presence of the ligand **6** gives a Pd(I) dimer **50**, which promotes Suzuki coupling by providing an active 12-electron Pd(0).²² Our preliminary results⁴¹ showed that the known “solvent-saturated” Pd(I)–Pd(I) complex $[\text{Pd}_2(\text{CH}_3\text{CN})_6][\text{SbF}_6]_2$ (**51**) is active in promot-

ing Suzuki cross-coupling reactions of aryl bromides with various aryl boronic acids at room temperature in a $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ mixture giving good to excellent yields. The use of Cs_2CO_3 as a base with the $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ solvent mixture gives the best yields. The anion also has an effect on the catalyst activity with the less basic and much bigger SbF_6^- performing better than BF_4^- . X-ray single-crystal crystallographic analysis revealed a “naked” Pd–Pd bonded dimer stabilized only by the solvent ligand CH_3CN . This appears to be the ideal source for unsaturated Pd(I) and other exquisite forms of Pd materials.⁴² These results have posed a range of unanswered questions. (1) Is it prerequisite for Pd(I) to reduce to Pd(0) before it can enter the catalytic cycle? (2) Could a radical pathway be proposed for Suzuki or other related cross-coupling reactions? (3) Can Suzuki coupling proceed through a Pd(I)/Pd(III) instead of Pd(0)/Pd(II) cycle? (4) Instead of oxidative addition and then reductive elimination at the metal center, can we benefit from oxidative addition across a Pd–Pd bond and reductive elimination to regenerate the Pd–Pd bond? Such knowledge could help in the design of the next generation of active catalysts.

5. Summary

Ferrocene-based hemilabile ligands functionalized with P,N and P,O donors are highly efficient and effective

in promoting Suzuki couplings. Their remarkable ability to stabilize low-valent and unsaturated metals has enabled us to isolate and study the structural features of species that are mechanistically significant. These ligands are electronically, sterically, and stereogeometrically tunable. Such tuning provides a fine balance between (thermodynamic) stability and (catalytic) activity, thereby allowing the syntheses of complexes that model catalytic intermediates. As we begin to understand better the synergic and complementary effect of hemilabile ligand and unsaturated metal, we can expect a larger variety of ligand designs, an example of which is given in our recent use of N,S-heterocyclic carbenes with functionalized side arms.⁴³ The use of imidazolylphosphines with supporting H-bonding is another example.^{29e} The benefits of hemilability in other catalytic systems are also beginning to emerge.⁴⁴

We thank the National University of Singapore and the Institute of Chemical and Engineering Sciences of Singapore for support and our co-workers, whose names appear in the references, for their intellectual and experimental contributions.

References

- (1) See, for instance: (a) de Meijere, A., Diederich, F., Eds. *Metal-Catalyzed Cross-Coupling Reactions*, Vol. 1&2, 2nd ed.; Wiley-VCH: Weinheim, Germany, 2004. (b) Beller, M.; Zapf, A. In *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E.-I., Ed.; Wiley: New York, 2002; Vol. 1, pp 1209–1222.
- (2) Hassan, J.; Sévignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. Aryl-Aryl Bond Formation One Century after the Discovery of the Ullmann Reaction. *Chem. Rev.* **2002**, *102*, 1359–1470.
- (3) Suzuki, A. Recent Advances in the Cross-Coupling Reactions of Organoboron Derivatives with Organic Electrophiles, 1995–1998. *J. Organomet. Chem.* **1999**, *576*, 147–168 and references therein.
- (4) Rouhi, A. M. Fine chemicals. *Chem. Eng. News* **2004**, *82*, 49–58.
- (5) Braunstein, P.; Nand, F. Hemilability of Hybrid Ligands and the Coordination Chemistry of Oxazoline-Based Systems. *Angew. Chem., Int. Ed.* **2001**, *40*, 680–699.
- (6) Alvarez, S. Bonding and Stereochemistry of Three-Coordinated Transition Metal Compounds. *Coord. Chem. Rev.* **1999**, *193–195*, 13–41.
- (7) Gan, K.-S.; Hor, T. S. A. *Ferrocenes: Homogeneous Catalysis, Organic Synthesis, Materials Science*; Togni, A., Hayashi, T., Eds.; VCH: Weinheim, Germany, 1995; Chapter 1, p 3.
- (8) See for example: Gstöttmayr, C. W. K.; Böhm, V. P. W.; Herdtweck, E.; Grosche, M.; Herrmann, W. A. A Defined N-Heterocyclic Carbene Complex for the Palladium-Catalyzed Suzuki Cross-Coupling of Aryl Chlorides at Ambient Temperatures. *Angew. Chem., Int. Ed.* **2002**, *41*, 1363–1365.
- (9) See for example: Grasa, G. A.; Viciu, M. S.; Huang, J.; Zhang, C.; Trudell, M. L.; Nolan, S. P. Suzuki-Miyaura Cross-Coupling Reactions Mediated by Palladium/Imidazolium Salt Systems. *Organometallics* **2002**, *21*, 2866–2873.
- (10) See for example: (a) Lysén, M.; Köhler, K. Palladium on Activated Carbon: A Recyclable Catalyst for Suzuki-Miyaura Cross-Coupling of Aryl Chlorides in Water. *Synthesis* **2006**, 692–698. (b) Cassez, A.; Ponchel, A.; Hapiot, F.; Monflier, E. Unexpected Multifunctional Effects of Methylated Cyclodextrins in a Palladium Charcoal-Catalyzed Suzuki-Miyaura Reaction. *Org. Lett.* **2006**, *8*, 4823–4826.
- (11) See for example: (a) Leadbeater, N. E.; Marco, M. Ligand-Free Palladium Catalysis of the Suzuki Reaction in Water Using Microwave Heating. *Org. Lett.* **2002**, *4*, 2973–2976. (b) Arvela, R. K.; Leadbeater, N. E.; Sangi, M. S.; Williams, V. A.; Granados, P.; Singer, R. D. A Reassessment of the Transition-Metal Free Suzuki-Type Coupling Methodology. *J. Org. Chem.* **2005**, *70*, 161–168. (c) Thathagar, M. B.; Beckers, J.; Rothenberg, G. Copper-Catalyzed Suzuki Cross-Coupling Using Mixed Nanocluster Catalysts. *J. Am. Chem. Soc.* **2002**, *124*, 11858–11859.
- (12) See for example: Zhang, J.; Zhao, L.; Song, M.; Mak, T. C. W.; Wu, Y. Highly Efficient Cyclopalladated Ferrocenylimine Catalyst for Suzuki Cross-Coupling Reaction of 3-Pyridylboronic Pinacol Ester with Aryl halides. *J. Organomet. Chem.* **2006**, *691*, 1301–1306.
- (13) See for example: (a) Alonso, D. A.; Nájera, C.; Pacheco, M. C. Highly Active Oxime-Derived Palladacycle Complexes for Suzuki-Miyaura and Ullmann-Type Coupling Reactions. *J. Org. Chem.* **2002**, *67*, 5588–5594. (b) Lee, S. 1,1'-Bis(oxazolonyl)ferrocene-Based Palladium Catalysts: Synthesis, X-ray Structures and Applications in Suzuki and Heck Coupling Reactions. *J. Organomet. Chem.* **2006**, *691*, 1347–1355.
- (14) (a) Grasa, G. A.; Hillier, A. C.; Nolan, S. P. Convenient and Efficient Suzuki-Miyaura Cross-Coupling Catalyzed by a Palladium/Diazabutadiene System. *Org. Lett.* **2001**, *3*, 1077–1080. (b) Li, J.-H.; Liu, W.-J. Dabco as an Inexpensive and Highly Efficient Ligand for Palladium-Catalyzed Suzuki-Miyaura Cross-Coupling Reaction. *Org. Lett.* **2004**, *6*, 2809–2811.
- (15) Mazet, C.; Gade, L. H. A Bis(oxazolonyl)pyrrole as a New Monoanionic Tridentate Supporting Ligand: Synthesis of a Highly Active Palladium Catalyst for Suzuki-Type C-C Coupling. *Organometallics* **2001**, *20*, 4144–4146.
- (16) Mohanty, S.; Punji, B.; Balakrishna, M. S. Synthesis and Reaction Kinetics of Pd(1,5-Cyclooctadiene)Cl₂ with N,N'-Methylene-bis(2-Aminopyridyl): An Efficient Catalyst for Suzuki-Cross-Coupling Reactions. *Polyhedron* **2006**, *25*, 815–820.
- (17) Weng, Z.; Koh, L. L.; Hor, T. S. A. Suzuki Cross-Coupling in Aqueous Media Catalyzed by a 1,1'-N-Substituted Ferrocenediyl Pd(II) Complex. *J. Organomet. Chem.* **2004**, *689*, 18–24.
- (18) Weng, Z.; Teo, S.; Koh, L. L.; Hor, T. S. A. Crystallographic Characterization of a Pd(II) Metallamacrocycle Supported by an Amino-Functionalized Ferrocene and Its Use as an Efficient Suzuki-Coupling Catalyst. *Organometallics* **2004**, *23*, 3603–3609.
- (19) See, for instance: (a) Litke, A. F.; Dai, C.; Fu, G. C. Versatile Catalysts for the Suzuki Cross-Coupling of Arylboronic Acids with Aryl and Vinyl Halides and Triflates under Mild Conditions. *J. Am. Chem. Soc.* **2000**, *122*, 4020–4028. (b) Kataoka, N.; Shelby, Q.; Stambuli, J. P.; Hartwig, J. F. Air Stable, Sterically Hindered Ferrocenyl Dialkylphosphines for Palladium-Catalyzed C-C, C-N, and C-O Bond-Forming Cross-Couplings. *J. Org. Chem.* **2002**, *67*, 5553–5566. (c) Dai, Q.; Gao, W.; Liu, D.; Kapes, L. M.; Zhang, X. Triazole-Based Monophosphine Ligands for Palladium-Catalyzed Cross-Coupling Reactions of Aryl Chlorides. *J. Org. Chem.* **2006**, *71*, 3928–3934. (d) Bedford, R. B.; Cazin, C. S. J.; Hazelwood, S. L. Simple Mixed Tricyclohexylphosphane-Triarylphosphite Complexes as Extremely High-Activity Catalysts for the Suzuki Coupling of Aryl Chlorides. *Angew. Chem., Int. Ed.* **2002**, *41*, 4120–4122.
- (20) (a) Herrmann, W. A. N-Heterocyclic Carbenes: A New Concept in Organometallic Catalysis. *Angew. Chem., Int. Ed.* **2002**, *41*, 1290–1309. (b) Viciu, M. S.; Nolan, S. P. The Use of N-Heterocyclic Carbenes as Ligands in Palladium-Mediated Catalysis. *Top. Organomet. Chem.* **2005**, *14*, 241–278.
- (21) Barder, T. E.; Walker, S. D.; Martinelli, J. R.; Buchwald, S. L. Catalysts for Suzuki-Miyaura Coupling Processes: Scope and Studies of the Effect of Ligand Structure. *J. Am. Chem. Soc.* **2005**, *127*, 4685–4696.
- (22) Weissman, H.; Shimon, L. J. W.; Milstein, D. Unsaturated Pd(0), Pd(I), and Pd(II) Complexes of a New Methoxy-Substituted Benzyl Phosphine. Aryl-X (X = Cl, I) Oxidative Addition, C-O Cleavage, and Suzuki-Miyaura Coupling of Aryl Chlorides. *Organometallics* **2004**, *23*, 3931–3940.
- (23) (a) Kwong, F. Y.; Lam, W. H.; Yeung, C. H.; Chan, K. S.; Chan, A. S. C. A Simple and Highly Efficient P,O-type Ligand for Suzuki-Miyaura Cross-Coupling of Aryl Halides. *Chem. Commun.* **2004**, 1922–1923. (b) Dai, W.-M.; Zhang, Y. A Family of Simple Amide-Derived Air-Stable P,O-ligands for Suzuki Cross-Coupling of Unactivated Aryl Chlorides. *Tetrahedron Lett.* **2005**, *46*, 1377–1381. (c) Bei, X.; Turner, H. W.; Weinberg, W. H.; Guram, A. S.; Petersen, J. L. Palladium/P,O-Ligand-Catalyzed Suzuki Cross-Coupling Reactions of Arylboronic Acids and Aryl Chlorides. Isolation and Structural Characterization of (P,O)-Pd(dba) Complex. *J. Org. Chem.* **1999**, *64*, 6797–6803. (d) Jensen, J. F.; Johannsen, M. New Air-Stable Planar Chiral Ferrocenyl Monophosphine Ligands: Suzuki Cross-Coupling of Aryl Chlorides and Bromides. *Org. Lett.* **2003**, *5*, 3025–3028. (e) Atkinson, R. C. J.; Gibson, V. C.; Long, N. J.; White, A. J. P.; Williams, D. J. Synthesis, Coordination Chemistry, and Catalytic Application of a Novel Unsymmetrical P/O Ferrocenediyl Ligand. *Organometallics* **2004**, *23*, 2744–2751.
- (24) (a) Old, D. W.; Wolfe, J. P.; Buchwald, S. L. A Highly Active Catalyst for Palladium-Catalyzed Cross-Coupling Reactions: Room-Temperature Suzuki Couplings and Amination of Unactivated Aryl Chlorides. *J. Am. Chem. Soc.* **1998**, *120*, 9722–9723. (b) Kočovský, P.; Vyskočil, Š.; Čisáková, I.; Sejbál, J.; Smrčina, M.; Lloyd-Jones, G. C.; Stephen, S. C.; Butts, C. P.; Murray, M.; Langer, V. Palladium(II) Complexes of 2-Dimethylamino-2'-diphenylphosphino-1,1'-

- binaphthyl (MAP) with Unique P,C_o-Coordination and Their Catalytic Activity in Allylic Substitution, Hartwig-Buchwald Amination, and Suzuki Coupling. *J. Am. Chem. Soc.* **1999**, *121*, 7714–7715. (c) Zapf, A.; Jackstell, R.; Rataboul, F.; Riermeier, T.; Monsees, A.; Fuhrmann, C.; Shaikh, N.; Dingerdissen, U.; Beller, M. Practical Synthesis of New and Highly Efficient Ligands for the Suzuki Reaction of Aryl Chlorides. *Chem. Commun.* **2004**, 38–39. (d) Liang, L.-C.; Chien, P.-S.; Huang, M.-H. Catalytic Suzuki Coupling Reactions by Amido Phosphine Complexes of Palladium. *Organometallics* **2005**, *24*, 353–357.
- (25) (a) Zhang, W.; Shi, M. A Novel P,S-Heterodonor Ligand and Palladium(0) Complex Catalyzed Suzuki Cross-Coupling Reaction. *Tetrahedron Lett.* **2004**, *45*, 8921–8924. (b) Gibson, V. C.; Long, N. J.; White, A. J. P.; Williams, C. K.; Williams, D. J.; Fontani, M.; Zanello, P. Synthesis, Characterisation and Catalytic Activity of Metal Complexes of Neutral, Unsymmetrical P/S Ferrocenediyl Ligands. *J. Chem. Soc., Dalton Trans.* **2002**, 3280–3289. (c) Scrivanti, A.; Beghetto, V.; Matteoli, U.; Antonaroli, S.; Marini, A.; Crociani, B. Catalytic Activity of η^2 -(Olefin)palladium(0) Complexes with Iminophosphine Ligands in the Suzuki-Miyaura Reaction. Role of the Olefin in the Catalyst Stabilization. *Tetrahedron* **2005**, *61*, 9752–9758. (d) Guo, M.; Jian, F.; He, R. The Air-Stable and Highly Efficient P,N-Chelated Palladium(II) Complexes as Catalysts for the Suzuki Cross-Coupling Reaction at Room Temperature. *Tetrahedron Lett.* **2006**, *47*, 2033–2036. (e) Piechaczyk, O.; Doux, M.; Ricard, L.; Le Floch, P. Synthesis of 1-Phosphabarrelene Phosphine Sulfide Substituted Palladium(II) Complexes: Application in the Catalyzed Suzuki Cross-Coupling Process and in the Allylation of Secondary Amines. *Organometallics* **2005**, *24*, 1204–1213. (f) Punji, B.; Mague, J. T.; Balakrishna, M. S. Synthesis of Neutral (Pd^{II}, Pt^{II}), Cationic (Pd^{II}), and Water-Induced Anionic (Pd^{II}) Complexes Containing New Mesocyclic Thioether-Aminophosphonite Ligands and Their Application in the Suzuki Cross-Coupling Reaction. *Inorg. Chem.* **2006**, *45*, 9454–9464.
- (26) Weng, Z.; Teo, S.; Koh, L. L.; Hor, T. S. A. Efficient Suzuki-Coupling of Aryl Chlorides Catalyzed by Palladium(0) with P,N-heteroligand and Isolation of Unsaturation Intermediates. *Organometallics* **2004**, *23*, 4342–4345.
- (27) (a) Walker, S. D.; Barder, T. E.; Martinelli, J. R.; Buchwald, S. L. A Rationally Designed Universal Catalyst for Suzuki-Miyaura Coupling Processes. *Angew. Chem., Int. Ed.* **2004**, *43*, 1871–1876. (b) Bedford, R. B.; Hazelwood (née Welch), S. L.; Limmert, M. E.; Albisson, D. A.; Draper, S. M.; Scully, P. N.; Coles, S. J.; Hursthouse, M. B. Orthopalladated and -platinated Bulky Triarylphosphite Complexes: Synthesis, Reactivity and Application as High-Activity Catalysts for Suzuki and Stille Coupling Reactions. *Chem.—Eur. J.* **2003**, *9*, 3216–3227.
- (28) Teo, S.; Weng, Z.; Hor, T. S. A. 1,1'-P/O Ferrocenyl Ligands in Pd-Catalyzed Suzuki Coupling of Aryl Chlorides. *Organometallics* **2006**, *25*, 1199–1205.
- (29) (a) Otsuka, S.; Yoshida, T.; Matsumoto, M.; Nakatsu, K. Bis(tertiary phosphine)palladium(0) and -Platinum(0) Complexes: Preparations and Crystal and Molecular structures. *J. Am. Chem. Soc.* **1976**, *98*, 5850–5858. (b) Paul, F.; Patt, J.; Hartwig, J. F. Structural Characterization and Simple Synthesis of {Pd[P(o-Tol)₃]₂}. Spectroscopic Study and Structural Characterization of the Dimeric Palladium(II) Complexes Obtained by Oxidative Addition of Aryl Bromides and Their Reactivity with Amines. *Organometallics* **1995**, *14*, 3030–3039. (c) Mann, G.; Incarvito, C.; Rheingold, A. L.; Hartwig, J. F. Palladium-Catalyzed C-O Coupling Involving Unactivated Aryl Halides. Sterically Induced Reductive Elimination To Form the C-O Bond in Diaryl Ethers. *J. Am. Chem. Soc.* **1999**, *121*, 3224–3225. (d) Reid, S. M.; Boyle, R. C.; Mague, J. T.; Fink, M. J. A Dicoordinate Palladium(0) Complex with an Unusual Intramolecular η^1 -Arene Coordination. *J. Am. Chem. Soc.* **2003**, *125*, 7816–7817. (e) Grotjahn, D. B.; Gong, Y.; Zakharov, L.; Golen, J. A.; Rheingold, A. L. Changes in Coordination of Sterically Demanding Hybrid Imidazolylphosphine Ligands on Pd(0) and Pd(II). *J. Am. Chem. Soc.* **2006**, *128*, 438–453.
- (30) Amatore, C.; Jutand, A. Role of dba in the Reactivity of Palladium(0) Complexes Generated in Situ from Mixtures of Pd(dba)₂ and Phosphines. *Coord. Chem. Rev.* **1998**, *178–180*, 511–528.
- (31) Yin, J.; Rainka, M. P.; Zhang, X.-X.; Buchwald, S. L. A Highly Active Suzuki Catalyst for the Synthesis of Sterically Hindered Biaryls: Novel Ligand Coordination. *J. Am. Chem. Soc.* **2002**, *124*, 1162–1163.
- (32) Christmann, U.; Vilar, R.; White, A. J. P.; Williams, D. J. Synthesis of Two Novel Dinuclear Palladium(I) Complexes and Studies of Their Catalytic Activity in Amination Reactions. *Chem. Commun.* **2004**, 1294–1295.
- (33) Jalón, F. A.; Manzano, B. R.; Torre, F. G.; López-Agenjo, A. M.; Rodríguez, A. M.; Weissensteiner, W.; Sturm, T.; Mahía, J.; Maestro, M. NMR Study on the Coordination of Dibenzylideneacetone to Chiral Palladium(0) Units. Fluxional Behaviour Including an Intramolecular Double Bond Exchange. *J. Chem. Soc., Dalton Trans.* **2001**, 2417–2424.
- (34) See, for instance: Reid, S. M.; Mague, J. T.; Fink, M. J. Synthetic, Structural, and Dynamic NMR Studies of (Bisphosphine)palladium(0) Complexes of Dibenzylideneacetone. *J. Organomet. Chem.* **2000**, *616*, 10–18.
- (35) Farina, V. High-Turnover Palladium Catalysts in Cross-Coupling and Heck Chemistry: A Critical Overview. *Adv. Synth. Catal.* **2004**, *346*, 1553–1582.
- (36) Fong, S.-W. A.; Vittal, J. J.; Hor, T. S. A. Isolation of a Stable Precursor for [Pt⁰(P-P)] and Its Reductive Addition to Ru₃(CO)₉(μ_3 -S)₂ Giving Square Heterometallic Clusters Supported by Two Face-Capping μ_4 -Sulfides [P-P = (C₅H₄PPh₂)₂M; M = Fe, Ru]. *Organometallics* **2000**, *19*, 918–924.
- (37) See, for instance: Braga, A. A. C.; Ujaque, G.; Maseras, F. A DFT Study of the Full Catalytic Cycle of the Suzuki-Miyaura Cross-Coupling on a Model System. *Organometallics* **2006**, *25*, 3662–3673.
- (38) Bei, X.; Uno, T.; Norris, J.; Turner, H. W.; Weinberg, W. H.; Guram, A. S.; Petersen, J. L. Phenyl Backbone-Derived P,O- and P,N-Ligands for Palladium/Ligand-Catalyzed Aminations of Aryl Bromides, Iodides, and Chlorides. Syntheses and Structures of (P,O)_n-Palladium(II)Aryl(Br) Complexes. *Organometallics* **1999**, *18*, 1840–1853.
- (39) Stambuli, J. P.; Incarvito, C. D.; Bühl, M.; Hartwig, J. F. Synthesis, Structure, Theoretical Studies, and Ligand Exchange Reactions of Monomeric, T-Shaped Arylpalladium(II) Halide Complexes with an Additional, Weak Agostic Interaction. *J. Am. Chem. Soc.* **2004**, *126*, 1184–1194.
- (40) (a) Stambuli, J. P.; Kuwano, R.; Hartwig, J. F. Unparalleled Rates for the Activation of Aryl Chlorides and Bromides: Coupling with Amines and Boronic Acids in Minutes at Room Temperature. *Angew. Chem., Int. Ed.* **2002**, *41*, 4746–4748. (b) Prashad, M.; Mak, X. Y.; Liu, Y.; Repi, O. Palladium-Catalyzed Amination of Aryl Bromides with Hindered N-Alkyl-Substituted Anilines Using a Palladium(I) Tri-tert-butylphosphine Bromide Dimer. *J. Org. Chem.* **2003**, *68*, 1163–1164. (c) Christmann, U.; Pantazis, D. A.; Benet-Buchholz, J.; McGrady, J. E.; Maseras, F.; Vilar, R. Experimental and Theoretical Investigations of New Dinuclear Palladium Complexes as Precatalysts for the Amination of Aryl Chlorides. *J. Am. Chem. Soc.* **2006**, *128*, 6376–6390. (d) Barder, T. E. Synthesis, Structural, and Electron Topographical Analyses of a Dialkylbiaryl Phosphine/Arene-Ligated Palladium(II) Dimer: Enhanced Reactivity in Suzuki-Miyaura Coupling Reactions. *J. Am. Chem. Soc.* **2006**, *128*, 898–904.
- (41) Han, X.; Weng, Z.; Hor, T. S. A. *Suzuki Coupling Catalyzed by a Cationic Pd(II) Dimer*; 37th International Conference on Coordination Chemistry, Aug 13–18, 2006, Cape Town, South Africa; Abstract 427.
- (42) Tatsumi, Y.; Shirato, K.; Murahashi, T.; Ogoshi, S.; Kurosawa, H. Sandwich Complexes Containing Bent Palladium Chains. *Angew. Chem., Int. Ed.* **2006**, *45*, 5799–5803.
- (43) (a) Yen, S. K.; Koh, L. L.; Hahn, F. E.; Huynh, H. V.; Hor, T. S. A. Convenient Entry to Mono- and Dinuclear Palladium(II) Benzothiazolin-2-ylidene Complexes and Their Activities toward Heck Coupling. *Organometallics* **2006**, *25*, 5105–5112. (b) Yen, S. K.; Cavelli, K. J.; Huynh, H. V.; Hor, T. S. A. *Preparation, Characterisation and Redox Activities of (Hyrido)metal(carbene) Complexes*; 6th International Symposium for Chinese Inorganic Chemists, Dec 17–20, 2006, Singapore; Poster Abstract 302.
- (44) (a) Huynh, H. V.; Yeo, C. H.; Tan, G. K. Hemilabile Behavior of a Thioether-Functionalized N-Heterocyclic Carbene Ligand. *Chem. Commun.* **2006**, 3833–3835. (b) Chisholm, M. H.; Gallucci, J. C.; Yaman, G. A tris-Pyrazolylborate Ligand with Hemilabile O-Donor Groups. Examples of η^3 , η^5 , η^6 and Bridging Modes of Bonding to Li⁺, Na⁺, K⁺, Tl⁺ and Ca²⁺ Ions. *Chem. Commun.* **2006**, 1872–1874.

AR600003H