

Enantiomerically Enriched Allylic Alcohols and Allylic Amines via C–C Bond-Forming Hydrogenation: Asymmetric Carbonyl and Imine Vinylation

EDUARDAS SKUCAS, MING-YU NGAI, VENUKRISHNAN KOMANDURI, AND MICHAEL J. KRISCHE*

Department of Chemistry and Biochemistry, University of Texas at Austin, 1 University Station, Welch Hall (A5300), Austin, Texas 78712-1167

Received May 5, 2007

ABSTRACT

Hydrogenation of alkynes in the presence of carbonyl compounds and imines using cationic rhodium(I) and iridium(I) precatalysts enables the formation of allylic alcohols and allylic amines, respectively. Through the use of hydrogenation catalysts modified by chiral ligands, allylic alcohols and allylic amines may be generated in highly optically enriched forms. Hydrogenative fragment couplings of this type circumvent the use of preformed organometallic reagents and avoid the generation of stoichiometric byproducts.

Introduction to Carbonyl and Imine Vinylation

The synthetic utility of allylic alcohols and allylic amines has driven efforts toward the development of catalytic enantioselective protocols for the vinylation of carbonyl compounds and imines. The majority of work in this area stems from the seminal studies of Oguni^{1a} and Noyori^{1b} on the enantioselective addition of dialkylzinc reagents to aldehydes.² The first asymmetric aldehyde vinylations of this type were reported by Oppolzer^{3a,4} and involve the generation of vinylzinc reagents via alkyne hydroboration, followed by transmetalation of the resulting vinyl boron

Eduardas Skucas obtained a B.S. degree in chemistry from Temple University, Philadelphia, PA, where he conducted undergraduate research under the tutelage of Professor Franklin Davis. In Fall 2005, he initiated graduate studies at the University of Texas at Austin under the supervision of Professor Michael J. Krische, where he is involved in the discovery and development of new C–C bond-forming hydrogenations.

Ming-Yu Ngai obtained a B.S. degree in chemistry at the University of Hong Kong (HKU), where he conducted undergraduate research under the tutelage of Professor Chi-Ming Che. During his undergraduate study, Ming-Yu Ngai spent 1 year abroad at the University of California, San Diego, under the supervision of Professor Michael VanNieuwenhze. In Fall 2004, he initiated graduate studies at the University of Texas at Austin under the supervision of Professor Michael J. Krische, where he is involved in the discovery and development of new C–C bond-forming hydrogenations.

Venukrishnan Komanduri obtained a B.S. degree in chemistry from Andhra Loyola College, Vijayawada, India, and an M.S. degree from the Indian Institute of Technology (IIT), Chennai, India. In Fall 2002, he initiated graduate studies at Wayne State University, under the tutelage of Professor Maarten H. D. Postema. In Fall 2004, he continued graduate studies at the University of Texas at Austin under the supervision of Professor Michael J. Krische, where he is involved in the discovery of new C–C bond-forming hydrogenations.

reagent to zinc using $ZnMe_2$. A related strategy for asymmetric aldehyde vinylation is reported by Wipf and involves alkyne hydrozirconation–transmetalation en route to vinylzinc reagents.^{3d} Finally, after catalytic enantioselective additions of alkylzinc and arylzinc reagents to ketones^{2b–d} described by Yus^{5a,b} and Fu,^{5c} respectively, catalytic asymmetric ketone vinylations were devised by Walsh.⁶

Parallel efforts toward the development of enantioselective imine additions using preformed metallic reagents reveal an additional set of challenges.⁷ Pursuant to Soai's seminal report,^{8a} several organocatalysts for the enantioselective addition of organozinc reagents to imines emerged.⁸ Because conventional imines are generally less reactive than aldehydes with respect to organocatalyzed organozinc addition, these studies employ *N*-acyl and *N*-(diphenylphosphinoyl) imines. To address the issue of reactivity, enantioselective metal-catalyzed organozinc additions to imines were developed,⁹ as first described by Tomioka in the case of copper.^{9a} Early transition-metal catalysts (Ti, Zr, and Hf)¹⁰ and late transition-metal catalysts (Rh)¹¹ also have been found to promote highly enantioselective organozinc additions to imines.


Beyond organozinc reagents, the enantioselective addition of organolithium reagents to imines catalyzed by chiral Lewis basic chelating agents has been described.¹² Additionally, under the conditions of rhodium catalysis, organotin,^{13a,b} organotitanium,^{13c} and organoboron^{13d–h} reagents have served in catalytic asymmetric imine additions. Despite considerable effort, highly enantioselective vinyl transfer to imines employing preformed organometallic reagents has not been achieved.^{14,15}

The aforementioned approaches to enantioselective carbonyl and imine vinylation are highly effective in terms of directing enantioselection. However, these strategies uniformly exploit preformed organometallic reagents, which are generally prepared via transmetalation from a “primary organometallic reagent”, which itself may require preparation through metalation of a precursor. For example, organoboron reagents are generally prepared from the corresponding organolithium reagents, which in turn

* To whom correspondence should be addressed. E-mail: mkrische@mail.utexas.edu.

Michael Krische obtained a B.S. degree in chemistry from the University of California at Berkeley, where he performed research under the tutelage of Professor Henry Rapoport. After 1 year of study abroad as a Fulbright Fellow, he initiated graduate study at Stanford University under the mentorship of Professor Barry Trost as a Veatch Graduate Fellow. Following his Ph.D. degree, he engaged in postdoctoral studies with Jean-Marie Lehn at the Université Louis Pasteur as an NIH Postdoctoral Fellow. In Fall 1999, Professor Krische was appointed as Assistant Professor of Chemistry at the University of Texas at Austin. He was promoted directly to Full Professor in Fall 2004. His research on hydrogenative C–C coupling has received recognition in the form of numerous awards: Elias J. Corey Award (2007), Dowpharma Prize (2007), Presidential Green Chemistry Award (2007), Solvias Ligand Prize (2006), Society of Synthetic Chemistry Japan Lectureship (2005), Johnson & Johnson Focused Giving Award (2005), Dreyfus Teacher Scholar Award (2003), Alfred P. Sloan Research Fellowship (2003), Cottrell Scholar Award (2002), Frasc Foundation Award in Chemistry (2002), Lilly Grantee Award (2002), and National Science Foundation–CAREER Award (2000).

are prepared by metal halogen exchange using butyl lithium. Here, three preformed organometallic reagents are used stoichiometrically in advance of C–C coupling. Such protocols mandate the generation of multiple stoichiometric byproducts and require the handling of multiple air- and moisture-sensitive materials (Scheme 1).




Direct metal-catalyzed reductive coupling circumvents the successive use of stoichiometrically preformed organometallic reagents.¹⁶ Of direct relevance to this Account are catalysts enabling direct alkyne–aldehyde reductive coupling.¹⁷ The first catalytic process of this type was reported by Ojima and involves the rhodium-catalyzed reductive cyclization of acetylenic aldehydes mediated by silane.¹⁸ Corresponding titanocene-catalyzed cyclizations were disclosed by Crowe,¹⁹ and nickel-catalyzed cyclizations were reported by Montgomery.^{20a–c,e} Finally, intermolecular reductive alkyne–aldehyde coupling was achieved using nickel-based catalysts, as described by Jamison,²¹ Takai,²² and Montgomery.^{20d} Reductive couplings of this type involve the capture of organometallics that arise transiently under catalytic conditions, signaling a departure from the stoichiometric preformation of organometallic reagents. However, the aforementioned methods employ terminal reductants, such as hydrosilanes, hydrostannanes, organozinc reagents, organoboron reagents, or chromium(II) chloride, which ultimately produce molar equivalents of chemical byproducts.

Completely atom economical and, hence, byproduct-free reductive C–C couplings are potentially achieved under the conditions of catalytic hydrogenation. Despite the fact that catalytic hydrogenation of organic compounds has been known for over a century,²³ use of elemental hydrogen as a terminal reductant in metal-catalyzed C–C coupling has only been explored in connection with processes involving carbon monoxide insertion, as exemplified by the Fischer–Tropsch²⁴ reaction and alkene hydroformylation.²⁵ Despite the impact of these prototypical hydrogen-mediated C–C bond formations, systematic efforts toward hydrogenative C–C couplings

that apply to conventional electrophilic partners in the form of carbonyl compounds and imines were absent from the literature prior to our work.^{26–29}

In this Account, we present a summary of our work on the hydrogenative coupling of alkynes to carbonyl compounds and imines to furnish allylic alcohols and allylic amines. Related hydrogenative couplings of alkenes to carbonyl compounds and imines, as exemplified by the hydrogen-mediated reductive aldol and Mannich reaction, and the hydrogen-mediated coupling of olefins to anhydrides are the subject of recent reviews.^{26,30} The collective hydrogenative fragment couplings that we report constitute a broad new class of catalytic C–C bond formations that circumvent byproduct generation and the requirement of stoichiometrically preformed organometallic reagents in certain C=X (X = O, NR) addition processes.




Hydrogenative Coupling of Alkynes to Carbonyl Compounds and Imines


Proof of concept studies toward the development of first-generation catalysts for hydrogenative C–C coupling, along with preliminary mechanistic studies corroborating their proposed mode of action, are described in prior accounts.²⁶ Salient features are summarized below. The collective data are consistent with a general mechanism involving alkyne–C=X (X = O, NR) oxidative coupling followed by direct or Brønsted acid co-catalyzed hydrogenolysis of the resulting metallacyclic intermediate. To mitigate competitive conventional substrate hydrogenation, cationic complexes of rhodium and iridium are required. A plausible explanation is as follows. Unlike corresponding neutral complexes, hydrogen activation is generally slower for cationic rhodium precatalysts,^{31,32} thus providing a greater kinetic window for entry into alkyne–C=X (X = O, NR) oxidative coupling manifolds. Oxidative coupling pathways are promoted further by the increased coordinative unsaturation of cationic complexes. As supported by theoretical studies,³³ Brønsted acid co-catalysts are postulated to circumvent highly energetic four-centered transition structures “A” for σ -bond metathesis, as required for direct hydrogenolysis of metallacyclic intermediates, with six-centered transition structures “C” for hydrogenolysis of rhodium carboxylates derived upon metallacycle protonolysis, which itself may occur through six-centered transition structure “B”. The collective data suggest that the oxarhodacyclopentene is the catalyst resting state and that hydrogenolysis of the oxarhodacyclopentene is the turnover limiting step in the catalytic mechanism (Scheme 1).

When conjugated alkynes are simply hydrogenated in the presence of vicinal dicarbonyl compounds, glyoxalates, and pyruvates, at ambient pressure using chirally modified cationic rhodium catalysts, reductive C–C coupling occurs to furnish α -hydroxy esters in highly optically enriched form.^{34a,b} Under nearly identical conditions, conjugated

Scheme 1. General Catalytic Mechanism for Hydrogenative Alkyne–Carbonyl Coupling




Scheme 2. Rhodium-Catalyzed Hydrogenative C–C Coupling of Alkynes to Carbonyl Compounds and Imines




alkynes hydrogenatively couple to heterocyclic aromatic aldehydes and ketones that are isoelectronic with respect to the vicinal dicarbonyl motif, thereby providing access to highly optically enriched heteroaryl-substituted secondary and tertiary alcohols.^{34c} Hydrogenation of 1,3-enynes in the presence of optically enriched ethyl (*N*-

sulfonyl)iminoacetates delivers novel nonproteogenic amino acid esters.^{34d} Finally, catalytic hydrogenation of acetylenic aldehydes using chirally modified rhodium catalysts provides products of reductive carbocyclization, again with excellent levels of asymmetric induction.^{34e} Brønsted acid co-catalysts were found to enhance the rate and con-

Scheme 3. Enantioselective (*Z*)-Butadienylation via Hydrogenative Coupling of Acetylene to Aldehydes and Imines

Scheme 4. Enantioselective Iridium-Catalyzed Hydrogenative Coupling of 2-Butyne to Aldehydes and Imines



version in all couplings examined. Couplings to ethyl (*N*-sulfinyl)iminoacetates were studied prior to the discovery of the Brønsted acid co-catalyst effect (Scheme 2).

Intermolecular hydrogenative couplings of alkynes to carbonyl compounds under the conditions of rhodium catalysis require highly activated electrophilic partners, such as vicinal dicarbonyl compounds and their structural relatives. It was found that gaseous acetylene couples to conventional aldehydes under hydrogenation conditions to furnish products of *Z*-butadienylation.^{35a} Isotopic labeling and electrospray ionization mass spectrometry (ESI-MS) analysis are consistent with a catalytic mechanism involving oxidative dimerization of acetylene to form a cationic rhodacyclopentadiene³⁶ followed by carbonyl insertion and Brønsted-acid-assisted hydrogenolysis of the

resulting oxarhodacycloheptadiene to provide the carbonyl addition product and cationic rhodium(I) to close the cycle. More recently, corresponding couplings to aldimines have been achieved.^{35b} For both aldehyde and imine couplings, chirally modified rhodium catalysts deliver highly optically enriched allylic alcohols and allylic amines, respectively (Scheme 3).

Attempted imine vinylation under the conditions of rhodium catalysis using 1,2-dialkylsubstituted alkynes is not an efficient process. Our rationale for addressing this limitation is as follows. In hydrogenative alkyne-carbonyl coupling, our collective studies suggest that a key feature of the catalytic mechanism involves oxidative coupling of the π -unsaturated reactants to furnish oxametallacyclic intermediates. Hydrogenolytic cleavage of this species via

σ -bond metathesis furnishes the coupling product with concomitant regeneration of the catalyst. It is plausible that π -backbonding in the metal–alkyne complex, as described by the Dewar–Chatt–Duncanson model,³⁷ facilitates the oxidative coupling event by conferring nucleophilic character to the bound alkyne. For rhodium, a comparatively weak π donor, conjugated alkynes are required, possibly because of the fact that they embody lower lying unoccupied molecular orbitals (LUMOs). Iridium is a stronger π donor than rhodium because of relativistic effects.^{38,39} These data suggest that iridium complexes may catalyze the hydrogenative coupling of nonconjugated alkynes that embody higher lying LUMOs.

The veracity of this analysis was challenged through experiment. Gratifyingly, it was found that using [Ir(cod)₂]-BARF as a precatalyst, hydrogenation of 2-butyne in the presence of α -benzyloxy acetaldehyde provides an 81% isolated yield of the fully saturated reductive coupling product as an equimolar mixture of diastereomers. Here, the initially formed allylic alcohol is subject to further hydrogenation to deliver the saturated alcohol. In contrast, iridium-catalyzed hydrogenation of 2-butyne in the presence of various *N*-arylsulfonyl imines provides the corresponding trisubstituted allylic amines with complete levels of *E/Z* selectivity ($\geq 95:5$) and, using catalysts modified by (*R*)-Cl₂MeO-BIPHEP, exceptional levels of asymmetric induction.⁴⁰ Notably, over-reduction of the unsaturated allylic amines are not observed. These results are consistent with the role of hydroxyl and sulfonamide moieties as active and inactive volumes, respectively, in iridium-catalyzed hydrogenation.⁴¹ The nonsymmetric alkynes 4-methyl-2-pentyne and 2-hexyne couple to *N*-arylsulfonyl imines with excellent regioselectivity proximal to the more highly substituted alkyne terminus to provide the corresponding allylic amines with uniformly high levels of enantiomeric excess (Scheme 4).

Future Directions

The prototypical C–C bond-forming hydrogenations, the Fischer–Tropsch reaction and alkene hydroformylation, were reported over half a century ago and are practiced on an enormous scale.^{24,25} While these processes are restricted to the use of carbon monoxide as a coupling partner, studies from our laboratory extend hydrogenative C–C coupling to the use of conventional electrophiles in the form of carbonyl compounds and imines. The generation and capture of nonstabilized carbanion equivalents under the essentially neutral conditions of hydrogenation dispenses with the requirement of stoichiometrically preformed organometallic reagents in certain C=X (X = O, NR) addition processes, enabling C–C bond formation in the absence of any stoichiometric byproducts. These studies open a new chapter in the area of catalytic hydrogenation and raise numerous possibilities *vis-à-vis* development of related byproduct-free C–C couplings.

References

- (1) (a) Oguni, N.; Omi, T. Enantioselective Addition of Diethylzinc to Benzaldehyde Catalyzed by a Small Amount of Chiral 2-Amino-1-Alcohols. *Tetrahedron Lett.* **1984**, *25*, 2823–2824. (b) Kitamura, M.; Suga, S.; Kawai, K.; Noyori, R. Catalytic Asymmetric Induction. Highly Enantioselective Addition of Dialkylzincs to Aldehydes. *J. Am. Chem. Soc.* **1986**, *108*, 6071–6072.
- (2) For reviews encompassing catalytic enantioselective addition of organozinc reagents to carbonyl compounds, see (a) Pu, L.; Yu, H.-B. Catalytic Asymmetric Organozinc Additions to Carbonyl Compounds. *Chem. Rev.* **2001**, *101*, 757–824. (b) Betancort, J. M.; Garcia, C.; Walsh, P. J. Development of the First Practical Catalyst for the Asymmetric Addition of Alkyl- and Arylzinc Reagents to Ketones. *Synlett* **2004**, 749–760. (c) Ramón, D. J.; Yus, M. Chiral Tertiary Alcohols Made by Catalytic Enantioselective Addition of Unreactive Zinc Reagents to Poorly Electrophilic Ketones. *Angew. Chem., Int. Ed.* **2004**, *43*, 284–287. (d) Garcia, C.; Martin, V. S. Asymmetric Addition to Ketones: Enantioselective Formation of Tertiary Alcohols. *Curr. Org. Chem.* **2006**, *10*, 1849–1889. (e) Ramón, D. J.; Yus, M. In the Arena of Enantioselective Synthesis, Titanium Complexes Wear the Laurel Wreath. *Chem. Rev.* **2006**, *106*, 2126–2208.
- (3) For enantioselective catalytic addition of vinylzinc reagents to aldehydes, see (a) Oppolzer, W.; Radinov, R. Catalytic Asymmetric Synthesis of Secondary (*E*)-Allyl Alcohols from Acetylenes and Aldehydes via (1-Alkenyl)zinc Intermediates. *Helv. Chim. Acta* **1992**, *75*, 170–173. (b) Oppolzer, W.; Radinov, R. Synthesis of (*R*)-(-)-Muscone by an Asymmetrically Catalyzed Macrocyclization of an ω -Alkynal. *J. Am. Chem. Soc.* **1993**, *115*, 1593–1594. (c) Soai, K.; Takahashi, K. J. Asymmetric Alkenylation of Chiral and Prochiral Aldehydes Catalysed by Chiral or Achiral Amino Alcohols: Catalytic Diastereoselective Synthesis of Protected *erythro*-Sphingosine and Enantioselective Synthesis of Chiral Diallyl Alcohols. *J. Chem. Soc., Perkin Trans. 1* **1994**, 1257–1258. (d) Wipf, P.; Xu, W. Preparation of Allylic Alcohols by Alkene Transfer from Zirconium to Zinc. *Tetrahedron Lett.* **1994**, *35*, 5197–5200. (e) Oppolzer, W.; Radinov, R. N.; De Brabander, J. Total Synthesis of the Macrolide (+)-Aspicilin by an Asymmetrically Catalyzed Macrocyclization of an ω -Alkynal ester. *Tetrahedron Lett.* **1995**, *36*, 2607–2610. (f) Wipf, P.; Ribe, S. Zirconocene–Zinc Transmetalation and in Situ Catalytic Asymmetric Addition to Aldehydes. *J. Org. Chem.* **1998**, *63*, 6454–6455. (g) Oppolzer, W.; Radinov, R. N.; El-Sayed, E. Catalytic Asymmetric Synthesis of Macrocyclic (*E*)-Allylic Alcohols from ω -Alkynals via Intramolecular 1-Alkenylzinc/Aldehyde Additions. *J. Org. Chem.* **2001**, *66*, 4766–4770. (h) Dahmen, S.; Bräse, S. [2,2]Paracyclophane-Based *N,O*-Ligands in Alkenylzinc Additions to Aldehydes. *Org. Lett.* **2001**, *3*, 4119–4122. (i) Chen, Y. K.; Lurain, A. E.; Walsh, P. J. A General, Highly Enantioselective Method for the Synthesis of *D* and *L*- α -Amino Acids and Allylic Amines. *J. Am. Chem. Soc.* **2002**, *124*, 12225–12231. (j) Ji, J.-X.; Qiu, L.-Q.; Yip, C. W.; Chan, A. S. C. A Convenient, One-Step Synthesis of Optically Active Tertiary Aminonaphthol and Its Applications in the Highly Enantioselective Alkenylations of Aldehydes. *J. Org. Chem.* **2003**, *68*, 1589–1590. (k) Lurain, A. E.; Walsh, P. J. A Catalytic Asymmetric Method for the Synthesis of γ -Unsaturated α -Amino Acid Derivatives. *J. Am. Chem. Soc.* **2003**, *125*, 10677–10683. (l) Ko, D.-H.; Kang, S.-W.; Kim, K. H.; Chung, Y.; Ha, D.-C. Asymmetric Alkenylzinc Additions to Aldehydes Catalyzed by a Binaphthyl Based *N,O*-Ligand. *Bull. Korean Chem. Soc.* **2004**, *25*, 35–36. (m) Sprout, C. M.; Richmond, M. L.; Seto, C. T. Solid-Phase Synthesis of Chiral *N*-Acylethylenediamines and Their Use as Ligands for the Asymmetric Addition of Alkylzinc and Alkenylzinc Reagents to Aldehydes. *J. Org. Chem.* **2004**, *69*, 6666–6673. (n) Jeon, S.-J.; Chen, Y. K.; Walsh, P. J. Catalytic Asymmetric Synthesis of Hydroxy Enol Ethers: Approach to a Two-Carbon Homologation of Aldehydes. *Org. Lett.* **2005**, *7*, 1729–1732. (o) Lauterwasser, F.; Gall, J.; Hoefener, S.; Bräse, S. Second-Generation *N,O*-[2,2]Paracyclophane Ketimine Ligands for the Alkenylzinc Addition to Aliphatic and Aromatic Aldehydes: Scope and Limitations. *Adv. Synth. Catal.* **2006**, *348*, 2068–2074. (p) Jeon, S.-J.; Fisher, E. L.; Carroll, P. J.; Walsh, P. J. Direct, Stereospecific Generation of (*Z*)-Disubstituted Allylic Alcohols. *J. Am. Chem. Soc.* **2006**, *128*, 9618–9619.
- (4) For reviews encompassing catalytic enantioselective aldehyde vinylation using organozinc reagents, see (a) Wipf, P.; Kendall, C. Novel Applications of Alkenyl Zirconocenes. *Chem.—Eur. J.* **2002**, *8*, 1778–1784. (b) Wipf, P.; Nunes, R. L. Selective Carbon–Carbon Bond Formations with Alkenylzirconocenes. *Tetrahedron* **2004**, *60*, 1269–1279.
- (5) (a) Ramón, D. J.; Yus, M. First Enantioselective Addition of Dialkylzinc to Ketones Promoted by Titanium(IV) Derivatives. *Tetrahedron Lett.* **1998**, *39*, 1239–1242. (b) Ramón, D. J.; Yus, M. First Enantioselective Addition of Diethylzinc and Dimethylzinc to Prostereogenic Ketones Catalyzed by Camphorsulfonamide-

- Titanium Alkoxide Derivatives. *Tetrahedron* **1998**, *54*, 5651–5666. (c) Dosa, P. I.; Fu, G. C. Catalytic Asymmetric Addition of $ZnPh_2$ to Ketones: Enantioselective Formation of Quaternary Stereocenters. *J. Am. Chem. Soc.* **1998**, *120*, 445–446.
- (6) For catalytic enantioselective ketone vinylation using organozinc reagents, see (a) Li, H.; Walsh, P. J. Catalytic Asymmetric Vinylation of Ketones. *J. Am. Chem. Soc.* **2004**, *126*, 6538–6539. (b) Li, H.; Walsh, P. J. Catalytic Asymmetric Vinylation and Dienylation of Ketones. *J. Am. Chem. Soc.* **2005**, *127*, 8355–8361. (c) Jeon, S.-J.; Li, H.; Garcia, C.; La Rochelle, L. K.; Walsh, P. J. Catalytic Asymmetric Addition of Alkylzinc and Functionalized Alkylzinc Reagents to Ketones. *J. Org. Chem.* **2005**, *70*, 448–455.
- (7) For reviews encompassing the catalytic enantioselective addition of organozinc reagents to imines, see (c) Bloch, R. Additions of Organometallic Reagents to C=N Bonds: Reactivity and Selectivity. *Chem. Rev.* **1998**, *98*, 1407–1438. (d) Kobayashi, S.; Ishitani, H. Catalytic Enantioselective Addition to Imines. *Chem. Rev.* **1999**, *99*, 1069–1094. (e) Vilaivan, T.; Bhanthumnavin, W.; Sritana-Anant, Y. Recent Advances in Catalytic Asymmetric Addition to Imines and Related C=N Systems. *Curr. Org. Chem.* **2005**, *9*, 1315–1392. (f) Schmidt, F.; Stemmler, R. T.; Rudolph, J.; Bolm, C. Catalytic Asymmetric Approaches Toward Enantiomerically Enriched Diarylmethanols and Diarylmethylamines. *Chem. Soc. Rev.* **2006**, *35*, 454–470.
- (8) For enantioselective organocatalytic addition of organozinc reagents to imines, see (a) Soai, K.; Hatanaka, T.; Miyazawa, T. Highly Enantioselective Alkylation of Carbon–Nitrogen Double Bonds. Catalytic and Stoichiometric Asymmetric Synthesis of Optically Active Amines by the Enantioselective Addition of Dialkylzinc Reagents to *N*-Diphenylphosphinoylimines. *J. Chem. Soc., Chem. Commun.* **1992**, 1097–1098. (b) Katritsky, A. R.; Harris, P. A. Enantioselective Ethylation of *N*-(Amidobenzyl)benzotriazoles Catalyzed by Chiral Amino Alcohols. *Tetrahedron Asymmetry* **1992**, *3*, 437–442. (c) Andersson, P. G.; Guijarro, D.; Tanner, D. Simple Aziridino Alcohols as Chiral Ligands Enantioselective Additions of Diethylzinc to *N*-Diphenylphosphinoylimines. *Synlett* **1996**, 727–728. (d) Andersson, P. G.; Guijarro, D.; Tanner, D. Preparation and Use of Aziridino Alcohols as Promoters for the Enantioselective Addition of Dialkylzinc Reagents to *N*-(Diphenylphosphinoyl) Imines. *J. Org. Chem.* **1997**, *62*, 7364–7375. (e) Guijarro, D.; Pinho, P.; Andersson, P. G. Enantioselective Addition of Dialkylzinc Reagents to *N*-(Diphenylphosphinoyl) Imines Promoted by 2-Azanorbonylmethanols. *J. Org. Chem.* **1998**, *63*, 2530–2535. (f) Brandt, P.; Hedberg, C.; Lawonn, K.; Pinho, P.; Andersson, P. G. A Theoretical and Experimental Study of the Asymmetric Addition of Dialkylzinc to *N*-(Diphenylphosphinoyl)benzalimine. *Chem.—Eur. J.* **1999**, *5*, 1692–1699. (g) Jimeno, C.; Reddy, S.; Sola, L.; Moyano, A.; Pericas, M. A.; Riera, A. Highly Enantioselective Addition of Diethylzinc to Diphenylphosphinoyl Imines under Dual Amino Alcohol/Halosilane Mediation. *Org. Lett.* **2000**, *2*, 3157–3159.
- (9) For enantioselective copper catalyzed addition of organozinc reagents to imines, see (a) Fujihara, H.; Nagai, K.; Tomioka, K. Copper–Amidophosphine Catalyst in Asymmetric Addition of Organozinc to Imines. *J. Am. Chem. Soc.* **2000**, *122*, 12055–12056. (b) Nagai, K.; Fujihara, H.; Kuriyama, M.; Yamada, K.-I.; Tomioka, K. Efficient Chiral Amidophosphine Ligand for Copper-Catalyzed Asymmetric Addition of Diethylzinc to *N*-Sulfonylimines. *Chem. Lett.* **2002**, 8–9. (c) Shi, M.; Wang, C.-J. Chiral Binaphthylthiophosphoramidate Cu(II) Catalyzed Asymmetric Addition of Diethylzinc to *N*-Sulfonylimines. *J. Org. Chem.* **2003**, *68*, 6229–6237. (d) Soeta, T.; Nagai, K.; Fujihara, H.; Kuriyama, M.; Tomioka, K. Asymmetric Alkylation of *N*-Toluenesulfonylimines with Dialkylzinc Reagents Catalyzed by Copper–Chiral Amidophosphine. *J. Org. Chem.* **2003**, *68*, 9723–9727. (e) Shi, M.; Zhang, W. Chiral C_1 -Symmetric Diaminothiophosphoramidate–Cu(II) Catalyzed Asymmetric Addition of Diethylzinc to *N*-Sulfonylimines. *Tetrahedron Asymmetry* **2003**, *14*, 3407–3414. (f) Shi, M.; Wang, C.-J. The Catalytic Asymmetric Addition of Diethylzinc to *N*-(Diphenylphosphinoyl) Imines Catalyzed by $Cu(OTf)_2$ -Chiral *N*-(Binaphthyl-2-yl)thiophosphoramidate Ligands. *Adv. Synth. Catal.* **2003**, *345*, 971–973. (g) Boezio, A. A.; Charette, A. B. Catalytic Enantioselective Addition of Dialkylzinc to *N*-Diphenylphosphinoylimines. A Practical Synthesis of α -Chiral Amines. *J. Am. Chem. Soc.* **2003**, *125*, 1692–1693. (h) Boezio, A. A.; Pytkowicz, J.; Côté, A.; Charette, A. B. Asymmetric, Catalytic Synthesis of α -Chiral Amines Using a Novel Bis(phosphine) Monoxide Chiral Ligand. *J. Am. Chem. Soc.* **2003**, *125*, 14260–14261. (i) Côté, A.; Boezio, A. A.; Charette, A. B. Catalytic Asymmetric Addition of Diorganozinc Reagents to *N*-Phosphinoylalkylamines. *Proc. Nat. Acad. Sci. U.S.A.* **2004**, *101*, 5405–5410. (j) Côté, A.; Boezio, A. A.; Charette, A. B. Evidence for the Structure of the Enantioactive Ligand in the Phosphine–Copper-Catalyzed Addition of Diorganozinc Reagents to Imines. *Angew. Chem., Int. Ed.* **2004**, *43*, 6525–6528. (k) Wang, M.-C.; Xu, C.-L.; Zou, Y.-X.; Liu, H.-M.; Wang, D.-K. New Chiral Ferrocenyl Amidophosphine Ligand for Remarkable Improvement of Enantioselectivities in Copper-Catalyzed Addition of Diethylzinc to *N*-Sulfonylimines. *Tetrahedron Lett.* **2005**, *46*, 5413–5416. (l) Desrosiers, J.-N.; Côté, A.; Charette, A. B. Enantioselective Synthesis of β -Amino Alcohols and α -Amino Acids via a Copper Catalyzed Addition of Diorganozinc Reagents to *N*-Phosphinoylimines. *Tetrahedron* **2005**, *61*, 6186–6192. (m) Wang, M.-C.; Liu, L.-T.; Hua, Y.-Z.; Zhang, J.-S.; Shi, Y.-Y.; Wang, D.-K. Novel Chiral Copper Complexes of *N,P*-Ferrocenyl Ligands with Central and Planar Chirality as Efficient Catalyst for Asymmetric Addition of Diethylzinc to Imines. *Tetrahedron Asymmetry* **2005**, *16*, 2531–2534. (n) Kim, B.-S.; Kang, S.-W.; Kim, K. H.; Ko, D.-H.; Chung, Y.; Ha, D. C. Catalytic Enantioselective Diethylzinc Addition to *N*-Phosphinoylimine Using Thiophosphoramidate of Homologated Binaphthyl *N,O*-Ligand. *Bull. Korean Chem. Soc.* **2005**, *26*, 1501–1502. (o) Lauzon, C.; Charette, A. B. Catalytic Asymmetric Synthesis of α,α,α -Trifluoromethylamines by the Copper-Catalyzed Nucleophilic Addition of Diorganozinc Reagents to Imines. *Org. Lett.* **2006**, *8*, 2743–2745. (p) Wang, M.-C.; Xu, C.-L.; Cheng, F.; Ding, X. Chiral Ferrocenyl Amidophosphine Ligand for Highly Enantioselective Addition of Diethylzinc to *N*-Diphenylphosphinoylimines. *Tetrahedron* **2006**, *62*, 12220–12226. (q) Shi, M.; Lei, Z.-Y.; Xu, Q. Asymmetric Addition of Diethylzinc to Diphenylphosphinoyl-Imines Catalyzed by Copper(II) Trifluoromethanesulfonate-Chiral (2-Ethylamino-[1,1]binaphthalenyl-2-yl)-thiophosphoramidate Acid *O,O*-Diaryl Ester Ligands. *Adv. Synth. Catal.* **2006**, *348*, 2237–2242.
- (10) For enantioselective zirconium, titanium, and hafnium catalyzed addition of organozinc reagents to imines, see (a) Porter, J. R.; Traverse, J. F.; Hoveyda, A. H.; Snapper, M. L. Enantioselective Synthesis of Arylamines through Zr-Catalyzed Addition of Dialkylzinc to Imines. Reaction Development by Screening of Parallel Libraries. *J. Am. Chem. Soc.* **2001**, *123*, 984–985. (b) Porter, J. R.; Traverse, J. F.; Hoveyda, A. H.; Snapper, M. L. Three-Component Catalytic Asymmetric Synthesis of Aliphatic Amines. *J. Am. Chem. Soc.* **2001**, *123*, 10409–10410. (c) Akullian, L. C.; Snapper, M. L.; Hoveyda, A. H. Three-Component Enantioselective Synthesis of Propargylamines through Zr-Catalyzed Additions of Alkyl Zinc Reagents to Alkynylimines. *Angew. Chem., Int. Ed.* **2003**, *42*, 4244–4247. (d) Akullian, L. C.; Porter, J. R.; Traverse, J. F.; Snapper, M. L.; Hoveyda, A. H. Asymmetric Synthesis of Acyclic Amines Through Zr and Hf Catalyzed Enantioselective Alkylzinc Reagents to Imines. *Adv. Synth. Catal.* **2005**, *347*, 417–425. (e) Basra, S.; Fennie, M. W.; Kozlowski, M. C. Catalytic Asymmetric Addition of Dialkylzinc Reagents to α -Aldiminoesters. *Org. Lett.* **2006**, *8*, 2659–2662.
- (11) For enantioselective rhodium catalyzed addition of organozinc reagents to imines, see Nishimura, T.; Yasuhara, Y.; Hayashi, T. Asymmetric Addition of Dimethylzinc to *N*-Tosylarylimines Catalyzed by a Rhodium–Diene Complex toward the Synthesis of Chiral 1-Arylethylamines. *Org. Lett.* **2006**, *8*, 979–981.
- (12) For reviews encompassing catalytic enantioselective addition of organolithium reagents to carbonyl compounds and imines, see (a) Tomioka, K. Asymmetric Synthesis Utilizing External Chiral Ligands. *Synthesis* **1990**, 541–549. (b) Berrisford, D. J. Catalytic Asymmetric C–C Bond Formation: New Enolato- and Organolithium Chemistry. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 178–180. (c) Denmark, S. E.; Nicaise, O. J.-C. Ligand Mediated Addition of Organometallic Reagents to Azomethine Functions. *Chem. Commun.* **1996**, 999–1004. (d) Enders, D.; Reinhold, U. Asymmetric Synthesis of Amines by Nucleophilic 1,2-Addition of Organometallic Reagents to the CN-Double Bond. *Tetrahedron Asymmetry* **1997**, *8*, 1895–1946.
- (13) For enantioselective rhodium-catalyzed addition of organometallic reagents to imines, see (a) Organotin reagents: Hayashi, T.; Ishigedani, M. Rhodium Catalyzed Asymmetric Arylation of Imines with Organostannanes. Asymmetric Synthesis of Diarylmethylamines. *J. Am. Chem. Soc.* **2000**, *122*, 976–977. (b) Hayashi, T.; Ishigedani, M. Rhodium Catalyzed Asymmetric Arylation of α,β -Unsaturated Imines with Arylstannanes. Catalytic Asymmetric Synthesis of Allylic Amines. *Tetrahedron* **2001**, *57*, 2589–2595. (c) Organotitanium reagents: Hayashi, T.; Kawai, M.; Tokunaga, N. Asymmetric Synthesis of Diarylmethyl Amines by Rhodium Catalyzed Asymmetric Addition of Aryl Titanium Reagents to Imines. *Angew. Chem., Int. Ed.* **2004**, *43*, 6125–6128. (d) Organoboron reagents: Kuriyama, M.; Soeta, T.; Hao, X.; Chen, Q.; Tomioka, K. *N*-Boc-L-Valine-Connected Amidomonophosphane Rhodium(I) Catalyst for Asymmetric Arylation of *N*-Tosylarylimines with Arylboroxines. *J. Am. Chem. Soc.* **2004**, *126*, 8128–8129. (e) Tokunaga, N.; Otomaru, Y.; Okamoto, K.; Ueyama, K.; Shintani, R.; Hayashi, T. C_2 -Symmetric Bicyclo[2.2.2]octadienes as Chiral Ligands: Their High Performance in Rhodium Catalyzed Asymmetric Arylation of *N*-Tosylarylimines. *J. Am. Chem. Soc.* **2004**, *126*, 13584–13585. (f) Weix, D. J.; Shi, Y.; Ellman, J. A. Diastereoselective and Enantioselective Rh(II)-Catalyzed Additions of Arylboronic Acids to *N*-tert-Butanesulfinyl and *N*-Diphenylphosphinoyl Aldimines. *J. Am.*

- Chem. Soc.* **2005**, *127*, 1092–1093. (g) Otomaru, Y.; Tokunaga, N.; Shintani, R.; Hayashi, T. C_2 -Symmetric Bicyclo[3.3.1]nonadiene as a Chiral Ligand for Rhodium-Catalyzed Asymmetric Arylation of *N*-(4-Nitrobenzenesulfonyl)arylimines. *Org. Lett.* **2005**, *7*, 307–310. (h) Duan, H.-F.; Jia, Y.-X.; Wang, L.-X.; Zhou, Q.-L. Enantioselective Rh-Catalyzed Arylation of *N*-Tosylarylimines with Arylboronic Acids. *Org. Lett.* **2006**, *8*, 2567–2569.
- (14) Catalyzed addition of vinylzirconocenes to imines is known, but enantioselective variants have not been developed. (a) Kakuuchi, A.; Taguchi, T.; Hanzawa, Y. Rh(I)-Catalyzed Addition of Alkenylzirconocene Chlorides to Aldimine Derivatives. *Tetrahedron Lett.* **2003**, *44*, 923–926. (b) Wipf, P.; Kendall, C.; Stephenson, C. R. J. Dimethylzinc-Mediated Additions of Alkenylzirconocenes to Aldimines. New Methodologies for Allylic Amine and *C*-Cyclopropyl-alkylamine Syntheses. *J. Am. Chem. Soc.* **2003**, *125*, 761–768.
- (15) Enantioselective Ni-catalyzed alkyne, imine, triethylborane three-component coupling has been reported, but modest selectivities (51–89% ee) are observed. For this method, vinylation is accompanied by ethyl transfer. Patel, S. J.; Jamison, T. F. Asymmetric Catalytic Coupling of Organoboranes, Alkynes, and Imines with a Removable (Trialkylsilyloxy)ethyl Group—Direct Access to Enantiomerically Pure Primary Allylic Amines. *Angew. Chem., Int. Ed.* **2004**, *43*, 3941–3944.
- (16) A compilation of reviews on the subject of metal-catalyzed reductive C–C coupling appears in volume 279 of *Top. Curr. Chem.*
- (17) For selected reviews encompassing intra- and intermolecular direct reductive coupling of alkynes to carbonyl partners, see (a) Ojima, I.; Tzamanioudaki, M.; Li, Z.; Donovan, R. J. Transition Metal-Catalyzed Carbocyclizations in Organic Synthesis. *Chem. Rev.* **1996**, *96*, 635–662. (b) Montgomery, J. Nickel-Catalyzed Cyclizations, Couplings, and Cycloadditions Involving Three Reactive Components. *Acc. Chem. Res.* **2000**, *33*, 467–473. (c) Montgomery, J.; Amarashinghe, K. K. D.; Chowdhury, S. K.; Oblinger, E.; Seo, J.; Savchenko, A. V. Nickel-Catalyzed Cyclizations of Alkynyl Enones and Alkynyl Enals. *Pure Appl. Chem.* **2002**, *74*, 129–133. (d) Ikeda, S.-I. Nickel-Catalyzed Coupling of Carbonyl Compounds and Alkynes or 1,3-Dienes: An Efficient Method for the Preparation of Allylic, Homoallylic, and Bishomoallylic Alcohols. *Angew. Chem., Int. Ed.* **2003**, *42*, 5120–5122. (e) Miller, K. M.; Molinaro, C.; Jamison, T. F. Catalytic Reductive Carbon–Carbon Bond-Forming Reactions of Alkynes. *Tetrahedron Asymmetry* **2003**, 3619–3625. (f) Montgomery, J. Nickel-Catalyzed Reductive Cyclizations and Couplings. *Angew. Chem., Int. Ed.* **2004**, *43*, 3890–3908.
- (18) Ojima, I.; Tzamanioudaki, M.; Tsai, C.-Y. Extremely Chemoselective Silylformylation and Silylcarbocyclization of Alkynals. *J. Am. Chem. Soc.* **1994**, *116*, 3643–3644.
- (19) (a) Crowe, W. E.; Rachita, M. J. Titanium-Catalyzed Reductive Cyclization of δ,ϵ -Unsaturated Ketones and Aldehydes. *J. Am. Chem. Soc.* **1995**, *117*, 6787–6788. (b) For an aligned study, see Kablaoui, N. M.; Buchwald, S. L. Reductive Cyclization of Enones by a Titanium Catalyst. *J. Am. Chem. Soc.* **1995**, *117*, 6785–6786.
- (20) (a) Oblinger, E.; Montgomery, J. A New Stereoselective Method for the Preparation of Allylic Alcohols. *J. Am. Chem. Soc.* **1997**, *119*, 9065–9066. (b) Tang, X.-Q.; Montgomery, J. Nickel Catalysis in the Stereoselective Preparation of Quinolidine, Pyrrolizidine, and Indolizidine Alkaloids: Total Synthesis of (+)-Allopumiliotoxin 267A. *J. Am. Chem. Soc.* **1999**, *121*, 6098–6099. (c) Tang, X.-Q.; Montgomery, J. Nickel-Catalyzed Preparation of Bicyclic Heterocycles: Total Synthesis of (+)-Allopumiliotoxin 267A, (+)-Allopumiliotoxin 339A, and (+)-Allopumiliotoxin 339B. *J. Am. Chem. Soc.* **2000**, *122*, 6950–6954. (d) Mahandru, G. M.; Liu, G.; Montgomery, J. Ligand-Dependent Scope and Divergent Mechanistic Behavior in Nickel-Catalyzed Reductive Couplings of Aldehydes and Alkynes. *J. Am. Chem. Soc.* **2004**, *126*, 3698–3699. (e) Knapp-Reed, B.; Mahandru, G. M.; Montgomery, J. Access to Macrocyclic Endocyclic and Exocyclic Allylic Alcohols by Nickel-Catalyzed Reductive Cyclization of Ynals. *J. Am. Chem. Soc.* **2005**, *127*, 13156–13157.
- (21) (a) Huang, W.-S.; Chan, J.; Jamison, T. F. Highly Selective Catalytic Intermolecular Reductive Coupling of Alkynes and Aldehydes. *Org. Lett.* **2000**, *2*, 4221–4223. (b) Miller, K. M.; Huang, W.-S.; Jamison, T. F. Catalytic Asymmetric Reductive Coupling of Alkynes and Aldehydes: Enantioselective Synthesis of Allylic Alcohols and α -Hydroxy Ketones. *J. Am. Chem. Soc.* **2003**, *125*, 3442–3443. (c) Miller, K. M.; Jamison, T. F. Highly Regioselective Catalytic Asymmetric Reductive Coupling of 1,3-Enynes and Ketones. *Org. Lett.* **2005**, *7*, 3077–3080.
- (22) Takai, K.; Sakamoto, S.; Isshiki, T. Regioselective Reductive Coupling of Alkynes and Aldehydes Leading to Allylic Alcohols. *Org. Lett.* **2003**, *5*, 653–655.
- (23) The first reported catalytic hydrogenations were developed by Paul Sabatier and constitute the basis for his receipt of the Nobel Prize in 1912. (a) Sabatier, P.; Senderens, J.-B. Action of Nickel on Ethylene. Ethane Synthesis. *C. R. Hebd. Seances Acad. Sci.* **1897**, *124*, 1358–1361. (b) Sabatier, P.; Senderens, J.-B. Hydrogenation of Acetylene in the Presence of Nickel. *C. R. Hebd. Seances Acad. Sci.* **1899**, *128*, 1173–1176. (c) Sabatier, P.; Senderens, J.-B. Preparation of Hexahydrobenzene. *C. R. Hebd. Seances Acad. Sci.* **1901**, *132*, 210–212. (d) For a biographical sketch of Paul Sabatier, see Lattes, A. From Catalytic Hydrogenation to the Chemical Theory of the Catalysis: Paul Sabatier, Genius Chemist, Decentralization Apostle. *C. R. Acad. Sci., Ser. Ilc: Chim.* **2000**, *3*, 705–709.
- (24) (a) Fischer, F.; Tropsch, H. The Preparation of Synthetic Oil Mixtures (Synthol) from Carbon Monoxide and Hydrogen. *Brennst. Chem.* **1923**, *4*, 276–285. (b) Fischer, F.; Tropsch, H. Synthesis of Higher Members of the Aliphatic Series from Carbon Monoxide. *Chem. Ber.* **1923**, *56B*, 2428–2443.
- (25) For recent reviews on alkene hydroformylation, see (a) Cornils, B.; Herrmann, W. A.; Kohlpaintner, C. W. Otto Roelen as the Pioneer of Industrial Homogeneous Catalysis. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 2144–2163. (b) Beller, M.; Cornils, B.; Frohning, C. D.; Kohlpaintner, C. W. Progress in Hydroformylation and Carbonylation. *J. Mol. Catal.* **1995**, *104*, 17–85. (c) Eilbracht, P.; Barfacker, L.; Buss, C.; Hollmann, C.; Kitsos-Rzychon, B. E.; Kranemann, C. L.; Rische, T.; Roggenbuck, R.; Schmidt, A. Tandem Reaction Sequences under Hydroformylation Conditions: New Synthetic Applications of Transition Metal Catalysis. *Chem. Rev.* **1999**, *99*, 3329–3365. (d) Nozaki, K. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer-Verlag: Berlin, Germany, 1999; Vol. 1, pp 381–413. (e) Breit, B. Synthetic Aspects of Stereoselective Hydroformylation. *Acc. Chem. Res.* **2003**, *36*, 264–275.
- (26) For selected reviews on hydrogen-mediated C–C coupling, see (a) Ngai, M.-Y.; Krische, M. J. Lessons in Green Chemistry from the Smallest of Molecules: C–C Bond Formation via Catalytic Hydrogenation. *Chim. Oggi/Chem. Today* **2006**, *24*, 12–14. (b) Ngai, M.-Y.; Kong, J.-R.; Krische, M. J. Hydrogen-Mediated C–C Bond Formation: A Broad New Concept in Catalytic C–C Coupling. *J. Org. Chem.* **2007**, *72*, 1063–1072. (c) Iida, H.; Krische, M. J. Catalytic Reductive Coupling of Alkenes and Alkynes to Carbonyl Compounds and Imines Mediated by Hydrogen. *Top. Curr. Chem.* **2007**, *279*, 77–104.
- (27) Prior to our work, two isolated examples of hydrogen-mediated C–C bond formations not involving carbon monoxide as a coupling partner were reported. (a) Molander, G. A.; Hoberg, J. O. Organotrium-Catalyzed Cyclization of Substituted 1,5- and 1,6-Dienes. *J. Am. Chem. Soc.* **1992**, *114*, 3123–3125. (b) Kokube, K.; Miura, M.; Nomura, M. Rhodium-Catalyzed Reaction of Benzoic Anhydride with Styrene under Molecular Hydrogen. *Organometallics* **1995**, *14*, 4521–4524.
- (28) Subsequent to our work, two isolated examples of hydrogen-mediated C–C bond formation not involving carbon monoxide as a coupling partner were reported. (a) Smith, D. M.; Pulling, M. E.; Norton, J. R. Tin-Free and Catalytic Radical Cyclizations. *J. Am. Chem. Soc.* **2007**, *129*, 770–771. (b) Jung, I. G.; Seo, J.; Lee, S. I.; Choi, S. Y.; Cheung, Y. K. Reductive Cyclization of Dienes and Enynes Catalyzed by Allyl Platinum *N*-Heterocyclic Carbene Complexes. *Organometallics* **2006**, *25*, 4240–4242.
- (29) Side products of reductive C–C bond formation have been observed in catalytic hydrogenation on rare occasions. (a) Moyes, R. B.; Walker, D. W.; Wells, P. B.; Whan, D. A.; Irvine, E. A. Catalysis and Surface Characterisation (Special Publication). *R. Soc. Chem.* **1992**, *114*, 207–212. (b) Bianchini, C.; Meli, A.; Peruzzini, M.; Vizzi, F.; Zanobini, F.; Frediani, P. A. Homogeneous Iron(II) System Capable of Selectivity Catalyzing the Reduction of Terminal Alkynes to Alkenes and Buta-1,3-Dienes. *Organometallics* **1989**, *8*, 2080–2082.
- (30) Garner, S.; Krische, M. J. Metal Catalyzed Reductive Aldol Coupling. In *Modern Reductions*; Andersson, P., Munslow, I., Eds.; Wiley-VCH: Weinheim, Germany, 2007, in press.
- (31) For mechanistic studies of alkene hydrogenation catalyzed by neutral Rh(I) complexes, see (a) Tolman, C. A.; Meakin, P. Z.; Lindner, D. L.; Jesson, J. P. Triarylphosphine, Hydride, and Ethylene Complexes or Rhodium(I) Chloride. *J. Am. Chem. Soc.* **1974**, *96*, 2762–2774. (b) Halpern, J.; Wong, C. S. Hydrogenation of Tris(triphenylphosphine)chlororhodium(I). *Chem. Commun.* **1973**, 629–630. (c) Halpern, J.; Okamoto, T.; Zakhariyev, A. Mechanism of the Chlorotris(triphenylphosphine) Rhodium(I)-Catalyzed Hydrogenation of Alkenes. The Reaction of Chlorodihydrodtris(triphenylphosphine)rhodium(III) with Cyclohexene. *J. Mol. Catal.* **1977**, *2*, 65–68.
- (32) For reviews encompassing the mechanism of asymmetric hydrogenation catalyzed by cationic rhodium complexes, see (a) Halpern, J. Asymmetric Catalytic Hydrogenation: Mechanism and Origin of Enantioselection. *Asymmetric Synth.* **1985**, *5*, 41–69. (b) Landis, C. R.; Brauch, T. W. Probing the Nature of H_2 Activation in Catalytic Asymmetric Hydrogenation. *Inorg. Chim. Acta* **1998**, *270*, 285–297.

- (33) Musashi, Y.; Sakaki, S. Theoretical Study of Rhodium(III)-Catalyzed Hydrogenation of Carbon Dioxide into Formic Acid. Significant Differences in Reactivity among Rhodium(III), Rhodium(I), and Ruthenium(II) Complexes. *J. Am. Chem. Soc.* **2002**, *124*, 7588–7603.
- (34) (a) Kong, J.-R.; Ngai, M.-Y.; Krische, M. J. Highly Enantioselective Direct Reductive Coupling of Conjugated Alkynes and α -Ketoesters via Rhodium-Catalyzed Asymmetric Hydrogenation. *J. Am. Chem. Soc.* **2006**, *128*, 718–719. (b) Hong, Y.-T.; Cho, C.-W.; Skucas, E.; Krische, M. J. Enantioselective Reductive Coupling of 1,3-Enynes to Glyoxalates Mediated by Hydrogen: Asymmetric Synthesis of β,γ -Unsaturated α -Hydroxy Esters. *Org. Lett.* **2007**, *9*, in press. (c) Komanduri, V.; Krische, M. J. Enantioselective Reductive Coupling of 1,3-Enynes to Heterocyclic Aromatic Aldehydes and Ketones via Rhodium-Catalyzed Asymmetric Hydrogenation: Mechanistic Insight into the Role of Brønsted Acid Additives. *J. Am. Chem. Soc.* **2006**, *128*, 16448–16449. (d) Kong, J.-R.; Cho, C.-W.; Krische, M. J. Hydrogen-Mediated Reductive Coupling of Conjugated Alkynes with Ethyl (*N*-Sulfinyl)iminoacetates: Synthesis of Unnatural α -Amino Acids via Rhodium-Catalyzed C–C Bond Forming Hydrogenation. *J. Am. Chem. Soc.* **2005**, *127*, 11269–11276. (e) Rhee, J.-U.; Krische, M. J. Highly Enantioselective Reductive Cyclization of Acetylenic Aldehydes via Rhodium Catalyzed Asymmetric Hydrogenation. *J. Am. Chem. Soc.* **2006**, *128*, 10674–10675.
- (35) (a) Kong, J.-R.; Krische, M. J. Catalytic Carbonyl (*Z*)-Dienylation via Multicomponent Reductive Coupling of Acetylene to Aldehydes and α -Ketoesters Mediated by Hydrogen: Carbonyl Insertion into Cationic Rhodocyclopentadienes. *J. Am. Chem. Soc.* **2006**, *128*, 16040–16041. (b) Skucas, E.; Kong, J.-R.; Krische, M. J. Enantioselective Reductive Coupling of Acetylene to *N*-Arylsulfonyl Imines via Rhodium Catalyzed C–C Bond Forming Hydrogenation: (*Z*)-Dienyl Allylic Amines. *J. Am. Chem. Soc.* **2007**, *129*, 7242–7243.
- (36) Bianchini, C.; Caulton, K. G.; Chardon, C.; Eisenstein, O.; Foltling, K.; Johnson, T. J.; Meli, A.; Peruzzini, M.; Rauscher, D. J.; Streib, W. E.; Vizza, F. An η^4 -Benzene Species Mediates Acetylene Cyclo-trimerization. *J. Am. Chem. Soc.* **1991**, *113*, 5127–5129, and references cited therein.
- (37) (a) Dewar, M. J. S. A Review of the π -Complex Theory. *Bull. Soc. Chim. Fr.* **1951**, *18*, C71–C79. (b) Chatt, J.; Duncanson, L. A. Olefin Co-ordination Compounds. III. Infrared Spectra and Structure: Attempted Preparation of Acetylene Compounds. *J. Chem. Soc.* **1953**, 2939–2947. (c) Dewar, M. J. S.; Ford, G. P. Relationship Between Olefinic π -Complexes and Three-Membered Rings. *J. Am. Chem. Soc.* **1979**, *101*, 783–791.
- (38) (a) Vaska, L.; Peone, J., Jr. Simple Synthetic Route to d8 Complexes of Rhodium and Iridium. Electronegativity Scale for Anionic Ligands. *Chem. Commun.* **1971**, 418–419. (b) Haynes, A.; McNish, J.; Pearson, J. M. Cis–Trans Isomerism in $[M(CO)_2L_4]^-$ ($M = Rh, Ir$): Kinetic, Mechanistic and Spectroscopic Studies. *J. Organomet. Chem.* **1998**, *551*, 339–347. (c) Grotjohn, D. B.; Collins, L. S. B.; Wolpert, M.; Bikzhanova, G. A.; Lo, H. C.; Combs, D.; Hubbard, J. L. First Direct Structural Comparison of Complexes of the Same Metal Fragment to Ketenes in Both C,C- and C,O-Bonding Mode. *J. Am. Chem. Soc.* **2001**, *123*, 8260–8270.
- (39) Li, J.; Schreckenbach, G.; Ziegler, T. A Reassessment of the First Metal-Carbonyl Dissociation Energy in $M(CO)_4$ ($M = Ni, Pd, Pt$), $M(CO)_5$ ($M = Fe, Ru, Os$), and $M(CO)_6$ ($M = Cr, Mo, W$) by a Quasirelativistic Density Functional Method. *J. Am. Chem. Soc.* **1995**, *117*, 486–494.
- (40) (a) Ngai, M.-Y.; Barchuk, A.; Krische, M. J. Allylic Amines via Iridium Catalyzed C–C Bond Forming Hydrogenation: Imine Vinylation in the Absence of Stoichiometric Byproducts or Metallic Reagents. *J. Am. Chem. Soc.* **2007**, *129*, 8432–8433. (b) Barchuk, A.; Ngai, M.-Y.; Krische, M. J. Optically Enriched Allylic Amines via Alkyne–Imine Reductive Coupling Under Hydrogenation Conditions: Enantioselective Iridium Catalyzed Imine Vinylation. **2007**, submitted for publication.
- (41) Brown, J. M. Directed Homogeneous Hydrogenation. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 190–203.

AR7001123