

Drugs of Abuse That Mediate Advanced Glycation End Product Formation: A Chemical Link to Disease Pathology

JENNIFER B. TREWEEK,[†] TOBIN J. DICKERSON,^{‡,⊥} AND
KIM D. JANDA^{*,†,⊥}


[†]Departments of Chemistry and Immunology of The Skaggs Institute for Chemical Biology, [‡]Department of Chemistry, and [⊥]Worm Institute for Research and Medicine (WIRM), The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037

RECEIVED ON NOVEMBER 2, 2008

CON SPECTUS

Nicotine and methamphetamine are frequently abused in modern society, despite the increasing evidence of their addictive, neuropharmacological, and toxic effects. Tobacco, the most widely abused substance, is the leading cause of preventable death in the United States, killing nearly half a million Americans annually. A methamphetamine epidemic has also spread during the past decade; severe neurotoxicity and addictiveness contribute to the drug's notoriety. Although the majority of research on these two drugs is of pharmacological and neurobiological motivation, further study of these molecules from a chemical perspective may provide novel mechanistic insight into either their addictive potential or their pathological effects. For example, nicotine and methamphetamine share a common structural feature, a secondary amine, suggesting that these molecules could possess similar (or analogous) *in vivo* reactivity. Discoveries concerning the synthetic requirements for aqueous aldol catalysis and the feasibility of the enamine mechanism under physiological conditions have given rise to the hypothesis that ingested molecules, such as abused drugs, could participate in reactions utilizing an enamine intermediate *in vivo*.

The chemical reactivity of exogenous drugs with amine functionalities was initially examined in the context of the Maillard reaction, or nonenzymatic browning. The heating of reducing sugars with amino acids yields a brown solution; studies of this reaction were originally applied to food chemistry for the production of distinct flavors and aromas. Further research has since revealed numerous instances in which the *in vivo* production of advanced glycation end products (AGEs) through the Maillard reaction contribute to the pathology of disease states. Specifically, the modification of long-lived proteins by glycation and glycoxidation and the accumulation of these AGEs compromise the original function of such proteins and change the mechanical properties of affected tissue. In this Account, we summarize our investigations into the capacity for exogenous compounds to initiate the Maillard reaction and the corresponding physiological and immunological impact of the drug-conjugated AGEs that form. Many of the pathological components of diabetes, atherosclerosis, cancer, macular degeneration, Alzheimer's disease, and even the normal aging process are attributable to AGEs and their potential for aggregate formation in the vasculature. A deeper understanding of AGEs, and particularly glycated proteins, will provide fundamental mechanistic insight into disease origins.



Introduction

Substance addiction, which continues to inflict a large economic and social burden on modern society, involves a complex sequence of brain

adaptations. The development of drug sensitization and addictive behaviors, which involves neuroplasticity in brain regions subserving learning and memory, is often examined through models of nicotine and psychostimulant abuse. Whereas




FIGURE 1. Structures of L-proline, nornicotine, dopamine, and (+)-methamphetamine.

amphetamines directly activate the mesolimbic dopaminergic circuits, which govern the learned reward aspect of addiction, nicotine binds nicotinic acetylcholine receptors whose downstream effect includes the modulation of dopamine reactivity within the nucleus accumbens. Thus, both drugs possess a similar capacity to induce severe drug dependence in the user.

The quest to elucidate the altered psychological and physical pathology of drug addicts and correspondingly to develop better addiction therapies motivates the majority of research on abused substances. Many drugs, nicotine and amphetamines notwithstanding, possess inherently reactive chemical functionalities that would allow them to participate in biologically relevant chemical reactions; however the potential for this *in vivo* reactivity to induce certain pathologies in addicts is often overlooked in neurobiological research. By contrast, the examination of drug reactivity has been undertaken by chemists, and a classic example is the finding that specific sedatives and depressants promote alkaloid formation *in vivo* by inhibiting the oxidation of aldehydes.¹ Their metabolism generates acetaldehyde, which readily condenses with biogenic amines or amino acids to form imines, and subsequent ring closure may lead to the formation of potentially psychoactive or toxic alkaloids through Schiff base formation by the Pictet–Spengler reaction. We have postulated that nornicotine and methamphetamine (Figure 1), psychoactive compounds possessing a secondary amine moiety, may mediate complementary enamine-based chemistry *in vivo* and that this reactivity of abused drugs may provide a mechanistic explanation for certain pathologies observed in addicts.

The Enamine Mechanism



Revered by chemists as an essential method to form carbon–carbon bonds with great stereoselective control, the aldol reaction also has prominence within the field of biology with its application to metabolic processes employing aldolases as glycolytic enzymes, the formation of DNA adducts of acetaldehyde, and prebiotic catalysis.^{2,3} Both natural aldolase and aldolase mimetics, which include both catalytic antibodies and compounds with aldolase activity, have been studied extensively in order to gain improved efficiency, stereoselectivity, and mechanistic understanding of the aldol reaction. Thus, the discovery that a chiral biomolecule, pro-

line, could catalyze asymmetric aldol reactions represented an important contribution to the field of organocatalysis, particularly with respect to iminium-based and enamine-based mechanisms for catalysis.^{4,5} This chemical framework was successfully applied to research in aldolase enzyme evolution and small oligopeptide-mediated catalysis of the asymmetric formation of sugars under prebiotic conditions.⁶ However, consideration of this enamine-based chemistry during investigations into the mechanistic basis of disease pathologies or drug effects *in vivo* was minimal until small-molecule-based aqueous aldol catalysis was proven operable under physiologically relevant conditions.⁷

Our research into the reactivity of nicotine metabolites led us to discover that nornicotine, a minor tobacco alkaloid, possessed organocatalytic activity under buffered aqueous conditions, and opposite to the biomolecule proline, such catalysis was lost in common organic solvents (Scheme 1).⁷ Furthermore, as proven through capturing the hypothesized enamine nucleophile in trapping experiments, nornicotine catalyzed the aqueous aldol reaction involving an activated aldehyde acceptor, while the biomolecule proline failed to mediate such catalysis at physiological pH and temperature. Whereas the nornicotine-derived enamine was relatively stable at pH 7.5–8, the proline-derived enamine was vulnerable to hydrolysis, thus precluding catalysis.⁸ Contrary to previous studies with proline, nornicotine-based catalysis illustrated the physiological relevance of the enamine mechanism to the aldol reaction (Scheme 2A) and was suggestive of the more general operability of this mechanism *in vivo* (Scheme 2B).

The Maillard Reaction

The Maillard reaction, which was first described as the pathway responsible for the color and aroma of certain cooked foods, is initiated by the reversible coupling between a free amine and an aldehyde or ketone moiety of a reducing sugar to form Schiff bases.⁹ Amadori rearrangement of the Schiff base intermediates and then degradation, fragmentation, or oxidation steps lead to the formation of advanced glycation end products (AGE, Figure 2A).¹⁰ Its scientific relevance has since extended beyond its application to food chemistry to the study of its reaction mechanism and role in disease states. With respect to the latter, the diverse array of glycated proteins are potentially toxic, long-lived, and deleterious to the

SCHEME 1. Enamine-Based Reaction Cycle for the Covalent Catalysis of the Aqueous Aldol Reaction by Nornicotine**SCHEME 2.** Catalysis by Nornicotine^a

^a The enamine mechanism is relevant to (A) the aqueous aldol reaction and (B) the Maillard reaction.

endogenous role of unmodified protein.¹¹ Specifically, the cross-linking of collagen, a protein with a slow rate of turnover, may precipitate endothelial dysfunction as AGE deposits accumulate on vasculature. Fiber stiffness, thermal denaturation temperature, and enzyme resistance comprise a few of the physical properties of proteins compromised by intra- and intermolecular cross-linking.¹² Indeed, many of the

pathological components of diabetes, atherosclerosis, cancer, macular degeneration, Alzheimer's disease (AD), and even the normal aging process are attributable to AGE and aggregate formation.^{10,13–16} A second major impact of the Maillard reaction is the side-chain modification of reacting proteins. The altered charge distribution of specific amino acids can abolish the interaction or recognition of the effected molecule with




FIGURE 2. Advanced glycation end product formation: (A) scheme of the Maillard reaction, depicted by the reaction of glucose and a protein-derived amine; (B) structures of common advanced glycation end products (AGEs), in which the ϵ -amino group of lysine serves as the protein-derived amine.

other endogenous biomolecules, a scenario important to degradative enzymes, cell surface receptors, and the cell–collagen interactions mediated by integrin recognition of arginine residues.

It should be noted that the Maillard products detected *in vivo* and implicated in disease states are not formed exclusively by glycation. Indeed, several physiologically relevant pathways contribute to the wide array of immunochemically distinct AGEs. The dehydration and rearrangement of Amadori products may form α -dicarbonyls and the classic AGEs *N* $^{\epsilon}$ -(carboxymethyl)lysine (CML), pentosidine, and pyrraline (Figure 2B).¹⁷ However, Schiff base fragmentation to generate glycolaldehyde and rearrangement of Amadori products

and glyceraldehyde 3-phosphate also produces α -oxoaldehydes, the most notable of which are the reactive dicarbonyl 3-deoxyglucosone (3-DG) and methylglyoxal.¹⁸ Autoxidation reactions of reducing sugars and polyunsaturated fatty acids such as in the formation of glyoxal from glucose emphasize the ability of various other carbohydrate or lipid-dependent metabolic processes to generate α -oxoaldehydes.¹⁹ A distinguishing feature of these Maillard reaction intermediate processes is the role of oxidative reactions (i.e., glycooxidation and nonoxidative glycation).²⁰ Regardless, all intermediates subsequently react with the free amino groups of proteins, phospholipids, and nucleic acids and thereby contribute to the diversity of late-stage AGEs.

In Vivo Effects of AGE Formation

The complex kinetics and thermodynamics of the *in vivo* pathways generating AGEs have impeded the study of AGE formation. Schiff base condensation is a rapid and reversible process amenable to pharmacodynamic modeling. By contrast, the irreversible pathway by which Amadori products rearrange and form cross-links to generate AGEs is complicated by its sensitivity to reaction conditions and by the imperfect characterization of the numerous chemical entities and reaction intermediates produced en route. Even though the Amadori products form within hours to days after the initial reaction of the first sugar, the intermediate products, though measurable, are presumably too short-lived to have dramatic physiological ramifications in contrast to the end products of subsequent glycation and oxidation reactions.²¹ Nevertheless, detection of *in vitro* and *in vivo* prepared glycation and oxidation products holds merit in elucidating the kinetics of the early stages of the Maillard reaction.²² The reactivity of the monosaccharide, in which the open-chain conformer of the reducing sugar is conducive to Schiff base condensation, and the concentration of the reducing sugars, which governs the probability that the sugar will encounter a reactive amine, tend to dominate the reaction kinetics.^{23,24} It follows that hyperglycemic conditions may encourage the forward reaction toward Schiff base and Amadori product formation.²⁵

To elucidate the chemical mechanism by which AGEs arise from Amadori products is nontrivial because it involves analysis of a complex, heterogeneous mixture of AGEs forming from an already diverse array of Amadori products. To bypass this hurdle, scientists have focused on identifying pathologies derived from AGE formation and further elucidating how the *in vivo* occurrence of the Maillard reaction is causally related to them. AGE formation appears to induce a few main biological effects via a distinct progression of events: AGE attachment to and cross-linking of proteins, AGE deposition and subsequent induction of oxidative stress, activation of receptors for AGE (RAGEs), and induction of an immune response against the antigenic AGE-modified proteins.^{26,27} With respect to the latter, more common AGE motifs such as CML may serve as major immunological epitopes of anti-AGE autoantibodies of broader scope. However, all members of the heterogeneous array of AGEs may elicit a unique polyclonal response, which greatly complicates their characterization. AGE-modified proteins are detected through various methods including the use of antibodies to known AGE motifs, visualization of AGE deposits on specific tissues such as vasculature and lens collagen, measurement of the altered flu-

orescence produced by certain Maillard-type cross-links, and detection of nonfluorescent AGE species through other spectrographic, chromatographic, or immunological methods. Alternatively, RAGE activation, which represents a cell surface receptor dependent effect of *in vivo* AGE generation, contributes largely to the pathogenesis of vasculature dysfunction, and its expression is more readily quantified than heterologous AGEs.²⁸

Nornicotine-Based Glycation

The characterization of protein adducts and AGEs formed *in vivo* has illuminated the physiological importance of the Maillard reaction. In conjunction with our elucidation of the enamine mechanism for nornicotine-based aqueous aldol catalysis, we were exploring the propensity of commonly administered drugs containing an amine moiety to participate in other enamine-based reactions such as the Maillard reaction (Scheme 2).⁷ The reactive glycation products or glycotoxins present in tobacco and tobacco smoke extracts were previously demonstrated to mediate AGE formation and protein cross-linking *in vitro*.²⁹ This study also verified that cigarette smoke-derived glycotoxins induced AGE modification of serum proteins of smokers, illustrating the causal link between smoking, tobacco-associated AGE, and vascular disease.²⁹ However, from a mechanistic standpoint, no distinction was made between AGE formation through glycotoxins preformed in tobacco (e.g., during the tobacco curing process) or through tobacco-derived reactive amines that initiated the Maillard reaction *in vivo* after their inhalation.

To test whether nornicotine would initiate the Maillard reaction *in vitro*, the Amadori product of nornicotine was prepared through the incubation of nornicotine and glucose under physiological conditions as well as through synthetic methods (Scheme 2B).³⁰ To verify that this intermediate would react further to generate the glycated proteins associated with smoking, the incubation of nornicotine and glucose was repeated in the presence of protein [ribonuclease A (RNase A), bovine serum albumin (BSA), or human serum albumin (HSA)]. Detection of nornicotine-modified AGEs was accomplished through enzyme-linked immunosorbent assay (ELISA) and Western blot analysis using mAb NIC6C12 (Figure 3). Containing a 3-pyrrolidine-2-yl-pyridine nucleus and an alkyl linker attached to the pyrrolidine nitrogen, the hapten used to generate mAb NIC6C12 was designed to elicit anti-nicotine antibodies with superior recognition for the pyrrolidine nitrogen and the pyrrolidyl methyl group (Figure 3A).³¹ Even though mAb NIC6C12 was originally prepared for use as a passive vaccine, the conservation of the alkaloid nucleus during nic-




FIGURE 3. (A) Haptent used to elicit anti-nicotine mAbs and (B) detection of normicotine-modified proteins using anti-nornicotine mAb NIC6C12 and normicotine glycosylated protein; SDS/PAGE and Western blots: (a) BSA (lane 1); BSA + normicotine and glucose (lane 2); HSA (lane 3); HSA + normicotine and glucose (lane 4); (b) HSA + normicotine and glucose (lane 1); nonsmoker plasma sample A (lanes 2–4; 2.5, 5.0, and 10 mg/mL total protein concentration); nonsmoker plasma sample B (lanes 5–7); smoker plasma sample C (lanes 8–10); smoker plasma sample D (lanes 11–13). Reproduced with permission from ref 30. Copyright 2002 National Academy of Sciences, U.S.A.

otine metabolism to normicotine as well as during normicotine-mediated glycation proved fortuitous in that mAb NIC6C12 displayed good affinity for both normicotine and normicotine-modified AGEs.³¹


Though normicotine represents a minor metabolite of nicotine (~8% of peripherally metabolized nicotine), its extended half-life of 8 h permits its accumulation in the serum of smokers.³² Given this and the *in vitro* stability of experimentally generated normicotine glycation products upon extended incubations (>7 months), we hypothesized that normicotine-based AGEs would reach detectable levels in serum samples of smokers if the mechanism of normicotine-based glycation was operable *in vivo*. Plasma samples of smokers and nonsmokers were assessed by ELISA with mAb NIC6C12, and gratifyingly, normicotine-modified protein levels were substantially elevated in the plasma of smokers.³⁰

Whereas the formation of normicotine-modified AGEs validated a hypothesis as to the reactivity of amine-containing compounds, the biological significance of this chemical prediction had yet to be realized. Thus, we embarked upon an investigation into the *in vivo* ramifications of this normicotine-mediated glycation. Specifically, the propensity of normicotine to target lysine during protein glycation and cross-linking resonated with the current research on Alzheimer's disease (AD). Whereas initial investigations into the therapeutic potential of nicotine were based on the upregulation of nicotinic acetylcholine receptors (nAChR) to counteract the nAChR deficiency in the AD brain, alternative mechanisms to explain the documented nicotine-derived protection from AD were still considered.³³ The formation of neurotoxic intraneuronal neu-

rofibrillary tangles and amyloid plaques have been shown to be directly proportional to the development of neurological deficits; however the mechanism driving their formation is still debated.³⁴ It follows that a molecule able to prevent aberrant protein aggregation and more specifically avert the "amyloid-initiated cascade" may substantially delay disease progression.³⁵ Thus, to explore an alternative explanation for the nicotine-mediated antagonism of A β -peptide toxicity, Salomon et al. examined the effect of nicotine on the different conformations of A β -peptide in solution and concluded that nicotine blocked α -helix to β -sheet conversion through binding the histidine residues of the α -helix.³³ The A β -peptide normally interconverts between random coil, monomeric α -helical and oligomeric β -sheet structures, but a shift in this equilibrium favoring the β -sheet structure was hypothesized to cause aggregation and precipitation of β -sheet structures as amyloid. This process would then encourage further α -helix to β -sheet conversion. Salomon et al. surmised that the N-CH₃ and 5'-CH₂ pyrrolidine moieties of nicotine mediated the interaction with the α -helix structure, implying that many of the major alkaloidic metabolites of nicotine would behave similarly. This model was subsequently refuted when NMR studies revealed that neither the α -helical nor the random coil conformations of A β -peptide were bound by nicotine to an appreciable extent, suggesting that nicotine might interact with the small soluble A β -sheet aggregates to attenuate amyloidosis.³³

The folding of A β -pleated sheets, the structural building block of A β aggregates, relies on the specific KLVFF amino acid sequence.³⁶ The requirement of lysine in A β -peptide mis-

SCHEME 3. Methamphetamine Protein Glycation as Initiated by Methamphetamine and Glucose



folding provoked the hypothesis that normicotine-based glycation could target lysine residues of $A\beta$ in a manner reminiscent of glycation.^{7,30} Through assays using thioflavin T staining of fibrils and diffusion NMR experiments to monitor the progress of $A\beta$ aggregation in incubations of normicotine, glucose, and either $A\beta^{1-40}$ or $A\beta^{12-28}$, we demonstrated that normicotine is indeed capable of covalently modifying exposed lysine residues such as the Lys-16 of amyloid peptides and soluble $A\beta$ aggregates and that this modification blocks subsequent peptide aggregation.³⁷

Whereas AGE formation and protein cross-linking have previously been implicated in disease pathologies, the hypothesis that exogenous molecules may participate in and even initiate protein glycation with potentially detrimental effects represents an unexplored application of glycation chemistry. Thus, we postulated that our initial research regarding normicotine could be extrapolated to other abused molecules containing a secondary amine.

Methamphetamine-Based AGEs

Nicotine and methamphetamine (Figure 1), two widely abused drugs joined in their high addictive potential and reactive amine moiety, have well-known toxicity regardless of their ability to mediate protein glycation. Upon exposure, methamphetamine spurs excessive release and signaling of all monoamine neurotransmitters within the CNS through a variety of molecular mechanisms.^{38,39} Acute methamphetamine binges produce diffuse neuronal damage, which compromises dopaminergic signaling; however, the ramifications of chronic,

low dose exposure and the processes through which methamphetamine causes damage to the cardiovascular and periphery are ambiguous.^{38,40,41} Of particular interest is how the long-term pathological consequences of methamphetamine abuse mirror the cardiovascular complications arising from AGE generation. Not only does the amine functionality of methamphetamine permit its participation in the Maillard reaction (Scheme 3), but also its long serum half-life ($t_{1/2} = 12$ h) in comparison to other psychostimulants and its frequent administration by addicts support the hypothesis that methamphetamine-derived AGEs are generated at experimentally detectable and pathologically relevant levels.

The AGEs formed in disease states such as macular degeneration and diabetes mellitus have been linked to autoantibody titers against the glycosylated proteins, and indeed, the process of attaching nonimmunogenic chemical moieties to a carrier protein is reminiscent of hapten preparation in a research setting for vaccination.^{27,42} Before chronic methamphetamine abuse could be linked to the appearance of AGE-related pathologies, the generation of methamphetamine-modified AGEs *in vivo* was illustrated through the detection of a polyclonal response against methamphetamine glycosylated serum albumin. Protein glycation by methamphetamine was conducted *in vitro* through the incubation of methamphetamine with glucose at physiological conditions; the Amadori rearrangement product was detected after 12 h by LC/MS. Addition of BSA to the starting reaction caused the methamphetamine-derived albumin glycation products to form after 2 weeks, as assessed by ELISA and dot blot. Immunization of




FIGURE 4. Quantification of ELISA results for presence of autoantibodies against methamphetamine-glycated MSA in serum samples of rats self-administering methamphetamine. Bars represent ratio of ELISA signal above negative control background. * denotes $p < 0.05$; ** denotes $p < 0.0002$; comparison to drug naïve. Reproduced with permission from ref 45. Copyright 2007 National Academy of Sciences, U.S.A.

mice with experimentally prepared methamphetamine-glycated mouse serum albumin (MSA) elicited serum antibodies against methamphetamine-conjugated AGEs (METH-AGEs) that were detectable via ELISA. To verify that this response was indeed specific to the METH-AGEs and distinct from any potentially immunogenic effect of MSA-AGE injected in mice, competition studies with MSA and MSA-AGE were employed to show that serum antibodies failed to recognize these substrates. Furthermore, the polyclonal sera measurably bound METH-AGEs alone ($K_{d,app} \approx 10$ nM), suggesting that continued immune system priming with methamphetamine and the sustained production of METH-AGE would serve to augment the polyclonal response as well as initiate an immunoinactivated or inflammatory state.

To investigate the potential significance of these results with respect to methamphetamine addiction in humans, a chronic schedule of methamphetamine self-administration was implemented in a rat model.^{43,44} Two groups of methamphetamine-addicted rats were allowed to self-administer methamphetamine for approximately 3 months in daily drug sessions lasting either 1 h (short access rats, ShA) or 6 h (long access rats, LgA). Gratifyingly, the level of drug intake by the different groups directly correlated with the serum antibody titers against methamphetamine glycosylated proteins (Figure 4). Moreover, comparison of ELISA signals between LgA, ShA, and drug naïve (DN) serum samples showed that there was no statistically significant difference in serum antibody binding to either serum albumin or methamphetamine-unmodified glycosylated protein ELISA substrates. The methamphetamine dose dependence and relative specificity of the LgA and ShA serum antibodies confirmed that the polyclonal response was attributable to methamphetamine self-administration and subsequent methamphetamine-mediated glycation.

The magnitude of the polyclonal response measured in LgA and ShA rats is suggestive of the *in vivo* significance of methamphetamine-mediated glycation. Namely, previous work into the neuronal and behavioral repercussions of methamphetamine in rodent models have relied on acute binge doses of drug, oftentimes experimenter-delivered, to evoke the measured phenotype.⁴⁰ Recently, such studies have attracted scrutiny as scientists have realized the value of translational research in providing insight into the human condition.⁴⁶ Our dosing regimen, in which LgA rats administered an average drug dose of 7 mg/kg per 6-h session over a prolonged period of time, was designed to approximate the chronic abuse and level of methamphetamine intake documented for human addicts.⁴⁶ While the 6-h drug exposure coupled with the half-life of methamphetamine in rats ($t_{1/2} \approx 1$ h) translates to continual immune challenge by methamphetamine relative to 1-h ShA sessions, the low methamphetamine intake of ShA rats was still sufficient to elicit measurable antibody titers, a testament to both the repercussions of more recreational methamphetamine abuse and the relevance of METH-AGE immunogenicity.

Immunomodulation By AGEs

Previous work has elucidated the immunosuppressive properties of methamphetamine, which is consistent with many abused drugs, while research into the inflammation or immunoinactivation caused by methamphetamine focuses on microglialosis, blood–brain barrier disruption, and neurotoxicity from oxidative stress and inflammatory gene upregulation.^{47,48} We speculated that METH-AGE production would spur a constellation of effects including the corresponding polyclonal response against METH-AGE, METH-AGE deposition on vasculature, and receptor for AGE (RAGE) activation and that these effects would induce altered cytokine expression levels in the periphery. With respect to the latter effect, the receptor for AGE (RAGE), a member of the immunoglobulin superfamily expressed on most cell types, precipitates the expression of inflammatory and prothrombotic genes upon AGE ligation.^{26,28} Whereas the impact of RAGE and inflammatory responses have been studied within the context of the chronic vascular disease seen in other disease states (e.g., diabetes, Alzheimer's disease, macular degeneration, nephropathy, retinopathy), this AGE–RAGE hypothesis of immunomodulation and vascular disease has not been investigated as a causative factor of human and nonhuman methamphetamine abuse pathologies.^{49,50}

To attribute discrete physiological consequences to a chemical mechanism, namely, methamphetamine-mediated glyca-




FIGURE 5. Modulation of relevant cytokine and chemokine levels in response to methamphetamine intake. Sera samples analyzed were obtained from ShA rats (gray) and LgA rats (black) self-administering methamphetamine 1 h daily and 6 h daily, respectively, for a period 87 days. Data is expressed as mean fold-increase over drug naïve levels \pm SEM: * denotes $p < 0.05$; ** denotes $p < 0.01$; significance in change compared with drug naïve levels. † denotes $P < 0.05$, H_0 , LgA > ShA. Reproduced with permission from ref 45. Copyright 2007 National Academy of Sciences, U.S.A.

tion, immunomodulation was assessed through measuring the levels of several major cytokines and chemokines in sera from DN, ShA, and LgA rats (Figure 5).⁴⁵ Both tumor necrosis factor- α (TNF- α) and interleukin-1 β (IL-1 β), two major proinflammatory cytokines that mediate macrophage activation and vascular permeance, were increased above basal levels in LgA rats. In particular, the substantial 5-fold elevation of TNF- α aligns with its known roles in promoting antibody production as well as RAGE expression.⁵¹ Methamphetamine self-administration resulted in the dose-dependent upregulation of other cytokine molecules specifically related to AGE exposure. Most notably, the level of vascular endothelial growth factor (VEGF) exhibited a 2-fold increase in ShA sera and a 6-fold increase in LgA sera compared with DN sera, an effect not previously reported alongside methamphetamine abuse. VEGF, expression of which is stimulated by AGE production, has been demonstrated to mediate many of the angiogenic activities of AGEs and to play a major role in the development of diabetic retinopathy and nephropathy.⁵²

In conjunction with the above major cytokines, the levels of several chemokines were also altered, an indication of the magnitude of the proinflammatory response initiated by chronic drug-taking behavior. This finding mirrors the release of MCP-1 upon AGE exposure and RAGE activation.⁵³ Other chemokines displayed a biphasic behavior with regard to drug dose, a pattern already observed in the varying expression of several cytokines. In sum, the fluctuation in both cytokine and

chemokine levels was sensitive to the duration of methamphetamine self-administration, akin to the variance in antibody titers.

Conclusion

Our research into the chemical reactivity of exogenous drug molecules highlights how the formation of drug-glycated proteins *in vivo* becomes a marker of a developing disease pathology. Whereas both endogenous and drug-modified AGEs have been shown to serve as immunological epitopes *in vivo*, the accumulation of drug-modified AGEs gains additional relevance to the chronic drug addict in its ability to vaccinate the addict against the unmodified drug molecule.

Though the mechanism behind the pathogenesis of certain diseases observed in addicts is undoubtedly multifaceted, the correlation between the generation of AGEs, the polyclonal response to AGEs, and the development of a detectable pathology illustrates the value in turning to chemistry for mechanistic explanations of biological phenomena. Despite potential kinetic and thermodynamic hurdles to the formation of drug-AGEs such as drug availability and conjugation of AGEs to long-lived proteins, there is strong evidence that drug-mediated glycation processes are operable and relevant *in vivo*. Thus, future work will strive to clarify the causative link between AGE formation and the downstream immunomodulatory and pathological effects. A better understanding of the pharmacodynamics of protein glycation when mediated by exogenous amine-containing compounds is necessary in order for this AGE hypothesis to influence the therapeutic treatment of pathologies attributed to AGEs.

We gratefully acknowledge the financial support of the Skaggs Institute for Chemical Biology and the National Institute on Drug Abuse (Grant DA 21939 to J.B.T.; Grant DA 98590 to K.D.J.).

Note Added after ASAP. This paper was posted to the web on March 10, 2009 with an error in Figure 1. The revised version was posted on March 20, 2009.

BIOGRAPHICAL INFORMATION

Jennifer B. Treweek received her B.S. degree in chemistry from Caltech (2004) and conducted research with Prof. Richard Roberts. In 2005, she joined the laboratory of Professor Janda as a graduate student at TSRI where she received a Pfeiffer Research Foundation Grant (2006) and an NIH NRSA Predoctoral Fellowship (2007). Her current research is focused on neuroscience and immunopharmacotherapy.

Tobin J. Dickerson received his B.S. degree in chemistry from the University of Virginia (1999) and Ph.D. from The Scripps

Research Institute (2004). In 2005, he was promoted to Assistant Professor in the Department of Chemistry at TSRI and WIRM. His research interests include the chemical reactivity of drugs of abuse, the interplay between small molecules and the immune system, approaches for the treatment of filarial infections, and technologies that modulate BoNT toxicity.

Kim D. Janda obtained a B.S. degree from the University of South Florida (1980) and a Ph.D. (1984) from the University of Arizona. He is currently a Professor in the Chemistry and Immunology Departments at TSRI, where he is also a Skaggs Scholar, the Ely R. Callaway, Jr. Chair, and Director of WIRM. Among his awards are the Alfred P. Sloan Fellow (1993) and the Arthur Cope Scholar (1999). He was a founder of CombiChem and Drug Abuse Sciences.

FOOTNOTES

*To whom correspondence should be addressed. E-mail: kdjanda@scripps.edu. Mailing address: The Scripps Research Institute, BCC-582, 10550 North Torrey Pines Road, La Jolla, CA 92037. Telephone: (858) 784-2516. Fax: (858) 784-2595.

REFERENCES

- Davis, V. E.; Walsh, M. J.; Yamanaka, Y. Augmentation of alkaloid formation from dopamine by alcohol and acetaldehyde *in vitro*. *J. Pharmacol. Exp. Ther.* **1970**, *174*, 401–412.
- Wong, C.-H.; Whitesides, G. M. *Enzymes in Synthetic Organic Chemistry*; Pergamon: Oxford, U.K., 1994.
- Fessner, W.-D.; Walter, C. *Top. Curr. Chem.* **1996**, *184*, 98–183.
- List, B.; Lerner, R. A.; Barbas, C. F., III. Proline-catalyzed direct asymmetric aldol reactions. *J. Am. Chem. Soc.* **2000**, *122*, 2395–2396.
- Mukherjee, S.; Yang, J. W.; Hoffmann, S.; List, B. Asymmetric enamine catalysis. *Chem. Rev.* **2007**, *107*, 5471–5569.
- List, B. Biocatalysis and organocatalysis: Asymmetric synthesis inspired by nature. *Asymmetric Synth.* **2007**, 161–165.
- Dickerson, T. J.; Janda, K. D. Aqueous aldol catalysis by a nicotine metabolite. *J. Am. Chem. Soc.* **2002**, *124*, 3220–3221.
- Chu, F. L.; Yaylayan, V. A. FTIR monitoring of oxazolidinone formation and decomposition in a glycoaldehyde-phenylalanine model system by isotope labeling techniques. *Carbohydr. Res.* 2008, in press.
- Maillard, L. C.; Gautier, M. Action des acides amines sur les sucres: formation des melanoidines par voie methodique. *C. R. Seances Acad. Sci. III* **1912**, *154*, 66–68.
- Brownlee, M.; Cerami, A.; Vlassara, H. Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications. *N. Engl. J. Med.* **1988**, *318*, 1315–1321.
- Monnier, V. M.; Cerami, A. Nonenzymatic browning in vivo: Possible process for aging of long-lived proteins. *Science* **1981**, *211*, 491–493.
- Vlassara, H.; Brownlee, M.; Manogue, K. R.; Dinarello, C. A.; Pasagian, A. Cachectin/TNF and IL-1 induced by glucose-modified proteins: role in normal tissue remodeling. *Science* **1988**, *240*, 1546–1548.
- Monnier, V. M.; Kohn, R. R.; Cerami, A. Accelerated age-related browning of human collagen in diabetes mellitus. *Proc. Natl. Acad. Sci. U.S.A.* **1984**, *91*, 583–587.
- Stopper, H.; Schinzel, R.; Sebekova, K.; Heidland, A. Genotoxicity of advanced glycation end products in mammalian cells. *Cancer Lett.* **2003**, *190*, 151–160.
- Vitek, M. P.; Bhattacharya, K.; Glendening, J. M.; Stopa, E.; Vlassara, H.; Bucala, R.; Manogue, K.; Cerami, A. Advanced glycation end products contribute to amyloidosis in Alzheimer disease. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 4766–4770.
- Ishibashi, T.; Murata, T.; Hangai, M.; Nagai, R.; Horiuchi, S.; Lopez, P. F.; Hinton, D. R.; Ryan, S. J. Advanced glycation end products in age-related macular degeneration. *Arch. Ophthalmol.* **1998**, *116*, 1629–1632.
- Bierhaus, A.; Hofmann, M. A.; Ziegler, R.; Nawroth, P. P. AGE and their interaction with AGE-receptors in vascular disease and diabetes. I. The AGE concept. *Cardiovasc. Res.* **1998**, *37*, 586–600.
- Lo, T. W.; Westwood, M. E.; McLellan, A. C.; Selwood, T.; Thornalley, P. J. Binding and modification of proteins by methylglyoxal under physiological conditions. A kinetic and mechanistic study with N alpha-acetylarginine, N alpha-acetylcysteine, and N alpha-acetyllysine, and bovine serum albumin. *J. Biol. Chem.* **1994**, *269*, 32299–32305.
- Wells-Knecht, K. J.; Zyzak, D. V.; Litchfield, J. E.; Thorpe, S. R.; Baynes, J. W.

- Identification of glyoxal and arabinose as intermediates in the autoxidative modification of proteins by glucose. *Biochemistry* **1995**, *34*, 3702–3709.
- Singh, R.; Barden, A.; Mori, T.; Beilin, L. Advanced glycation end-products: a review. *Diabetologia* **2001**, *44*, 129–146.
- Monnier, V. M.; Sell, D. R.; Dai, Z.; Nemet, I.; Collard, F.; Zhang, J. The role of the Amadori product in the complications of diabetes. *Ann. N.Y. Acad. Sci.* **2008**, *1126*, 81–88.
- Khalifah, R. G.; Todd, P.; Booth, A. A.; Yang, S. X.; Mott, J. D.; Hudson, B. G. Kinetics of nonenzymatic glycation of ribonuclease A leading to advanced glycation end products. Paradoxical inhibition by ribose leads to facile isolation of protein intermediate for rapid post-Amadori studies. *Biochemistry* **1996**, *35*, 4645–4654.
- Bunn, H. F.; Higgins, P. J. Reaction of monosaccharides with proteins: Possible evolutionary significance. *Science* **1981**, *213*, 222–224.
- Booth, A. A.; Khalifah, R. G.; Todd, P.; Hudson, B. G. *In vitro* kinetic studies of formation of antigenic advanced glycation end products (AGEs). Novel inhibition of post-Amadori glycation pathways. *J. Biol. Chem.* **1997**, *272*, 5430–5437.
- McPherson, J. D.; H., S. B.; Walton, D. J. Role of fructose in glycation and cross-linking of proteins. *Biochemistry* **1988**, *27*, 1901–1907.
- Yan, S. D.; Schmidt, A. M.; Anderson, G. M.; Zhang, J.; Brett, J.; Zou, Y. S.; Pinsky, D.; Stern, D. Enhanced cellular oxidant stress by the interaction of advanced glycation end products with their receptors/binding proteins. *J. Biol. Chem.* **1994**, *269*, 9889–9897.
- Turk, Z.; Ljubic, S.; Turk, N.; Benko, B. Detection of autoantibodies against advanced glycation endproducts and AGE-immune complexes in serum of patients with diabetes mellitus. *Clin. Chim. Acta* **2001**, *303*, 105–115.
- Kislinger, T.; Fu, C.; Huber, B.; Qu, W.; Taguchi, A.; Du Yan, S.; Hofmann, M.; Yan, S. F.; Pischetsrieder, M.; Stern, D.; Schmidt, A. M. N(epsilon)-(carboxymethyl)lysine adducts of proteins are ligands for receptor for advanced glycation end products that activate cell signaling pathways and modulate gene expression. *J. Biol. Chem.* **1999**, *274*, 31740–31749.
- Cerami, C.; Founds, H.; Nicholl, I.; Mitsuhashi, T.; Giordano, D.; Vanpatten, S.; Lee, A.; Al-Abed, Y.; Vlassara, H.; Bucala, R.; Cerami, A. Tobacco smoke is a source of toxic reactive glycation products. *Proc. Natl. Acad. Sci. U.S.A.* **1997**, *94*, 13915–13920.
- Dickerson, T. J.; Janda, K. D. A previously undescribed chemical link between smoking and metabolic disease. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 15084–15088.
- Isomura, S.; Wirsching, P.; Janda, K. D. An immunotherapeutic program for the treatment of nicotine addiction: Hapten design and synthesis. *J. Org. Chem.* **2001**, *66*, 4115–4121.
- Kyerematen, G. A.; Morgan, M.; Chattopadhyay, B.; DeBethizy, J. D.; Vesell, E. S. Disposition of nicotine and eight metabolites in smokers and nonsmokers: Identification in smokers of two metabolites that are longer lived than cotinine. *Clin. Pharmacol. Ther.* **1998**, *48*, 641–651.
- Salomon, A. R.; Marciniowski, K. J.; Friedland, R. P.; Zagorski, M. G. Nicotine inhibits amyloid formation by the beta-peptide. *Biochemistry* **1996**, *35*, 13568–13578.
- Sisodia, S. S.; Price, D. J. Role of the beta-amyloid protein in Alzheimer's disease. *FASEB J.* **1995**, *9*, 366–370.
- Zeng, H.; Zhang, Y.; Peng, L.; Shao, H.; Menon, N. K.; Yang, J.; Salomon, A. R.; Freidland, R. P.; Zagorski, M. G. Nicotine and amyloid formation. *Biol. Psychiatry* **2001**, *49*, 248–257.
- Tjernberg, L. O.; Näslund, J.; Lindqvist, F.; Johansson, J.; Karlström, A. R.; Thyberg, J.; Terenius, L.; Nordstedt, C. Arrest of β -amyloid fibril formation by a pentapeptide ligand. *J. Biol. Chem.* **1996**, *271*, 8545–8548.
- Dickerson, T. J.; Janda, K. D. Glycation of the amyloid β -protein by a nicotine metabolite: A fortuitous chemical dynamic between smoking and Alzheimer's disease. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 8182–8187.
- Barr, A. M.; Panenka, W. J.; MacEwan, G. W.; Thornton, A. E.; Lang, D. J.; Honer, W. G.; Lecomte, T. The need for speed: An update on methamphetamine addiction. *J. Psychiatry Neurosci.* **2006**, *31*, 301–313.
- Saunders, C.; Ferrer, J. V.; Shi, L.; Chen, J.; G., M.; Lamb, M. E.; Leeb-Lundberg, L. M.; Carvelli, L.; Javitch, J. A.; Galli, A. Amphetamine-induced loss of human dopamine transporter activity: An internalization-dependent and cocaine-sensitive mechanism. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 6850–6855.
- Itzhak, Y.; Achat-Mendes, C. Methamphetamine and MDMA (ecstasy) neurotoxicity: "Of mice and men". *IUBMB Life* **2004**, *56*, 249–255.
- Melega, W. P.; Jorgensen, M. J.; acutean, G. L.; Way, B. M.; Pham, J.; Morton, G.; Cho, A. K.; Fairbanks, L. A. Long-term methamphetamine administration in the Vervet monkey models aspects of a human exposure: Brain neurotoxicity and behavioral profiles. *Neuropsychopharmacology* **2008**, *33*, 1441–1452.
- Gu, X.; Meer, S. G.; Miyagi, M.; Rayborn, M. E.; Hollyfield, J. G.; Crabb, J. W.;

- Salomon, R. G. Carboxyethylpyrrole protein adducts and autoantibodies, biomarkers for age-related macular degeneration. *J. Biol. Chem.* **2003**, *278*, 42027–42035.
- 43 Kitamura, O.; Wee, S.; Specio, S. E.; Koob, G. F.; Pulvirenti, L. Escalation of methamphetamine self-administration in rats: A dose-effect function. *Psychopharmacology (Berl)*. **2006**, *186*, 48–53.
- 44 Wee, S.; Wang, Z.; Woolverton, W. L.; Pulvirenti, L.; Koob, G. F. Effect of aripiprazole, a partial dopamine D2 receptor agonist, on increased rate of methamphetamine self-administration in rats with prolonged session duration. *Neuropsychopharmacology* **2007**, *32*, 2238–2247.
- 45 Treweek, J.; Wee, S.; Koob, G. F.; Dickerson, T. J.; Janda, K. D. Self-vaccination by methamphetamine glycation products chemically links chronic drug abuse and cardiovascular disease. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, *104*, 11580–11584.
- 46 Segal, D. S.; Kuczenski, R.; O'Neil, M. L.; Melega, W. P.; Cho, A. K. Escalating dose methamphetamine pretreatment alters the behavioral and neurochemical profiles associated with exposure to a high-dose methamphetamine binge. *Neuropsychopharmacology* **2003**, *28*, 1730–1740.
- 47 Asanuma, M.; Miyazaki, I.; Higashi, Y.; Tsuji, T.; Ogawa, N. Specific gene expression and possible involvement of inflammation in methamphetamine-induced neurotoxicity. *Ann. N.Y. Acad. Sci.* **2004**, *1025*, 69–75.
- 48 In, S.-W.; Son, E.-W.; Rhee, D.-K.; Pyo, S. Methamphetamine administration produces immunomodulation in mice. *J. Toxicol. Environ. Health A*. **2005**, *68*, 2133–2145.
- 49 He, S. Y.; Matoba, R.; Fujitani, N.; Sodesaki, K.; Onishi, S. Cardiac muscle lesions associated with chronic administration of methamphetamine in rats. *Am. J. Forensic Med. Pathol.* **1996**, *17*, 155–62.
- 50 Schaiberger, P. H.; Kennedy, T. C.; Miller, F. C.; Gal, J.; Petty, T. L. Pulmonary hypertension associated with long-term inhalation of "crank" methamphetamine. *Chest* **1993**, *104*, 614–616.
- 51 Tanaka, N.; Yonekura, H.; Yamagishi, S.; Fujimori, H.; Yamamoto, Y.; Yamamoto, H. The receptor for advanced glycosylation end products is induced by the glycation product themselves and TNF-alpha through nuclear factor kappa B and by 17-beta-estradiol through Sp1 in human vascular endothelial cells. *J. Biol. Chem.* **2000**, *275*, 25781–25790.
- 52 Treins, C.; Giorgetti-Peraldi, S.; Murdaca, J.; Van Obberghen, E. Regulation of vascular endothelial growth factor expression by advanced glycation end products. *J. Biol. Chem.* **2001**, *276*, 43836–43841.
- 53 Yamagishi, S.-i.; Inagaki, Y.; Okamoto, T.; Amano, S.; Koga, K.; Takeuchi, M.; Makita, Z. Advanced glycation end product-induced apoptosis and overexpression of vascular endothelial growth factor and monocyte chemoattractant protein-1 in human-cultured mesangial cells. *J. Biol. Chem.* **2002**, *277*, 20309–20315.