

Table 3. Enthalpy changes accompanying the formation of adducts between SbCl_5 and given donors in ethylene chloride solution. ΔH -values are expressed in kcal per mole.

Donor	$-\Delta H_1$	$-\Delta H_2$	$-\Delta H$
$\text{Me}_2\text{N}(\text{Me})\text{CO}$	28.71	0.91	27.80 ± 0.08
$(\text{Me}_2\text{N})_2\text{CO}$	30.69	1.05	29.64 ± 0.03

Tetramethylurea, b.p. 64.5° at 13 mm Hg, n_D^{25} 1.4493, d_4^{25} 0.9619. (Found: C 51.7; H 10.4. Calc. for $\text{C}_5\text{H}_{12}\text{N}_2\text{O}$: C 51.7; H: 10.4).

Units of measurements. The results of the calorimetric experiments are expressed in terms of the defined calorie equal to 4.1840 abs. joules, and refer to the isothermal process at 25° and to the true mass. The molecular weights were computed from the 1961 table of atomic weights.⁵

Results. The experimental results are summarized in Tables 1–3. Two different systems were used, I and II, with heat equivalents 3996 ± 4 and 3302 ± 2 cal per unit of $\log R_i/R_f$, respectively. In the heat of reaction measurements (Table 1) the first column gives the name of the donor together with a sum correction, q , compensating for a number of small heat effects.² Columns 2 and 3 give the amounts of donor and acceptor. The SbCl_5 was added in excess prior to the experiment to take care of traces of water inside the calorimeter and in the solvent. Column 4 gives the corrected temperature rise expressed in units of $10^4 \log R_i/R_f$, where R_i and R_f are the extrapolated initial and final thermistor resistances. The last column gives the molar enthalpy change. The heat of solution experiments (Table 2) were performed in pure ethylene chloride. The data obtained are given in Table 3 together with the calculated values of the enthalpy change for the formation of the adducts in solution.

Acknowledgement. A grant from the Faculty of Mathematics and Natural Sciences, University of Uppsala, is gratefully acknowledged.

- Zackrisson, M. *Acta Chem. Scand.* **15** (1961) 1784.
- Olofsson, G., Lindqvist, I. and Sunner, S. *Acta Chem. Scand.* **17** (1963) 259.
- Olofsson, G. *Acta Chem. Scand.* **18** (1964) 11.

- Lindqvist, I. *Inorganic Adduct Molecules of Oxo-compounds* Springer-Verlag, Berlin 1963.
- Cameron, A. E. and Wichers, E. *J. Am. Chem. Soc.* **84** (1962) 4175.

Received May 5, 1964.

Spectroscopic Studies on Metal Carbonyls

II. Generalized Mean-Square Amplitudes of Chromium and Molybdenum Hexacarbonyls

S. J. CYVIN and J. BRUNVOLL

Institute of Theoretical Chemistry, Technical University of Norway, Trondheim, Norway

In this work the mean-square parallel and perpendicular amplitudes, and mean cross products¹ of $\text{Cr}(\text{CO})_6$ and $\text{Mo}(\text{CO})_6$ have been calculated. Because of the high symmetry of the appropriate model (O_h), all mean cross products vanish except for one type of distances. Numerical results are given in Table 1, where the atom pairs, which define the types of distances, refer to the numbering previously adopted.² As to the orientation of the perpendicular amplitudes, all Δy have been referred to the XZ plane, except for the 1–8 and 7–8 type distances, where the appropriate plane is 1–4–6. These quantities as referred to somewhat arbitrary orientations, are not of much interest. But so are the quantities obtained as the sum $\langle \Delta x \rangle^2 + \langle \Delta y \rangle^2$, which is invariant with respect to the individual orientations. This sum enters

Table 1. Mean-square parallel and perpendicular amplitudes (\AA^2) for chromium and molybdenum hexacarbonyls*

Atom pair	Parallel $\langle \Delta z^2 \rangle$	Perpendicular $\langle \Delta x^2 \rangle$	Perpendicular $\langle \Delta y^2 \rangle$
Chromium hexacarbonyl			
7-13 (Cr-C)	(0.0024) 0.0031	(0.0050) 0.0150	(0.0050) 0.0150
1-7 (C-O)	(0.0012) 0.0012	(0.0057) 0.0122	(0.0057) 0.0122
1-13 (Cr.C.O)**	(0.0024) 0.0033	(0.0092) 0.0401	(0.0092) 0.0401
1-10 (CCrCO)**	(0.0038) 0.0050	(0.0092) 0.0349	(0.0092) 0.0349
7-10 (C.Cr.C)**	(0.0037) 0.0048	(0.0069) 0.0182	(0.0069) 0.0182
1-4 (OCCrCO)**	(0.0038) 0.0051	(0.0103) 0.0454	(0.0103) 0.0454
7-8 (C...C)	(0.0075) 0.0224	(0.0062) 0.0158	(0.0088) 0.0291
1-2 (O...O)	(0.0133) 0.0563	(0.0087) 0.0342	(0.0161) 0.0789
1-8 (C...O)	(0.0092) 0.0321	(0.0086) 0.0323	(0.0126) 0.0540
Molybdenum hexacarbonyl			
7-13 (Mo-C)	(0.0025) 0.0044	(0.0054) 0.0189	(0.0054) 0.0198
1-7 (C-O)	(0.0012) 0.0012	(0.0061) 0.0137	(0.0061) 0.0137
1-13 (Mo.C.O)**	(0.0026) 0.0046	(0.0097) 0.0501	(0.0097) 0.0501
1-10 (CMoCO)**	(0.0041) 0.0056	(0.0102) 0.0435	(0.0102) 0.0435
1-10 (C.Mo.C)**	(0.0040) 0.0055	(0.0078) 0.0238	(0.0078) 0.0238
7-4 (OCMoCO)**	(0.0041) 0.0058	(0.0114) 0.0561	(0.0114) 0.0561
7-8 (C...C)	(0.0092) 0.0339	(0.0076) 0.0249	(0.0102) 0.0393
1-2 (O...O)	(0.0154) 0.0764	(0.0103) 0.0491	(0.0181) 0.0999
1-8 (C...O)	(0.0111) 0.0465	(0.0101) 0.0465	(0.0142) 0.0696

* Values at 298°K, and (in parentheses) at absolute zero temperature.

** Atoms on straight line.

into the calculation of Bastiansen—Morino shrinkage effects.³

The only non-vanishing mean cross product is $\langle \Delta z \Delta x \rangle$ for the 1-8 type distance, viz. -0.0024 and -0.0027 \AA^2 at $T = 0$, and -0.0143 and -0.0184 \AA^2 at $T = 298 \text{ °K}$ for $\text{Cr}(\text{CO})_6$ and $\text{Mo}(\text{CO})_6$, respectively.

1. Morino, Y. and Hirota, E. *J. Chem. Phys.* **23** (1955) 737.

2. Brunvoll, J. and Cyvin, S. J. Part I of this series (to be published); see also: Pistorius, C. W. F. T. and Haarhoff, P. C. *J. Mol. Spectry.* **3** (1959) 671.

3. Morino, Y., Cyvin S. J., Kuchitsu, K. and Iijima, T. *J. Chem. Phys.* **36** (1962) 1109; Cyvin, S. J. *Tidsskr. Kjemi, Bergvesen Met.* **22** (1962) 44, 74, and references cited therein.

Received April 14, 1964,