






as do other reactive nitrile oxides. 3-Alkoxyisoxazolines **4**, which are masked  $\beta$ -hydroxy esters, are available by cycloaddition of dihaloformaldehyde oxime to alkenes and subsequent dehaloalkoxylation at C-3<sup>3</sup> [eqn. (3)]. It is thus possible to shorten the routes to **4** and **5** by one



step. The exchange of Cl for OR at C-3 in the aromatic isoxazoles required drastic conditions with use of strong bases.

## Results and discussion

*Alkyl formhydroximates.* A simplified procedure for the preparation of alkyl formhydroximates was required. It has been reported that treatment of thioesters with hydroxylamine give alkyl hydroximates in good yields.<sup>4a</sup> However, it was also reported that some aromatic thioesters give thiohydroxamic acids.<sup>4b</sup> We found that the thioformates, obtained by acid<sup>5</sup> or  $\text{FeCl}_3$ <sup>6</sup> catalyzed thionation of orthoesters<sup>7</sup> with hydrogen sulfide, reacted with hydroxylamine to give **2a,b,d,e** in modest to good yields [eqn. (4)]. The reaction conditions are critical



**4a**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{CH}_2\text{NHCOO-t-Bu}$

**4b**  $\text{R}^1 = \text{CH}_2\text{C}_6\text{H}_5$ ;  $\text{R}^2 = \text{COOC}_2\text{H}_5$

**4c**  $\text{R}^1 = \text{CH}_2\text{C}_6\text{H}_5$ ;  $\text{R}^2 = \text{C}_6\text{H}_5$

**4d**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{COOC}_2\text{H}_5$

**4e**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = n\text{-Bu}$

**4f**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{CH(OC}_2\text{H}_5)_2$


**4g**  $\text{R}^1 = \text{CH}_2\text{C}_6\text{H}_5$ ;  $\text{R}^2 = \text{CH}_2\text{NHCOO-t-Bu}$

**4h**  $\text{R}^1 = \text{C}_2\text{H}_5$ ;  $\text{R}^2 = \text{COOC}_2\text{H}_5$

**4i**  $\text{R}^1 = \text{C}_2\text{H}_5$ ;  $\text{R}^2 = \text{Bu}$

**4j**  $\text{R}^1 = \text{C}_2\text{H}_5$ ;  $\text{R}^2 = \text{COCH}_3$

**4k**  $\text{R}^1 = \text{C}_2\text{H}_5$ ;  $\text{R}^2 = \text{COC}_6\text{H}_5$



**5a**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{R}^3 = \text{COOCH}_3$

**5b**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{R}^3 = \text{C}_6\text{H}_5$

**5c**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{C}_6\text{H}_5$ ;  $\text{R}^3 = \text{H}$


**5d**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{Bu}_3\text{Sn}$ ;  $\text{R}^3 = \text{H}$

**5e**  $\text{R}^1 = \text{CH}_2\text{C}_6\text{H}_5$ ;  $\text{R}^2 = \text{C}_6\text{H}_5$ ;  $\text{R}^3 = \text{H}$


**5f**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = (\text{CH}_3)_3\text{Si}$ ;  $\text{R}^3 = \text{H}$

**5g**  $\text{R}^1 = \text{C}_2\text{H}_5$ ;  $\text{R}^2 = \text{C}_6\text{H}_5$ ;  $\text{R}^3 = \text{H}$


**5h**  $\text{R}^1 = \text{CH}_2\text{C}_6\text{H}_5$ ;  $\text{R}^2 = \text{Bu}_3\text{Sn}$ ;  $\text{R}^3 = \text{H}$




is a naturally occurring antibiotic.<sup>8</sup> Commercially available dimethylaminothioformamide was heated under reflux with hydroxylamine in aqueous ethanol to give *N,N*-dimethylamino-*N*-hydroxyformamidine (dimethylaminoformamide oxime), **8**.<sup>10</sup> Chlorination of **8** with NCS and subsequent dehydrochlorination in the presence of ethyl acrylate did not give the desired isoxazoline **9**, [eqn. (6)]. No further reactions were performed with **8**.




*3-Alkoxyisoxazolines 4 and 3-alkoxyisoxazoles 5.* The chlorination of alkyl formhydroximates **2b,d,e** was conveniently carried out with NCS. Use of *tert*-butyl hypochlorite did not offer any advantages. The intermediate blue-green nitroso compounds rearranged to the oximes and the end of the chlorination was determined by the disappearance of the blue-green colour. Potassium hydrogen carbonate was the most suitable base for the dehydrochlorination of **3**. It slowly released the nitrile oxides, preventing their dimerization to furoxans and improved the yield of the cyclization products **4**. Occasionally triethylamine was used as base. The furoxan **10**



**10**



**9**




**11a**  $\text{R}^1 = \text{C}_2\text{H}_5$ ;  $\text{R}^2 = \text{CH}_3$ ;  $\text{R}^3 = \text{CH(OC}_2\text{H}_5)_2$

**11b**  $\text{R}^1 = \text{C}_2\text{H}_5$ ;  $\text{R}^2 = \text{CH(OC}_2\text{H}_5)_2$ ;  $\text{R}^3 = \text{CH}_3$

**11c**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{CH}_3$ ;  $\text{R}^3 = \text{COOCH}_3$

**11d**  $\text{R}^1 = i\text{-Pr}$ ;  $\text{R}^2 = \text{COOCH}_3$ ;  $\text{R}^3 = \text{CH}_3$



**12a**

for the outcome of the reaction. Compounds **2a-d** are crystalline solids, which decompose and liquify within a few days at 25°C but are stable for several months in the freezer. Then benzyl derivative **2e** is stable at room temperature. We were not successful in transforming triphenyl orthoformate<sup>7</sup> into **2f**.

*Attempted preparation of C-alkoxynitrones and dimethylaminonitrile oxide. Preparation of thioformin, 6.*<sup>8</sup> The reaction of thioformates with *N*-substituted hydroxylamines took a different course. The corresponding *N*-hydroxythioformamides<sup>9</sup> were obtained. The desired nitrone **7** was not observed, [eqn. (5)]. Compound **6**

was isolated as a crystalline solid in the absence of a substrate. The isoxazolines **4** or isoxazoles **5** were obtained most conveniently in a one-pot reaction where **2**, NCS, potassium hydrogen carbonate and the olefin or acetylene were mixed in ethyl acetate as the solvent and reacted at ca. 48°C for 24 h, Table 1. Monosubstituted olefins and acetylenes gave practically only the 5-substituted

Table 1. Synthesis of 3-alkoxyisoxazolines and 3-alkoxyisoxazoles.

Compound No.	Ratio oxime: alkene or alkyne	Base, Temp., Reaction time	Yield (%)	M.p./°C or (b.p./mmHg)	Eluent TLC, silica	Alkene or alkyne	Selected <sup>1</sup> H, <sup>13</sup> C NMR, δ (CDCl <sub>3</sub> ); IR ν <sub>max</sub> /cm <sup>-1</sup> ; UV, λ <sub>max</sub> /nm; MS, m/z data.
4a	1 : 0.6	KHCO <sub>3</sub> , 25°C, 70 h	85	Oil	CHCl <sub>3</sub> -CH <sub>3</sub> OH 93 : 7	N-Boc-allyl-amine	<sup>1</sup> H: 1.27 (6 H, d, J = 6 Hz), 1.39 (9 H, s), 2.67 (1 H, dd, J = 8.5 and 17.5 Hz), 2.91 (1 H, dd, J = 10.0 and 17.5 Hz), 3.3 (2 H, m), 4.60 (1 H, m), 4.73 (1 H, septet, J = 6 Hz), 4.93 (1 H, br s). <sup>13</sup> C: 21.9, 28.5, 36.0, 43.6, 74.1, 79.7, 79.9, 156.7, 166.8. IR (film): 1720, 1628. MS: 258 (M <sup>+</sup> ), 203, 185, 160, 143, 128, 110.
4b	1 : 1.2	KHCO <sub>3</sub> , 25°C, 48 h KHCO <sub>3</sub> , 48°C, 38 h Et <sub>3</sub> N, 48°C, 1 h	51 64 49	Oil	CHCl <sub>3</sub>	Ethyl acrylate	<sup>1</sup> H: 1.27 (3 H, t, J = 7.0 Hz), 3.23 (2 H, m), 4.22 (2 H, q, J = 7.0 Hz), 5.0 (1 H, dd, J = 9 and 10 Hz), 5.12 (2 H, s), 7.34 (5 H, s). <sup>13</sup> C: 14.3, 36.4, 62.2, 72.8, 78.7, 128.7, 128.9, 129.1, 135.8, 166.8, 170.6. IR (film): 1760, 1640. MS: 249 (M <sup>+</sup> ), 176, 146, 91.
4c	1 : 0.9	KHCO <sub>3</sub> , 48°C, 34 h	78	Oil	CH <sub>2</sub> Cl <sub>2</sub> -Et <sub>2</sub> O 9 : 1	Styrene	<sup>1</sup> H: 3.05 (1 H, dd, J = 9.0 and 16.5 Hz), 3.35 (1 H, dd, J = 10.5 and 16.5 Hz), 5.22 (2 H, s), 5.66 (1 H, dd, J = 9.0 and 10.5 Hz), 7.4 (5 H, m). <sup>13</sup> C: 40.9, 72.4, 83.1, 167.3. MS: 254 (M <sup>+</sup> + 1), 91.
4d	1 : 1.2	KHCO <sub>3</sub> , 25°C, 48 h KHCO <sub>3</sub> , 48°C, 34 h Et <sub>3</sub> N, 48°C, 1 h	82 91 92	Oil	CHCl <sub>3</sub> -CH <sub>3</sub> OH 99 : 1	Ethyl acrylate	<sup>1</sup> H: 1.20 (3 H, t, J = 7.0 Hz), 1.23 (6 H, d, J = 6.5 Hz), 3.08 (1 H, dd, J = 7.5 and 16.5 Hz), 3.16 (1 H, dd, J = 10.5 and 16.5 Hz), 4.13 (2 H, q, J = 7.0 Hz), 4.70 (1 H, septet, J = 6.5 Hz), 4.87 (1 H, dd, J = 7.5 and 10.5 Hz). <sup>13</sup> C: 14.2, 21.8, 36.9, 62.0, 74.6, 77.9, 165.9, 170.8. IR (film): 1745, 1630. MS: 201 (M <sup>+</sup> ), 186, 160, 159, 128, 127.
4e	1 : 1.25	KHCO <sub>3</sub> , 48°C, 34 h Et <sub>3</sub> N, 48°C, 1 h	54 44	Oil	CHCl <sub>3</sub> -CH <sub>3</sub> OH 99 : 1	1-Butene	<sup>1</sup> H: 0.88 (3 H, t, J = 6 Hz), 1.28 (6 H, d, J = 6 Hz), 1.2-1.8 (6 H, m), 2.57 (1 H, dd, J = 9 and 16.5 Hz), 2.88 (1 H, dd, J = 9 and 16.5 Hz), 4.51 (1 H, m), 4.78 (1 H, septet, J = 6 Hz). <sup>13</sup> C: 14.1, 22.0, 22.7, 27.8, 34.8, 38.8, 73.7, 81.3, 166.8. IR (film): 1625. MS: 185 (M <sup>+</sup> ), 170, 143, 111, 86.
4f	1 : 1.25	Et <sub>3</sub> N, 48°C, 1 h	55	Oil	CH <sub>2</sub> Cl <sub>2</sub> -CH <sub>3</sub> OH 98 : 2	Acrolein diethylacetal	<sup>13</sup> C: 15.4, 21.8, 34.3, 63.6, 64.6, 73.8, 80.9, 102.7, 166.6. IR (film): 1630. MS: 186 (M <sup>+</sup> - OEt), 140, 116, 103.
4g	1 : 0.6	KHCO <sub>3</sub> , 25°C, 72 h	65	Oil	CH <sub>2</sub> Cl <sub>2</sub>	N-Boc-allylamine	<sup>1</sup> H: 1.40 (9 H, s), 2.75 (1 H, dd, J = 8.5 and 16.5 Hz), 2.96 (1 H, dd, J = 9.5 and 16.5 Hz), 3.3 (2 H, m), 4.64 (1 H, m), 5.15 (1 H, br s), 7.3 (5 H, m). MS: 306 (M <sup>+</sup> ).

<b>4h</b>	1 : 1.1	KHCO <sub>3</sub> , 47°C, 48 h	83	Oil	CH <sub>2</sub> Cl <sub>2</sub> -EtOAc 1 : 1	Ethyl acrylate	<sup>1</sup> H: 1.26 (3 H, t, J = 7.5 Hz), 1.30 (3 H, t, J = 7.5 Hz), 3.17 (1 H, dd, J = 7.5 and 16.5 Hz), 3.21 (1 H, dd, J = 10 and 16.5 Hz), 4.16 (2 H, q, J = 7.5 Hz), 4.21 (2 H, q, J = 7.5 Hz), 4.97 (1 H, dd, J = 7.5 and 10 Hz). <sup>13</sup> C: 14.2, 14.5, 36.4, 62.1, 66.8, 78.2, 166.8, 170.7. IR (film): 1735, 1630. MS: 187 (M <sup>+</sup> ).
<b>4i</b>	1 : 1.25	KHCO <sub>3</sub> , 47°C, 45 h	55	Oil (130/0.2)	1 - Hexene	<sup>1</sup> H: 0.88 (3 H, t, J = 7 Hz), 1.28 (3 H, t, J = 7 Hz), 1.4-1.8 (6 H, m), 2.58 (1 H, dd, J = 9 and 16 Hz), 2.90 (1 H, dd, J = 9 and 16 Hz), 4.13 (2 H, q, J = 7.0 Hz), 4.52 (1 H, m). IR (film): 1625. MS: 171 (M <sup>+</sup> ).	
<b>4j</b>	1 : 1.5	KHCO <sub>3</sub> , 25°C, 72 h	82	Oil	Diethyl ether-ethyl acetate 1 : 9	Methyl vinyl ketone	<sup>1</sup> H: 1.30 (3 H, t, J = 6.9 Hz), 2.29 (3 H, s), 3.09 (1 H, dd, J = 10.3 and 16.6 Hz), 3.17 (1 H, dd, J = 6.9 and 16.6 Hz), 4.14 (2 H, q, J = 6.9 Hz), 4.81 (1 H, dd, J = 6.9 and 10.3 Hz). <sup>13</sup> C: 14.5, 26.4, 34.7, 66.8, 84.5, 167.4, 208.5. MS: 157 (M <sup>+</sup> ), 130, 114. IR: 1720, 1625.
<b>4k</b>	1 : 1	KHCO <sub>3</sub> , 25°C, 72 h	77	71	CH <sub>2</sub> Cl <sub>2</sub>	Phenyl vinyl ketone	<sup>13</sup> C: 14.5, 33.7, 66.8, 81.4, 129.2, 130.1, 134.4, 134.8, 167.6, 194.4. IR: 1690, 1625. MS: 219 (M <sup>+</sup> ), 190, 114, 105, 77.
<b>5a</b>	1 : 1	KHCO <sub>3</sub> , 40 h	35	Oil	CH <sub>2</sub> Cl <sub>2</sub> -Et <sub>2</sub> O 4 : 1	Dimethyl acetylenedicarboxylate	<sup>1</sup> H: 1.37 (6 H, d, J = 6.3 Hz), 3.84 (3 H, s), 3.93 (3 H, s), 4.95 (1 H, septet, J = 6.3 Hz). <sup>13</sup> C: 21.7, 52.7, 53.4, 75.5, 107.6, 157.1, 160.4, 160.8, 168.5. IR (film): 1735, 1625. MS: 201 (M <sup>+</sup> - C <sub>3</sub> H <sub>6</sub> ), 170, 169, 111. UV (EtOH): 222.
<b>5b</b>	1 : 0.75	KHCO <sub>3</sub> , 48°C, 36 h	33	86-87	Heptane-Et <sub>2</sub> O 4 : 1	Diphenylacetylene	<sup>1</sup> H: 1.40 (6 H, d, J = 6.3 Hz), 5.03 (1 H, septet, J = 6.3 Hz), 7.3-7.6 (10 H, m). IR (KBr): 1640, 1505. UV (EtOH): 223, 270. MS: 279 (M <sup>+</sup> ), 237, 209, 208.
<b>5c</b>	1 : 0.8	KHCO <sub>3</sub> , 48°C, 40 h	65	Oil	CH <sub>2</sub> Cl <sub>2</sub>	Phenylacetylene	<sup>1</sup> H: 1.40 (6 H, d, J = 6.3 Hz), 4.92 (1 H, septet, J = 6.3 Hz), 6.08 (1 H, s), 7.4 (3 H, m), 7.7 (2 H, m). IR (film): 1620, 1505. UV (EtOH): 258. MS: 203 (M <sup>+</sup> ), 161, 105.
<b>5d<sup>a</sup></b>	1 : 0.6	KHCO <sub>3</sub> , 48°C, 40 h	78	Oil	CH <sub>2</sub> Cl <sub>2</sub> on basic alumina	Ethynyltributyltin	<sup>1</sup> H: 0.8-1.7 (Sn-Bu, m), 1.33 (6 H, d, J = 6 Hz), 4.82 (1 H, septet, J = 6 Hz), 5.87 (1 H, s).
<b>5e</b>	1 : 1	KHCO <sub>3</sub> , 48°C, 48 h	90	65	CH <sub>2</sub> Cl <sub>2</sub> -Et <sub>2</sub> O 4 : 1	Phenylacetylene	<sup>1</sup> H: 5.37 (2 H, s), 6.20 (1 H, s), 7.4 and 7.75 (10 H, m). MS: 251 (M <sup>+</sup> ).
<b>5f</b>	1 : 1	KHCO <sub>3</sub> , 48°C, 34 h	56	(55/2) 39-40	Ethynyltrimethylsilane	<sup>1</sup> H: 0.27 (9 H, s), 1.33 (6 H, d, J = 6 Hz), 4.82 (1 H, septet, J = 6 Hz), 5.94 (1 H, s). <sup>13</sup> C: -2.1, 22.0, 73.9, 104.9, 171.6, 180.4.	
<b>5g</b>	1 : 1.1	KHCO <sub>3</sub> , 47°C, 45 h	21	Oil	CHCl <sub>3</sub>	Phenylacetylene	<sup>1</sup> H: 1.40 (3 H, t, J = 6.5 Hz), 4.33 (2 H, q, J = 6.5 Hz), 6.09 (1 H, s), 7.3-7.7 (5 H, m). MS: 189 (M <sup>+</sup> ).

Table continued

Table 1. Continued

Compound No.	Ratio oxime: alkene or alkyne	Base, Temp, Reaction time	Yield (%)	M.p./°C or (b.p./mmHg)	Eluent TLC, silica	Alkene or alkyne	Selected <sup>1</sup> H, <sup>13</sup> C NMR, δ (CDCl <sub>3</sub> ); IR ν <sub>max</sub> /cm <sup>-1</sup> ; UV, λ <sub>max</sub> /nm; MS, m/z data.
5h <sup>a</sup>	1 : 0.6	KHCO <sub>3</sub> , 48°C, 38 h	89 (crude)	Oil	Et <sub>2</sub> O–heptane 3 : 2, on basic alumina	Ethynyltributyltin	<sup>1</sup> H: 0.8–1.8 (Sn–Bu, m), 5.27 (2 H, s), 6.01 (1 H, s), 7.4 (5 H, m). <sup>13</sup> C: 1.35 (6 H, d, J = 6 Hz), 1.45 (6 H, d, J = 6 Hz), 4.83 (1 H, septet, J = 6 Hz), 5.05 (1 H, septet, J = 6 Hz), <sup>13</sup> C: 21.7, 22.0, 75.1, 79.9, 157.0, 178.8. IR: 1610. MS: 202, 186, 145.
10	—	KHCO <sub>3</sub> , 48°C, 24 h	16	180–190 (decomp.)	Heptane–Et <sub>2</sub> O 3 : 2	—	<sup>1</sup> H (11a): 3.04 (H <sup>4</sup> , dd, J = 5 and 8 Hz), <sup>1</sup> H (11b): 3.21 (H <sup>4</sup> , q, J = 7 Hz). <sup>1</sup> H (11c): 1.25 (3 H, d, J = 6.5 Hz), 3.47 (H <sup>4</sup> , d, J = 9.5 Hz), 4.67 (H <sup>5</sup> , dq, J = 9.5 and 6.5 Hz), <sup>1</sup> H (11d): 3.19 (H <sup>4</sup> , quintet, J = 7 Hz), 4.37 (H <sup>5</sup> , d, J = 7.5 Hz). <sup>1</sup> H (12a): 2.89 (1 H, ddd, J = 2.7, 8.5 and 12.0 Hz), 4.91 (1 H, dd, J = 2.7 and 8.4 Hz), <sup>1</sup> H (12b): 3.92 (1 H, ddd, J = 2.1, 3.6 and 10.1 Hz), 4.73 (1 H, ddd, J = 4.7, 9.9 and 9.9 Hz), <sup>1</sup> H (12c): 2.78 (2 H, t, J = 6.0 Hz), 3.88 (1 H, dd, J = 4.8 and 8.7 Hz), 5.12 (1 H, dd, J = 4.4 and 8.7 Hz). <sup>1</sup> H: 1.21 (6 H, d, J = 6 Hz), 1.32 (3 H, s), 1.83 (2 H, m), 2.58 (1 H, d, J = 16 Hz), 2.80 (1 H, d, J = 16 Hz), 2.9 (1 H, br d), 3.68 (2 H, m), 4.68 (1 H, septet, J = 6 Hz). <sup>13</sup> C: 21.9, 25.6, 41.7, 44.2, 59.0, 73.6, 86.2, 166.7. MS: 187 (M <sup>+</sup> ), 145, 115, 113, 100, 95. IR: 3500, 1620.
11a,b <sup>b</sup>	1 : 1	KHCO <sub>3</sub> , 25°C, 72 h	30	Oils	CHCl <sub>3</sub> –CH <sub>3</sub> OH 99 : 1	Crotonaldehyde diethylacetal	<sup>1</sup> H: 1.25 (6 H, m), 1.58 (3 H, s), 2.80 (1 H, d, J = 16.5 Hz), 3.38 (1 H, d, J = 16 Hz), 4.15 (4 H, m). <sup>13</sup> C: 14.1, 14.4, 23.4, 42.1, 62.1, 66.3, 85.9, 166.6, 172.6. IR (film): 1740, 1625.
11c,d <sup>c</sup>	1 : 1	KHCO <sub>3</sub> , 48°C, 48 h	83 (crude)	Oil	CHCl <sub>3</sub> –CH <sub>3</sub> OH 99 : 1	Ethyl crotonate	
12a–c <sup>d</sup>	1 : 1.4	KHCO <sub>3</sub> , 25°C, 68 h	28 <sup>f</sup>	12a <sup>e</sup> 41–43 12b,c oils	CH <sub>2</sub> Cl <sub>2</sub> –CH <sub>3</sub> OH 99.5 : 0.5	Cycloheptatriene	
16	1 : 1.2	KHCO <sub>3</sub> , 48°C, 38 h	85	Oil	CHCl <sub>3</sub> –CH <sub>3</sub> OH 95 : 5	Isopentenyl alcohol	
17	1 : 1.1	KHCO <sub>3</sub> , 47°C, 48 h	83	Oil (120/0.08)	CH <sub>2</sub> Cl <sub>2</sub> –EtOAc 1 : 1	Ethyl methacrylate	


<sup>a</sup>Decomp. on silica. Contaminated by 3-alkoxyisoxazole formed by metallation. <sup>b</sup>11a : 11b ~ 1 : 1. <sup>c</sup>11c : 11d ~ 1 : 1. <sup>d</sup>12a : 12b : 12c ~ 6 : 2.5 : 1. <sup>e</sup>The isomers were separated by HPLC, SiO<sub>2</sub>, normal phase, diethyl ether–heptane, 1 : 5. <sup>f</sup>The corresponding 3-isopropoxy derivatives were obtained in a total yield of 65% and in approx. the same ratio.

derivatives. Disubstituted olefins gave mixtures of regioisomers with low regioselectivity. The relative stereostructure at C-4,5 depends on the *cis*- or *trans*-structure of the starting olefins (cf. compounds **11a–d** and **12a–c**). An excess of nitrile oxide was used for less active olefins or acetylenes in order to increase the yield of the isoxazole derivatives.

Lower temperature required longer reaction times. Triethylamine liberates the nitrile oxides very fast and the reaction is over within a few minutes. For less reactive olefins, the formation of furoxans from two moles of nitrile oxide was observed. The alkoxy nitrile oxides reacted more efficiently with olefins and acetylenes than did the halonitrile oxides, i.e., the carboxy-hydroxylation reaction via the alkoxy nitrile oxide route was preferred; it reduced the number of steps by one, gave higher yields and avoided the used of toxic haloform oximes.


Cycloaddition to cycloheptatriene gave three products **12a–c** with the isomer **12a** as the major product as predicted by MO theory. The isomer **12d**, formed by a [6 + 4] cycloaddition, was not observed. The structures **12a–c** were proved by two-dimensional H,H correlation spectra.

**Reduction.** Synthesis of  $\gamma$ -amino- $\beta$ -hydroxybutyric acid (*Gabob*), mevalonic acid, malic acid and citramalic acid derivatives. Catalytic reduction over Raney-Ni was the method of choice when other reducible groups, e.g., nitro, epoxy, acetylenic or olefinic groups were absent. Reduction by  $Ti^{3+}$  supplemented the catalytic reduction well. The usefulness of the reaction described here is demonstrated for the synthesis of a DL-amino acid derivative **13**, (a *Gabob* derivative) from Boc-allylamine via **4a** or **4g**, isopropyl mevalonate **14** from 3-methyl-3-butenol via **16**, the diethyl ester of citramalic acid **15**



from ethyl methacrylate via **17** and dialkyl malates **19a,b** from ethyl acrylate via **4d,b**. Citramalic acid has been synthesized previously via another isoxazoline route.<sup>12</sup> Reduction of **12a** ( $R^1 = C_2H_5$ ) with titanous ions gave **18** in a yield of ca. 30%.

**Attempted preparation of tetric acids.** Vinyl ketones gave **4j,k** in high yields and reductive ring opening of **4j,k** gave **20a,b**. Attempted acid-catalysed enol rearrangement of **20a,b** under various conditions combined with lactonization did not lead to the desired  $\gamma$ -methyl- and  $\gamma$ -phenyl-tetric acids **21a,b**, [eqn. (7)].



## Experimental

The  $^1H$  and  $^{13}C$  NMR spectra were obtained with a Varian Gemini 200 spectrometer.  $Me_4Si$  was used as an internal standard. IR spectra were recorded with a Nicolet MX-S, UV spectra with a Unikon 860 and mass spectra with a Micromass 7070 F. Melting points were determined with an Electrothermal melting point apparatus and are uncorrected. Preparative TLC was performed on silica gel 60, PF<sub>254+360</sub>, layers on glass plates (0.2 × 20 × 20 cm). Column chromatography on Kieselgel 60 (0.063–0.20 mm) Merck.

**Triisopropyl orthoformate** was prepared by acid-catalysed transesterification of trimethyl orthoformate with 4 mol of isopropyl alcohol at reflux temperature for ca. 6 h.<sup>7a</sup> The acid was neutralized with metallic sodium and the product was distilled over an efficient fractionating column. The yield was ca. 79%, b.p. 66°C/18 mmHg.  $^1H$  NMR ( $CDCl_3$ ):  $\delta$  5.22 (1 H, s, C–H). Fractions containing mixed methyl–isopropyl orthoformates were reused for further transesterifications.

**Tribenzyl orthoformate** was prepared according to the method of Alexander and Busch.<sup>7b</sup> The crude product was heated to 130°C/0.2 mmHg to remove by-products. The residue in the flask consisted of nearly pure tribenzyl orthoformate, which was used directly for further reactions.  $^1H$  NMR ( $CDCl_3$ ):  $\delta$  5.62 (1 H, s, C–H).

**Methyl formhydroximate, 2a.** Trimethyl orthoformate (5.3 g, 0.05 mol) and 0.35 g of 70% perchloric acid were reacted with  $H_2S$  in a tube-shaped reaction flask fitted with a sintered inlet tube and an outlet connected to  $H_2S$  traps containing conc. NaOH. The space above the orthoformate was first flushed with  $H_2S$  and then the inlet tube was lowered into the orthoformate.  $H_2S$  was introduced in a rapid stream at room temperature. After ca. 5–10 min the solution was poured into

a stirred aqueous solution of hydroxylamine (0.05 mol) at 0°C. The hydroxylamine solution was prepared from hydroxylamine hydrochloride (3.5 g, 0.05 mol), sodium hydroxide (2.0 g, 0.05 mol) and potassium carbonate (2.8 g, 0.02 mol) in water (5 ml). The evolution of H<sub>2</sub>S and CO<sub>2</sub> stopped after ca. 0.5 min. A 1 : 1 mixture of MgSO<sub>4</sub>·K<sub>2</sub>CO<sub>3</sub> (20 g) and dichloromethane (50 ml) were added with vigorous stirring at 0°C. The solid was filtered off and carefully washed with dichloromethane (100 ml). Evaporation of the solvent yielded methyl formhydroximate, **2a**, 1.3 g, 35%, white crystals, m.p. 75–77°C. Recrystallization from carbon tetrachloride raised the m.p. to 85–86°C (lit.<sup>2</sup> 90°C). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 3.87 (3 H, s), 6.55 (1 H, s), 6.55 (1 H, s). The product contained some methyl formate. Methyl formhydroximate **2a** was unstable at room temperature and liquified after a few days. It could be stored in the freezer for a number of months.

*Ethyl formhydroximate*, **2b**, was obtained in a yield of ca. 65% from triethyl orthoformate by applying the procedure described for **2a**. Compound **2b** melted at 81–83°C (from CCl<sub>4</sub>), white needles, (lit.<sup>2</sup> 80°C). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.36 (3 H, t, *J* = 7.5 Hz), 4.11 (2 H, q, *J* = 7.5 Hz), 6.62 (1 H, s). The compound decomposed overnight at 25°C, but could be stored in the freezer.

*Isopropyl formhydroximate*, **2d**, was prepared as described for **2a**, yield 78%, m.p. 85–86°C from carbon tetrachloride. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.33 (6 H, d, *J* = 6.5 Hz), 4.28 (1 H, septet, *J* = 6.5 Hz), 6.66 (1 H, s). MS: *m/z* = 103 (*M*<sup>+</sup>); IR (KBr): 1680 cm<sup>-1</sup> (C=N). The compound was stored in the freezer.

*Benzyl formhydroximate*, **2e**. Tribenzyl orthoformate (16.7 g, 0.05 mol) was reacted with H<sub>2</sub>S at 25°C in the presence of ZnCl<sub>2</sub> (0.3 g). The procedure described for **2a** was then followed but the work-up was changed. The slurry obtained from the oximation was poured onto the top of a silica (65 g) chromatography column (φ = 45 mm). Benzyl alcohol was first eluted with diethyl ether–heptane (1 : 4, 500 ml) and **2e** with diethyl ether–dichloromethane (1 : 1, 500 ml). Evaporation of the solvent *in vacuo* gave **2e** (5.1 g, 68%), white crystals, m.p. 129–130°C. Crystallization from carbon tetrachloride gave white needles, m.p. 130–132°C, <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 5.09 (2 H, s), 6.70 (1 H, s), 7.37 (5 H, s); MS: *m/z* = 151 (*M*<sup>+</sup>); IR (KBr): 1680 cm<sup>-1</sup> (C=N). The compound was stable for several weeks at 25°C.

*General procedure for the preparation of 3-alkoxyisoxazolines and 3-alkoxyisoxazoles, 4,5.* *Method A.* NCS (*N*-chlorosuccinimide, 0.02 mol) was reacted with the alkyl formhydroximate (0.02 mol) at 25°C in chloroform (20 ml) with a few drops of pyridine as a catalyst. The chlorination was complete after ca. 15 min as observed by the disappearance of suspended NCS and of the blue–green colour indicating rearrangement of the intermediate

chloro-nitroso compound into the chloro-oxime. For larger batches it is advisable to cool the reaction flask with tap water. The alkene or alkyne (0.02 mol) was added and finally triethylamine (0.021 mol in 3 ml of CHCl<sub>3</sub>) was added dropwise at 48°C. The ratio dipolarophile : nitrile oxide could be adjusted according to the reactivity, availability, price, etc. of the dipolarophile. The reaction was complete after ca. 30 min. The suspension was filtered, the filtrate was evaporated and carbon tetrachloride (50 ml) was added. The succinimide was filtered off and evaporation of the solution gave the crude 3-alkylisoxazole derivative, which was purified by TLC, column chromatography on silica or basic alumina or distillation at low pressure (< 1 mmHg).

*Method B.* NCS (0.021 mol), alkyl formhydroximate (0.02 mol), alkene or alkyne (0.02 mol), potassium hydrogen carbonate (10 g) and a few drops of water were stirred in ethyl acetate at 48°C for 1–2 days or at 25°C for 2–3 days. The reaction mixture was filtered, evaporated and suspended in carbon tetrachloride (50 ml). Filtration and evaporation of the solution gave the crude isoxazole derivatives. The yields were modest to good. Method B gave somewhat better yields but took longer to perform.

*Catalytic reduction* of the 3-alkoxyisoxazolines was carried out in aqueous methanol in the presence of boric acid using Raney-Ni as the catalyst. The reduction was complete within ca. 1–2 h. The solution was filtered through a thin layer of Celite, evaporated *in vacuo* to a small volume and extracted with chloroform. Drying (MgSO<sub>4</sub>) and evaporation of the solvent gave the crude product, which was purified chromatographically on silica.

*Thioformin, 6.* To ethyl thioformate, prepared from triethyl orthoformate (3.7 g) according to the method described above, were added methylhydroxylamine (from CH<sub>3</sub>NHOH·HCl) (1.74 g), NaOH (1.0 g) and K<sub>2</sub>CO<sub>3</sub> (1.5 g) in water (5 ml) at 0°C and the mixture was stirred for 3 min. Dichloromethane (40 ml), K<sub>2</sub>CO<sub>3</sub> (10 g) and MgSO<sub>4</sub> (16 g) were added with stirring and the suspension was filtered and evaporated *in vacuo*. **6**, 1.54 g, 69%, was obtained. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 3.57 (3 H, s), 8.57 (1 H, s). <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 42.6, 167.9. MS (*m/z*) 91 (*M*<sup>+</sup>). UV (λ<sub>max</sub>): 270 nm in agreement with data reported.<sup>8</sup>

*Furoxan 10* was isolated as a crystalline solid when **1d** was generated according to method B in the absence of a substrate. The crude oily product was purified by preparative TLC (SiO<sub>2</sub>, heptane–Et<sub>2</sub>O, 3 : 2) to give **10** as a solid in a yield of 16%. It decomposed at ca. 180–190°C.

Compound **13a** was obtained as an oil in a yield of 79% by catalytic reduction over Raney-Ni. The crude product was purified by preparative TLC (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>–Et<sub>2</sub>O, 98 : 2). <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 22.0, 28.6, 39.1, 45.7, 68.8, 79.9, 157.1, 172.7. MS (*m/z*): 261 (*M*<sup>+</sup>).

**13b**, yield 71%, TLC (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>-Et<sub>2</sub>O, 9 : 1). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.40 (9 H, s), 2.51 (2 H, m), 3.10 (1 H, ddd, *J* = 14.7, 6.5 and 6.5 Hz), 3.3 (1 H, m), 4.11 (1 H, m), 4.96 (1 H, br s), 5.13 (2 H, s), 7.32 (5 H, br s). MS (*m/z*): 309 (*M*<sup>+</sup>).

**14**, crude yield 78%, TLC (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.24 (6 H, d, *J* = 6.5 Hz), 1.30 (3 H, s), 1.75 (2 H, m), 2.40 (1 H, d, *J* = 15 Hz), 2.57 (1 H, d, *J* = 15 Hz), 3.83 (2 H, m), 5.06 (1 H, septet, *J* = 6.5 Hz).

DL-Diethyl citramalate, **15**, was obtained by catalytic reduction of **17** in a crude yield of 82%. It was practically pure. <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 14.3, 26.5, 44.5, 61.1, 62.3, 72.8, 171.6, 176.2.

**18**. The isoxazoline **12a** (60 mg) in ethanol (1.5 ml) was reduced with aqueous titanium trichloride (1.5 M solution, 150% excess) for 3 days under N<sub>2</sub> with stirring. Powdered Zn was added to maximize the full reducing power of the solution and it was neutralized to pH ca. 3 with sodium carbonate. The solution slowly became colourless. Water (3 ml) was added and the solution was extracted with chloroform. The organic phase was dried over MgSO<sub>4</sub> and evaporated. The oily residue consisted mainly of **18**, 30%, contaminated by a small amount of **12a**. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.25 (3 H, t, *J* = 7.3 Hz), 2.5–2.9 (3 H, m), 4.15 (2 H, q, *J* = 7.3 Hz), 4.62 (1 H, br s), 5.75–6.1 (4 H, m). <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 14.3, 27.4, 48.8, 61.2, 70.3, 125.5, 125.8, 133.3, 133.7, 174.9.

**19a**, yield, 86%, TLC (SiO<sub>2</sub>, CHCl<sub>3</sub>), oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.10 (6 H, d, *J* = 6.5 Hz), 1.17 (3 H, q, *J* = 7 Hz), 2.65 (2 H, m), 3.45 (1 H, br s), 4.12 (2 H, q, *J* = 7 Hz), 4.35 (1 H, t, *J* = 5.5 Hz), 4.91 (1 H, septet, *J* = 6.5 Hz).

**19b**, yield, 74%, TLC, (SiO<sub>2</sub>, CHCl<sub>3</sub>), oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.16 (3 H, t, *J* = 7.5 Hz), 2.80 (2 H, m), 4.15 (2 H, q, *J* = 7.5 Hz), 4.48 (1 H, t, *J* = 6 Hz), 5.10 (2 H, s), 7.30 (5 H, s).

**20a**. The reduction was carried out in aqueous ethanol as the solvent and was stopped when one equivalent of hydrogen had been absorbed. The pH was adjusted to ca. 4 with 2 M HCl. The solution was evaporated to a small volume and extracted several times with ethyl acetate. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated *in vacuo*. The crude yield of **20a** was ca. 90%. The compound was unstable and could not be purified on TLC. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.25 (3 H, t, *J* = 7.5 Hz), 2.28 (3 H, s), 2.72 (1 H, dd, *J* = 1.35 and 6.0 Hz), 2.86 (1 H, dd,

*J* = 13.5 and 4.4 Hz), 4.14 (2 H, q, *J* = 7.5 Hz), 4.36 (1 H, dd, *J* = 6.0 and 4.4 Hz).

**20b**, yield, 63%, TLC (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>-EtOAc, 1 : 1). The reduction was stopped when one equivalent of hydrogen had been absorbed. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.20 (3 H, t, *J* = 7.5 Hz), 2.57 (1 H, dd, *J* = 16.1 and 7.6 Hz), 2.83 (1 H, dd, *J* = 16.1 and 3.5 Hz), 4.11 (2 H, q, *J* = 7.5 Hz), 5.37 (1 H, dd, *J* = 7.6 and 3.5 Hz), 7.3–8.0 (5 H, m).

*Acknowledgments.* We thank Statens Naturvidenskabelige Forskningsråd and Århus Universitets Forskningsfond, Denmark, for financial support.

## References

- In the sequel this name is used in accordance with the naming of other nitrile oxides, e.g., halonitrile oxides.
- Houben, J. *J. Prakt. Chem.* 105 (1922) 7.
- (a) Chiarino, D., Napolitano, M. and Sala, A. *Tetrahedron Lett.* (1986) 3181; *J. Heterocycl. Chem.* 24 (1987) 43; *Synth. Commun.* 18 (1988) 1171; (b) Caldirola, P., De Amici, M. and De Micheli, C. *Tetrahedron Lett.* (1986) 4651; (c) Halling, K., Thomsen, I. and Torsell, K. B. G. *Liebigs Ann. Chem.* (1989) 985 and references therein.
- (a) Brion, J. D., Reynaud, P. and Kirkiacharian, S. *Synthesis* (1983) 220; (b) Mizukami, S. and Nagata, K. *Chem. Pharm. Bull. Jpn.* 14 (1966) 1249.
- (a) Stowell, J. C., Ham, B. M., Esslinger, M. A. and Duplantier, A. J. *J. Org. Chem.* 54 (1989) 1212; (b) Lü Lei and Senning, A. *Sulfur Lett.* 11 (1990) 201.
- Ohno, A., Koizumi, T. and Tsuchihashi, G. *Tetrahedron Lett.* (1968) 2083.
- (a) De Wolfe, R. H. *Synthesis* (1974) 153; (b) Alexander, E. R. and Busch, H. M. *J. Am. Chem. Soc.* 74 (1952) 554; (c) Scheeren, J. W., Van Roy, J. J. H. and Van der Veek, A. P. M. *Recl. Trav. Chim. Pays-Bas* 90 (1971) 745.
- Egawa, Y., Umino, K., Ito, Y. and Okuda, T. *J. Antibiot.* 24 (1971) 124.
- (a) Coates, R. M. and Firsan, S. J. *J. Org. Chem.* 51 (1986) 5198; (b) Walter, W. and Schaumann, E. *Synthesis* (1971) 111.
- See, e. g., Metzger, H. in Houben-Weyl, *Methoden der Organischen Chemie*, Vol. 10 : 4, p. 209, Thieme, Stuttgart 1968; for a review of this class of compound.
- (a) Timms, G. H. and Wildschmidt, E. *Tetrahedron Lett.* (1971) 195; (b) Andersen, S. H., Das, N. B., Jørgensen, R. D., Kjeldsen, G., Knudsen, J. S., Sharma, S. C. and Torsell, K. B. G. *Acta Chem. Scand., Ser. B* 36 (1982) 1.
- (a) Brandi, A., De Sarlo, F., Guarna, A. and Speroni, G. *Synthesis* (1982) 719; (b) Yang, S., Hayden, W., Faber, K. and Gringl, H. *Synthesis* (1992) 365.

Received February 15, 1993.