

Acta Cryst. (1979). A35, 248

Absorption corrections for neutron diffraction. By A. W. HEWAT, *Institute Laue-Langevin, BP 156X, Centre de Tri, 38042 Grenoble, France*

(Received 19 June 1978; accepted 6 September 1978)

Abstract

The transmission factors tabulated by Rouse, Cooper, York & Chakera [*Acta Cryst.* (1970), A26, 682–691] for $\mu r < 1$ can be expressed as the product of two factors: a scale factor and a Debye–Waller factor. In the case of a sphere or a cylinder, the absorption corrections then reduce to simple adjustments of these parameters at the end of the refinement, instead of tedious corrections to the data itself. The results are of particular use for powder data. A printing error in the original paper is also corrected.

Absorption corrections are usually made by reference to a tabulation of the transmission factor A_{hkl} , or by the use of computer programs to correct the raw data. For example, Rouse, Cooper, York & Chakera (1970) have produced a table particularly suited to neutron diffraction, where usually $\mu r < 1$. Of course the use of such tables is tedious, and when computer programs are available it is not always clear what the effect will be of errors in the measurement of the absorption coefficient μ , the crystal radius r , or even the different approximations used to calculate A_{hkl} .

Rouse *et al.* also give an analytical approximation for A_{hkl} , where the error does not exceed 0.0035 for $\mu r < 1$ ($1 > A_{hkl} > 0.1965$):

$$A_{hkl} = \exp[-(a_1 + b_1 \sin^2 \theta)\mu r - (a_2 + b_2 \sin^2 \theta)(\mu r)^2],$$

where the coefficients are:

	Cylinder	Sphere
a_1	1.7133	1.5108
b_1	-0.0368	-0.0315
a_2	-0.0927	-0.0951
b_2	-0.3750	-0.2898.

0567-7394/79/010248-01\$01.00 © 1979 International Union of Crystallography

Acta Cryst. (1979). A35, 248–250

Errors in coordinates, bond lengths, and bond angles. By G. B. CARPENTER, *Metcalf Chemical Laboratories, Brown University, Providence, Rhode Island 02912, USA*

(Received 20 June 1978; accepted 12 September 1978)

Abstract

Standard deviations in bond lengths and bond angles are related to standard deviations in atomic coordinates according to published equations [Cruickshank (1959). *International Tables for X-ray Crystallography*. Vol. II, pp. 331–332]. These equations were derived for an idealized model in

0567-7394/79/010248-03\$01.00

(Note that there is a printing error for b_2 in the original paper.) Clearly this expression can be rewritten as

$$\begin{aligned} A_{hkl} &= k \exp[-\Delta B (\sin \theta / \lambda)^2], \\ k &= \exp[-a_1 \mu r + a_2 (\mu r)^2], \\ \Delta B &= \lambda^2 [b_1 \mu r + b_2 (\mu r)^2], \end{aligned}$$

where k is a scale factor and ΔB is an overall Debye–Waller factor.

Then, unless absolute measurements are being made, the absorption correction for spherical or cylindrical samples can be performed at the end of the refinement simply by adding $\Delta B \cos \gamma^*$ to all the B_{ij} , where γ^* is the angle between the reciprocal-lattice axes i and j .

The result is of particular use for neutron powder diffraction where the product μr can be obtained simply by measuring the transmission through a fine slit placed in front of the sample. For example, a value of $\mu r = 0.579$ was obtained for a 16 mm diameter sample of CsPbCl_3 , giving $\Delta B = 0.27$. Even for a weakly absorbing sample of deuterio-naphthalene, we found $\mu r = 0.274$ and hence $\Delta B = 0.07$. In most cases, absorption corrections are not made for neutron powder diffraction, and this results in significant underestimations of the Debye–Waller factors, even for apparently weakly absorbing materials. Given the ease with which the above formulae can be applied, there is now no excuse for not making these corrections.

The concurrence of M. J. Cooper with these comments on his original paper is acknowledged.

Reference

ROUSE, K. D., COOPER, M. J., YORK, E. J. & CHAKERA, A. (1970). *Acta Cryst.* A26, 682–691.

which the distribution of coordinate errors is isotropic. Tests show that typical structures exhibit only moderate deviations from this model, and so the calculated standard deviations are accurate. Furthermore, the standard deviation in a bond angle φ (in degrees) can be well approximated by the expression $\sigma(\varphi) \sim 81 [\sigma(R)/R]_{\text{r.m.s.}}$, where the quantity in square brackets is the root-mean-square value of $\sigma(R)/R$ for

© 1979 International Union of Crystallography