

SHORT COMMUNICATIONS

Acta Cryst. (1994). A50, 406–408

Magnetic physical-property tensors. By DANIEL B. LITVIN, *Department of Physics, The Pennsylvania State University, Penn State Berks Campus, PO Box 7009, Reading, PA 19610-6009, USA*

(Received 6 July 1993; accepted 18 October 1993)

Abstract

A method is presented to determine the form of magnetic physical-property tensors from the form of non-magnetic physical-property tensors.

1. Introduction

The derivation and tabulation of physical-property tensors invariant under the non-magnetic crystallographic point groups have been considered by many authors (Jahn, 1949; Nye, 1957; Birss, 1964; Kopsky, 1979*a*; Sands, 1982; Brandmuller & Winter, 1985; and references contained in these sources). Tables of a wide variety of physical-property tensors invariant under non-magnetic point groups have been given by Sirotnin & Shaskolskaya (1975) and Brandmuller, Bross, Bauer & Winter (1986) and will appear in the forthcoming Volume D of *International Tables for Crystallography* (1994).

The derivation and tabulation of physical-property tensors invariant under magnetic groups have been considered by Birss (1964) and Kopsky (1979*b*). An analysis of the relationships among the forms of tensors invariant under space inversion, time inversion, space–time inversion and all three inversions has been given by Grimmer (1991). What we present below is a two-step algorithm to read from tables of the form of physical-property tensors invariant under the non-magnetic point groups the form of tensors invariant under all magnetic point groups. This method has already been implemented by Litvin & Litvin (1991) in tabulating the form of rank 0, 1, 2 and 3 magnetic physical-property tensors and in work on non-ferroelastic magnetoelectric twin laws (Litvin, Janovec & Litvin, 1994).

2. Determining the form of magnetic physical-property tensors

Let V denote a polar vector tensor and $V^n = V \times V \times \dots \times V$ the n th ranked product of V ; e and a denote zero-rank tensors that change sign under spatial inversion $\bar{1}$ and time inversion $1'$, respectively. The form of magnetic physical-property tensors is determined by the invariance of the tensors under the elements of the magnetic point group. For example, the invariance of a tensor V^n and the transformation of its components under a pure rotation p gives

$$pV_{ij\dots k}^n = V_{ij\dots k}^n = D^{1-}(p)_{ii} \cdot D^{1-}(p)_{jj} \dots D^{1-}(p)_{kk} \cdot V_{i'j' \dots k'}^n,$$

where the first equality denotes the invariance and the second gives the corresponding constraints on the components of the tensor. For typographical simplicity, we write the above as $pV^n = V^n = D(p)V^n$. In Table 1, we give

Table 1. Invariance and constraints

n odd		n even	
$pV^n = V^n$	$= D(p)V^n$	$pV^n = V^n$	$= D(p)V^n$
$\bar{p}V^n = V^n$	$= -D(p)V^n$	$\bar{p}V^n = V^n$	$= D(p)V^n$
$p'V^n = V^n$	$= D(p)V^n$	$p'V^n = V^n$	$= D(p)V^n$
$\bar{p}'V^n = V^n$	$= -D(p)V^n$	$\bar{p}'V^n = V^n$	$= D(p)V^n$
$p(eV^n) = eV^n$	$= D(p)(eV^n)$	$p(eV^n) = eV^n$	$= D(p)(eV^n)$
$\bar{p}(eV^n) = eV^n$	$= D(p)(eV^n)$	$\bar{p}(eV^n) = eV^n$	$= -D(p)(eV^n)$
$p'(eV^n) = eV^n$	$= D(p)(eV^n)$	$p'(eV^n) = eV^n$	$= D(p)(eV^n)$
$\bar{p}'(eV^n) = eV^n$	$= D(p)(eV^n)$	$\bar{p}'(eV^n) = eV^n$	$= -D(p)(eV^n)$
$p(aV^n) = aV^n$	$= D(p)(aV^n)$	$p(aV^n) = aV^n$	$= D(p)(aV^n)$
$\bar{p}(aV^n) = aV^n$	$= -D(p)(aV^n)$	$\bar{p}(aV^n) = aV^n$	$= D(p)(aV^n)$
$p'(aV^n) = aV^n$	$= -D(p)(aV^n)$	$p'(aV^n) = aV^n$	$= -D(p)(aV^n)$
$\bar{p}'(aV^n) = aV^n$	$= D(p)(aV^n)$	$\bar{p}'(aV^n) = aV^n$	$= -D(p)(aV^n)$
$p(aeV^n) = aeV^n$	$= D(p)(aeV^n)$	$p(aeV^n) = aeV^n$	$= D(p)(aeV^n)$
$\bar{p}(aeV^n) = aeV^n$	$= D(p)(aeV^n)$	$\bar{p}(aeV^n) = aeV^n$	$= -D(p)(aeV^n)$
$p'(aeV^n) = aeV^n$	$= -D(p)(aeV^n)$	$p'(aeV^n) = aeV^n$	$= -D(p)(aeV^n)$
$\bar{p}'(aeV^n) = aeV^n$	$= -D(p)(aeV^n)$	$\bar{p}'(aeV^n) = aeV^n$	$= D(p)(aeV^n)$

Table 2. Number of the column of Table 3 to be used in determining the non-magnetic point group is given at the intersection of the row denoting the magnetic physical-property tensor and the column denoting the parity of the rank of the tensor

	n odd, V^n	n even, eV^n
V^n	2	1
eV^n	1	2
aV^n	4	3
aeV^n	3	4

the invariance and corresponding constraints on the components of tensors V^n , eV^n , aV^n and aeV^n under elements p , \bar{p} , p' and \bar{p}' of a magnetic point group. The two columns of Table 1 correspond to tensors of odd and even rank. These conditions are applicable not only to general tensors of rank n but also to tensors with additional intrinsic symmetry.

From Table 1, we see that there are only two types of sets of constraints: (tensor) = $D(p)$ (tensor) and (tensor) = $-D(p)$ (tensor). Consequently, for odd- (even-) ranked tensors, the form of any magnetic physical-property tensor invariant under a magnetic group M is identical with the form of the physical-property tensor V^n (eV^n) invariant under a non-magnetic point group G . The non-magnetic point group G corresponding to the magnetic group M is determined by replacing those elements p , \bar{p} , p' , \bar{p}' that give rise to the constraint (tensor) = $D(p)$ (tensor) with the element p and those elements that give rise to the constraint (tensor) = $-D(p)$ (tensor) with the element \bar{p} . The resulting non-magnetic groups G are determined from Tables 2 and 3. In Table 2, at the intersection of the row

Table 3. *Magnetic-to-non-magnetic point-group conversion table*

		1	2	3	4			1	2	3	4
1	11'	1	1	$\bar{1}$	$\bar{1}$	46	$\bar{3}1'$	3	$\bar{3}$	$\bar{3}$	$\bar{3}$
2	$\bar{1}1'$	1	$\bar{1}$	$\bar{1}$	$\bar{1}$	47	$\bar{3}'$	3	$\bar{3}$	$\bar{3}$	3
3	$\bar{1}$	1	$\bar{1}$	$\bar{1}$	1	48	321'	32	32	$\bar{3}m$	$\bar{3}m$
4	21'	2	2	2/m	2/m	49	32'	32	32	$\bar{3}m$	$\bar{3}m$
5	2'	2	2	m	m	50	3m1'	32	3m	$\bar{3}m$	$\bar{3}m$
6	m1'	2	m	2/m	2/m	51	3m'	32	3m	3m	32
7	m'	2	m	m	m	52	$\bar{3}m1'$	32	$\bar{3}m$	$\bar{3}m$	$\bar{3}m$
8	2/m1'	2	2/m	2/m	2/m	53	$\bar{3}'m$	32	$\bar{3}'m$	$\bar{3}'m$	$\bar{3}'m$
9	2'/m	2	2/m	2/m	m	54	$\bar{3}m'$	32	3m	3m	$\bar{3}m$
10	2/m'	2	2/m	2/m	2	55	$\bar{3}'m'$	32	$\bar{3}'m$	$\bar{3}'m$	32
11	2'/m'	2	2/m	m	2/m	56	61'	6	6	6/m	6/m
12	2221'	222	222	mmm	mmm	57	6'	6	6	$\bar{6}$	$\bar{6}$
13	2'2'2	222	222	mm2	mm2	58	$\bar{6}1'$	6	$\bar{6}$	6/m	6/m
14	mm21'	222	mm2	mmm	mmm	59	$\bar{6}'$	6	$\bar{6}$	$\bar{6}$	6
15	m'm2'	222	mm2	m2m	2mm	60	6/m1'	6	6/m	6/m	6/m
16	m'm'2	222	mm2	mm2	222	61	6'/m	6	6/m	6/m	$\bar{6}$
17	mmm1'	222	mmm	mmm	mmm	62	6'/m'	6	6/m	6/m	6
18	m'mm	222	mmm	mmm	2mm	63	6'/m'	6	6/m	$\bar{6}$	6/m
19	m'm'm	222	mmm	mm2	mmm	64	6221'	622	622	6/mmm	6/mmm
20	m'm'm'	222	mmm	mmm	222	65	6'2'2	622	622	$\bar{6}m2$	$\bar{6}m2$
21	41'	4	4	4/m	4/m	66	62'2'	622	622	6mm	6mm
22	4'	4	4	$\bar{4}$	$\bar{4}$	67	6mm1'	622	6mm	6/mmm	6/mmm
23	$\bar{4}1'$	4	$\bar{4}$	4/m	4/m	68	6'm'm	622	6mm	$\bar{6}m2$	$\bar{6}2m$
24	$\bar{4}'$	4	$\bar{4}$	$\bar{4}$	4	69	6m'm'	622	6mm	6mm	622
25	4/m1'	4	4/m	4/m	4/m	70	6m21'	622	$\bar{6}m2$	6/mmm	6/mmm
26	4'/m	4	4/m	$\bar{4}$	4/m	71	$\bar{6}'m'2$	622	$\bar{6}m2$	$\bar{6}m2$	622
27	4/m'	4	4/m	4/m	4	72	$\bar{6}'m'2'$	622	$\bar{6}m2$	$\bar{6}2m$	6mm
28	4'/m'	4	4/m	4/m	$\bar{4}$	73	$\bar{6}m'2'$	622	$\bar{6}m2$	6mm	$\bar{6}2m$
29	4221'	422	422	4/mmm	4/mmm	74	6/mmm1'	622	6/mmm	6/mmm	6/mmm
30	4'22'	422	422	$\bar{4}2m$	$\bar{4}2m$	75	6'/m'm	622	6/mmm	6/mmm	6mm
31	42'2'	422	422	4mm	4mm	76	6'/m'm'm	622	6/mmm	6/mmm	$\bar{6}2m$
32	4mm1'	422	4mm	4/mmm	4/mmm	77	6'/m'm'm	622	6/mmm	$\bar{6}m2$	6/mmm
33	4'm'm	422	4mm	$\bar{4}m2$	$\bar{4}2m$	78	6'/m'm'm'	622	6/mmm	6mm	6/mmm
34	4m'm'	422	4mm	4mm	422	79	6/m'm'm'	622	6/mmm	6/mmm	622
35	$\bar{4}2m1'$	422	$\bar{4}2m$	4/mmm	4/mmm	80	231'	23	23	$m\bar{3}$	$m\bar{3}$
36	$\bar{4}'2'm$	422	$\bar{4}2m$	$\bar{4}m2$	4mm	81	$m\bar{3}1'$	23	$m\bar{3}$	$m\bar{3}$	$m\bar{3}$
37	$\bar{4}'2m'$	422	$\bar{4}2m$	$\bar{4}2m$	422	82	$m'3'$	23	$m\bar{3}$	$m\bar{3}$	23
38	$\bar{4}'2'm'$	422	$\bar{4}2m$	4mm	$\bar{4}m2$	83	4321'	432	432	$m\bar{3}m$	$m\bar{3}m$
39	4/mmm1'	422	4/mmm	4/mmm	4/mmm	84	4'32'	432	432	43m	43m
40	4'/m'm	422	4/mmm	4/mmm	4mm	85	$\bar{4}3m1'$	432	$\bar{4}3m$	$m\bar{3}m$	$m\bar{3}m$
41	4'/m'm'm	422	4/mmm	$\bar{4}m2$	4/mmm	86	$\bar{4}'3m'$	432	$\bar{4}3m$	$\bar{4}3m$	432
42	4'/m'm'm'	422	4/mmm	4/mmm	$\bar{4}2m$	87	$m\bar{3}m1'$	432	$m\bar{3}m$	$m\bar{3}m$	$m\bar{3}m$
43	4/mn'm'	422	4/mmm	4mm	4/mmm	88	$m'3'm$	432	$m\bar{3}m$	$m\bar{3}m$	43m
44	4/m'm'm'	422	4/mmm	4/mmm	422	89	$m\bar{3}m'$	432	$m\bar{3}m$	$\bar{4}3m$	$m\bar{3}m$
45	31'	3	3	$\bar{3}$	$\bar{3}$	90	$m'3'm'$	432	$m\bar{3}m$	$m\bar{3}m$	432

corresponding to the type of magnetic physical-property tensor and the column corresponding to either the odd or even rank of the tensor, is a number denoting a column of Table 3. In Table 3, at the intersection of this column and the row corresponding to M , is the required non-magnetic group G .

For example, consider the magnetoelectric effect tensor, which is of the type aeV^2 , invariant under the magnetic group $M = \bar{4}'2'm$. From Table 2, we have that the form of this tensor is the same as the form of the tensor eV^2 invariant under the non-magnetic group G in the fourth column of Table 3 on the row corresponding to the magnetic group $M = \bar{4}'2'm$. From Table 3, we have $G = 4mm$. The form of the physical-property tensor aeV^2 invariant under $M = \bar{4}'2'm$ is then the same as the form of the tensor eV^2 invariant under $G = 4mm$. The latter is given by Sirotn & Shaskolskaya (1975) as

$$\begin{bmatrix} 0 & A & 0 \\ -A & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

which agrees with the form of the magnetoelectric physical-property tensor invariant under $\bar{4}'2'm$ given by Birss (1964).

This work was supported by the National Science Foundation under grants DMR-9100418 and DMR-9305825.

References

- BIRSS, R. R. (1964). *Symmetry and Magnetism*. Amsterdam: North-Holland.
- BRANDMULLER, J., BROSS, H., BAUER, W. G. & WINTER, F. X. (1986). Tables to accompany Brandmuller & Winter (1985), 2nd ed.
- BRANDMULLER, J. & WINTER, F. X. (1985). *Z. Kristallogr.* **172**, 191–231.
- GRIMMER, H. (1991). *Acta Cryst.* **A47**, 226–232.
- International Tables for Crystallography* (1994). Vol. D: *Physical Properties of Crystals*, edited by A. AUTHIER. In preparation.
- JAHN, H. A. (1949). *Acta Cryst.* **2**, 30–33.

KOPSKY, V. (1979a). *Acta Cryst.* **A35**, 83–95.

KOPSKY, V. (1979b). *Acta Cryst.* **A35**, 95–101.

LITVIN, D. B., JANOVEC, V. & LITVIN, S. Y. (1994). *Ferroelectrics*. In the press.

LITVIN, S. Y. & LITVIN, D. B. (1991). *Acta Cryst.* **A47**, 290–292

NYE, J. F. (1957). *Physical Properties of Crystals*. Oxford: Clarendon Press.

SANDS, D. E. (1982). *Vectors and Tensors in Crystallography*. Reading, MA: Addison-Wesley.

SIROTIN, YU. & SHASKOLSKAYA, M. P. (1975). *Osnovi Kristallogfiziki*. Moscow: Nauka.

Acta Cryst. (1994). **A50**, 408–409

Normalization factors for Kubic harmonic density functions. By ZHENGWEI SU and PHILIP COPPENS, *Department of Chemistry, State University of New York at Buffalo, Buffalo, New York 14214-3094, USA*

(Received 19 August 1993; accepted 22 November 1993)

Abstract

Normalization factors N_{lj} for the Kubic harmonics K_{lj} defined by

$$\int_{\theta=0}^{\pi} \int_{\varphi=0}^{2\pi} N_{lj} |K_{lj}| \sin \theta \, d\theta \, d\varphi = 2 - \delta_{l0}$$

have been evaluated numerically for $l \leq 10$.

Introduction

For cubic site symmetries, the symmetry-adapted functions for describing the angular dependence of the atomic charge density are the Kubic harmonics K_{lj} (Kurki-Suonio, 1977; Dawson, 1967). However, the normalization factors available in the literature, defined by

$$(1/M_l^2) \int_{\theta=0}^{\pi} \int_{\varphi=0}^{2\pi} K_{lj}^2 \sin \theta \, d\theta \, d\varphi = 1 \quad (1)$$

(Von der Lage & Bethe, 1947; Kurki-Suonio, 1977), apply to the normalization of wave functions. For density functions, the normalization

$$N_{lj} \int_{\theta=0}^{\pi} \int_{\varphi=0}^{2\pi} |K_{lj}| \sin \theta \, d\theta \, d\varphi = 2 - \delta_{l0} \quad (2)$$

is appropriate. With this normalization, a population coefficient $P_{lj} = 1$ implies, for $l = 0$, a population of one electron and, for $l > 0$, that one electron has been transferred from the negative to the positive lobes of the function $R(r)N_{lj}K_{lj}$, where $R(r)$ is a normalized radial function.

The unnormalized Kubic harmonics K_{lj} have been expressed in different forms (Kurki-Suonio, 1977; Dawson, 1967; Von der Lage & Bethe, 1947). We adopt the expressions given by Kurki-Suonio (1977).

The calculation

Table 1 lists the definitions of K_{lj} up to $l = 10$, together with the N_{lj} values. Except for K_0 ($N_0 = 1/4\pi$) and K_3 ($N_3 = 1/15$), the values have been calculated by numerical integration using the Gaussian quadrature (Press, Flannery,

Table 1. *Normalization factors for Kubic harmonics*

$u_{lm\pm}$ is given here as $P_l^m(\cos\theta) \frac{\cos(m\varphi)}{\sin(m\varphi)}$	Unnormalized $K_{lj}(0, 0)$	Normalization factor N_{lj}
	$K_0 = u_{00+} = 1$	$1/4\pi = 0.07957747$
	$K_1 = u_{12-}$	$2/30 = 1/15 = 0.06666667$
	$K_4 = u_{40+} + (1/168)u_{44+}$	$2/4.602568 = 0.4345400$
	$K_{6,1} = u_{60+} - (1/360)u_{64+}$	$2/7.930299 = 0.2521973$
	$K_{6,2} = u_{62+} - (1/792)u_{66+}$	$2/96.00000 = 0.02083333$
	$K_7 = u_{72-} + (1/1560)u_{76-}$	$2/136.8711 = 0.01461229$
	$K_8 = u_{80+} + (1/5940)[u_{84+} + (1/672)u_{88+}]$	$2/3.552877 = 0.5629241$
	$K_{9,1} = u_{92-} - (1/2520)u_{96-}$	$2/335.7424 = 0.005956948$
	$K_{9,2} = u_{94-} - (1/4080)u_{98-}$	$2/13513.50 = 0.0001480001$
	$K_{10,1} = u_{10,0+} - (1/5460)[u_{10,4+} + (1/4320)u_{10,8+}]$	$2/5.480931 = 0.3649015$
	$K_{10,2} = u_{10,2+} + (1/43680)[u_{10,6+} - (1/456)u_{10,10+}]$	$2/210.1611 = 0.009516509$

Table 2. *Some high-order $y_{lm\pm}$ [$z = \cos(\theta)$]*

u_{80+}	$(1/128)(6435z^8 - 12012z^6 + 6930z^4 - 1260z^2 + 35)$
u_{84+}	$(1/8)(10395)(z^2 - 1)^2(65z^4 - 26z^2 + 1)\cos(4\varphi)$
u_{88+}	$2027025(z^2 - 1)^4\cos(8\varphi)$
u_{92-}	$(495/16)z(1 - z^2)(221z^6 - 273z^4 + 91z^2 - 7)\sin(2\varphi)$
u_{94-}	$(135135/8)z(z^2 - 1)^2(17z^4 - 10z^2 + 1)\sin(4\varphi)$
u_{96-}	$(675675/2)z(1 - z^2)^3(17z^2 - 3)\sin(6\varphi)$
u_{98-}	$34459425z(z^2 - 1)^4\sin(8\varphi)$
$u_{10,0+}$	$(1/256)(46189z^{10} - 109395z^8 + 90090z^6 - 30030z^4 + 3465z^2 - 63)$
$u_{10,2+}$	$(495/128)(1 - z^2)(4199z^8 - 6188z^6 + 2730z^4 - 364z^2 + 7)$
$u_{10,4+}$	$(45045/16)(z^2 - 1)^2(323z^6 - 255z^4 + 45z^2 - 1)\cos(4\varphi)$
$u_{10,6+}$	$(675675/8)(1 - z^2)^3(323z^4 - 102z^2 + 3)\cos(6\varphi)$
$u_{10,8+}$	$(34459425/2)(z^2 - 1)^4(19z^2 - 1)\cos(8\varphi)$
$u_{10,10+}$	$654729075(1 - z^2)^5\cos(10\varphi)$

Teukolsky & Vetterling, 1986). Satisfactory results have been obtained by dividing the intervals of integration for θ and φ into about 200 parts, evaluating the integral in each small solid angle by the 10-point Gaussian formula and subsequently summing the integrals. Finer divisions did not change the values to seven significant figures. For example, the numerical result for N_3 is 0.999999997/15 for both a 200-by-200 and a 300-by-300 division of $\theta\varphi$ space.