

of monomeric units arranged in layers, as previously proposed (Martell & Calvin, 1959). Dimerization is achieved by the sharing of one oxygen atom from each 'salen' ligand by two cobalt atoms. Thus the cobalt atom is five coordinated in a distorted rectangular based pyramid with an axial bond Co-O of 2.25 Å. The close approach of the two halves of the dimer requires the two planes of the 'salen' ligand to bend away from each other. The Co-N (mean value 1.88 Å) and Co-O (1.88 Å) bond lengths agree with the mean values of 1.88 and 1.86 Å respectively, found in some organo-cobalt derivatives of bis(acetylacetonate)-ethylendiimine (BAEH₂), (Brückner, Calligaris, Nardin & Randaccio, 1968a). On the other hand, the Co-O value of 1.95 Å agrees with the mean value of 1.92 Å found in vinyl-aquo(BAE)Co (Brückner, Calligaris, Nardin & Randaccio, 1968b). In the last compound the lengthening of Co-O bonds has been ascribed to the intermolecular hydrogen bonds in which these oxygen atoms are involved. Analogously the lengthening found here can be ascribed to the additional coordination bond of 2.25 Å in which the oxygen atom is involved.

The torsional angle around the CH₂-CH₂ bond is 42.0° and the angle between the planes passing through the two halves of the 'salen' ligand is 22.2°. The close stacking of the dimeric units does not introduce voids in the structure. Thus the crystal packing of the inactive form is such that it may be regarded as 'self-protecting' towards oxygen absorption.

The dimeric nature of this compound suggests a possible explanation of the two-to-one stoichiometry of the oxygenated adduct. If one assumes that the crystals of the active form consist of dimers packed in an open way, the formation of an adduct [(cosalen)₂O₂]_n seems likely when the oxygen molecules penetrate into the voids of the structure. On the contrary, in the model of equally displaced monomeric units, proposed by Calvin, the 2:1 and 1:1 ratios would both be possible.

However this conjecture requires further experimental information to support it.

We acknowledge financial assistance from the Consiglio Nazionale delle Ricerche, Roma, Italy. We are grateful to Professor A. Ripamonti for helpful discussions.

References

- ALBANO, V., DOMENICANO, A. & VACIAGO, A. (1966). *Gazz. Chim. Ital.* **96**, 922.
 BAILES, R. H. & CALVIN, M. (1947). *J. Amer. Chem. Soc.* **69**, 1886.
 BAYER, E. & SCHRETZMANN, P. (1967). *Structure and Bonding*, **2**, 181.
 BERGHUIS, J., HAANAPPEL, IJ. M., POTTERS, M., LOOPSTRA, B. O., MACGILLAVRY, C. H. & VEENENDAAL, A. L. (1955). *Acta Cryst.* **8**, 478.
 BRÜCKNER, S., CALLIGARIS, M., NARDIN, G. & RANDACCIO, L. (1968a). *Inorg. Chim. Acta*, **2**, 386.
 BRÜCKNER, S., CALLIGARIS, M., NARDIN, G. & RANDACCIO, L. (1968b). *Inorg. Chim. Acta*, **2**, 416.
 HALL, D. & WATERS, T. N. (1960). *J. Chem. Soc.* p. 2644.
 HAMILTON, W. C., ROLLETT, J. S. & SPARKS, R. A. (1965). *Acta Cryst.* **18**, 129.
 HANSON, H. P., HERMAN, F., LEA, J. D. & SKILLMAN, S. (1964). *Acta Cryst.* **17**, 1040.
International Tables for X-ray Crystallography (1962). Vol. III. Birmingham: Kynoch Press.
 MARTELL, A. E. & CALVIN, M. (1959). *Chemistry of the Metal Chelate Compounds*. Englewood Cliffs, New Jersey: Prentice Hall Inc.
 PHILLIPS, D. C. (1954). *Acta Cryst.* **7**, 746.

Notes and News

Announcements and other items of crystallographic interest will be published under this heading at the discretion of the Editorial Board. The notes (in duplicate) should be sent to the General Secretary of the International Union of Crystallography (G. Boom, Laboratorium voor Fysische Metaalkunde der Rijksuniversiteit, Universiteitscomplex Paddepoel, Groningen 8002, The Netherlands). Publication of an item in a particular issue cannot be guaranteed unless the draft is received 8 weeks before the date of publication.

Crystallographic programs for the IBM 1130 computer

Colleague crystallographers may be interested to know that we have available a collection of crystallographic programs suitable for use on an IBM 1130 computer with 8 K core and one magnetic disc storage:

1. Lorentz-polarization corrections for Weissenberg data.
2. Lorentz-polarization corrections for precession data.
3. Wilson plot and E-calculation.
4. Temperature-sharpened Patterson map calculation.
5. Centrosymmetric structure factor calculation, and Fourier summation (2- and 3-dimensional).
6. Acentric structure factor calculation, and Fourier summation (2- and 3-dimensional).

7. Interatomic-distance and bond-angle calculation.
8. All-distance calculation.
9. Projection down any direction.
10. Molecular translation search programs for plane groups *p2*, *pgg*, *pmg*, *pg*, *pm*.

These programs are not original, but have been collected from several sources and modified for this model computer.

On request, a FORTRAN listing and operating instructions will be supplied.

MICHAEL LAING

*Department of Chemistry
 Arthur Sale, Computer Centre
 University of Natal
 Durban
 South Africa*