



Fig. 1. The absolute configuration of centaurepensin.

with a slope of 1.01 passing through the origin. The correlation coefficient and  $\chi^2$  of the fit are 0.99 and 0.01, respectively, indicating effectively identical results.\*

\* Positional and anisotropic thermal parameters for non-hydrogen atoms, bond distances, bond angles, torsion angles and lists of structure factors have been deposited with the British Library Lending Division as Supplementary Publication No. SUP 33565 (18 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

The present absolute configuration, shown in Fig. 1, corresponds to the enantiomorph of the one previously reported (Mason, Hewson, Kennard & Pettersen, 1972; Hewson, Pettersen & Kennard, 1972). This was confirmed after refinement of both enantiomers to give  $R = 0.044$  and  $0.054$  ( $R_w$  being 0.057 and 0.076, respectively). Finally a comparison of the 30 most relevant Bijvoet pairs gave the following discrepancy indices: averaged Bijvoet difference of 0.72 (3.11 for the enantiomorph) and averaged Bijvoet ratio of 0.05 (0.49).

We thank the staff of the Centro de Proceso de Datos del MEC of Spain for facilities on the Univac 1108 computer.

#### References

- ABRAHAMS, S. C. & KEVE, E. T. (1971). *Acta Cryst.* **A27**, 157–165.  
 GONZALEZ, A. G., BERMEJO, J., BRETÓN, J. L., MASSANET, G. M., DOMINGUEZ, B. & AMARO, J. M. (1976). *J. Chem. Soc. Perkin Trans.* **1**, pp. 1663–1666.  
 HEWSON, A. T., PETERSEN, R. C. & KENNARD, O. (1972). *Cryst. Struct. Commun.* **1**, 383–388.  
 MASON, J. H., HEWSON, A. T., KENNARD, O. & PETERSEN, R. C. (1972). *Chem. Commun.* pp. 460–461.  
 STEWART, J. M., KUNDELL, F. A. & BALDWIN, J. C. (1970). The XRAY 70 system. Computer Science Center, Univ. of Maryland, College Park, Maryland.

*Acta Cryst.* (1978). **B34**, 2670

**4,5-Dichloro-1,3,6,8-tetrafluoro-2,7-naphthyridine: erratum.** By D. M. W. VAN DEN HAM and G. J. VAN HUMMEL, *Twente University of Technology, Chemical Physics Laboratory, PO Box 217, Enschede, The Netherlands*

(Received 22 March 1978; accepted 6 April 1978)

In the discussion of the experimental results the Cl–Cl distance of the title compound was erroneously given as 2.307 Å in the paper by van den Ham & van Hummel [*Acta Cryst.* (1977), **B33**, 3866–3868]. This value should be 3.109 Å. As this value is still smaller than twice the van der Waals radius, this error does not affect the conclusion.

All the relevant information is given in the Abstract.

*Acta Cryst.* (1978). **B34**, 2670–2671

**Experimental and theoretical difference densities for cyanuric acid. A simple refinement of density distributions of bonding electrons VI. Erratum.** By C. SCHERINGER, A. KUTOGLU and E. HELLNER, *Institut für Mineralogie der Universität Marburg, D-3550 Marburg/Lahn, Federal Republic of Germany*

(Received 27 June 1978)

Owing to an error in the data transmission with respect to the form factor curve of oxygen, Fig. 1(a,b) of the paper by Scheringer, Kutoglu, Hellner, Hase, Schulte & Schweig [*Acta Cryst.* (1978). **B34**, 2162–2165] – hereafter referred to as SKHHSS – was incorrectly drawn. Similarly, the peak heights of the experimental investigation in Table 1 of SKHHSS are not correct. The corrected Fig. 1(a,b) and the full Table 1 are given. The good agreement between the experimental and theoretical peak heights mentioned in SKHHSS no longer holds for the lone-pair peaks of the oxygen atoms. The experimental peaks are considerably lower and in better agreement with the  $X - N$  map of Coppens & Vos [*Acta Cryst.* (1971), **B27**, 146–158].