

**Literatur**

- BOSWIJK, K. H., VAN DER HEIDE, J., VOS, A. & WIEBENGA, E. H. (1956). *Acta Cryst.* **9**, 274–277.
- BOSWIJK, K. H. & WIEBENGA, E. H. (1953). *Acta Cryst.* **7**, 417–423.
- BURNHAM, C. W. (1966). *Am. Mineral.* **51**, 159–164.
- CARPENTER, G. B. & RICHARDS, S. M. (1962). *Acta Cryst.* **15**, 360–364.
- HASSEL, O. & RØMMING, C. (1967). *Acta Chem. Scand.* **21**, 2659–2662.
- International Tables for X-ray Crystallography* (1968). Bd. III, 2. Aufl., S. 202–203, 211, 215–216. Birmingham: Kynoch Press.
- NAKAMOTO, K. (1960). *J. Phys. Chem.* **64**, 1420–1425.
- RØMMING, C. (1972). *Acta Chem. Scand.* **26**, 1555–1560.
- RUNDLE, R. E. (1962). *Rec. Chem. Prog.* **23**, 206–215.
- SOLED, S. & CARPENTER, G. B. (1973). *Acta Cryst.* **B29**, 2104–2109.
- STEWART, J. M. (1976). The XRAY 76 system. Tech. Rep. TR-446. Computer Science Center, Univ. of Maryland, College Park, Maryland.
- WEIDNER, R. T. (1947). *Phys. Rev.* **72**, 1268–1269.
- YAGI, Y. & POPOV, A. I. (1967). *J. Inorg. Nucl. Chem.* **29**, 2223–2230.

**SHORT COMMUNICATIONS**

*Contributions intended for publication under this heading should be expressly so marked; they should not exceed about 1000 words; they should be forwarded in the usual way to the appropriate Co-editor; they will be published as speedily as possible.*

*Acta Cryst.* (1979), **B35**, 1933

**Produit de cyclisation: la dihydro-5,6 (oxo-2 propyl)-6 4H-pyrrolo[1,2-a]thiéno[3,2-f][diazépine-1,4]one-4. Errata.** Par NGUYEN-HUY-DUNG, Laboratoire de Chimie Minérale et Structurale, UER des Sciences Pharmaceutiques de Caen, 1 rue Vaubanard, 14032 Caen CEDEX, France et SYLVAIN RAULT et MAX ROBBA, Laboratoire de Pharmacie Chimique, UER des Sciences Pharmaceutiques de Caen, 1 rue Vaubanard, 14032 Caen CEDEX, France

(Received 14 June 1979)

**Abstract**

In the paper by Nguyen-Huy-Dung, Rault & Robba [*Acta Cryst.* (1979), **B35**, 1290–1293] the values given for  $M_r$ ,

0567-7408/79/081933-01\$01.00

$D_x$  and  $\mu$  are incorrect. The correct values are:  $M_r = 260.31$ ,  $D_x = 1.40 \text{ Mg m}^{-3}$ ,  $\mu = 0.256 \text{ mm}^{-1}$ .

Le résumé contient tous les détails.

© 1979 International Union of Crystallography

*Acta Cryst.* (1979), **B35**, 1933–1934

**Some comments on refinement in a space group of unnecessarily low symmetry.\*** By VERNER SCHOMAKER, Department of Chemistry, University of Washington, Seattle, Washington 98105, USA and RICHARD E. MARSH, Arthur Amos Noyes Laboratory of Chemical Physics, California Institute of Technology, Pasadena, California 91125, USA

(Received 24 January 1979; accepted 12 April 1979)

**Abstract**

Releasing the constraint of a center of symmetry causes a well known singularity, which, however, often goes unrecognized. Deleting an element of translational symmetry (transition to a supercell) also causes singularity. In contrast,

deleting any element other than a center or a pure translation (to give a lower Laue group) is uneventful.

Lee (1971) has suggested that 'in cases where the space group is not absolutely certain, from a structural point of view it is safest (and cannot be wrong) to assume the lower symmetry'. In rebuttal, Donohue (1971) pointed out that the logical import of Lee's suggestion is that all structures should

\* Contribution No. 5962 from the Arthur Amos Noyes Laboratory.

0567-7408/79/081933-02\$01.00

© 1979 International Union of Crystallography