

SHORT COMMUNICATIONS

Acta Cryst. (1997). B53, 188**Neutron powder diffraction study of α -Ti(HPO₄)₂·H₂O and α -Hf(HPO₄)₂·H₂O; H-atom positions. Erratum**

MIGUEL A. SALVADÓ,^a PILAR PERTIERRA,^a SANTIAGO GARCÍA-GRANDA,^a JOSÉ R. GARCÍA,^b JULIO RODRÍGUEZ^b AND M. T. FERNÁNDEZ-DÍAZ^c at ^a*Departamento de Química Física y Analítica* and ^b*Area de Química Inorgánica, Facultad de Química, Universidad de Oviedo, 33006 Oviedo, Spain,* and ^c*Institut Laue-Langevin, BP 156, F-38042 Grenoble CEDEX 9, France.*
 E-mail: sgg@dwarf1.quimica.uniovi.es

(Received 30 November 1996)

Fractional atomic coordinates and isotropic displacement parameters (Å^2) for H atoms

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
Abstract				
The H-atom coordinates were omitted from Tables 2 and 3 on p. 897 [Salvadó, Pertierra, García-Granda, García, Rodríguez & Fernández-Díaz, <i>Acta Cryst.</i> (1996), B52, 896–898]. The values are now listed.				
α -TiP				
H1	0.4133 (17)	0.245 (3)	0.7235 (9)	0.0415 (17)
H2	0.1440 (19)	0.943 (3)	0.7289 (9)	0.0415 (17)
H3	0.1948 (17)	0.390 (3)	0.7548 (11)	0.0415 (17)
H4	0.3083 (18)	0.233 (4)	0.8353 (8)	0.0415 (17)
α -HfP				
H1	−0.100 (2)	0.811 (4)	0.7167 (15)	0.052 (3)
H2	0.659 (3)	0.076 (4)	0.7282 (16)	0.052 (3)
H3	0.294 (3)	0.079 (4)	0.763 (2)	0.052 (3)
H4	0.212 (2)	0.221 (5)	0.6681 (10)	0.052 (3)