

Structures of mono-unsaturated triacyl-glycerols. II. The β_2 polymorph. Corrigendum

Jan B. van Mechelen,* Rene Peschar and Henk Schenk

University of Amsterdam, HIMS/FNWI/Kristallografie, Valckenierstraat 65, 1018 XE Amsterdam, The Netherlands

Correspondence e-mail: janvm@science.uva.nl

A replacement Fig. 3 to the paper by van Mechelen *et al.* (2006), *Acta Cryst. B62*, 1131–1138 is given.

Fig. 3 in the article by van Mechelen *et al.* [(2006b), *Acta Cryst. B62*, 1131–1138] is actually a duplicate of Fig. 3 from van Mechelen *et al.* [(2006a), *Acta Cryst. B62*, 1121–1130]. The correct figure is given here.

References

- Mechelen, J. B. van, Peschar, R. & Schenk, H. (2006a). *Acta Cryst. B62*, 1121–1130.
 Mechelen, J. B. van, Peschar, R. & Schenk, H. (2006b). *Acta Cryst. B62*, 1131–1138.




Figure 3

Higher-angle part of Fig. 2 showing the differences between the β_2 patterns. (1, k , -1) peaks are labeled (k) and (0, k , 2) peaks are labeled underscore k .