Fig. 2. Molecular packing viewed down the b axis.

The glycosyl torsion angle $C(2)-N(1)-C(1')-O(4')$ is $-74.2(2)^\circ$. The sugar pucker is in a twisted form, $C(4')\text{-endo}-C(3')\text{-exo}$, with pseudorotation angle $P = 220.9^\circ$ as defined by Altona & Sundaralingam (1972). A similar conformation is also observed in 8,2'-*O*-cycloadenosine ($P = 217.5^\circ$) and in 8,2'-*S*-cyclo-5'-AMP ($P = 232.7^\circ$).

The molecular packing is illustrated in Fig. 2. A hydrogen bond is observed between $O(5')$ and $N(3)$ in the neighbouring molecule related by the 2_1 screw axis.

References

- ALTONA, C. & SUNDARALINGAM, M. (1972). *J. Am. Chem. Soc.* **94**, 8205–8212.
 ASHIDA, T. (1979). *HBL SV. The Universal Crystallographic Computing System—Osaka*. Library of Programs, Computing Center, Osaka Univ.
International Tables for X-ray Crystallography (1974). Vol. IV. Birmingham: Kynoch Press.
 JOHNSON, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee.
 KITAMURA, K., MIZUNO, H., SUGIO, S., OKABE, N., IKEHARA, M. & TOMITA, K. (1983). *Acta Cryst. C* **39**, 1551–1553.
 MAIN, P., HULL, S. E., LESSINGER, L., GERMAIN, G., DECLERCQ, J.-P. & WOOLFSON, M. M. (1978). *MULTAN78. A System of Computer Programs for the Automatic Solution of Crystal Structures from X-ray Diffraction Data*. Univer. of York, England, and Louvain, Belgium.
 NEIDLE, S., TAYLOR, G. L. & COWLING, P. C. (1979). *Acta Cryst. B* **35**, 708–712.
 TANAKA, K., FUJII, S., FUJIWARA, T. & TOMITA, K. (1979). *Acta Cryst. B* **35**, 929–933.

Acta Cryst. (1984). **C40**, 188–190

Structure of 1-(*p*-Ethoxyphenyl)-1,3-dihydro-3-phenyl-2*H*-benzimidazole-2-thione, $C_{21}H_{18}N_2OS$


BY A. CRIADO, A. CONDE AND R. MÁRQUEZ

Departamento de Óptica y Sección de Física del Centro Coordinado de CSIC, Universidad de Sevilla, Spain

(Received 18 April 1983; accepted 21 September 1983)

Abstract. $M_r = 346.4$, orthorhombic, $P2_12_12_1$, $a = 7.600(1)$, $b = 11.132(2)$, $c = 20.767(3)\text{ \AA}$, $V = 1756.9(5)\text{ \AA}^3$, $Z = 4$, $D_x = 1.31\text{ Mg m}^{-3}$, $\lambda(\text{Cu }K\alpha) = 1.5418\text{ \AA}$, $\mu = 1.67\text{ mm}^{-1}$, $F(000) = 728$, $T = 300\text{ K}$. Final $R = 0.049$ for 1531 observed independent reflections. The benzimidazole bicyclic is quasi-planar, the dihedral angle between the two fused rings being $1.6(2)^\circ$. The unsubstituted phenyl ring is planar while the phenyl ring with the ethoxy substituent deviates significantly from the expected planar conformation.

The title compound (I) has been obtained (Fernández-Bolaños, Fuentes-Mota & Fernández-Bolaños Guzmán, 1983a) from the *p*-toluenesulfonate of 4-(β -D-erythrofuranosyl)-1-(*p*-ethoxyphenyl)-1,3-dihydro-3-phenyl-2*H*-imidazole-2-thione (Fernández-Bolaños, Fuentes-Mota & Fernández-Bolaños Guzmán, 1983b) using a reaction similar to that of Tipson & Cohen (1965). This synthesis is the first case of nucleoside aromatization.



Introduction. The crystal structure of the title compound has been determined as part of a systematic structural analysis of imidazole *C*-nucleosides synthesized in the Organic Chemistry Department of this University. Recently, we have reported (Criado, Conde & Márquez, 1983) the structure of the first disubstituted (at both N atoms) imidazole *C*-nucleoside.

Experimental. Needle-shaped colorless crystals, $0.05 \times 0.06 \times 0.14$ mm, kindly supplied by Professor Fernández-Bolaños of the Organic Chemistry Department of this University; preliminary studies showed orthorhombic symmetry, systematic absences consistent with $P2_12_12_1$. Philips PW 1100 four-circle computer-controlled diffractometer, graphite-monochromated radiation. Unit-cell parameters from least-squares refinement of θ values of 33 reflections. 1729 reflections with $\theta < 65^\circ$ ($h < 9$, $k < 14$, $l < 25$) measured, $\omega-2\theta$ scan mode, 1531 with $I > 2\sigma(I)$ considered observed. Two reference reflections monitored periodically, intensity changes less than 2%. Lorentz and polarization corrections, none for absorption or extinction effects. Weighted tangent formula refinement (*MULTAN78*, Main, Hull, Lessinger, Germain, Declercq & Woolfson, 1978) of 180 reflections with $E > 1.5$. Full-matrix least-squares refinement of 281 parameters based on F_o over all observed reflections (*CRYLSQ* of XRAY70, Stewart, Kundell & Baldwin, 1970). Fourier difference synthesis up to $\sin\theta/\lambda = 0.7 \text{ \AA}^{-1}$ gave all H-atom positions; a least-squares refinement in a mixed mode including H atoms, each one with an isotropic temperature factor equal to that of the attached skeleton atom, gave $R = R_w = 0.049$. Last-cycle parameter shifts less than 0.3σ , average 0.1σ . Final difference Fourier map excursions $-0.25 \leq \Delta\rho \leq 0.35 \text{ e \AA}^{-3}$. Atomic scattering factors from *International Tables for X-ray Crystallography* (1974). Weighting scheme in the form $1/\sigma^2(I)$ based on counting statistics.

Discussion. Final atomic coordinates and equivalent isotropic temperature factors (Hamilton, 1959) are given in Table 1.* Bond lengths and angles for the non-H atoms are given in Fig. 1. The C—H distances range from 0.95 to 1.19 Å, with a mean value of 1.08 (6) Å.

Benzimidazole ring. Bond distances and angles in the imidazole ring are in good agreement with the reported values for imidazole-2-thione compounds (Conde, López-Castro & Márquez, 1978; Criado *et al.*, 1983) showing no appreciable distortion due to the bicyclic formation. Only the C(2)—C(3) bond distance is somewhat long in comparison with the mean value [1.345 (5) Å] of the aforementioned compounds. The ring keeps its planar feature and the atomic deviations from the least-squares plane through the five atoms are within the standard deviations ($\chi^2 = 0.72$). The benzene ring, on the contrary, shows an appreciable distortion from planarity ($\chi^2 = 27.39$), with atoms whose distance

to the least-squares plane is greater than three standard deviations. The average C-C bond length in the ring is 1.389 (7) Å and C-C-C angle is 120.0 (5)°. The whole bicycle is quasi-planar, the dihedral angle between the imidazole and benzene least-squares planes being 1.6 (2)°.

Table 1. Positional parameters ($\times 10^5$) and equivalent isotropic thermal parameters ($\times 10^4$) for the non-H atoms with e.s.d.'s in parentheses

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{eq}}(\text{\AA}^2)$
S	52527 (18)	10175 (13)	-23129 (6)	450 (4)
O(1)	-4101 (48)	-27848 (31)	-11181 (18)	501 (12)
N(1)	52808 (54)	3744 (34)	-10470 (19)	365 (12)
N(2)	73068 (55)	16914 (35)	-13045 (19)	365 (13)
C(1)	59232 (63)	10275 (44)	-15567 (22)	371 (14)
C(2)	62185 (64)	6115 (41)	-4873 (22)	351 (15)
C(3)	75071 (67)	14402 (41)	-6487 (23)	362 (15)
C(4)	86847 (76)	18824 (47)	-2034 (25)	475 (18)
C(5)	85689 (85)	14255 (53)	4187 (27)	549 (20)
C(6)	72880 (88)	5597 (52)	5804 (26)	542 (20)
C(7)	61161 (75)	1292 (45)	1287 (24)	430 (16)
C(8)	37658 (65)	-4036 (42)	-10742 (23)	372 (15)
C(9)	37768 (70)	-14158 (46)	-14696 (25)	432 (16)
C(10)	23521 (75)	-21796 (45)	-14697 (26)	447 (17)
C(11)	9032 (68)	-19503 (44)	-10830 (24)	394 (15)
C(12)	8414 (67)	-9217 (47)	-6999 (23)	411 (15)
C(13)	23083 (72)	-1425 (45)	-7045 (25)	404 (16)
C(14)	-20360 (76)	-25485 (55)	-7782 (31)	568 (21)
C(15)	-32549 (77)	-35714 (56)	-9505 (32)	577 (21)
C(16)	84187 (65)	24958 (46)	-16586 (23)	377 (15)
C(17)	101124 (72)	21513 (47)	-18020 (26)	465 (18)
C(18)	112177 (73)	29465 (54)	-21276 (28)	542 (20)
C(19)	105956 (72)	40733 (53)	-23039 (28)	536 (19)
C(20)	88844 (83)	43990 (52)	-21639 (32)	608 (22)
C(21)	77897 (77)	36129 (48)	-18433 (30)	531 (20)




Fig. 1. Bond lengths (\AA) and angles ($^\circ$) in the molecule. Standard deviations are below 0.009\AA and 0.6° .

* Lists of structure factors, anisotropic thermal parameters, H-atom coordinates and least-squares-planes' data have been deposited with the British Library Lending Division as Supplementary Publication No. SUP 38889 (12 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

The dihedral angles formed by the imidazole ring and the two bound phenyl rings are $73.7(2)^\circ$ for the N(2)-attached ring and $61.4(2)^\circ$ for the N(1)-attached ring, both values being similar to those encountered in the analogous compounds studied previously, ranging from 60 to 80° . The torsional angles describing the bonding are $C(1)-N(1)-C(8)-C(9) = 63.5(6)$ and $C(1)-N(2)-C(16)-C(21) = 75.2(7)^\circ$, respectively.

Phenyl rings. The N(2)-attached phenyl ring is planar, with atomic deviations from the mean plane within the standard deviations ($\chi^2 = 4.52$). The average C-C bond length is $1.385(8)$ Å and the mean C-C-C angle is $120.0(5)^\circ$. The N(1)-attached phenyl ring, on the contrary, deviates significantly from the expected planar conformation ($\chi^2 = 29.07$), with an average C-C bond length of $1.390(7)$ Å and C-C-C angle of $120.0(5)^\circ$.

The ethoxy group exhibits asymmetry in the $C(10)-C(11)-O(1) = 115.1(5)$ and $C(12)-C(11)-O(1) = 124.4(5)^\circ$ angles, a common feature of the phenoxy group (Domiano, Nardelli, Balsamo, Macchia & Macchia, 1979).

Fig. 2 shows a view of the unit cell along the b axis. No evidence for hydrogen bonds exists and no intermolecular contacts appreciably shorter than the sum of the van der Waals radii have been detected.

The authors thank Professor Fernández-Bolaños for supplying the crystals and the staff of the Instituto 'Rocasolano' of CSIC (Madrid) for collecting the diffractometer data. The present work forms part of a wider research project supported by the Government through the 'Comisión Asesora de Investigación Científica y Técnica'.




Fig. 2. Unit cell viewed along the b axis.

References

- CONDE, A., LÓPEZ-CASTRO, A. & MÁRQUEZ, R. (1978). *Rev. Iberoam. Cristalogr. Miner. Metalogen.* **1**, 23–26.
- CRİADO, A., CONDE, A. & MÁRQUEZ, R. (1983). *Acta Cryst.* **C39**, 122–125.
- DOMIANO, P., NARDELLI, M., BALSAMO, A., MACCHIA, B. & MACCHIA, F. (1979). *Acta Cryst.* **B35**, 1363–1372.
- FERNÁNDEZ-BOLAÑOS, J., FUENTES-MOTA, J. & FERNÁNDEZ-BOLAÑOS GUZMÁN, J. (1983a). *Carbohydr. Res.* To be published.
- FERNÁNDEZ-BOLAÑOS, J., FUENTES-MOTA, J. & FERNÁNDEZ-BOLAÑOS GUZMÁN, J. (1983b). *An. Quím.* To be published.
- HAMILTON, W. C. (1959). *Acta Cryst.* **12**, 609–610.
- International Tables for X-ray Crystallography (1974). Vol. IV. Birmingham: Kynoch Press.
- MAIN, P., HULL, S. E., LESSINGER, L., GERMAIN, G., DECLERCQ, J.-P. & WOOLFSON, M. M. (1978). *MULTAN78. A System of Computer Programs for the Automatic Solution of Crystal Structures from X-ray Diffraction Data*. Univs. of York, England, and Louvain, Belgium.
- STEWART, J. M., KUNDELL, F. A. & BALDWIN, J. C. (1970). The XRAY70 system. Computer Science Center, Univ. of Maryland, College Park, Maryland.
- TIPSON, R. S. & COHEN, A. (1965). *Carbohydr. Res.* **1**, 338–340.

Acta Cryst. (1984). **C40**, 190–193

2-(1,3-Dithiol-2-ylidene)-4-cyclopentene-1,3-dione, $C_8H_4O_2S_2$, and 2-(1,3-Benzodithiol-2-ylidene)cyclohexanone, $C_{13}H_{12}OS_2$

BY WILLIAM H. WATSON, JEAN GALLOY AND DAVID A. GROSSIE

FASTBIOS Laboratory, Department of Chemistry, Texas Christian University, Fort Worth, TX 76129, USA

AND JUZO NAKAYAMA

Department of Chemistry, Faculty of Science, Saitama University, Urawa, Saitama 338, Japan

(Received 22 July 1983; accepted 22 September 1983)

Abstract. $C_8H_4O_2S_2$ (2), $M_r = 196.25$, orthorhombic, $Pnma$, $a = 18.080(2)$, $b = 11.102(1)$, $c = 3.8904(4)$ Å, $V = 780.9(1)$ Å 3 , $Z = 4$, $D_x = 1.67$ Mg m $^{-3}$, $Cu K\alpha$, $\lambda = 1.54178$ Å, $\mu = 5.549$ mm $^{-1}$, $F(000) = 400$, room temperature, $R = 0.046$ based on 597 unique reflections. $C_{13}H_{12}OS_2$ (3), M_r

= 248.24, monoclinic, $P2_1/c$, $a = 6.459(1)$, $b = 14.263(3)$, $c = 13.514(4)$ Å, $\beta = 112.16(2)^\circ$, $V = 1153.0(6)$ Å 3 , $Z = 4$, $D_x = 1.43$ Mg m $^{-3}$, $Cu K\alpha$, $\mu = 3.90$ mm $^{-1}$, $F(000) = 520$, room temperature, $R = 0.074$ based on 1429 unique reflections. Compound (2) is planar except for a slight 2° fold along the S-S line.