

- EMSLEY, J. (1980). *Chem. Soc. Rev.* pp. 91–124, and references therein.
- FAIR, C. K. & SCHLEMPER, E. O. (1978). *Acta Cryst. B34*, 436–442.
- HADZI, D. (1965). *Pure Appl. Chem. 11*, 435–437.
- HANANIA, G. I. H. & IRVINE, D. H. (1962). *J. Chem. Soc. Dalton Trans.* pp. 2745–2749.
- HEDWIG, G. R. & POWELL, H. K. (1974). *J. Chem. Soc. Dalton Trans.* pp. 47–52.
- HUSSAIN, M. S. & HAQUE, M. (1983). *Acta Cryst. C39*, 292–294.
- HUSSAIN, M. S. & SCHLEMPER, E. O. (1979). *Inorg. Chem. 18*, 1116–1121, 2275–2282, and references therein.
- MAIN, P., HULL, S. E., LESSINGER, L., GERMAIN, G., DECLERCQ, J.-P. & WOOLFSON, M. M. (1978). *MULTAN78. A System of Computer Programs for the Automatic Solution of Crystal Structures from X-ray Diffraction Data*. Univs. of York, England, and Louvain, Belgium.
- SCHLEMPER, E. O., HUSSAIN, M. S. & MURMANN, R. K. (1981). *Acta Cryst. B37*, 234–237.
- SHELDICK, G. M. (1976). *SHELX76. Program for crystal structure determination*. Univ. of Cambridge, England.
- STEWART, R. F., DAVIDSON, E. R. & SIMPSON, W. T. (1965). *J. Chem. Phys. 42*, 3175–3187.
- VASSIAN, E. G. & MURMANN, R. K. (1967). *Inorg. Chem. 6*, 2043–2047.

Acta Cryst. (1984). **C40**, 816–818

Structures of the *cis* and *trans* Isomers of 2,4,6-Tri-*tert*-butyl-2,4,6-trifluorocyclotrisilazane, $C_{12}H_{30}F_3N_3Si_3$

BY WILLIAM CLEGG, GEORGE M. SHELDICK AND DIETMAR STALKE

Institut für Anorganische Chemie der Universität, Tammannstrasse 4, D-3400 Göttingen, Federal Republic of Germany

(Received 17 October 1983; accepted 19 December 1983)

Abstract. $M_r = 357.6$; *cis* isomer (1): trigonal, $P\bar{3}$, $a = 10.873$ (1), $c = 10.040$ (1) Å, $U = 1027.9$ Å 3 , $Z = 2$, $D_x = 1.155$ Mg m $^{-3}$, Mo $K\alpha$, $\lambda = 0.71069$ Å, $\mu = 0.25$ mm $^{-1}$, $F(000) = 384$, $T = 291$ K, $R = 0.054$ for 1232 observed reflections; *trans* isomer (2): monoclinic, $P2_1/n$, $a = 18.077$ (4), $b = 10.569$ (2), $c = 22.015$ (6) Å, $\beta = 104.67$ (3) $^\circ$, $U = 4069.0$ Å 3 , $Z = 8$, $D_x = 1.167$ Mg m $^{-3}$, $\mu = 0.25$ mm $^{-1}$, $F(000) = 1536$, $T = 291$ K, $R = 0.071$ for 4421 observed reflections. (1) has crystallographic threefold rotation symmetry and a virtually planar Si_3N_3 ring [r.m.s. deviation = 0.017 (2) Å]. The two crystallographically independent molecules of (2) lie in general positions and deviations from planarity of the ring are greater [r.m.s. $\Delta = 0.086$ (3) and 0.128 (3) Å].

Introduction. A number of chain and ring silazanes built up from $-HNSiBuF-$ units have recently been synthesized by controlled condensation reactions of $BuSiF_3$ and $LiNH_2$ [$Bu = tert$ -butyl, $(CH_3)_3Cl$] (Klingebiel & Vater, 1983). The cyclotrisilazane ($BuSiFNH$) $_3$ exists in two stable non-interconverting isomeric forms, the separation of which has been accomplished by fractional crystallization: the *cis* isomer is less soluble in non-polar solvents and can be selectively crystallized from a mixed solution in CH_2Cl_2 by addition of *n*-hexane.

We have investigated the structures of both isomers in order to assess the influence of the geometrical substituent distribution on the expected planarity of the Si_3N_3 ring.

Experimental. Crystals obtained from CH_2Cl_2/n -hexane (1) or *n*-hexane by cooling (2), $0.4 \times 0.5 \times 0.15$ mm (1), $0.3 \times 0.3 \times 0.45$ mm (2), mounted in capillaries, Stoe-Siemens AED diffractometer, unit-cell parameters from 2θ values of 36 (1) and 32 (2) reflections ($20 < 2\theta < 25^\circ$), 1826 reflections with $2\theta < 55^\circ$ and $h,k \geq 0$ (1) and 8179 reflections with $2\theta < 50^\circ$ and $k \geq 1$ (2), profile analysis (Clegg, 1981), no significant intensity variation for 3 standard reflections, no absorption corrections, $R_{int} = 0.027$ (1) and 0.037 (2), 1581 (1) and 7144 (2) unique reflections, 1232 (1) and 4421 (2) with $F > 4\sigma(F)$; automatic multisolution direct methods, blocked-cascade refinement on F , $w^{-1} = \sigma^2(F) + gF^2$, $g = 0.00068$ (1), 0.00041 (2), Bu H atoms constrained to give $C-H = 0.96$ Å, $H-C-H = 109.5^\circ$, $U(H) = 1.2U_{eq}(C)$, anisotropic thermal parameters for non-H atoms, no extinction corrections, scattering factors from *International Tables for X-ray Crystallography* (1974). (1): N–H refined freely, 77 parameters, $R = 0.054$, $wR = 0.068$, slope of normal probability plot = 1.51, max. $\Delta/\sigma = 0.013$, mean = 0.001, largest peak in final difference map = 0.25 e Å $^{-3}$, largest hole = -0.33 e Å $^{-3}$; (2): minor disorder component for one $BuSiF$ group with site occupation factor 0.119 (2) for Si and F; disorder of Bu group not resolved; N–H constrained to lie on Si–N–Si external bisector with $N-H = 0.62$ Å [value obtained from free refinement of (1)], $U(H) = 1.2U_{eq}(N)$, 398 parameters, $R = 0.071$, $wR = 0.073$, slope = 1.61, max. $\Delta/\sigma = 0.029$, mean = 0.005, largest peak = 0.63 e Å $^{-3}$, largest hole =

Table 1. Atomic coordinates ($\times 10^4$) and equivalent isotropic thermal parameters ($\text{\AA}^2 \times 10^4$) for (1)

	x	y	z	U
Si	6802 (1)	5046 (1)	6339 (1)	417 (3)
N	5283 (2)	3450 (2)	6374 (3)	482 (11)
H(1)	4753 (30)	3515 (32)	6374 (29)	491 (97)
F	6862 (2)	5818 (2)	4963 (2)	622 (9)
C(1)	6907 (3)	6310 (3)	7646 (3)	587 (14)
C(2)	5599 (4)	6492 (4)	7591 (4)	893 (26)
C(3)	6986 (5)	5766 (4)	9023 (3)	946 (27)
C(4)	8243 (6)	7746 (4)	7408 (6)	1341 (29)

 $U = \frac{1}{3}$ (trace of the orthogonalized U_{ij} matrix).

Table 2. Bond lengths (Å) and angles (°) for (1)

Si—F	1.602 (2)	Si—C(1)	1.862 (3)
N—Si	1.695 (2)	N—Si ^l	1.692 (3)
N—H(1)	0.615 (38)	C(1)—C(2)	1.532 (7)
C(1)—C(3)	1.523 (5)	C(1)—C(4)	1.529 (4)
N—Si—N ^{ll}	107.0 (2)	F—Si—N	108.5 (1)
F—Si—N ^{ll}	108.6 (1)	C(1)—Si—N	114.2 (1)
C(1)—Si—N ^{ll}	113.9 (1)	F—Si—C(1)	104.5 (1)
Si—N—Si ^l	133.0 (2)	Si—N—H(1)	111.9 (28)
Si ^l —N—H(1)	115.1 (28)	Si—C(1)—C(2)	110.6 (2)
Si—C(1)—C(3)	110.3 (3)	Si—C(1)—C(4)	108.7 (3)
C(2)—C(1)—C(3)	108.7 (3)	C(2)—C(1)—C(4)	109.4 (4)
C(3)—C(1)—C(4)	109.2 (3)		

Symmetry operators: (i) $1-y, x-y, z$; (ii) $1-x+y, 1-x, z$.

$-0.32 \text{ e \AA}^{-3}.$ * Programs: *SHELXTL* (Sheldrick, 1978), diffractometer control program by WC.

Discussion. Atomic coordinates, bond lengths and angles are given in Tables 1–4. The molecular structures are shown in Figs. 1–3. The more symmetrical *cis* isomer (1) forms much better quality crystals in a higher-symmetry space group; it was possible to refine freely the H atom attached to N. Thermal motion in (2) is rather greater (although the crystal density is slightly higher), and free refinement of six independent N—H groups proved unsuccessful. There is also disorder of one of the BuSiF groups in the second molecule (Fig. 3); derived geometry within and around this group is unreliable. It was not possible to resolve the disorder of the Bu group or of the neighbouring NH groups.

Corresponding bond lengths and angles for the two isomers are insignificantly different; for (1) and for (2) respectively, mean values for the most important parameters (ignoring disorder) are: Si—N 1.695 (2), 1.696 (7); Si—F 1.602 (2), 1.600 (5); Si—C 1.862 (3), 1.851 (9) Å; N—Si—N 107.0 (2), 106.5 (8); Si—N—Si 133.0 (2), 131.5 (22)°. They are also similar to those of the simple cyclotrisilazane *cis*-(PhSiFNMe)₃ (Clegg, Noltemeyer, Sheldrick & Vater, 1980). As in this case,

Table 3. Atomic coordinates ($\times 10^4$) and equivalent isotropic thermal parameters ($\text{\AA}^2 \times 10^4$) for (2)

	x	y	z	U
Si(1)	2385 (1)	-870 (1)	-459 (1)	587 (5)
F(1)	2186 (2)	-679 (3)	-1203 (1)	771 (11)
C(11)	2362 (3)	-2610 (5)	-333 (2)	707 (21)
C(12)	2921 (4)	-3277 (6)	-619 (4)	1317 (38)
C(13)	1566 (3)	-3106 (6)	-627 (4)	1276 (36)
C(14)	2558 (5)	-2890 (7)	353 (3)	1636 (47)
N(1)	1739 (2)	-35 (4)	-188 (2)	728 (16)
Si(2)	1829 (1)	1012 (1)	407 (1)	591 (5)
F(2)	1676 (2)	323 (3)	1007 (1)	915 (13)
C(21)	1117 (2)	2302 (4)	234 (2)	651 (19)
C(22)	309 (3)	1788 (6)	69 (3)	956 (27)
C(23)	1225 (4)	3053 (6)	-348 (3)	1260 (35)
C(24)	1230 (4)	3189 (6)	803 (3)	1476 (39)
N(2)	2751 (2)	1544 (4)	577 (2)	674 (16)
Si(3)	3517 (1)	1079 (1)	321 (1)	644 (5)
F(3)	4182 (1)	680 (3)	922 (1)	807 (11)
C(31)	3944 (3)	2339 (5)	-72 (2)	708 (20)
C(32)	4074 (3)	3547 (5)	327 (3)	1011 (27)
C(33)	4712 (3)	1897 (7)	-170 (3)	1137 (32)
C(34)	3404 (3)	2650 (6)	-712 (3)	1059 (28)
N(3)	3252 (2)	-221 (4)	-124 (2)	824 (18)
Si(4)	77 (1)	-82 (1)	7896 (1)	539 (5)
F(4)	330 (1)	-559 (3)	8606 (1)	761 (11)
C(41)	8 (3)	-1507 (5)	7393 (3)	715 (22)
C(42)	808 (3)	-2012 (6)	7401 (3)	1087 (31)
C(43)	-407 (4)	-1174 (7)	6705 (3)	1210 (33)
C(44)	-430 (3)	-2550 (6)	7623 (3)	1158 (34)
N(4)	-764 (2)	713 (3)	7797 (2)	610 (15)
Si(5)	-1068 (1)	2054 (1)	7395 (1)	543 (5)
F(5)	-1726 (1)	1724 (3)	6774 (1)	709 (11)
C(51)	-1516 (2)	3175 (4)	7834 (2)	620 (19)
C(52)	-1732 (4)	4408 (5)	7475 (3)	1153 (33)
C(53)	-2251 (3)	2612 (6)	7949 (3)	997 (29)
C(54)	-963 (3)	3461 (6)	8466 (3)	1180 (31)
N(5)	-321 (2)	2621 (4)	7143 (2)	725 (17)
Si(6)	608 (1)	2163 (2)	7243 (1)	586 (6)
F(6)	737 (2)	1668 (4)	6588 (1)	871 (15)
C(61)	1320 (3)	3439 (5)	7436 (2)	770 (22)
C(62)	1202 (3)	4389 (6)	6919 (3)	906 (29)
C(63)	2124 (3)	2908 (7)	7583 (3)	975 (30)
C(64)	1222 (4)	4167 (7)	8035 (3)	1181 (38)
N(6)	738 (2)	967 (4)	7781 (2)	704 (16)
Si(6x)	634 (5)	2675 (9)	7746 (4)	411 (35)
F(6x)	778 (12)	3526 (21)	8335 (10)	684 (90)

the *cis* isomer (1) contains a virtually planar Si₃N₃ ring, with a very slight puckering towards a chair conformation, and the electron-withdrawing fluorine substituents cause a reduction in Si—N bond length compared with other simple cyclotrisilazanes, such as (Bu₂SiNH)₃ [1.727 (2) Å] (Clegg, Sheldrick & Stalke, 1984).

There is no significant difference between the two crystallographically independent molecules of the *trans* isomer (2) (ignoring the disorder). Deviations from planarity of the Si₃N₃ rings are greater than in (1), but are still very small compared with those observed when bulky substituents are attached to N, as in (Me₂SiNSiMe₃)₃, which has a boat conformation (Adamson & Daly, 1970).

We may conclude that a planar Si₃N₃ ring is normal for cyclotrisilazanes, except when the steric bulk of the N substituents or ring fusion render this impossible (Clegg, Sheldrick & Stalke, 1984). Bulky Si substituents do not greatly affect this planar conformation, and crystal packing produces only very minor perturbations.

* Lists of structure factors, anisotropic thermal parameters and H-atom parameters have been deposited with the British Library Lending Division as Supplementary Publication No. SUP 39132 (66 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

We thank Dr U. Klingebiel and Dr N. Vater for supplying crystals, and the Verband der Chemischen Industrie for financial support.




Fig. 1. Molecular structure of the *cis* isomer (1). H atoms are omitted from the *tert*-butyl groups.




Fig. 2. Molecule 1 of the *trans* isomer (2). H atoms are omitted.




Fig. 3. Molecule 2 of the *trans* isomer (2), showing both disorder components. H atoms are omitted.

Table 4. Bond lengths (Å) and angles (°) for (2)

Si(1)–N(1)	1.689 (4)	Si(4)–N(4)	1.702 (4)
Si(1)–N(3)	1.698 (4)	Si(4)–N(6)	1.695 (4)
Si(2)–N(1)	1.691 (4)	Si(5)–N(4)	1.688 (4)
Si(2)–N(2)	1.707 (4)	Si(5)–N(5)	1.693 (4)
Si(3)–N(2)	1.695 (4)	Si(6)–N(5)	1.706 (4)
Si(3)–N(3)	1.685 (4)	Si(6)–N(6)	1.708 (4)
Si(1)–F(1)	1.598 (3)	Si(4)–F(4)	1.596 (3)
Si(2)–F(2)	1.591 (3)	Si(5)–F(5)	1.607 (2)
Si(3)–F(3)	1.602 (3)	Si(6)–F(6)	1.604 (4)
Si(1)–C(11)	1.862 (5)	Si(4)–C(41)	1.855 (5)
Si(2)–C(21)	1.848 (5)	Si(5)–C(51)	1.842 (5)
Si(3)–C(31)	1.859 (6)	Si(6)–C(61)	1.839 (5)
C(11)–C(12)	1.496 (9)	C(41)–C(42)	1.537 (8)
C(11)–C(13)	1.515 (7)	C(41)–C(43)	1.550 (7)
C(11)–C(14)	1.492 (8)	C(41)–C(44)	1.517 (9)
C(21)–C(22)	1.512 (6)	C(51)–C(52)	1.523 (7)
C(21)–C(23)	1.561 (9)	C(51)–C(53)	1.534 (8)
C(21)–C(24)	1.537 (8)	C(51)–C(54)	1.524 (7)
C(31)–C(32)	1.533 (8)	C(61)–C(62)	1.493 (8)
C(31)–C(33)	1.531 (8)	C(61)–C(63)	1.515 (7)
C(31)–C(34)	1.532 (7)	C(61)–C(64)	1.575 (9)
Si(6x)–N(5)	1.893 (9)	Si(6x)–N(6)	1.815 (10)
Si(6x)–F(6x)	1.546 (23)	Si(6x)–C(61)	1.756 (11)
F(1)–Si(1)–C(11)	105.6 (2)	F(4)–Si(4)–C(41)	106.7 (2)
F(1)–Si(1)–N(1)	107.5 (2)	F(4)–Si(4)–N(4)	107.9 (2)
C(11)–Si(1)–N(1)	114.9 (2)	C(41)–Si(4)–N(4)	113.7 (2)
F(1)–Si(1)–N(3)	109.7 (2)	F(4)–Si(4)–N(6)	108.4 (2)
C(11)–Si(1)–N(3)	112.7 (2)	C(41)–Si(4)–N(6)	112.6 (2)
N(1)–Si(1)–N(3)	106.3 (2)	N(4)–Si(4)–N(6)	107.4 (2)
N(1)–Si(2)–F(2)	109.9 (2)	N(4)–Si(5)–F(5)	109.6 (2)
N(1)–Si(2)–C(21)	113.4 (2)	N(4)–Si(5)–C(51)	113.1 (2)
F(2)–Si(2)–C(21)	104.5 (2)	F(5)–Si(5)–C(51)	104.9 (2)
N(1)–Si(2)–N(2)	106.4 (2)	N(4)–Si(5)–N(5)	106.8 (2)
F(2)–Si(2)–N(2)	109.6 (2)	F(5)–Si(5)–N(5)	106.2 (2)
C(21)–Si(2)–N(2)	113.1 (2)	C(51)–Si(5)–N(5)	116.0 (2)
N(2)–Si(3)–F(3)	107.9 (2)	N(5)–Si(6)–F(6)	109.8 (2)
N(2)–Si(3)–C(31)	114.7 (2)	N(5)–Si(6)–C(61)	115.5 (2)
F(3)–Si(3)–C(31)	105.1 (2)	F(6)–Si(6)–C(61)	101.2 (2)
N(2)–Si(3)–N(3)	107.0 (2)	N(5)–Si(6)–N(6)	104.9 (2)
F(3)–Si(3)–N(3)	108.0 (2)	F(6)–Si(6)–N(6)	110.8 (2)
C(31)–Si(3)–N(3)	113.9 (2)	C(61)–Si(6)–N(6)	114.6 (2)
Si(1)–N(1)–Si(2)	132.5 (2)	Si(4)–N(4)–Si(5)	130.0 (2)
Si(2)–N(2)–Si(3)	131.8 (2)	Si(5)–N(5)–Si(6)	134.3 (2)
Si(1)–N(3)–Si(3)	132.6 (3)	Si(4)–N(6)–Si(6)	127.5 (2)
Si(1)–C(11)–C(12)	111.1 (4)	Si(4)–C(41)–C(42)	110.7 (3)
Si(1)–C(11)–C(13)	109.4 (3)	Si(4)–C(41)–C(43)	109.9 (4)
C(12)–C(11)–C(13)	109.1 (5)	C(42)–C(41)–C(43)	108.2 (5)
Si(1)–C(11)–C(14)	109.7 (4)	Si(4)–C(41)–C(44)	110.6 (4)
C(12)–C(11)–C(14)	108.7 (5)	C(42)–C(41)–C(44)	108.3 (4)
C(13)–C(11)–C(14)	108.7 (6)	C(43)–C(41)–C(44)	109.1 (4)
Si(2)–C(21)–C(22)	111.4 (3)	Si(5)–C(51)–C(52)	111.5 (4)
Si(2)–C(21)–C(23)	108.6 (4)	Si(5)–C(51)–C(53)	110.6 (3)
C(22)–C(21)–C(23)	107.4 (4)	C(52)–C(51)–C(53)	107.3 (4)
Si(2)–C(21)–C(24)	109.9 (3)	Si(5)–C(51)–C(54)	109.5 (3)
C(22)–C(21)–C(24)	109.6 (5)	C(52)–C(51)–C(54)	109.1 (4)
C(23)–C(21)–C(24)	109.9 (5)	C(53)–C(51)–C(54)	108.7 (5)
Si(3)–C(31)–C(32)	110.6 (4)	Si(6)–C(61)–C(62)	111.1 (3)
Si(3)–C(31)–C(33)	110.2 (4)	Si(6)–C(61)–C(63)	110.8 (4)
C(32)–C(31)–C(33)	108.7 (4)	C(62)–C(61)–C(63)	110.8 (5)
Si(3)–C(31)–C(34)	109.8 (4)	Si(6)–C(61)–C(64)	109.3 (4)
C(32)–C(31)–C(34)	108.5 (4)	C(62)–C(61)–C(64)	106.6 (5)
C(33)–C(31)–C(34)	109.0 (5)	C(63)–C(61)–C(64)	107.9 (4)
N(5)–Si(6x)–N(6)	93.8 (4)	C(61)–Si(6x)–F(6x)	94.2 (10)
N(5)–Si(6x)–F(6x)	122.4 (10)	N(6)–Si(6x)–F(6x)	123.3 (10)
N(5)–Si(6x)–C(61)	110.5 (5)	N(6)–Si(6x)–C(61)	113.5 (6)
Si(5)–N(5)–Si(6x)	116.3 (4)	Si(4)–N(6)–Si(6x)	126.0 (4)

References

- ADAMSON, G. W. & DALY, J. J. (1970). *J. Chem. Soc. A*, pp. 2724–2728.
- CLEGG, W. (1981). *Acta Cryst. A* **37**, 22–28.
- CLEGG, W., NOLTEMEYER, M., SHELDICK, G. M. & VATER, N. (1980). *Acta Cryst. B* **36**, 2461–2462.
- CLEGG, W., SHELDICK, G. M. & STALKE, D. (1984). *Acta Cryst. C* **40**, 433–434.
- International Tables for X-ray Crystallography (1974). Vol. IV, pp. 99, 149. Birmingham: Kynoch Press.
- KLINGEBIEL, U. & VATER, N. (1983). *Chem. Ber.* **116**, 3277–3282.
- SHELDICK, G. M. (1978). SHELXTL. An Integrated System for Solving, Refining and Displaying Crystal Structures from Diffraction Data. Univ. of Göttingen.