

3.28 (7) Å; C(43)—H(432)···Pd 119 (3)°]. Short intermolecular contacts include the interaction H(442)···I [1-x, ½+y, ½-z] 2.90 (1); C(44)···I 3.926 (8) Å; C(44)—H(442)···I 158.8 (6)°.

A slight skeletal deformation of the benzene rings of the triphenylphosphine ligands in the title compound is observed. It may be described in terms of an average decrease of 2.3° from 120° of the endocyclic bond angle, opposite to the P atom. Domenicano, Vaciago & Coulson (1975) explain this in terms of hybridization effects and valence-shell electron-pair repulsions.

The X-ray data were kindly collected by Dr A. J. M. Duisenberg. Part of this work (ALS) was supported by the Netherlands Foundation for Chemical Research (SON) with financial aid from the Netherlands Organization for the Advancement of Pure Research (ZWO).

*Acta Cryst.* (1987). **C43**, 463–465

## Structure of {2,6-Bis[(dimethylamino)methyl]phenyl}iodoplatinum(II)

BY W. J. J. SMEETS, A. L. SPEK\* AND A. J. M. DUISENBERG

*Vakgroep Algemene Chemie, Afdeling Kristal- en Structuurchemie, University of Utrecht, Padualaan 8, 3508 TB Utrecht, The Netherlands*

AND J. A. M. VAN BEEK AND G. VAN KOTEN

*Anorganisch Chemisch Laboratorium, University of Amsterdam, Nieuwe Achtergracht 166, 1018 WV Amsterdam, The Netherlands*

(Received 2 October 1986; accepted 21 October 1986)

**Abstract.** [PtI(C<sub>12</sub>H<sub>19</sub>N<sub>2</sub>)],  $M_r = 513.28$ , monoclinic,  $I2/a$ ,  $a = 17.044$  (6),  $b = 9.259$  (3),  $c = 18.44$  (1) Å,  $\beta = 96.65$  (4)°,  $V = 2890$  (2) Å<sup>3</sup>,  $Z = 8$ ,  $D_x = 2.359$  g cm<sup>-3</sup>, Mo  $K\alpha$ ,  $\lambda = 0.71073$  Å,  $\mu = 119.1$  cm<sup>-1</sup>,  $F(000) = 1888$ ,  $T = 294$  K,  $R = 0.0301$  for 2635 observed reflections with  $I > 2\sigma(I)$ . Pt is square-planar (within 0.013 Å) coordinated by C(1), I, N(1) and N(2). The molecule has approximate twofold axial symmetry along the Pt—I axis.

**Introduction.** Recently it appeared that the title compound [=PtI(NCN)] readily forms a very stable complex with I<sub>2</sub> (van Beek, van Koten, Smeets & Spek, 1986). The resulting PtI(NCN)( $\eta^1$ -I<sub>2</sub>) compound is the first example of a complex mimicking the initial stage of the oxidative addition of halogens at d<sup>8</sup> metal centres.

- ### References
- BONDI, A. (1964). *J. Chem. Phys.* **68**, 441–451.  
 CROMER, D. T. & LIBERMAN, D. (1970). *J. Chem. Phys.* **53**, 1891–1898.  
 CROMER, D. T. & MANN, J. B. (1968). *Acta Cryst.* **A24**, 321–324.  
 DOMENICANO, A., VACIAGO, A. & COULSON, C. A. (1975). *Acta Cryst.* **B31**, 221–234.  
 KOBAYASHI, Y., IITAKA, Y., YAMAZAKI, H. (1972). *Acta Cryst.* **B28**, 899–906.  
 MCCANDLISH, L. E., STOUT, G. H. & ANDREWS, L. C. (1975). *Acta Cryst.* **A31**, 245–249.  
 NEGISHI, E. (1978). *Aspects of Mechanism in Organometallic Chemistry*, edited by J. H. BREWSTER, pp. 283–316. New York: Plenum.  
 OLMSTEAD, M. M., FARR, J. P. & BALCH, A. L. (1981). *Inorg. Chim. Acta*, **52**, 47–53.  
 SHELDRICK, G. M. (1976). *SHELX76*. Program for crystal structure determination. Univ. of Cambridge, England.  
 SPEK, A. L. (1982). *The EUCLID Package*. In *Computational Crystallography*, edited by D. SAYRE, p. 528. Oxford: Clarendon Press.  
 WALKER, N. & STUART, D. (1983). *Acta Cryst.* **A39**, 158–166.

PtI(NCN)( $\eta^1$ -I<sub>2</sub>) has a distorted square-pyramidal coordination with an end-on bonded I<sub>2</sub> molecule occupying the apical position. Two of the NMe groups are located on the opposite side of the  $\eta^1$  bonded I<sub>2</sub> molecule and seem to shield that side from further attack by I<sub>2</sub>.

In order to study this aspect in more detail the structure of the parent compound PtI(NCN) was determined.

**Experimental.** Transparent yellow prismatic crystal 0.48 × 0.29 × 0.13 mm glued on top of a glass fibre. Enraf–Nonius CAD-4F diffractometer, Zr-filtered Mo  $K\alpha$  radiation, cell constants from the angular settings of 12 reflections with 11.5 <  $\theta$  < 14.0°. Intensity data of 3512 unique reflections were collected within a quarter of the reflection sphere;  $-21 \leq h \leq 22$ ,  $0 \leq k \leq 12$ ,  $-23 \leq l \leq 0$ ;  $(\sin\theta/\lambda)_{\max} = 0.649$  Å<sup>-1</sup>;

\* Author to whom correspondence should be addressed.

$\omega-2\theta$  scan mode with  $\Delta\omega = (0.55 + 0.35\tan\theta)^\circ$ . Three reference reflections [400 (e.s.d. = 0.8%), 040 (e.s.d. = 0.7%), 004 (e.s.d. = 1.0%)], measured every hour showed a linear decay of 1.6% during the 67.1 h of X-ray exposure time. The intensities were corrected for this decay, Lorentz and polarization effects and for absorption using a Gaussian integration method (grid:  $8 \times 6 \times 12$ ; corrections in the range 2.728 to 6.703), resulting in 2636 observed reflections with  $I > 2\sigma(I)$ . Standard deviations in the intensities were increased according to an analysis of the variance of the reference reflections:  $\sigma^2(I) = \sigma^2(I)_{cs} + (0.017I)^2$  (McCandlish, Stout & Andrews, 1975). Space group  $I2/a$  derived from systematic absences [ $hkl$  ( $h + k + l = 2n + 1$ ) and  $h0l$  ( $h, l = 2n + 1$ )] and discriminated from  $Ia$  during the structure determination process. Structure solved by Patterson (*SHELX84*; Sheldrick, 1984) and subsequent difference Fourier methods. H atoms introduced on calculated positions ( $C-H = 0.98 \text{ \AA}$ ) and included in weighted, blocked full-matrix refinement (on  $F$ ) riding on their carrier atom. All non-H atoms were refined with anisotropic thermal parameters, H atoms refined with one isotropic temperature factor. The 002 reflections with a  $\Delta F/\sigma(F)$  ratio of 24.9 was excluded from the final refinement cycles. Convergence with 159 parameters was reached at  $R = 0.0301$ ,  $wR = 0.0310$ ,  $w = 1/\sigma^2(F)$ ,  $S = 1.90$ ,  $(\Delta/\sigma)_{\max} = 0.094$ , max. and min. residual densities 0.88 and  $-1.46 \text{ e \AA}^{-3}$  near the heavy atoms. The crystal exhibited some secondary extinction for which the  $F_c$  values were corrected by refinement of an empirical isotropic extinction parameter  $X[=1.0(1) \times 10^{-8}]$ :  $F_c' = F_c[1 - (XF^2/\sin\theta)]$ .

Scattering factors from Cromer & Mann (1968); anomalous-dispersion factors from Cromer & Liberman (1970). Calculations performed with *SHELX76* (Sheldrick, 1976) and the *EUCLID* package (geometry calculations and illustrations) (Spek, 1982) on the CDC Cyber-855 of the University of Utrecht.

**Discussion.** Fractional atomic coordinates and equivalent isotropic thermal parameters for the non-hydrogen atoms are given in Table 1.\* The molecular structure with the adopted atom numbering is shown in Fig. 1. Bond distances and angles are listed in Table 2.

The monoclinic unit cell contains eight discrete molecules of the title compound. The Pt atom has a distorted square-planar coordination by N(1), N(2), I and C(1). These atoms are coplanar with Pt within 0.013 Å. The distortion from regular square-planar coordination is illustrated by the deviation of the

\* Lists of structure factors, anisotropic thermal parameters, H-atom parameters, torsion angles and geometrical data concerning H atoms have been deposited with the British Library Document Supply Centre as Supplementary Publication No. SUP 43506 (25 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

Table 1. Final atomic coordinates and equivalent isotropic thermal parameters with e.s.d.'s in parentheses

$$U_{eq} = \frac{1}{3} \sum_i \sum_j U_{ij} a_i^* a_j^* a_i a_j$$

	x	y	z	$U_{eq}(\text{\AA}^2)$
Pt	0.16958 (1)	0.61977 (3)	0.37491 (1)	0.0369 (1)
I	0.07875 (3)	0.37817 (7)	0.37017 (3)	0.0746 (2)
N(1)	0.2762 (3)	0.5235 (6)	0.3541 (3)	0.041 (2)
N(2)	0.0840 (3)	0.7687 (6)	0.3981 (3)	0.047 (2)
C(1)	0.2339 (3)	0.7921 (7)	0.3788 (3)	0.041 (2)
C(2)	0.2055 (4)	0.9184 (7)	0.4033 (4)	0.049 (2)
C(3)	0.2519 (4)	1.0418 (8)	0.4051 (4)	0.060 (3)
C(4)	0.3270 (4)	1.0342 (9)	0.3849 (5)	0.065 (3)
C(5)	0.3562 (4)	0.9064 (8)	0.3596 (4)	0.061 (3)
C(6)	0.3089 (3)	0.7835 (8)	0.3558 (3)	0.047 (2)
C(7)	0.3266 (3)	0.6362 (8)	0.3254 (4)	0.050 (2)
C(8)	0.2691 (4)	0.4035 (8)	0.3015 (4)	0.064 (3)
C(9)	0.3144 (3)	0.4697 (8)	0.4255 (4)	0.049 (2)
C(10)	0.1260 (4)	0.9065 (8)	0.4305 (4)	0.063 (3)
C(11)	0.0298 (4)	0.7200 (9)	0.4506 (4)	0.061 (3)
C(12)	0.0369 (4)	0.808 (1)	0.3289 (4)	0.077 (3)

Table 2. Bond distances (Å) and angles (°) for the non-hydrogen atoms

Pt-I	2.716 (2)	N(2)-C(12)	1.472 (9)
Pt-N(1)	2.099 (5)	C(1)-C(2)	1.363 (9)
Pt-N(2)	2.087 (5)	C(1)-C(6)	1.395 (7)
Pt-C(1)	1.933 (6)	C(2)-C(10)	1.50 (1)
N(1)-C(7)	1.487 (9)	C(2)-C(3)	1.39 (1)
N(1)-C(8)	1.471 (9)	C(3)-C(4)	1.38 (1)
N(1)-C(9)	1.486 (9)	C(4)-C(5)	1.39 (1)
N(2)-C(10)	1.549 (9)	C(5)-C(6)	1.39 (1)
N(2)-C(11)	1.484 (9)	C(6)-C(7)	1.52 (1)
I-Pt-N(1)	98.4 (2)	C(10)-N(2)-C(12)	108.2 (6)
I-Pt-N(2)	98.1 (2)	C(11)-N(2)-C(12)	108.8 (5)
I-Pt-C(1)	179.6 (2)	Pt-C(1)-C(2)	119.9 (4)
N(1)-Pt-N(2)	163.4 (2)	Pt-C(1)-C(6)	118.5 (5)
N(1)-Pt-C(1)	81.8 (2)	C(2)-C(1)-C(6)	121.5 (6)
N(2)-Pt-C(1)	81.6 (2)	C(1)-C(2)-C(10)	114.6 (6)
Pt-N(1)-C(7)	108.4 (4)	C(1)-C(2)-C(3)	119.4 (6)
Pt-N(1)-C(8)	115.8 (4)	C(3)-C(2)-C(10)	125.9 (6)
Pt-N(1)-C(9)	106.3 (4)	C(2)-C(3)-C(4)	119.8 (7)
C(7)-N(1)-C(8)	107.6 (5)	C(3)-C(4)-C(5)	121.1 (7)
C(7)-N(1)-C(9)	109.8 (5)	C(4)-C(5)-C(6)	119.1 (6)
C(8)-N(1)-C(9)	109.0 (5)	C(1)-C(6)-C(5)	118.9 (6)
Pt-N(2)-C(10)	108.7 (4)	C(1)-C(6)-C(7)	113.1 (6)
Pt-N(2)-C(11)	115.8 (4)	C(5)-C(6)-C(7)	127.9 (5)
Pt-N(2)-C(12)	108.0 (4)	N(1)-C(7)-C(6)	110.6 (5)
C(10)-N(2)-C(11)	107.1 (5)	N(2)-C(10)-C(2)	109.0 (6)




Fig. 1. *PLUTO* (*EUCLID* version: Spek, 1982) drawing of the title compound with the adopted atom numbering.

relevant angles from 90 or 180°: N(1)—Pt—N(2) and I—Pt—C(1) being 163.4 (2) and 179.6 (2)° respectively with a range of 81.6 (2)—98.4 (2)° for the four remaining angles. The angle between the coordination plane and the phenyl ring is 9.5 (3)°. The molecule shows approximate twofold symmetry along the line through Pt and C(1); this is illustrated by the  $\Delta C_2$ [Pt—C(1)] asymmetry parameter for the eight-membered ring C(1)—C(2)—C(10)—N(2)—Pt—N(1)—C(7)—C(6) that amounts to 4.2 (6)° (Duax, Weeks & Rohrer, 1976). The molecular twofold axis is also present in the octahedral PtCl<sub>3</sub>(NCN) complex (Terheijden, van Koten, de Booy, Ubbels & Stam, 1983) in which the angle between the coordination plane and the phenyl ring is 16.3 (5)° and the  $\Delta C_2$ [Pt—C(1)] parameter is 4.9 (7)°. Some structural features of PtI(NCN) are comparable with those of the PtI(NCN)( $\eta^1$ -I<sub>2</sub>) complex (van Beek, van Koten, Smeets & Spek, 1986) which has a distorted square-pyramidal coordination with an I<sub>2</sub> molecule occupying the apical position, whereas the other coordinating atoms are the same as in the present structure. The geometry of the NCN ligand compares favourably with the geometries found in the studies of PtCl<sub>3</sub>(NCN) and PtI(NCN)( $\eta^1$ -I<sub>2</sub>). The Pt—C(1) distances are nearly equal: 1.933 (6) Å in the present PtI complex and 1.937 (9) Å in the PtI( $\eta^1$ -I<sub>2</sub>) structure. The Pt—I distance and the averages of the Pt—N distances are slightly different; 2.716 (2) and 2.093 (6) Å respectively in PtI(NCN) and 2.727 (1), 2.111 (8) Å in PtI(NCN)( $\eta^1$ -I<sub>2</sub>).

Characteristic differences between the PtI and PtI( $\eta^2$ -I<sub>2</sub>) complexes include the position of the Pt atom with respect to the coordination plane [I(1), N(1), N(2), C(1)] which is included in the plane in the present PtI compound and lifted 0.18 (2) Å out of this plane in the

square-pyramidal PtI( $\eta^1$ -I<sub>2</sub>) complex. The molecular symmetry of PtI( $\eta^1$ -I<sub>2</sub>) is also different, the molecule having an approximate mirror plane through the Pt—I—C(1) atoms, perpendicular to the phenyl ring. The different molecular symmetry can also be seen from the signs of the C—C—C—N torsion angles with values of -23.0 (7) and -20.5 (8)° for N(1) and N(2) in PtI(NCN) and -26 (1), 25 (1)° for the corresponding angles in the PtI( $\eta^1$ -I<sub>2</sub>) complex, indicating the different conformations of the side chains.

The investigation was supported in part (WJJS and ALS) by the Netherlands Foundation for Chemical Research (SON) with financial aid from the Netherlands Organization for Advancement of Pure Research (ZWO).

#### References

- BEEK, J. A. M. VAN, VAN KOTEN, G., SMEETS, W. J. J. & SPEK, A. L. (1986). *J. Am. Chem. Soc.* **108**, 5010–5011.  
 CROMER, D. T. & LIBERMAN, D. (1970). *J. Chem. Phys.* **53**, 1891–1898.  
 CROMER, D. T. & MANN, J. B. (1968). *Acta Cryst.* **A24**, 321–324.  
 DUAX, W. L., WEEKS, C. M. & ROHRER, D. C. (1976). *Topics in Stereochemistry*, Vol. 9, edited by E. L. ELIEL & N. ALLINGER, pp. 271–383. New York: John Wiley.  
 MCCANDLISH, L. E., STOUT, G. H., ANDREWS, L. C. (1975). *Acta Cryst.* **A31**, 245–249.  
 SHELDRIK, G. M. (1976). *SHELX76*. Program for crystal structure determination. Univ. of Cambridge, England.  
 SHELDRIK, G. M. (1984). *SHELX84*. Program for crystal structure determination. Univ of Göttingen, Federal Republic of Germany.  
 SPEK, A. L. (1982). *The EUCLID Package*. In *Computational Crystallography*, edited by D. SAYRE, p. 528. Oxford: Clarendon Press.  
 TERHEIJDEN, J., VAN KOTEN, G., DE BOOYS, J. L., UBBELS, H. J. C. & STAM, C. H. (1983). *Organometallics*, **2**, 1882–1883.

*Acta Cryst.* (1987). **C43**, 465–470

## Structure of the Organic Semiconducting Radical Cation Salt Bis[2,3:6,7-bis(methylenedioxydibenzofuran)]\* Hexafluoroarsenate

BY GUNNAR OLOVSSON

*Institute of Chemistry, University of Uppsala, Box 531, S-751 21 Uppsala, Sweden*

(Received 12 August 1986; accepted 21 October 1986)

**Abstract.** (C<sub>14</sub>H<sub>8</sub>O<sub>5</sub>)<sub>2</sub><sup>+</sup>.AsF<sub>6</sub><sup>-</sup>, *M<sub>r</sub>* = 701.33, monoclinic, *C*2/*c*, *a* = 9.564 (1), *b* = 13.899 (2), *c* = 19.364 (3) Å,  $\beta$  = 92.63 (1)°, *V* = 2571.4 (6) Å<sup>3</sup>, *Z* = 4, *D<sub>x</sub>* = 1.81 g cm<sup>-3</sup>, *Mo K $\alpha$* ,  $\lambda$  = 0.71069 Å,  $\mu$  = 15.1 cm<sup>-1</sup>, *F*(000) = 1404, *T* = 298 K. Final *R* = 0.035 for 2047

unique reflexions. Dimers consisting of (bMDODBF)<sub>2</sub><sup>+</sup> molecules (C<sub>14</sub>H<sub>8</sub>O<sub>5</sub> = bMDODBF) form stacks along the [110] and [1 $\bar{1}$ 0] directions and are surrounded by AsF<sub>6</sub><sup>-</sup> ions. Interplanar distances are 3.19 (2) Å within the dimers and 3.25 (2) Å between the dimers in the stacks. Adjacent dimers have poor molecular overlap, giving rise to semiconducting electronic properties.

\* 1',3'-Benzodioxolo[5',6':3,2]benzofuro[6,5-*d*][1,3]dioxole.