

**Structure of nucleoside analogues 3'-fluoro-2',3'-dideoxyuridine, 3'-fluoro-2',3'-dideoxy-5-bromouridine and 3'-azido-2',3'-dideoxy-5-bromouridine. Erratum.**

By D. H. EVERAERT, O. M. PEETERS, N. M. BLATON and C. J. DE RANTER, *Laboratorium voor Analytische Chemie en Medicinale Fysicochemie, Instituut voor Farmaceutische Wetenschappen, Katholieke Universiteit Leuven, Van Evenstraat 4, B-3000 Leuven, Belgium*, and A. VAN AERSCHOT and P. HERDEWIJN, *Laboratorium voor Farmaceutische Chemie, Rega Institute for Medical Research, Katholieke Universiteit Leuven, B-3000 Leuven, Belgium*

(Received 10 May 1991; accepted 21 May 1991)

**Abstract**

According to the IUPAC-IUB Joint Commission on Biochemical Nomenclature [*Pure Appl. Chem.* (1983), **55**, 1273–1280] guidelines, for compound (III) [Everaert, Peeters, Blaton, De Ranter, Van Aerschot & Herdewijn (1991). *Acta Cryst.* C47, 898–902] the C4'–C5' conforma-

tion for molecule *A* is *ap* (antiperiplanar) with  $\gamma = 172\cdot1(1)^\circ$  instead of *+sc* with  $\gamma = 172\cdot1(1)^\circ$  and for molecule *B* *+sc* with  $\gamma = 50\cdot5(7)^\circ$  instead of *ap* (antiperiplanar) with  $\gamma = 50\cdot5(7)^\circ$ .

---

All relevant information is given in the *Abstract*.

0108-2701/91/081776-01\$03.00

© 1991 International Union of Crystallography