

| | | | | |
|-------|-------------|--------------|-------------|-----------|
| F(7) | 0.0484 (7) | 0.2704 (14) | 0.1984 (3) | 0.169 (5) |
| F(8) | 0.1154 (6) | 0.5331 (12) | 0.1688 (4) | 0.142 (4) |
| F(9) | -0.0181 (6) | 0.3792 (13) | 0.1217 (3) | 0.143 (4) |
| F(10) | 0.2162 (5) | -0.3761 (8) | 0.0532 (3) | 0.096 (3) |
| F(11) | 0.2402 (6) | -0.2066 (10) | -0.0195 (3) | 0.103 (3) |
| F(12) | 0.0653 (5) | -0.2054 (9) | 0.0188 (3) | 0.104 (3) |
| O(1) | 0.5896 (5) | -0.0043 (8) | 0.1110 (2) | 0.044 (2) |
| O(2) | 0.5306 (5) | 0.3248 (8) | 0.1754 (2) | 0.045 (2) |
| O(3) | 0.2924 (5) | 0.2937 (8) | 0.1415 (2) | 0.049 (2) |
| O(4) | 0.3501 (5) | -0.0234 (8) | 0.0710 (2) | 0.048 (2) |
| N(1) | 0.4764 (5) | 0.3744 (10) | 0.0449 (2) | 0.038 (3) |
| C(1) | 0.6859 (8) | 0.0008 (12) | 0.1442 (4) | 0.043 (3) |
| C(2) | 0.7119 (8) | 0.1277 (13) | 0.1877 (3) | 0.050 (3) |
| C(3) | 0.6327 (9) | 0.2784 (13) | 0.2006 (3) | 0.047 (4) |
| C(4) | 0.7832 (10) | -0.1578 (16) | 0.1322 (5) | 0.066 (4) |
| C(5) | 0.6640 (11) | 0.4042 (19) | 0.2519 (5) | 0.083 (5) |
| C(6) | 0.1850 (9) | 0.2410 (13) | 0.1264 (3) | 0.045 (3) |
| C(7) | 0.1490 (8) | 0.0866 (13) | 0.0899 (4) | 0.051 (3) |
| C(8) | 0.2341 (9) | -0.0342 (13) | 0.0656 (3) | 0.044 (3) |
| C(9) | 0.0808 (10) | 0.3608 (20) | 0.1530 (5) | 0.078 (5) |
| C(10) | 0.1855 (10) | -0.2064 (16) | 0.0295 (5) | 0.066 (5) |
| C(11) | 0.5066 (7) | 0.5632 (13) | 0.0523 (3) | 0.043 (3) |
| C(12) | 0.4693 (7) | 0.3124 (11) | -0.0082 (4) | 0.041 (3) |

Table 2. Geometric parameters (Å, °)

| | | | |
|-------------------|------------|-------------------|------------|
| Cu(1)—O(1) | 1.963 (5) | Cu(1)—O(2) | 1.938 (5) |
| Cu(1)—O(3) | 1.956 (6) | Cu(1)—O(4) | 1.946 (5) |
| Cu(1)—N(1) | 2.248 (6) | O(1)—C(1) | 1.267 (10) |
| O(2)—C(3) | 1.263 (10) | O(3)—C(6) | 1.243 (11) |
| O(4)—C(8) | 1.249 (11) | N(1)—C(11) | 1.327 (11) |
| N(1)—C(12) | 1.326 (10) | C(1)—C(2) | 1.362 (12) |
| C(1)—C(4) | 1.536 (13) | C(2)—C(3) | 1.373 (12) |
| C(3)—C(5) | 1.508 (15) | C(6)—C(7) | 1.398 (12) |
| C(6)—C(9) | 1.546 (15) | C(7)—C(8) | 1.377 (13) |
| C(8)—C(10) | 1.521 (13) | C(11)—C(12A) | 1.378 (12) |
| C(12)—C(11A) | 1.378 (12) | | |
| O(1)—Cu(1)—O(2) | 91.3 (2) | O(1)—Cu(1)—O(3) | 163.9 (2) |
| O(2)—Cu(1)—O(3) | 85.9 (2) | O(1)—Cu(1)—O(4) | 87.4 (2) |
| O(2)—Cu(1)—O(4) | 168.5 (2) | O(3)—Cu(1)—O(4) | 92.1 (2) |
| O(1)—Cu(1)—N(1) | 99.1 (2) | O(2)—Cu(1)—N(1) | 96.8 (2) |
| O(3)—Cu(1)—N(1) | 97.0 (2) | O(4)—Cu(1)—N(1) | 94.7 (2) |
| Cu(1)—O(1)—C(1) | 124.5 (5) | Cu(1)—O(2)—C(3) | 125.0 (5) |
| Cu(1)—O(3)—C(6) | 124.6 (5) | Cu(1)—O(4)—C(8) | 125.7 (5) |
| Cu(1)—N(1)—C(11) | 122.8 (5) | Cu(1)—N(1)—C(12) | 121.6 (5) |
| C(11)—N(1)—C(12) | 115.6 (7) | O(1)—C(1)—C(2) | 128.0 (8) |
| O(1)—C(1)—C(4) | 114.2 (7) | C(2)—C(1)—C(4) | 117.8 (8) |
| C(1)—C(2)—C(3) | 122.1 (8) | O(2)—C(3)—C(2) | 127.9 (8) |
| O(2)—C(3)—C(5) | 113.2 (8) | C(2)—C(3)—C(5) | 118.9 (8) |
| O(3)—C(6)—C(7) | 128.0 (8) | O(3)—C(6)—C(9) | 114.5 (8) |
| C(7)—C(6)—C(9) | 117.5 (8) | C(6)—C(7)—C(8) | 122.3 (8) |
| O(4)—C(8)—C(7) | 127.2 (8) | O(4)—C(8)—C(10) | 114.5 (8) |
| C(7)—C(8)—C(10) | 118.2 (8) | N(1)—C(11)—C(12A) | 122.8 (7) |
| N(1)—C(12)—C(11A) | 121.6 (7) | | |

Acknowledgement is made to the Donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this research, as well as the Welch Foundation. We wish to thank F. Cervantes-Lee for collecting the intensity data.

Lists of structure factors, anisotropic thermal parameters, H-atom coordinates and geometry involving fluorine have been deposited with the British Library Document Supply Centre as Supplementary Publication No. SUP 71112 (8 pp.). Copies may be obtained through The Technical Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England. [CIF reference: HH1035]

References

- Belford, R. C. E., Fenton, D. E. & Truter, M. R. (1974). *J. Chem. Soc. Dalton Trans.* pp. 17–24.

- Dariet, J., Haddad, M. S., Duesler, E. N. & Hendrickson, D. N. (1979). *Inorg. Chem.* **18**, 2679–2682.
Haddad, M. S., Hendrickson, D. N., Cannady, J. P., Drago, R. S. & Bielska, D. S. (1979). *J. Am. Chem. Soc.* **101**, 898–906.
Kaim, W. (1983). *Angew. Chem. Int. Ed. Engl.* **22**, 171–258.
Losee, D. B., Richardson, H. W. & Hatfield, W. E. (1973). *J. Chem. Phys.* **59**, 3600–3603.
Richardson, H. W. & Hatfield, W. E. (1976). *J. Am. Chem. Soc.* **98**, 835–839.
Santoro, A., Mighell, A. D. & Reimann, C. W. (1970). *Acta Cryst.* **B26**, 979–984.
Sheldrick, G. M. (1990). *SHELXTL-Plus. An Integrated System for Solving, Refining and Displaying Crystal Structures from Diffraction Data.* Univ. of Göttingen, Germany.
Valentine, J. S., Silverstein, A. J. & Soos, Z. G. (1974). *J. Am. Chem. Soc.* **96**, 97–103.

Acta Cryst. (1993). **C49**, 1489–1492

The Monopotassium Salt of the Ligand of a Ziegler–Natta Catalyst

WILLIAM P. SCHAEFER,* W. DONALD COTTER AND JOHN E. BERCAW

Division of Chemistry and Chemical Engineering† and The Beckman Institute, Mail Code 139-74, California Institute of Technology, Pasadena, California 91125, USA

(Received 14 October 1993; accepted 16 February 1993)

Abstract

The anion of dipotassium (*tert*-butylamino)-dimethyl(tetramethylcyclopentadienyl)silanediide can coordinate to transition metals to form precursors for single-component Ziegler–Natta polymerization catalysts. The related monoanion derivative potassium (*tert*-butylamino)dimethyl(tetramethylcyclopentadienyl)silanide tetrahydrofuran solvate has been isolated as the first crystalline salt of this important ligand family. The tetramethylcyclopentadienyl group in the crystal has a potassium ion on each face, 2.8 Å from the center of the five-membered ring; each K atom is bonded to two rings and one tetrahydrofuran solvent molecule.

Comment

The organoscandium complexes $[(C_5Me_4)SiMe_2(NCMe_3)\{ScR\}_2(PMe_3)_x]$ ($R = H, CH_2CH_2CH_3$; $x = 2, 0$) are the first compounds which catalyze the homogeneous Ziegler–Natta polymerization of

† Contribution No. 8735

α -olefins without activation by a co-catalyst, such as alkylaluminum reagents or 'methylaluminumoxane' (Shapiro, Bunel, Schaefer & Bercaw, 1990). The synthesis of these complexes is hampered by retention of nonstoichiometric amounts of LiCl and tetrahydrofuran (THF) in the precursor $[(C_5Me_4)SiMe_2(NCMe_3)ScCl]_x-(LiCl)_y(THF)_z$, formed by treatment of $ScCl_3(THF)_3$ with the dilithium reagent $Li_2[(C_5Me_4)SiMe_2(NCMe_3)]$ in toluene. We have found that the dipotassium reagent $K_2[(C_5Me_4)SiMe_2(NCMe_3)]$ reacts with $ScCl_3(THF)_3$ in THF yielding cleanly $[(C_5Me_4)SiMe_2(NCMe_3)ScCl](THF)$, which is a suitable precursor for the preparation of the catalytic complexes (Cotter & Bercaw, 1993). However, the preparation of $K_2[(C_5Me_4)SiMe_2(NCMe_3)]$ itself is problematic; frequently the monopotassium derivative $K[(C_5Me_4)SiMe_2(N(H)CMe_3)]$ is obtained. In contrast to the lithium derivative, these potassium salts are highly crystalline and soluble in warm donor solvents.

In a dry box, $H_2[(C_5Me_4)SiMe_2(NCMe_3)]$ (26.06 g, 0.104 mol) was dissolved in THF (250 ml). KH (8.30 g, 0.207 mol) was added to the solution. The resulting slurry was refluxed overnight under argon on a Schlenk line. THF was removed *in vacuo*. The crude solid product was transferred to a large frit assembly and rinsed with petroleum ether, leaving behind a white powdery solid. A portion (10.81 g) of this material was purified *via* Soxhlet extraction in THF, with a great reduction in yield (5.48 g). Elemental analysis of this material is consistent with the formulation $K[(C_5Me_4)SiMe_2(N(H)CMe_3)]$ [calc. (found): C 62.87 (61.53); H 9.75 (9.52); N 4.84% (4.99%)]. Colorless rod-shaped crystals of this material were obtained by slow cooling of a hot THF solution and were placed in capillaries in a dry box. Outside a sealed environment, these crystals are sensitive to solvent loss, visibly losing crystallinity within minutes of their removal from the crystallization flask.

The asymmetric unit of this structure is shown in Fig. 1. Bond distances and angles are generally within normal ranges; the Si—C(cyclopentadienyl) bond is 1.827 (4) Å, much shorter than the Si—CH₃ bonds [average 1.873 (6) Å, compared to 1.86 Å expected (Allen, Kennard, Watson, Brammer, Orpen & Taylor, 1987)] and much shorter than others we have measured recently [average 1.867 (14) Å (Marsh, Schaefer, Coughlin & Bercaw, 1992; Schaefer, Köhn & Bercaw, 1992)]. The cyclopentadienyl ring shows evidence of localized double bonds, in that the C—C bonds to C1 average 1.426 (6) Å, C2—C3 and C4—C5 average 1.385 (16) and C3—C4 is 1.410 (7) Å. Thus, C1 has less double-bond character in its Cp ring bonds and could form a stronger bond to Si, as observed.




Fig. 1. An ORTEP drawing with 50% probability ellipsoids of one asymmetric unit showing the numbering system used. H atoms are not shown.

The ligand molecules are joined in chains by the potassium ions (Fig. 2). Each potassium ion is bonded to two cyclopentadienyl groups, and each cyclopentadienyl ring has a potassium ion on each side of it. The centroid of the five ring-carbon atoms in the cyclopentadienyl group, Cp*, is 2.81 Å from K⁺ on one side and 2.78 Å from another K⁺ on the other. The tetrahydrofuran solvent is attached to the potassium ion through oxygen: K—O 2.762 (5) Å. The two Cp* centers and one oxygen form a planar array about K⁺, with angles Cp*—K—O 108.4 and 106.1°; Cp*—K—Cp*' is 145.5°. The angle K—Cp*—K' is 176.2°, with the plane of the cyclopentadienyl group nearly perpendicular to the K—K' vector (the primes indicate a transformation by the twofold screw axis).




Fig. 2. An ORTEP drawing of the contents of a unit cell, with a unit cell outlined. The view is perpendicular to the *bc* plane, with the *c* axis horizontal. The potassium ion and the tetrahydrofuran molecule are shown without quadrant outlining. H atoms are not shown.

Experimental

Crystal data

K⁺.C₁₅H₂₈NSi⁻.C₄H₈OM_r = 361.68

Monoclinic

P2₁/n

a = 10.799 (3) Å

b = 10.676 (5) Å

c = 19.894 (4) Å

β = 97.13 (2)°

V = 2275.8 (13) Å³

Z = 4

D_x = 1.06 Mg m⁻³

Mo Kα radiation

λ = 0.71073 Å

Cell parameters from 25 reflections

θ = 6–17°

μ = 0.29 mm⁻¹

T = 294 K

Irregular parallelepiped

1.22 × 0.44 × 0.35 mm

Colorless

Data collection

Enraf-Nonius CAD-4 diffractometer

θ–2θ scans

Absorption correction: none

4870 measured reflections

2116 independent reflections

2116 observed reflections

R_{int} = 0.026θ_{max} = 20°

h = 0 → 10

k = -10 → 10

l = -19 → 19

3 standard reflections

frequency: 150 min

intensity variation: within counting statistics

Refinement

Refinement on F²

Final R = 0.064

wR = 0.013

S = 3.36

2116 reflections

212 parameters

H-atom parameters not refined (except NH)

w = 1/σ²(F_o²)(Δ/σ)_{max} = 0.13Δρ_{max} = 0.40 e Å⁻³Δρ_{min} = -0.24 e Å⁻³

Atomic scattering factors from Cromer & Waber (1974)

Table 1. Fractional atomic coordinates and equivalent isotropic thermal parameters (Å²)

$$U_{eq} = \frac{1}{3} \sum_i \sum_j U_{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j$$

| | x | y | z | U _{eq} |
|-----|-------------|-------------|------------|-----------------|
| K | 0.7457 (1) | 0.2949 (1) | 0.7913 (1) | 0.0821 (4) |
| C1 | 0.8612 (4) | 0.0482 (4) | 0.7601 (2) | 0.0583 (12) |
| C2 | 0.7919 (6) | 0.0801 (4) | 0.6969 (2) | 0.0631 (15) |
| C3 | 0.6665 (6) | 0.0659 (4) | 0.7006 (3) | 0.0670 (16) |
| C4 | 0.6509 (5) | 0.0243 (4) | 0.7663 (3) | 0.0676 (16) |
| C5 | 0.7696 (5) | 0.0134 (4) | 0.8026 (2) | 0.0605 (14) |
| C6 | 0.8433 (5) | 0.1234 (4) | 0.6334 (2) | 0.0987 (17) |
| C7 | 0.5606 (6) | 0.0843 (4) | 0.6440 (3) | 0.1129 (19) |
| C8 | 0.5292 (6) | -0.0058 (5) | 0.7919 (3) | 0.1142 (19) |
| C9 | 0.7970 (5) | -0.0306 (4) | 0.8754 (2) | 0.0886 (15) |
| Si | 1.0298 (1) | 0.0507 (1) | 0.7850 (1) | 0.0757 (4) |
| N | 1.0536 (5) | 0.1524 (4) | 0.8530 (2) | 0.0765 (13) |
| C10 | 1.0895 (5) | -0.1109 (5) | 0.8051 (3) | 0.129 (2) |
| C11 | 1.1252 (5) | 0.1032 (6) | 0.7174 (3) | 0.1172 (19) |
| C12 | 1.1652 (5) | 0.1926 (5) | 0.8970 (3) | 0.0882 (17) |
| C13 | 1.2500 (7) | 0.0863 (7) | 0.9211 (4) | 0.202 (3) |
| C14 | 1.2376 (7) | 0.2854 (8) | 0.8598 (4) | 0.189 (3) |
| C15 | 1.1281 (7) | 0.2597 (9) | 0.9566 (4) | 0.207 (4) |
| O | 0.7190 (6) | 0.2918 (4) | 0.9276 (2) | 0.1190 (15) |
| C16 | 0.7863 (7) | 0.3309 (7) | 0.9877 (5) | 0.149 (3) |
| C17 | 0.7038 (12) | 0.3424 (10) | 1.0361 (4) | 0.201 (4) |
| C18 | 0.5925 (10) | 0.2807 (10) | 1.0100 (6) | 0.217 (4) |
| C19 | 0.6030 (9) | 0.2590 (9) | 0.9413 (5) | 0.184 (4) |

Table 2. Geometric parameters (Å, °)

Cp* is the centroid of the five C atoms of the cyclopentadienyl ring.

| | | | |
|-----------|-----------|-------------|------------|
| K—Cp* | 2.808 | Si—C10 | 1.867 (6) |
| K—O | 2.762 (5) | Si—C11 | 1.878 (6) |
| C1—C2 | 1.423 (6) | Si—N | 1.729 (4) |
| C1—C5 | 1.428 (6) | N—C12 | 1.463 (7) |
| C2—C3 | 1.373 (7) | C12—C13 | 1.499 (10) |
| C2—C6 | 1.514 (7) | C12—C14 | 1.512 (10) |
| C3—C4 | 1.410 (7) | C12—C15 | 1.483 (10) |
| C3—C7 | 1.515 (7) | O—C16 | 1.385 (9) |
| C4—C5 | 1.396 (7) | O—C19 | 1.361 (11) |
| C4—C8 | 1.502 (8) | C16—C17 | 1.396 (13) |
| C5—C9 | 1.515 (7) | C17—C18 | 1.412 (15) |
| Si—C1 | 1.827 (4) | C18—C19 | 1.405 (15) |
| Cp*—K—O | 108.3 | C1—Si—N | 105.5 (2) |
| C5—C1—C2 | 105.0 (4) | C10—Si—C11 | 102.9 (3) |
| C3—C2—C1 | 109.9 (4) | C10—Si—N | 113.5 (2) |
| C6—C2—C1 | 127.1 (4) | C11—Si—N | 109.4 (2) |
| C6—C2—C3 | 123.0 (4) | Si—N—C12 | 133.1 (4) |
| C4—C3—C2 | 108.5 (4) | C13—C12—N | 113.2 (5) |
| C7—C3—C2 | 127.2 (5) | C14—C12—N | 109.7 (5) |
| C7—C3—C4 | 124.3 (5) | C15—C12—N | 109.6 (5) |
| C5—C4—C3 | 107.4 (4) | C14—C12—C13 | 108.7 (5) |
| C8—C4—C3 | 126.3 (5) | C15—C12—C13 | 109.0 (6) |
| C8—C4—C5 | 126.3 (5) | C15—C12—C14 | 106.4 (6) |
| C4—C5—C1 | 109.3 (4) | C19—O—C16 | 107.6 (6) |
| C9—C5—C1 | 125.3 (4) | C17—C16—O | 108.1 (7) |
| C9—C5—C4 | 125.4 (4) | C18—C17—C16 | 107.0 (9) |
| C1—Si—C10 | 110.5 (2) | C19—C18—C17 | 105.5 (9) |
| C1—Si—C11 | 115.3 (2) | C18—C19—O | 109.9 (8) |

Before averaging, 41 reflections were deleted from the data set because of shutter problems with the diffractometer. The goodness-of-fit for merging 1967 multiply-measured reflections was 1.25. The weights and variances [$\sigma^2(F_o^2)$] were derived from counting statistics plus an additional term, $(0.014I)^2$; variances of the merged data were obtained by propagation of error plus another additional term, $(0.014I)^2$. Values for f' were taken from Cromer (1974). The programs used were those of the *CRYM Crystallographic Computing System* (Duchamp, 1964) and *ORTEP* (Johnson, 1976). H atoms on C were placed by calculation (C—H 0.95 Å) assuming staggered configurations except on the Cp* group, where the atoms were placed at idealized positions based on difference maps calculated in the expected planes. H atoms were included at fixed positions, with B values corresponding to 1.2 times the isotropic equivalent U_{ij} value of the bonded atoms. They were repositioned once near the end of the refinement. The H atom on N was located in a difference map and its parameters were refined.

This work was supported by the USDOE Office of Basic Energy Sciences (grant No. DE-FG03-85ER113431) and by Exxon Chemical Americas.

Lists of structure factors, anisotropic thermal parameters, H-atom coordinates and complete geometry have been deposited with the British Library Document Supply Centre as Supplementary Publication No. SUP 71110 (16 pp.). Copies may be obtained through The Technical Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England. [CIF reference: HH1051]

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans.* pp. S1–S19.

- Cotter, W. D. & Bercaw, J. E. (1993). Unpublished results.
- Cromer, D. T. (1974). *International Tables for X-ray Crystallography*, Vol. IV, pp. 149–151. Birmingham: Kynoch Press. (Present distributor Kluwer Academic Publishers, Dordrecht.)
- Cromer, D. T. & Waber, J. T. (1974). *International Tables for X-ray Crystallography*, Vol. IV, pp. 99–101. Birmingham: Kynoch Press. (Present distributor Kluwer Academic Publishers, Dordrecht.)
- Duchamp, D. J. (1964). *Am. Crystallogr. Assoc. Meet., Bozeman, Montana*, Paper B14, p. 29.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Marsh, R. E., Schaefer, W. P., Coughlin, E. B. & Bercaw, J. E. (1992). *Acta Cryst. C* **48**, 1773–1776.
- Schaefer, W. P., Köhn, R. D. & Bercaw, J. E. (1992). *Acta Cryst. C* **48**, 251–253.
- Shapiro, P. J., Bunel, E., Schaefer, W. P. & Bercaw, J. E. (1990). *Organometallics*, **9**, 867–869.

Acta Cryst. (1993). **C49**, 1492–1493

A Second Crystal Form of Bis- μ -[bis(dimethylphosphino)methane]-bis(tricarbonylrhenium)(Re—Re), $\text{Re}_2(\text{dmpm})_2(\text{CO})_6$

DAVID R. TYLER AND TIMOTHY J. R. WEAKLEY

Department of Chemistry, University of Oregon, Eugene, OR 97403, USA

(Received 3 August 1992; accepted 15 February 1993)

Abstract

The second crystalline form, (II), of the title compound has the same space group ($P2_1/n$) as form (I) of known structure [Milder, Castellani, Weakley, Tyler, Miskowski, & Stiegman (1990). *J. Phys. Chem.* **94**, 6599–6603] but has a different packing. There is no disorder in (II) [in contrast to (I)] and the Re—Re bond length is significantly longer [3.126 (1) versus 3.105 (1) Å] despite the smaller (*ca* 1.3%) molecular volume.

Comment

The data crystal of form (II) came from the air-stable batch which earlier (Milder *et al.*, 1990) had afforded the crystal of form (I). Both forms have space group $P2_1/n$ with two molecules per cell, but they clearly differ in molecular packing. The unit cell of (II), and hence the molecular volume, is 1.3% smaller. Accurately centrosymmetric $\text{Re}_2(\text{dmpm})_2(\text{CO})_6$ molecules with Re—Re bonds, bridging dmpm ligands, *trans* P—Re—P geometry, and generally similar dimensions, are present in both forms. The principal points of difference are as follows. The Re—Re bond in (II)

[3.126 (1) Å] is significantly longer than in (I) [3.105 (1) Å]. Alternative positions for three of the five dmpm C atoms are discernible in (I), but (II) is free from disorder, a feature that accounts for the more efficient molecular packing in (II) and the fact that the C—P—C angles lie within the range 100.9–104.0 (6)° compared with 86–116 (1)° for the major conformer in (I).

A series of events led us to determine the unit cells of 20 crystals from a stable batch. Eight proved to be of form (II). The two forms, both amber prisms, cannot be readily distinguished under the polarizing microscope. We think it is unlikely that most workers routinely find the cell dimensions for several crystals from an apparently homogeneous sample if the first crystal proves suitable for data collection. We feel, therefore, that the presence of two crystalline forms in one batch may be more common than is generally supposed.




Fig. 1. View of the $\text{Re}_2(\text{dmpm})_2(\text{CO})_6$ molecule showing thermal ellipsoids.

Experimental

Crystal data

$[\text{Re}_2(\text{C}_5\text{H}_{14}\text{P}_2)_2(\text{CO})_6]$

$M_r = 812.70$

Monoclinic

$P2_1/n$

$a = 9.938$ (2) Å

$b = 8.579$ (3) Å

$c = 14.783$ (3) Å

$\beta = 93.53$ (2)°

Mo $K\alpha$ radiation

$\lambda = 0.71069$ Å

Cell parameters from 25 reflections

$\theta = 11.5$ – 16.1 °

$\mu = 10.0$ mm⁻¹

$T = 297$ K

Prism

$0.30 \times 0.25 \times 0.15$ mm