

*Acta Cryst.* (1996). **C52**, 1008–1010

## Bis(4-methoxyphenyl)methano[60]fullerene with a Methano Bridge at the 6–6-Ring Junction

YOSHIHIKO TEZUKA,<sup>a</sup> NOBUYUKI KAWASAKI,<sup>a</sup> HIROFUMI YAJIMA,<sup>a</sup> TADAIHIRO ISHII,<sup>a\*</sup> TOSHIYUKI OYAMA,<sup>b</sup> KAZUO TAKEUCHI,<sup>b</sup> AKIKO NAKAO<sup>c</sup> AND CHUJI KATAYAMA<sup>c</sup>

<sup>a</sup>Department of Applied Chemistry, Faculty of Science, Science University of Tokyo, 1-3 Kagurazaka, Shinjuku-ku, Tokyo 162, Japan, <sup>b</sup>The Institute of Physical and Chemical Research (RIKEN), Hirosawa, Wako-shi, Saitama 350-01, Japan, and <sup>c</sup>Research Laboratory, MAC SCIENCE Co. Ltd, 3-12-1 Nakamachidai, Tuzuki-ku, Yokohama 224, Japan.  
E-mail: taishii@ch.kagu.sut.ac.jp


(Received 7 July 1995; accepted 20 September 1995)

### Abstract

In the title compound, C<sub>75</sub>H<sub>14</sub>O<sub>2</sub>, a bis(4-methoxyphenyl)methylene group bridges a 6–6-ring junction of the C<sub>60</sub> sphere, forming a slightly deformed cyclopropane ring. The C—C bond length involved in the cyclopropane ring fusion is 1.635 (4) Å, more than 0.2 Å longer than that at the 6–6-ring junction of C<sub>60</sub>. The elongation of this C—C bond leads to a slight deformation of the C<sub>60</sub> framework, especially a shortening of two C—C bonds in equatorial positions.

### Comment

Since the discovery of the method for large-scale preparation of fullerenes, their chemical modifications have been investigated intensively for the purpose of functionalization. Addition of diazoalkanes or diazoacetates to C<sub>60</sub>-fullerene developed by Wudl and co-workers (Suzuki, Li, Khemani, Wudl & Almarsson, 1991; Wudl, 1992) is a promising method for the preparation of the functionalized fullerenes. Single addition of diazo compounds to C<sub>60</sub> gives a mixture of isomers with a methano-bridge at different positions, either at the 6–6-ring junction or the 5–6-ring junction of the C<sub>60</sub> sphere as determined by <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy (Isaacs, Wehrsig & Diederich, 1993). Recently, the 6–6-ring-bridged structure was confirmed by X-ray analysis for several methano-bridged fullerenes (Osterodt, Nieger & Vogtle, 1994; Anderson *et al.*, 1994; Paulus & Bingel, 1995). The X-ray structure is useful for the prediction of the reactivity of a C—C bond on the C<sub>60</sub> sphere and the estimation of intermolecular interactions. We report here the X-ray structure analysis of the title compound, (I), and compare its structure with those of previously reported methano-bridged fullerenes.



(I)

Fig. 1 shows the molecular structure of (I). The molecule has a mirror plane containing atoms C(8), C(9), C(10), C(25) and C(26), since both 4-methoxyphenyl groups are arranged symmetrically. The phenyl ring is essentially planar and the methoxy group belongs to this plane [deviations of the atoms from the least-squares plane through atoms C(1) and O(1) and the phenyl ring are less than 0.006 Å]. The phenyl ring is almost perpendicular to the plane through atoms C(5), C(8) and C(5') [symmetry code: (')  $x, \frac{1}{2} - y, z$ ; dihedral angle 88.3 (2)°]. The dihedral angle between the least-squares planes through the phenyl rings is 99.8 (3)°.




Fig. 1. An ORTEP view (Johnson, 1965) of (I) with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

The C(9)—C(10) bond involved in the cyclopropane ring fusion is 1.635 (4) Å long, which is the longest among the reported methano-bridged fullerenes. It is 0.12 Å longer than a C—C bond in cyclopropane (1.52 Å), leading to a slight deformation of the cyclopropane ring [angles C(9)—C(8)—C(10), C(8)—C(9)—C(10) and C(8)—C(10)—C(9) are 65.3 (2), 57.4 (2) and 57.3 (2)°, respectively]. The C(9)—C(10) bond length is more than 0.2 Å longer than that at the 6–6-ring junction of C<sub>60</sub> (1.40 Å) determined by NMR spectroscopy (Yannoni, Bernier, Bethune, Meijer & Salem, 1991). The elongation of a C—C bond leads to a slight deforma-

tion of the C<sub>60</sub> framework. The most affected bonds are those at the 6–6-ring junction in equatorial positions, C(17)—C(17') and C(18)—C(18'), which have lengths of 1.370 (5) and 1.374 (5) Å, respectively. They are the shortest two bonds on the C<sub>60</sub> sphere and thus have the highest double-bond character. This supports the theory, as demonstrated by theoretical and experimental methods, that the position of highest reactivity towards nucleophiles is at the equatorial position of singly methano-bridged fullerenes (Hirsch, Lamparth & Karfunkel, 1994; Hirsch, Lamparth, Grösser & Karfunkel, 1994).

Fig. 2 shows the crystal packing of (I). The packing is characterized by a layer parallel to the *ac* plane. The closest packed layers cut the *b* axis at  $y = \frac{1}{4}$  and  $\frac{3}{4}$ . The molecules are arranged in the same direction in a layer. A convenient molecular axis passing through atom C(8) and the midpoint of C(25)—C(26) forms an angle of 41.3 (1)° with the *a* axis. The adjacent layers are related by a twofold screw axis perpendicular to the *ac* plane. The shortest intermolecular distance between the phenyl ring and the C<sub>60</sub> sphere is 3.359 (3) Å, between atom C(2) of one molecule and atom C(37') of the next molecule in the same layer. This is close to the sum of the van der Waals radii, suggesting a weak interaction, as observed in (3,4-dimethoxyphenyl)phenylmethano[60]-fullerene (Osterodt, Nieger & Vogtle, 1994).




Fig. 2. A molecular packing diagram of (I) in the unit cell.

## Experimental

The title compound, (I), was synthesized according to the literature method of Shi, Khemani, Li & Wudl (1992). A

solution of 2 equiv. of bis(4-methoxyphenyl)diazomethane and one equiv. of C<sub>60</sub> in toluene was stirred at room temperature for 12 h and then refluxed for 24 h. The products were separated by flash chromatography on neutral alumina using toluene/hexane (1:1 v/v) solution as eluent. One of the fractional solutions yielded black crystals without any cooling or concentration. The crystals were heated to 430 K under reduced pressure to remove traces of solvent. The density *D<sub>m</sub>* was measured on a Berman density torsion balance.

### Crystal data

C<sub>75</sub>H<sub>14</sub>O<sub>2</sub>  
*M<sub>r</sub>* = 946.9  
 Monoclinic  
*P*2<sub>1</sub>/*m*  
*a* = 11.323 (2) Å  
*b* = 17.602 (4) Å  
*c* = 9.852 (2) Å  
 $\beta$  = 101.94 (2)°  
*V* = 1921.3 (7) Å<sup>3</sup>  
*Z* = 2  
*D<sub>x</sub>* = 1.63 Mg m<sup>-3</sup>  
*D<sub>m</sub>* = 1.63 Mg m<sup>-3</sup>

Mo *K*α radiation  
 $\lambda$  = 0.71073 Å  
 Cell parameters from 22 reflections  
 $\theta$  = 15–17.5°  
 $\mu$  = 0.097 mm<sup>-1</sup>  
*T* = 298 K  
 Prism  
 0.30 × 0.25 × 0.20 mm  
 Black

### Data collection

Mac Science MXC18K diffractometer  
 2θ/ω scans  
 Absorption correction: none  
 4980 measured reflections  
 4093 independent reflections  
 3003 observed reflections  
 [*F* > 3σ(*F*)]

*R*<sub>int</sub> = 0.03  
 $\theta_{\max}$  = 27.5°  
*h* = -14 → 14  
*k* = 0 → 22  
*l* = 0 → 12  
 3 standard reflections monitored every 100 reflections  
 intensity decay: 3%

### Refinement

Refinement on *F*  
*R* = 0.041  
*wR* = 0.038  
*S* = 0.756  
 3003 reflections  
 383 parameters  
 H atoms refined isotropically  
 Unit weights applied

( $\Delta/\sigma$ )<sub>max</sub> = 0.430  
 $\Delta\rho_{\max}$  = 0.17 e Å<sup>-3</sup>  
 $\Delta\rho_{\min}$  = -0.19 e Å<sup>-3</sup>  
 Extinction correction: none  
 Atomic scattering factors from *International Tables for X-ray Crystallography* (1974, Vol. IV)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (Å<sup>2</sup>)

$$U_{eq} = (1/3)\sum_i \sum_j U_{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j$$

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> <sub>eq</sub>
O(1)	0.5507 (1)	0.5009 (1)	0.8341 (2)	0.050 (1)
C(1)	0.6385 (3)	0.5341 (2)	0.7695 (4)	0.071 (2)
C(2)	0.4846 (2)	0.4420 (1)	0.7671 (2)	0.036 (1)
C(3)	0.4984 (2)	0.4125 (1)	0.6415 (2)	0.043 (1)
C(4)	0.4267 (2)	0.3522 (1)	0.5841 (2)	0.040 (1)
C(5)	0.3410 (2)	0.3210 (1)	0.6490 (2)	0.031 (1)
C(6)	0.3276 (2)	0.3518 (1)	0.7748 (2)	0.037 (1)
C(7)	0.3989 (2)	0.4115 (1)	0.8337 (2)	0.039 (1)
C(8)	0.2749 (2)	1/4	0.5914 (3)	0.031 (1)
C(9)	0.2125 (2)	1/4	0.4399 (3)	0.030 (1)
C(10)	0.1382 (2)	1/4	0.5656 (3)	0.030 (1)
C(11)	0.1906 (2)	0.1829 (1)	0.3467 (2)	0.035 (1)
C(12)	0.0597 (2)	0.1828 (1)	0.5675 (2)	0.036 (1)
C(13)	0.1829 (2)	0.2091 (1)	0.2063 (2)	0.044 (1)

C(14)	-0.0516 (2)	0.2092 (2)	0.6007 (2)	0.045 (1)
C(15)	-0.1159 (2)	0.1696 (2)	0.0940 (2)	0.055 (1)
C(16)	-0.1593 (2)	0.1691 (2)	0.5565 (2)	0.056 (1)
C(17)	0.0452 (2)	0.2111 (2)	-0.0227 (2)	0.060 (1)
C(18)	-0.2709 (2)	0.2110 (2)	0.5099 (2)	0.064 (2)
C(19)	-0.0664 (2)	0.1690 (2)	-0.0690 (2)	0.061 (2)
C(20)	-0.3414 (2)	0.1692 (2)	0.3936 (3)	0.063 (2)
C(21)	-0.1735 (2)	0.2091 (2)	-0.1143 (2)	0.055 (1)
C(22)	-0.4097 (2)	0.2089 (2)	0.2829 (3)	0.059 (1)
C(23)	-0.2837 (2)	0.1836 (2)	-0.0749 (2)	0.050 (1)
C(24)	-0.4129 (2)	0.1835 (2)	0.1423 (2)	0.050 (1)
C(25)	-0.3519 (3)	1/4	-0.0511 (3)	0.046 (2)
C(26)	-0.4152 (3)	1/4	0.0564 (3)	0.047 (2)
C(27)	0.1246 (2)	0.1206 (1)	0.3710 (2)	0.040 (1)
C(28)	0.0579 (2)	0.1204 (1)	0.4835 (2)	0.040 (1)
C(29)	0.0527 (2)	0.0799 (1)	0.2540 (3)	0.050 (1)
C(30)	-0.0546 (2)	0.0797 (1)	0.4344 (3)	0.052 (1)
C(31)	0.0499 (2)	0.1033 (2)	0.1185 (3)	0.056 (2)
C(32)	-0.1600 (2)	0.1031 (2)	0.4725 (3)	0.058 (2)
C(33)	-0.0639 (2)	0.1029 (2)	0.0168 (3)	0.061 (2)
C(34)	-0.2739 (2)	0.1029 (2)	0.3697 (3)	0.061 (2)
C(35)	-0.1684 (2)	0.0787 (2)	0.0555 (3)	0.058 (2)
C(36)	-0.2755 (2)	0.0788 (2)	0.2359 (3)	0.059 (2)
C(37)	-0.2809 (2)	0.1201 (2)	0.0082 (3)	0.053 (1)
C(38)	-0.3469 (2)	0.1200 (2)	0.1198 (3)	0.053 (1)
C(39)	-0.0570 (2)	0.0546 (1)	0.2941 (3)	0.056 (2)
C(40)	-0.1654 (3)	0.0539 (2)	0.1964 (3)	0.060 (2)

Table 2. Selected geometric parameters (Å, °)

C(8)—C(9)	1.514 (4)	C(18)—C(18')	1.374 (5)
C(8)—C(10)	1.516 (4)	C(21)—C(21')	1.440 (5)
C(9)—C(10)	1.635 (4)	C(25)—C(26)	1.396 (5)
C(13)—C(13')	1.438 (4)	C(39)—C(40)	1.395 (4)
C(17)—C(17')	1.370 (5)		
C(5)—C(8)—C(5')	112.1 (2)	C(8)—C(9)—C(10)	57.4 (2)
C(9)—C(8)—C(10)	65.3 (2)	C(8)—C(10)—C(9)	57.3 (2)
C(3)—C(4)—C(5)—C(8)	-173.1 (4)	C(8)—C(5)—C(6)—C(7)	172.5 (4)

Data collection: MAC Science MXC software. Cell refinement: MAC Science MXC software. Data reduction: MAC Science MXC software. Program(s) used to solve structure: *SIR* (Altomare *et al.*, 1994) in *CRYSTAN-GM* (Edwards, Gilmore, Mackay & Stewart, 1995). Program(s) used to refine structure: *LSQ* (Mallinson & Muir, 1985) in *CRYSTAN-GM*. Molecular graphics: *ORTEP* (Johnson, 1965) in *CRYSTAN-GM*. Software used to prepare material for publication: *CRYSTAN-GM*.

This work was partly supported by special coordination funds of the Science and Technology Agency of the Japanese Government.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: AS1208). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

## References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.  
 Anderson, H. L., Boudon, C., Diederich, F., Gisselbrecht, J.-P., Gross, M. & Seiler, P. (1994). *Angew. Chem. Int. Ed. Engl.* **33**, 1628–1632.  
 Edwards, C., Gilmore, C. J., Mackay, S. & Stewart, N. (1995). *CRYSTAN. A Computer Program for the Solution and Refinement of Crystal Structures*. MAC Science, Japan.  
 Hirsch, A., Lamparth, I., Grösser, T. & Karfunkel, H. R. (1994). *J. Am. Chem. Soc.* **116**, 9385–9386.

- Hirsch, A., Lamparth, I. & Karfunkel, H. R. (1994). *Angew. Chem. Int. Ed. Engl.* **33**, 437–438.  
 Isaacs, L., Wehrsig, A. & Diederich, F. (1993). *Helv. Chim. Acta*, **76**, 1231–1250.  
 Johnson, C. K. (1965). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee, USA.  
 Mallinson, P. R. & Muir, K. W. (1985). *J. Appl. Cryst.* **18**, 51–53.  
 Osterodt, J., Nieger, M. & Vogtle, F. (1994). *J. Chem. Soc. Chem. Commun.* pp. 1607–1608.  
 Paulus, E. F. & Bingel, C. (1995). *Acta Cryst.* **C51**, 143–146.  
 Shi, S., Khemani, K. C., Li, Q. C. & Wudl, F. (1992). *J. Am. Chem. Soc.* **114**, 10656–10657.  
 Suzuki, T., Li, Q., Khemani, K. C., Wudl, F. & Almarsson, O. (1991). *Science*, **254**, 1186–1188.  
 Wudl, F. (1992). *Acc. Chem. Res.* **25**, 157–161.  
 Yannoni, C. S., Bernier, P. P., Bethune, D. S., Meijer, G. & Salem, J. R. (1991). *J. Am. Chem. Soc.* **113**, 3190–3192.

*Acta Cryst.* (1996). **C52**, 1010–1012

## 3-Cyano-1-[4-(1,3-dithian-2-yl)butyl]-1,4,5,6-tetrahydropyridine

ZAHID HUSSAIN,<sup>a</sup> FRASER F. FLEMING,<sup>a</sup> RICHARD E. NORMAN<sup>a</sup> AND SHIH-CHI CHANG<sup>b</sup>

<sup>a</sup>Department of Chemistry and Biochemistry, Duquesne University, Pittsburgh, PA 15282, USA, and <sup>b</sup>Department of Physics, Duquesne University, Pittsburgh, PA 15282, USA. E-mail: renorman@duq3.cc.duq.edu

(Received 17 July 1995; accepted 2 October 1995)

## Abstract

The title nitrile {1-[4-(1,3-dithian-2-yl)butyl]-1,4,5,6-tetrahydropyridine-3-carbonitrile, C<sub>14</sub>H<sub>22</sub>N<sub>2</sub>S<sub>2</sub>} was prepared as part of our investigations into the conjugate addition reactions of unsaturated nitriles [Fleming & Pak (1995). *J. Org. Chem.* **60**, 4299–4301]. The tetrahydropyridine ring adopts a half-chair conformation and is appended by a four-carbon chain to the dithiane ring which adopts a chair conformation.

## Comment

Conjugate addition to unsaturated nitriles is often problematic with conventional nucleophiles (Lipshutz, Wilhelm & Kozlowski, 1984; House & Umen, 1973). We sought to promote the conjugate addition reaction by tethering a nucleophilic dithiane anion to an unsaturated nitrile and therefore prepared the title compound, (I), as a suitable substrate for this reaction. (I) is readily prepared by coupling 3-cyanotetrahydropyridine (Kikugawa, Kuramoto, Saito & Yamada, 1973) with 2-(4-chlorobutyl)-1,3-dithiane (Seebach, Jones & Corey, 1968). Concentration of the crude reaction mix-