

Refinement

Refinement on F	$(\Delta/\sigma)_{\max} = 0.0072$
$R = 0.0370$	$\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$
$wR = 0.0424$	$\Delta\rho_{\min} = -0.15 \text{ e } \text{\AA}^{-3}$
$S = 1.913$	Extinction correction: none
1095 reflections	Atomic scattering factors
100 parameters	from <i>International Tables</i>
H-atom parameters not refined	for <i>X-ray Crystallography</i>
$w = 4F^2/\sigma^2(F^2)$	(1974, Vol. IV)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (\AA^2)

$$U_{\text{eq}} = (1/3)\sum_i \sum_j U_{ij} a_i^* a_j \cdot a_j$$

	x	y	z	U_{eq}
O(1)	0.1941 (2)	-0.1448 (2)	0.68535 (6)	0.0435 (3)
O(2)	-0.0589 (2)	0.0916 (2)	0.68345 (7)	0.0554 (4)
O(3)	-0.0157 (2)	0.0357 (2)	0.60615 (6)	0.0474 (4)
N(1)	-0.2140 (4)	-0.4225 (3)	0.5738 (1)	0.0811 (7)
C(1)	-0.0464 (4)	-0.3452 (3)	0.5674 (1)	0.0540 (6)
C(2)	0.1650 (3)	-0.2412 (3)	0.5580 (1)	0.0483 (5)
C(3)	0.1907 (3)	-0.0758 (2)	0.61182 (9)	0.0382 (4)
C(4)	0.3927 (3)	0.0512 (3)	0.6000 (1)	0.0509 (5)
C(5)	0.3898 (3)	0.2202 (3)	0.6529 (1)	0.0580 (6)
C(6)	0.3273 (3)	0.1604 (3)	0.7298 (1)	0.0537 (6)
C(7)	0.1349 (3)	0.0195 (3)	0.72563 (10)	0.0467 (5)

Table 2. Selected geometric parameters (\AA , $^\circ$)

O(1)—C(3)	1.408 (2)	C(1)—C(2)	1.459 (3)
O(1)—C(7)	1.414 (2)	C(2)—C(3)	1.515 (2)
O(2)—O(3)	1.478 (2)	C(3)—C(4)	1.509 (2)
O(2)—C(7)	1.435 (2)	C(4)—C(5)	1.522 (3)
O(3)—C(3)	1.445 (2)	C(5)—C(6)	1.508 (3)
N(1)—C(1)	1.136 (2)	C(6)—C(7)	1.503 (3)
C(3)—O(1)—C(7)	102.0 (1)	O(3)—C(3)—C(4)	109.6 (1)
O(3)—O(2)—C(7)	103.8 (1)	C(2)—C(3)—C(4)	114.9 (1)
O(2)—O(3)—C(3)	104.8 (1)	C(3)—C(4)—C(5)	110.0 (1)
N(1)—C(1)—C(2)	178.3 (2)	C(4)—C(5)—C(6)	111.5 (2)
C(1)—C(2)—C(3)	111.9 (1)	C(5)—C(6)—C(7)	110.5 (2)
O(1)—C(3)—O(3)	103.3 (1)	O(1)—C(7)—O(2)	103.0 (1)
O(1)—C(3)—C(2)	109.5 (1)	O(1)—C(7)—C(6)	111.1 (1)
O(1)—C(3)—C(4)	110.9 (1)	O(2)—C(7)—C(6)	112.0 (2)
O(3)—C(3)—C(2)	108.0 (1)		

In the weighting scheme $\sigma^2(F^2) = S^2(C + R^2B) + (0.04F^2)^2/Lp^2$, where S = scan rate, C = total integrated peak count, R = ratio of scan to background counting times, B = total background count and Lp = Lorentz-polarization factor.

Data collection: *MSCIAFC Diffractometer Control Software* (Molecular Structure Corporation, 1992a). Cell refinement: *MSCIAFC Diffractometer Control Software*. Data reduction: *TEXSAN* (Molecular Structure Corporation, 1992b). Program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994). Program(s) used to refine structure: *TEXSAN*. Software used to prepare material for publication: *TEXSAN*.

Financial support from the Jacob A. and Frieda M. Hunkele Charitable Trust and the Kresge Foundation is gratefully acknowledged.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: BK1178). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Altomare, A., Cascarano, M., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435–436.
- Buckleton, J. S., Clark, G. R. & Rickard, C. E. F. (1995). *Acta Cryst.* **C51**, 494–495.
- Bunnelle, W. H. & Isbell, T. A. (1992). *J. Org. Chem.* **57**, 729–740.
- Bunnelle, W. H., Isbell, T. A., Barnes, C. L. & Qualls, S. (1991). *J. Am. Chem. Soc.* **113**, 8168–8169.
- Bunnelle, W. H. & Lee, S. (1992). *J. Am. Chem. Soc.* **114**, 7577–7578.
- Cummins, W. J., Drew, M. G. B., Mann, J. & Walsh, E. B. (1983). *J. Chem. Soc. Perkin Trans. 1*, pp. 167–171.
- Jerzykiewicz, L. B., Dziewońska-Baran, D., Baran, J. & Lis, T. (1993). *Acta Cryst.* **C49**, 400–402.
- Karban, J., McAtee, J. L. Jr, Belew, J. S., Mullica, D. F., Milligan, W. O. & Korp, J. (1978). *J. Chem. Soc. Chem. Commun.* pp. 729–730.
- Miura, M., Ikegami, A., Nojima, M., Kusabayashi, S., McCullough, K. J. & Nagase, S. (1983). *J. Am. Chem. Soc.* **105**, 2414–2426.
- Molecular Structure Corporation (1992a). *MSCIAFC Diffractometer Control Software*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Molecular Structure Corporation (1992b). *TEXSAN. Crystal Structure Analysis Package*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.

Acta Cryst. (1996). **C52**, 1014–1017

Two Schiff Base Ligands Derived from 1,2-Diaminocyclohexane

JASON C. CANNADINE, JONATHAN P. CORDEN, WILLIAM ERRINGTON, PETER MOORE AND MALCOLM G. H. WALLBRIDGE

Department of Chemistry, University of Warwick, Coventry CV4 7AL, England. E-mail: msrpq@csv.warwick.ac.uk

(Received 17 July 1995; accepted 10 October 1995)


Abstract

The crystal structures of the free ligands 2,2'-(1,2-cyclohexanediyl)bis(nitrilomethylidene)bisphenol, C₂₀H₂₂N₂O₂, (I), and 2,2'-(1,2-cyclohexanediyl)bis(nitriloethylidene)bisphenol, C₂₂H₂₆N₂O₂, (II), have been determined. In both molecules the N—O distances are indicative of intramolecular hydrogen bonding. In compound (I), the two aromatic rings are inclined at an angle of 56.5 (1)° and the O···O separation is 6.082 (3) Å; in compound (II) the corresponding values are 83.15 (8)° and 5.544 (5) Å. Thus, it is evident that the methyl groups in (II) have a very significant effect upon the overall conformation.

Comment

Relatively few crystal structures have been reported for quadridentate Schiff base ligands (Corden, Errington,

Moore & Wallbridge, 1996). Investigations within these laboratories into the coordination chemistry of the ligands (I) and (II) have prompted us to determine the structures of these free ligands, so that subsequent changes resulting from the coordination of a metal ion may be identified.



The molecular structures of these two ligands are represented in Figs. 1 and 2. The bond lengths and angles, while unexceptional, clearly indicate that the enolimine tautomer is favoured. Furthermore, the shortest N—O distances [average 2.604 (3) Å for (I); average 2.515 (3) Å for (II)] are indicative of intramolecular hydrogen bonding. Although structure (I) is non-centrosymmetric, both enantiomers are present; structure (II) is centrosymmetric.

Perhaps the most interesting comparative aspect of these two structures involves the influence of the methyl groups upon the molecular conformation. While several of the corresponding torsion angles are very similar, a large difference is noted between C7—N1—C9—C14 and C7—N1—C8—C13; the cumulative effect of the differences is that the orientation of the two aromatic rings is 56.5 (1)° in (I), but 83.15 (8)° in (II). An alternative way of expressing the significant differences between the conformations of the two molecules is to




Fig. 1. View of molecule (I) showing the atomic numbering. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. View of molecule (II) showing the atomic numbering. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 3. Superposition of the two molecules with respect to the cyclohexane rings. The atomic numbering refers to structure (II).

note that the O atoms are 6.082 (3) Å apart in (I), but only 5.544 (5) Å apart in (II). The overall situation is represented in Fig. 3 in which the cyclohexane rings have been superimposed for the two molecules to show the lack of coincidence in the 'wings' of the molecules.

Experimental

Compound (I) was prepared by the condensation of salicylaldehyde (150 mmol) with *trans*-1,2-diaminocyclohexane in 100 ml of methanol. The reaction mixture was stirred at 298 K for 20 min and then placed in a freezer for 1 h. The yellow precipitate was collected by filtration, washed with dry diethyl ether and dried *in vacuo*. Crystals of X-ray quality were obtained by the slow evaporation of a saturated methanolic solution.

Compound (II) was prepared by the condensation of 2-hydroxyacetophenone (74 mmol in 100 ml methanol) with *trans*-1,2-diaminocyclohexane (37 mmol in 40 ml methanol). The reaction mixture was stirred at 323 K for 1 h and then cooled to 253 K for 3 h. The bright yellow precipitate was washed and dried *in vacuo*. Crystals suitable for X-ray diffraction were obtained by slow evaporation from a saturated methanolic solution.

Compound (I)*Crystal data*C₂₀H₂₂N₂O₂ $M_r = 322.40$

Monoclinic

Cc

 $a = 16.013 (6) \text{ \AA}$ $b = 11.860 (6) \text{ \AA}$ $c = 9.643 (2) \text{ \AA}$ $\beta = 98.50 (2)^\circ$ $V = 1811.2 (12) \text{ \AA}^3$

Z = 4

 $D_x = 1.182 \text{ Mg m}^{-3}$ D_m not measured*Data collection*Delft Instruments FAST TV
area-detector diffractometer

Flat-plate detector

Absorption correction:
none

3417 measured reflections

1936 independent reflections

*Refinement*Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.0436$ $wR(F^2) = 0.1219$

S = 1.022

1932 reflections

217 parameters

H-atom parameters not
refined $w = 1/[\sigma^2(F_o^2) + (0.0729P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = -0.001$ Mo $K\alpha$ radiation $\lambda = 0.71073 \text{ \AA}$ Cell parameters from 250
reflections $\theta = 2.15\text{--}25.00^\circ$ $\mu = 0.077 \text{ mm}^{-1}$

T = 293 (2) K

Block

 $0.28 \times 0.16 \times 0.14 \text{ mm}$

Yellow

1728 observed reflections

 $[I > 2\sigma(I)]$ $R_{\text{int}} = 0.0461$ $\theta_{\max} = 25.02^\circ$ $h = -17 \rightarrow 16$ $k = -13 \rightarrow 12$ $l = -10 \rightarrow 9$ $\Delta\rho_{\max} = 0.153 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.141 \text{ e \AA}^{-3}$

Extinction correction: none

Atomic scattering factors

from *International Tables*
for *Crystallography* (1992),
Vol. C, Tables 4.2.6.8 and
6.1.1.4)

Absolute configuration:

Flack (1983) parameter

= 1.4 (18)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (\AA^2) for (I)
$$U_{\text{eq}} = (1/3)\sum_i\sum_j U_{ij}a_i^*a_j^* \mathbf{a}_i \cdot \mathbf{a}_j.$$

	x	y	z	U_{eq}
O1	1.0259 (2)	0.2372 (2)	-0.1151 (2)	0.0865 (7)
O2	0.8539 (2)	0.1559 (2)	0.3976 (2)	0.0845 (7)
N1	0.89210 (15)	0.1387 (2)	-0.0487 (2)	0.0556 (6)
N2	0.7976 (2)	0.2349 (2)	0.1494 (2)	0.0540 (6)
C1	1.0726 (2)	0.1683 (2)	-0.0234 (3)	0.0580 (7)
C2	1.1599 (2)	0.1739 (3)	-0.0062 (3)	0.0736 (9)
C3	1.2079 (2)	0.1053 (3)	0.0885 (4)	0.0784 (9)
C4	1.1704 (2)	0.0303 (3)	0.1688 (4)	0.0811 (10)
C5	1.0836 (2)	0.0227 (3)	0.1522 (3)	0.0709 (8)
C6	1.0326 (2)	0.0899 (2)	0.0559 (3)	0.0546 (7)
C7	0.9416 (2)	0.0787 (2)	0.0372 (3)	0.0539 (6)

C8	0.8011 (2)	0.1204 (2)	-0.0594 (3)	0.0542 (7)
C9	0.7620 (2)	0.1021 (3)	-0.2118 (3)	0.0676 (8)
C10	0.6676 (2)	0.0849 (3)	-0.2245 (4)	0.0795 (9)
C11	0.6249 (2)	0.1821 (3)	-0.1601 (3)	0.0750 (9)
C12	0.6650 (2)	0.2028 (3)	-0.0086 (3)	0.0700 (8)
C13	0.7596 (2)	0.2206 (2)	0.0021 (3)	0.0529 (7)
C14	0.8311 (2)	0.3288 (2)	0.1894 (3)	0.0498 (6)
C15	0.8746 (2)	0.3473 (2)	0.3306 (3)	0.0506 (7)
C16	0.9080 (2)	0.4527 (2)	0.3702 (3)	0.0587 (7)
C17	0.9563 (2)	0.4695 (3)	0.5003 (3)	0.0673 (8)
C18	0.9702 (2)	0.3800 (3)	0.5923 (3)	0.0685 (9)
C19	0.9359 (2)	0.2760 (3)	0.5576 (3)	0.0681 (9)
C20	0.8874 (2)	0.2588 (2)	0.4284 (3)	0.0574 (7)

Table 2. Selected geometric parameters (\AA , $^\circ$) for (I)

O1—C1	1.347 (4)	N2—C13	1.470 (4)
O2—C20	1.348 (3)	C6—C7	1.447 (4)
N1—C7	1.276 (3)	C8—C13	1.524 (4)
N1—C8	1.462 (4)	C14—C15	1.452 (4)
N2—C14	1.270 (3)		
C7—N1—C8	118.7 (2)	N1—C8—C13	110.2 (2)
C14—N2—C13	119.4 (2)	N2—C13—C8	108.9 (2)
O1—C1—C6	120.1 (2)	N2—C14—C15	122.6 (2)
N1—C7—C6	123.2 (2)	O2—C20—C15	120.9 (3)
O1—C1—C6—C7	2.4 (4)	N1—C8—C13—N2	-58.6 (3)
C8—N1—C7—C6	-180.0 (2)	C13—N2—C14—C15	-175.9 (2)
C1—C6—C7—N1	-1.1 (4)	N2—C14—C15—C20	4.8 (4)
C7—N1—C8—C13	110.0 (3)	C14—C15—C20—O2	-5.2 (4)
C14—N2—C13—C8	123.4 (2)		

Compound (II)*Crystal data*C₂₂H₂₆N₂O₂ $M_r = 350.45$

Monoclinic

 $P2_1/c$ $a = 11.832 (9) \text{ \AA}$ $b = 13.797 (5) \text{ \AA}$ $c = 12.954 (7) \text{ \AA}$ $\beta = 112.51 (4)^\circ$ $V = 1953.6 (20) \text{ \AA}^3$

Z = 4

 $D_x = 1.192 \text{ Mg m}^{-3}$ D_m not measured*Data collection*Delft Instruments FAST TV
area-detector diffractometer

Flat-plate detector

Absorption correction:
none

8708 measured reflections

2965 independent reflections

*Refinement*Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.0451$ $wR(F^2) = 0.2860$

S = 0.688

239 reflections

239 parameters

H-atom parameters not
refined $w = 1/[\sigma^2(F_o^2) + (0.0374P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$ Mo $K\alpha$ radiation $\lambda = 0.71073 \text{ \AA}$ Cell parameters from 250
reflections $\theta = 1.86\text{--}24.97^\circ$ $\mu = 0.076 \text{ mm}^{-1}$

T = 293 (2) K

Block

 $0.30 \times 0.30 \times 0.15 \text{ mm}$

Yellow

1134 observed reflections

 $[I > 2\sigma(I)]$ $R_{\text{int}} = 0.0882$ $\theta_{\max} = 24.97^\circ$ $h = -13 \rightarrow 13$ $k = -14 \rightarrow 15$ $l = -14 \rightarrow 11$ $(\Delta/\sigma)_{\max} = -0.001$ $\Delta\rho_{\max} = 0.185 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.186 \text{ e \AA}^{-3}$

Extinction correction: none

Atomic scattering factors

from *International Tables*
for *Crystallography* (1992),
Vol. C, Tables 4.2.6.8 and
6.1.1.4)

Table 3. Fractional atomic coordinates and equivalent isotropic displacement parameters (\AA^2) for (II)
$$U_{eq} = (1/3)\sum_i \sum_j U_{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j$$

	x	y	z	U_{eq}
O1	0.0234 (2)	0.1323 (2)	0.0819 (2)	0.0892 (7)
O2	-0.4654 (2)	0.23501 (15)	-0.1220 (2)	0.0873 (7)
N1	-0.0929 (2)	0.1855 (2)	-0.1170 (2)	0.0626 (6)
N2	-0.3216 (2)	0.1064 (2)	-0.1349 (2)	0.0649 (7)
C1	0.0817 (3)	0.2181 (2)	0.0971 (3)	0.0676 (8)
C2	0.1739 (3)	0.2354 (3)	0.2011 (3)	0.0827 (10)
C3	0.2380 (3)	0.3206 (3)	0.2212 (3)	0.0954 (11)
C4	0.2107 (3)	0.3905 (3)	0.1390 (3)	0.0971 (11)
C5	0.1189 (3)	0.3736 (2)	0.0370 (3)	0.0851 (10)
C6	0.0519 (3)	0.2878 (2)	0.0115 (3)	0.0615 (7)
C7	-0.0442 (3)	0.2702 (2)	-0.0994 (2)	0.0608 (8)
C8	-0.0800 (3)	0.3517 (2)	-0.1841 (2)	0.0866 (10)
C9	-0.1871 (2)	0.1569 (2)	-0.2258 (2)	0.0615 (8)
C10	-0.1288 (3)	0.1291 (2)	-0.3080 (2)	0.0754 (9)
C11	-0.2254 (3)	0.0974 (2)	-0.4190 (2)	0.0792 (9)
C12	-0.2993 (3)	0.0135 (2)	-0.4037 (2)	0.0885 (10)
C13	-0.3548 (3)	0.0377 (2)	-0.3175 (2)	0.0782 (9)
C14	-0.2588 (2)	0.0727 (2)	-0.2072 (2)	0.0632 (8)
C15	-0.3111 (2)	0.0674 (2)	-0.0414 (3)	0.0618 (8)
C16	-0.2319 (3)	-0.0184 (2)	0.0096 (3)	0.0910 (10)
C17	-0.3833 (3)	0.1107 (2)	0.0170 (2)	0.0616 (8)
C18	-0.3814 (3)	0.0734 (2)	0.1181 (3)	0.0791 (9)
C19	-0.4496 (3)	0.1143 (3)	0.1723 (3)	0.0923 (11)
C20	-0.5215 (3)	0.1930 (3)	0.1276 (3)	0.0947 (12)
C21	-0.5278 (2)	0.2323 (2)	0.0279 (3)	0.0819 (10)
C22	-0.4578 (3)	0.1927 (3)	-0.0265 (2)	0.0683 (9)

Table 4. Selected geometric parameters (\AA , $^\circ$) for (II)

O1—C1	1.346 (3)	C6—C7	1.471 (4)
O2—C22	1.340 (3)	C7—C8	1.514 (3)
N1—C7	1.284 (3)	C9—C14	1.512 (3)
N1—C9	1.477 (3)	C15—C17	1.467 (4)
N2—C15	1.288 (3)	C15—C16	1.497 (4)
N2—C14	1.475 (3)		
C7—N1—C9	122.4 (2)	N1—C9—C14	108.5 (2)
C15—N2—C14	125.1 (2)	N2—C14—C9	108.0 (2)
O1—C1—C6	121.7 (3)	N2—C15—C17	116.7 (3)
N1—C7—C6	116.8 (3)	N2—C15—C16	124.3 (3)
N1—C7—C8	124.4 (3)	O2—C22—C17	121.9 (3)
O1—C1—C6—C7	-0.4 (4)	N1—C9—C14—N2	63.1 (3)
C9—N1—C7—C6	-177.7 (2)	C14—N2—C15—C17	-178.8 (2)
C1—C6—C7—N1	-5.3 (3)	N2—C15—C17—C22	-1.7 (4)
C7—N1—C9—C14	-157.9 (2)	C15—C17—C22—O2	-0.5 (4)
C15—N2—C14—C9	-124.0 (3)		

H atoms were added at calculated positions and refined using a riding model. Anisotropic displacement parameters were used for all non-H atoms; each H atom was given an isotropic displacement parameter equal to 1.2 (or 1.5 for methyl and —OH H atoms) times the equivalent isotropic displacement parameter of the atom to which it is attached. For compound (II) the structure was solved by direct methods in *Pc* and then transformed into the space group *P2₁/c*.

For both compounds, data collection: *MADNES* (Pflugrath & Messerschmidt, 1992); cell refinement: *MADNES*; data reduction: *SHELXTL-Plus* (Sheldrick, 1990); program(s) used to solve structures: *SHELXTL-Plus*; program(s) used to refine structures: *SHELXL93* (Sheldrick, 1993); molecular graphics: *SHELXTL-Plus*; software used to prepare material for publication: *SHELXL93*.

We wish to thank Professor M. Hursthouse and the EPSRC X-ray Crystallographic Service (University of Wales, Cardiff) for collecting the diffraction data. We also wish to acknowledge the use of the EPSRC's

Chemical Database Service (Allen *et al.*, 1991) at Daresbury. Financial assistance was provided by BP (JPC) and ICI (JCC).

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: KA1152). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Allen, F. H., Davies, J. E., Galloy, J. J., Johnson, O., Kennard, O., Macrae, C. F., Mitchell, E. M., Mitchell, G. F., Smith, J. M. & Watson, D. G. (1991). *J. Chem. Inf. Comput. Sci.* **31**, 187–204.
- Corden, J. P., Errington, W., Moore, P. & Wallbridge, M. G. H. (1996). *Acta Cryst.* **C52**, 125–127.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Pflugrath, J. W. & Messerschmidt, A. (1992). *Munich Area Detector Systems*. Enraf-Nonius, Delft, The Netherlands.
- Sheldrick, G. M. (1990). *SHELXTL-Plus*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.

Acta Cryst. (1996). **C52**, 1017–1019

8-Hydroxy-7-nitroquinoline-5-sulfonic Acid Monohydrate

THAILAM PILLAI BALASUBRAMANIAN† AND PACKIANATHAN THOMAS MUTHIAH*

Department of Chemistry, Bharathidasan University, Tiruchirapalli 620 024, India

(Received 3 May 1995; accepted 23 August 1995)

Abstract

The title compound, $\text{C}_9\text{H}_6\text{N}_2\text{O}_6\text{S}\cdot\text{H}_2\text{O}$, is zwitterionic (*i.e.* 8-hydroxy-7-nitroquinolinium-5-sulfonate monohydrate) with a deprotonated sulfonic group and a protonated quinoline N atom which interacts with the hydroxy O atom. The H atom of the hydroxy group is distal to the N atom and is involved in an interaction with an O atom of the nitro group, which is oriented at $32.2(1)^\circ$ with respect to the quinoline ring system.

Comment

Derivatives of 8-hydroxyquinoline (oxine) are known for their anti-amoebic, antibacterial and antifungal activities, which can be correlated to their metal chelating

† Present address: Department of Physics, Nehru Memorial College, Puthanampatti 621 007, India.