

References

- Beurskens, P. T., Admiraal, G., Beurskens, G., Bosman, W. P., Garcia-Granda, S., Gould, R. O., Smits, J. M. M. & Smykalla, C. (1992). *The DIRDIF Program System*. Technical Report. Crystallography Laboratory, University of Nijmegen, The Netherlands.
- Cromer, D. T. & Waber, J. T. (1974). *International Tables for X-ray Crystallography*, Vol. IV, Table 2.2A, pp. 72–98. Birmingham: Kynoch Press. (Present distributor Kluwer Academic Publishers, Dordrecht.)
- Enraf–Nonius (1988). *Diffraction Control Software*. Enraf–Nonius, Delft, The Netherlands.
- Fong, L. K., Fox, J. R. & Cooper, N. J. (1987). *Organometallics*, **6**, 223–231.
- Gibson, D. H., Ding, Y., Mashuta, M. S. & Richardson, J. F. (1996). *Acta Cryst. C52*, 559–560.
- Hermann, G. S., Alt, H. G. & Thewalt, U. (1990). *J. Organomet. Chem.* **399**, 83–92.
- Hoffman, D. M., Chester, N. D. & Fay, R. C. (1983). *Organometallics*, **2**, 48–52.
- Huffman, J. C., Moloy, K. G., Marsella, J. A. & Caulton, K. G. (1980). *J. Am. Chem. Soc.* **102**, 3009–3014.
- Jessop, P. G., Ikariya, T. & Noyori, R. (1995). *Chem. Rev.* **95**, 259–272.
- Johnson, C. K. (1976). *ORTEP*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Molecular Structure Corporation (1993). *TEXSAN. Single Crystal Structure Analysis Software*. Version 1.6. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Wöhrle, T. & Thewalt, U. (1994). *J. Organomet. Chem.* **468**, C1–C3.
- Zachariassen, W. H. (1967). *Acta Cryst.* **23**, 558–564.

Acta Cryst. (1996). **C52**, 1616–1618

A Diphenylamido–Bromide Complex of Niobium: Dibromo[tris(diphenylamido)]-niobium(V) Tetrahydrofuran Solvate

DAVID M. HOFFMAN AND SRI PRAKASH RANGARAJAN

Department of Chemistry, University of Houston, Houston, Texas 77204, USA. E-mail: hoffman@uh.edu

(Received 13 November 1995; accepted 17 January 1996)


Abstract

NbBr₅ reacts with three equivalents of Me₃SiNPh₂ in ether to give the title compound, [NbBr₂(C₁₂H₁₀N)₃].·C₄H₈O. Nb(NPh₂)₃Br₂ has trigonal bipyramidal geometry with axial bromide ligands.

Comment

Several diethylamido–chloride and –fluoride complexes of niobium have been prepared from NbX₅ and trimethylsilyldiethylamine, including Nb(NEt₂)₃Cl₂, [Nb(NEt₂)₂Cl₃]₂, Nb(NEt₂)F₄ and Nb(NEt₂)₂F₃ (Chao,

Polson & Wigley, 1990; Fugglè, Sharp & Winfield, 1972; Fowles & Pleass, 1957). We were interested in preparing similar amido–bromide derivatives to obtain a selection of complexes available for reduction studies (Hoffman & Rangarajan, 1993; Bott, Hoffman & Rangarajan, 1995). In this paper we describe the preparation and structure of Nb(NPh₂)₃Br₂, (I).



NbBr₅ reacts with three equivalents of Me₃SiNPh₂ or LiNPh₂ in ether to give Nb(NPh₂)₃Br₂. The compound crystallizes from thf/CH₃CN as dark red needles, but it is difficult to isolate cleanly from one or two persistent unidentified contaminants. The preparation based on Me₃SiNPh₂ more consistently gives clean crystalline material than the one using LiNPh₂. Proton NMR spectra for Nb(NPh₂)₃Br₂ show only resonances arising from one kind of phenyl group, consistent with the solid-state structure.

Nb(NPh₂)₃Br₂ crystallizes with one molecule of thf in the lattice. The Nb(NPh₂)₃Br₂ molecules are trigonal bipyramidal with axial bromide ligands (Fig. 1). A crystallographic threefold axis passes through the Nb and Br atoms. The amido ligand NC₂ ‘blades’ deviate by less than 4° from the trigonal plane defined by the three N atoms. In this configuration the amido N *pπ*-donor orbitals can interact only with niobium *d_{xz,yz}* and not at all with *d_{xy,x2-y2}*. This is expected since *d_{xy}* and *d_{x2-y2}* are involved in Nb–N *σ* bonding and would be too high in energy for an effective interaction with the *pπ* orbitals of nitrogen.

The Nb–N bond lengths [1.935 (8) Å] are similar to those in Nb(NMe₂)₅ [1.977 (17)–2.044 (14) Å]




Fig. 1. View of Nb(NPh₂)₃Br₂ showing the labelling of the non-H atoms. Displacement ellipsoids are shown at the 40% probability level.

(Heath & Hursthouse, 1971; Bradley & Thomas, 1962). The Nb—Br bond lengths [2.637 (4) and 2.647 (4) Å] are within the range of terminal Nb—Br distances observed in other neutral complexes {e.g. [NbBr₄(μ-Br)₂] [2.403 (3), 2.462 (2) Å] (Müller & Klingelhöfer, 1983), NbBr₄(PMe₂Ph)₃ [2.541 (4)–2.621 (4) Å], NbBr₃(PMe₂Ph)₃ [2.554 (1)–2.573 (1) Å] (Cotton, Diebold & Roth, 1985) and Nb₂Br₆(SC₄H₈)₃ [2.604 (7)–2.648 (6) Å] (Templeton, Dorman, Clardy & McCarley, 1978)}.

Experimental

The synthesis was carried out under an inert atmosphere. To a dark brown solution of NbBr₅ (1.00 g, 2.03 mmol) in ether, Me₃SiNPh₂ (1.43 g, 6.09 mmol) was added slowly over a period of 10 min. The colour of the solution changed immediately to reddish brown. The reaction mixture was stirred for 48 h. The mixture was then stripped *in vacuo*, and the brown residue was extracted with 30 cm³ of toluene and the extracts filtered through celite. The red–brown filtrate was stripped *in vacuo*, and the residue was extracted with thf/CH₃CN (4:1 ratio; 15 cm³). The volume of the extract was reduced *in vacuo* to 5 cm³. The solution was stored for 1 d at 237 K. This produced dark red needles. Yield: 1.32 g, 88%. The sample for analysis was washed with 20 cm³ of hexane and dried *in vacuo*. Analysis: calculated for NbBr₂N₃C₃₆H₃₀ C 57.09, H 3.99, N 5.54%; found C 57.75, H 3.73, N 5.40%. ¹H NMR (C₆D₆): δ 7.65 (*d*, 6H, *p*-H), 6.64 (*t*, 12H, *m*-H), 6.49 *p.p.m.* (*t*, 12H, *o*-H).

Crystal data

[NbBr₂(C₁₂H₁₀N)₃].C₄H₈O
M_r = 829.54
 Hexagonal
*P*6₃
a = 13.671 (3) Å
c = 11.400 (3) Å
V = 1845 Å³
Z = 2
D_x = 1.49 Mg m⁻³
D_m not measured

Mo Kα radiation
 λ = 0.71073 Å
 Cell parameters from 38 reflections
 θ = 15–29°
 μ = 2.49 mm⁻¹
T = 223 K
 Hexagonal column
 0.70 × 0.20 × 0.20 mm
 Blood red

Data collection

Nicolet R3m/V diffractometer
 ω-scan technique
 Absorption correction: ψ scan (SHELXTL-Plus; Sheldrick, 1987)
T_{min} = 0.529, *T_{max}* = 0.565
 1153 measured reflections
 1153 independent reflections

728 observed reflections [*I* > 3σ(*I*)]
 θ_{max} = 25°
h = 0 → 15
k = 0 → 15
l = 0 → 13
 2 standard reflections monitored every 100 reflections
 intensity decay: none

Refinement

Refinement on *F*
R = 0.029
wR = 0.026
S = 2.2

(Δ/σ)_{max} = 0.3
 Δρ_{max} = 0.4 e Å⁻³
 Δρ_{min} = -0.2 e Å⁻³
 Extinction correction: none

728 reflections
 146 parameters
 Only H-atom *U*'s refined
w = 1/σ²(*F_o*)

Atomic scattering factors from SHELXTL-Plus
 Absolute configuration: Rogers (1981) parameter = 0.93

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (Å²)

$$U_{eq} = (1/3)\sum_i \sum_j U_{ij} a_i^* a_j^*$$

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{eq}</i>
Nb	1/3	2/3	0.3000	0.031 (1)
Br(1)	1/3	2/3	0.5313 (4)	0.044 (1)
Br(2)	1/3	2/3	0.0678 (4)	0.041 (1)
N	0.2776 (5)	0.7719 (5)	0.2986 (13)	0.038 (2)
C(1)	0.1609 (6)	0.7434 (6)	0.3056 (17)	0.043 (2)
C(2)	0.1195 (7)	0.7728 (8)	0.2048 (10)	0.053 (2)
C(3)	0.0077 (8)	0.7457 (9)	0.2038 (13)	0.076 (2)
C(4)	-0.0609 (7)	0.6888 (8)	0.2940 (16)	0.077 (2)
C(5)	-0.0218 (9)	0.6582 (8)	0.3941 (12)	0.077 (2)
C(6)	0.0918 (7)	0.6856 (7)	0.3976 (11)	0.056 (2)
C(7)	0.3578 (6)	0.8932 (6)	0.2988 (15)	0.041 (2)
C(8)	0.4268 (7)	0.9438 (7)	0.2038 (10)	0.048 (2)
C(9)	0.5015 (8)	1.0612 (7)	0.2073 (10)	0.055 (2)
C(10)	0.5072 (7)	1.1226 (7)	0.3071 (17)	0.060 (2)
C(11)	0.4409 (8)	1.0714 (8)	0.4010 (12)	0.058 (2)
C(12)	0.3642 (7)	0.9541 (7)	0.3979 (10)	0.044 (2)
C(13)†	1.014 (2)	0.956 (2)	0.528 (2)	0.102 (2)
C(14)†	0.955 (2)	0.958 (2)	0.3846 (12)	0.102 (2)
C(15)†	0.893 (2)	0.920 (2)	0.437 (2)	0.102 (2)
C(16)†	0.908 (2)	0.872 (2)	0.482 (2)	0.102 (2)
C(17)†	0.930 (3)	0.988 (3)	0.529 (3)	0.102 (2)
C(18)†	0.963 (3)	0.895 (2)	0.503 (3)	0.102 (2)

† Partial occupancy (see below).

Table 2. Selected geometric parameters (Å, °)

Nb—Br(1)	2.637 (4)	Nb—Br(2)	2.647 (4)
Nb—N	1.935 (8)	N—C(1)	1.444 (11)
N—C(7)	1.461 (8)		
Br(2)—Nb—N	89.5 (4)	Br(1)—Nb—N	90.5 (4)
Nb—N—C(1)	126.3 (4)	Nb—N—C(7)	119.5 (5)
C(1)—N—C(7)	114.0 (7)		

During the final stages of refinement, a diffuse area of residual electron density, characteristic of included solvent, was located around a threefold axis. On the basis of NMR data, the solvent in the crystals was assumed to be thf. As thf has so many possible configurations, however, it was impossible to refine it as a rigid body. Thus, the positions in the difference map having the highest electron density were each refined as C atoms having population factors of 0.333 (C13–C16) or 0.222 (C17 and C18). To determine the chirality of the molecule in the crystal, a coefficient multiplying Δ*f*' was refined, and this indicated that the absolute structure reported herein is correct.

Data collection: SHELXTL-Plus (Sheldrick, 1987). Cell refinement: SHELXTL-Plus. Data reduction: SHELXTL-Plus. Program(s) used to solve structure: SHELXTL-Plus. Program(s) used to refine structure: SHELXTL-Plus. Molecular graphics: SHELXTL-Plus. Software used to prepare material for publication: local program.

We acknowledge the Robert A. Welch Foundation and the American Chemical Society–Petroleum Research Fund for support of this work. We thank Dr James D. Korp for his technical assistance with the X-ray crystal structure determination. DMH is a 1992–1994 Alfred P. Sloan Research Fellow.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: SK1001). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Bott, S. G., Hoffman, D. M. & Rangarajan, S. P. (1995). *Inorg. Chem.* **34**, 4305–4310.
- Bradley, D. C. & Thomas, I. M. (1962). *Can. J. Chem.* **40**, 449–454.
- Chao, Y.-W., Polson, S. & Wigley, D. E. (1990). *Polyhedron*, **22**, 2709–2716.
- Cotton, F. A., Diebold, M. P. & Roth, W. J. (1985). *Inorg. Chim. Acta*, **105**, 41–49.
- Fowles, G. W. A. & Pleass, C. M. (1957). *J. Chem. Soc.* pp. 2078–2080.
- Fuggle, J. C., Sharp, D. W. A. & Winfield, J. M. (1972). *J. Chem. Soc. Dalton Trans.* pp. 1766–1768.
- Heath, C. & Hursthouse, M. B. (1971). *J. Chem. Soc. Chem. Commun.* pp. 143–144.
- Hoffman, D. M. & Rangarajan, S. P. (1993). *Polyhedron*, **12**, 2899–2900.
- Müller, U. & Klingelhöfer, P. (1983). *Z. Naturforsch. Teil B*, **38**, 559–561.
- Rogers, D. (1981). *Acta Cryst.* **A37**, 734–741.
- Sheldrick, G. M. (1987). *SHELXTL-Plus*. Release 3.4 for Nicolet R3m/V crystallographic system. Nicolet Instrument Corporation, Madison, Wisconsin, USA.
- Templeton, J. L., Dorman, W. C., Clardy, J. C. & McCarley, R. E. (1978). *Inorg. Chem.* **17**, 1263–1267.

Acta Cryst. (1996). **C52**, 1618–1621

Potassium Aqua(3,4-toluenediamine-*N,N,N',N'*-tetraacetato)ferrate(III)–Water (1/1.5)

M. HERNÁNDEZ-PADILLA,^a J. SANCHIZ,^a S. DOMÍNGUEZ,^a A. MEDEROS,^{a*} J. M. ARRIETA^b AND F. J. ZUÑIGA^c

^aDepartamento de Química Inorgánica, Universidad de La Laguna, Tenerife, Canary Islands, Spain, ^bDepartamento de Química Inorgánica, Universidad del País Vasco, Bilbao, Spain, and ^cDepartamento de Física de la Materia Condensada, Universidad del País Vasco, Bilbao, Spain

(Received 22 March 1995; accepted 26 February 1996)

Abstract


The ligand 3,4-toluenediamine-*N,N,N',N'*-tetraacetate in the title compound, K[Fe(C₁₅H₁₄N₂O₈)(H₂O)].1.5H₂O, is hexadentate, but the central iron(III) ion is seven-coordinate, with one coordinated water molecule. The complex has roughly pentagonal-bipyramidal geometry. The Fe—N bond lengths are 2.304(3) Å in the *para*

position and 2.349(3) Å in the *meta* position with respect to the methyl group. The presence of the CH₃ group increases the basicity of the N atom in the *para* position, shortening the length of the Fe—N bond.

Comment

The preparation of coordinating agents derived from aromatic diamines is of special interest since the use of N atoms for coordination to a single cation is directly related to their situation in *ortho*, *meta* or *para* positions. Thus, in the case of diaminetetramethylene-carboxylic acids, derived from *o*-phenylenediamines, the closer proximity of the N atoms permits the simultaneous coordination of both to the same metal cation, as has been established by the X-ray crystallographic determination of the structures of the complexes of *o*-phdta acid (*ortho*-phenylenediamine-*N,N,N',N'*-tetraacetic acid) with Co^{II} (McCandlish, Michael, Neal, Lingafelter & Rose, 1978), Mn^{II} (Nakasuka, Azuma, Katayama, Honda, Tanaka & Tanaka, 1985), Zn^{II} (Azuma, Nakasuka & Tanaka, 1986), Cu^{II} (Nakasuka, Azuma & Tanaka, 1986*a*), Cd^{II} (Nakasuka, Azuma & Tanaka, 1986*b*), Mg^{II} (Nakasuka & Shiro, 1989) and Fe^{III} (Mizuno, Funahashi, Nakasuka & Tanaka, 1991). Contrary to this is the behaviour of diaminetetramethylenecarboxylic acids derived from *meta*- or *para*-phenylenediamines, which can only coordinate one N atom of the ligand to the same metallic cation. Therefore, these ligands can bridge metal ions, as has been proven by the preparation of bimetallic species *M*₂*L* for *p*-phdta acid (*para*-phenylenediamine-*N,N,N',N'*-tetraacetic acid) (Ruíz-Pérez, Rodríguez, Rodríguez Romero, Mederos, Gili & Martín-Zarza, 1990) and dimer species *M*₂*L*₂ for *m*-phdta acid (*meta*-phenylenediamine-*N,N,N',N'*-tetraacetic acid) (Mederos *et al.*, 1990).

In the present work, the solid crystalline complex K[Fe(H₂O)(3,4-tdta)].1.5H₂O, (I), was prepared and characterized. 3,4-tdta (3,4-toluenediamine-*N,N,N',N'*-tetraacetic acid) is a potentially hexacoordinating ligand, analogous to *o*-phdta and edta.



The determination by X-ray diffraction of the structure of the complex [Fe(OH₂)(*o*-phdta)]⁻ (Mizuno, Funahashi, Nakasuka & Tanaka, 1991) shows that Fe^{III} is heptacoordinate, with roughly bipyramidal pentagonal