

C45	0.9165 (5)	-0.0075 (5)	0.3718 (4)	0.065 (2)
C46	0.8498 (4)	-0.0046 (4)	0.3299 (4)	0.055 (2)
C51	0.6887 (4)	-0.0115 (4)	0.2447 (3)	0.035 (2)
C52	0.7106 (4)	-0.0865 (4)	0.2373 (3)	0.051 (2)
C53	0.6559 (5)	-0.1433 (4)	0.2460 (4)	0.065 (2)
C54	0.5798 (5)	-0.1253 (5)	0.2593 (4)	0.068 (3)
C55	0.5571 (4)	-0.0526 (5)	0.2663 (4)	0.061 (2)
C56	0.6103 (5)	0.0035 (4)	0.2601 (3)	0.046 (2)
C61	0.7112 (3)	0.1503 (3)	0.2479 (3)	0.033 (2)
C62	0.6722 (4)	0.1907 (4)	0.1981 (3)	0.046 (2)
C63	0.6320 (4)	0.2561 (4)	0.2149 (4)	0.055 (2)
C64	0.6305 (4)	0.2797 (4)	0.2812 (5)	0.062 (2)
C65	0.6680 (5)	0.2397 (5)	0.3313 (4)	0.063 (2)
C66	0.7087 (4)	0.1753 (4)	0.3154 (3)	0.053 (2)

† Cp1 and Cp2 are the centroids of rings C11–C15 and C21–C25, respectively.

Table 2. Selected geometric parameters (Å, °)

Fe1—Cp1	1.646 (7)	C11—C12	1.424 (8)
Fe1—Cp2	1.655 (7)	C11—C15	1.425 (8)
P1—C1	1.845 (6)	C12—C13	1.436 (9)
P1—C41	1.788 (6)	C13—C14	1.384 (9)
P1—C51	1.794 (7)	C14—C15	1.402 (9)
P1—C61	1.790 (6)	C21—C22	1.380 (10)
O2—C2	1.217 (7)	C21—C25	1.397 (10)
C1—C2	1.545 (8)	C22—C23	1.403 (10)
C1—C11	1.504 (8)	C23—C24	1.389 (10)
C2—C31	1.498 (9)	C24—C25	1.380 (10)
Cp1—Fe1—Cp2	177.7 (3)	P1—C1—C2	107.9 (4)
C41—P1—C51	111.2 (3)	P1—C1—C11	114.6 (4)
C41—P1—C61	107.0 (3)	C2—C1—C11	110.4 (5)
C61—P1—C51	109.3 (3)	O2—C2—C1	119.6 (6)
C1—P1—C61	108.9 (3)	O2—C2—C31	120.7 (7)
C1—P1—C51	108.8 (3)	C1—C2—C31	119.7 (6)
C1—P1—C41	111.5 (3)		
C41—P1—C1—C11	-46.5 (6)	P1—C1—C2—O2	-67.4 (7)
C51—P1—C1—C11	-169.6 (5)	C11—C1—C2—C31	-123.4 (6)
C61—P1—C1—C11	71.4 (5)	P1—C1—C2—C31	110.7 (6)
C11—C1—C2—O2	58.6 (8)		

Examination of the structure with *PLATON* (Spek, 1995a) showed that there were no solvent-accessible voids in the crystal lattice.

Data collection: *CAD-4-PC* (Enraf–Nonius, 1992). Cell refinement: *SET4* and *CELDIM* (Enraf–Nonius, 1992). Data reduction: *DATRD2* in *NRCVAX94* (Gabe, Le Page, Charland, Lee & White, 1989). Program(s) used to solve structure: *NRCVAX94*. Program(s) used to refine structure: *NRCVAX94* and *SHELXL93* (Sheldrick, 1993). Molecular graphics: *NRCVAX94*, *ORTEPII* (Johnson, 1976), *PLATON*, *PLUTON* (Spek 1995b). Software used to prepare material for publication: *NRCVAX94*, *SHELXL93* and *WordPerfect* macro *PREPCIF*.

GF thanks NSERC (Canada) for research grants. SZA thanks CVCP (UK) and the University of St Andrews for financial support.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: CF1086). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

## References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.  
Alver, E. & Høltedahl, B. H. (1967). *Acta Chem. Scand.* **21**, 359–370.

- Archer, S. J., Modro, T. A. & Nassimbeni, L. R. (1981). *Phosphorus Sulfur*, **11**, 101–110.  
Bondi, A. (1984). *J. Phys. Chem.* **68**, 441–451.  
Enraf–Nonius (1992). *CAD-4-PC*. Version 1.1. Enraf–Nonius, Delft, The Netherlands.  
Ferguson, G., Gallagher, J. F., Glidewell, C. & Zakaria, C. M. (1994). *Acta Cryst.* **B50**, 146–150.  
Ferguson, G., Glidewell, C., Gosney, I., Lloyd, D., Metcalfe, S. & Lumbroso, H. (1988). *J. Chem. Soc. Perkin Trans. 2*, pp. 1829–1837.  
Ferguson, G., Glidewell, C., Lloyd, D., Metcalfe, S. & Ruhl, B. L. (1987). *J. Chem. Res. (S)*, p. 32; *J. Chem. Res. (M)*, pp. 458–473.  
Ferguson, G., Lough, A. J., McAlees, A. J. & McCrindle, R. (1993). *Acta Cryst.* **C49**, 573–577.  
Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.  
Glidewell, C., Zakaria, C. M., Ferguson, G. & Gallagher, J. F. (1994). *Acta Cryst.* **C50**, 233–238.  
Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.  
Mazhar-ul-Haque (1979). *Acta Cryst.* **B35**, 2601–2604.  
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.  
Pauson, P. L. & Watts, W. E. (1963). *J. Chem. Soc.* pp. 2990–2995.  
Penicaud, A., Perez-Benitez, A., Gleason, V. R., Munoz, P. E. & Escudro, R. (1993). *J. Am. Chem. Soc.* **115**, 10392–10393.  
Schweizer, E. E., Baldacchini, C. J. & Rheingold, A. L. (1989). *Acta Cryst.* **C45**, 1236–1239.  
Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.  
Spek, A. L. (1995a). *PLATON. Molecular Geometry Program*. July 1995 version. University of Utrecht, The Netherlands.  
Spek, A. L. (1995b). *PLUTON. Molecular Graphics Program*. July 1995 version. University of Utrecht, The Netherlands.  
Staples, R. J., Carlson, T., Wang, S. & Fackler, J. P. (1995). *Acta Cryst.* **C51**, 498–500.

*Acta Cryst.* (1996). **C52**, 1636–1638

## *trans*-Chloromethylbis(tri-*p*-tolylarsine)-platinum(II)

STEFANUS OTTO AND ANDREAS ROODT\*

Department of Chemistry, University of the Orange Free State, Bloemfontein 9300, South Africa. E-mail: char@uovsvml.uovs.ac.za


(Received 29 June 1995; accepted 15 January 1996)

## Abstract

The title compound, *trans*-[PtCl(CH<sub>2</sub>){As(C<sub>6</sub>H<sub>4</sub>CH<sub>3</sub>)<sub>3</sub>]<sub>2</sub>, is one of the few bis tertiary arsine complexes of platinum(II). The complex exhibits square-planar geometry with the two arsine ligands in a non-linear *trans* orientation with an As1—Pt—As2 bond angle of 175.66 (4)°. The large *trans* influence of the methyl substituent is manifested in the long Pt—Cl bond distance of 2.397 (3) Å.

### Comment

*trans*-Chloromethylbis(tri-*p*-tolylarsine)platinum(II), (I), has been structurally characterized and is one of the few bis(AsX<sub>3</sub>)-platinum(II) (X = alkyl or aryl) complexes isolated to date. The complex was investigated as part of a continued study of complexes of the form PtCl(CH<sub>3</sub>)(L)<sub>2</sub>, with L being different tertiary phosphine or arsine ligands. The *trans* influence of the methyl group is less pronounced in the case of the bis(AsPh<sub>3</sub>) complex compared with the bis(PPh<sub>3</sub>) complex. In the latter case, Pt—Cl bond distances of 2.431 (3) and 2.415 (5) Å were respectively found in two independent structural determinations for the PPh<sub>3</sub> compound (Bardi & Piazzesi, 1981; Otto, Roodt & Leipoldt, 1995), compared with the value of 2.410 (2) Å for the AsPh<sub>3</sub> compound obtained from a previous study (Roodt, Otto & Leipoldt, 1995). This is in agreement with the electronic influence PPh<sub>3</sub> > AsPh<sub>3</sub> (Cheeseman, Odell & Raethel, 1968). The As(*p*-Tol)<sub>3</sub> and AsPh<sub>3</sub> ligands have the same steric demand while the former is a considerably stronger base, an effect that is evident from the respective <sup>1</sup>H NMR spectra of the compounds. The



methyl resonance is shifted towards higher field for the As(*p*-Tol)<sub>3</sub> compound (0.04 p.p.m.) relative to the AsPh<sub>3</sub> compound (0.07 p.p.m.). This is in agreement with more effective electronic shielding of the methyl group, as to be expected from the larger electron-donating capability of the As(*p*-Tol)<sub>3</sub> ligand. The increase in electron density at the Pt centre is, however, not manifested in an increase in the Pt—Cl bond distance, *i.e.* 2.397 (3) Å found in (I) compared with 2.410 (2) Å for the AsPh<sub>3</sub> compound. The explanation for this phenomenon comes from the fact that an increase in the Pt—Me bond distance *trans* to the Cl atom, 2.111 (9) Å, compared with the Pt—Me distance of 2.073 (8) Å in the case of the AsPh<sub>3</sub> complex, is observed. This leads to a weaker effective *trans* influence of the methyl group and thus a decrease in the Pt—Cl bond distance. The larger steric demand of the methyl group compared with that of the chloro ligand is well illustrated in the platinum-coordination polyhedron bond angles which deviate significantly from normal square-planar values, *i.e.* 175.66 (4) and 173.7 (3)°, respectively, for the As1—Pt—As2 and C1—Pt—Cl1 angles. This is also evident from the C41—As2—Pt and C21—As1—Pt bond angles, which deviate by almost 10° from normal tetrahedral values.




Fig. 1. Structure showing numbering scheme and displacement ellipsoids (30% probability). The H atoms except for the methyl H atoms are omitted for clarity.

### Experimental

The title compound was prepared by the addition of an excess of As(C<sub>6</sub>H<sub>4</sub>CH<sub>3</sub>)<sub>3</sub> (1.05 g, 3 mmol) in acetone (20 ml) to *trans*-[PtCl(CH<sub>3</sub>){S(CH<sub>3</sub>)<sub>2</sub>}]<sub>2</sub> (Scott & Puddephatt, 1983) (500 mg, 1.35 mmol) in acetone (10 ml). The reaction mixture was stirred at room temperature for 1 h and evaporated to dryness to remove the liberated S(Me<sub>2</sub>)<sub>2</sub>. The residue was redissolved in acetone and filtered; slow evaporation of the solvent gave the desired product (1.06 g, yield >80%) as colourless plates. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ(CH<sub>3</sub>) 0.04 p.p.m., <sup>2</sup>J<sub>Pt—H</sub> 78 Hz.

### Crystal data

[PtCl(CH<sub>3</sub>)(C<sub>21</sub>H<sub>21</sub>As)<sub>2</sub>]  
*M<sub>r</sub>* = 942.17  
 Orthorhombic  
*Pna*2<sub>1</sub>  
*a* = 21.771 (4) Å  
*b* = 10.608 (2) Å  
*c* = 16.885 (3) Å  
*V* = 3899.5 (12) Å<sup>3</sup>  
*Z* = 4  
*D<sub>x</sub>* = 1.605 Mg m<sup>-3</sup>  
*D<sub>m</sub>* = 1.592 Mg m<sup>-3</sup>  
*D<sub>m</sub>* measured by flotation in NaI/H<sub>2</sub>O

Mo Kα radiation  
 λ = 0.71073 Å  
 Cell parameters from 25 reflections  
 θ = 17–22°  
 μ = 5.379 mm<sup>-1</sup>  
*T* = 293 (2) K  
 Thick plate  
 0.56 × 0.26 × 0.24 mm  
 Colourless

### Data collection

Enraf–Nonius CAD-4 diffractometer  
 θ/2θ scans  
 Absorption correction: numerical by integration from crystal shape (Akselrud, Grin, Zavalii, Petcharsky & Fundamentalsky, 1989)  
*T<sub>min</sub>* = 0.397, *T<sub>max</sub>* = 0.487

3964 measured reflections  
 2282 independent reflections  
 2282 observed reflections  
 [I > 2σ(I)]  
 θ<sub>max</sub> = 24.97°  
*h* = 0 → 25  
*k* = 0 → 12  
*l* = 0 → 20  
 3 standard reflections  
 frequency: 60 min  
 intensity decay: <1%

## Refinement

Refinement on  $F^2$   
 $R[F^2 > 2\sigma(F^2)] = 0.0226$   
 $wR(F^2) = 0.0545$   
 $S = 1.077$   
 2282 reflections  
 432 parameters  
 H atoms riding  
 $w = 1/[\sigma^2(F_o^2) + (0.0475P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = -0.083$

$\Delta\rho_{\max} = 0.914 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -1.153 \text{ e } \text{\AA}^{-3}$   
 Extinction correction: none  
 Atomic scattering factors  
 from *International Tables for Crystallography* (1992, Vol. C, Tables 4.2.6.8 and 6.1.1.4)  
 Absolute configuration:  
 Flack (1983) parameter  
 $= -0.004 (10)$

Cl—Pt—As1	91.0 (3)	C21—As1—Pt	119.3 (3)
Cl—Pt—As2	91.9 (3)	C31—As1—Pt	110.4 (3)
As1—Pt—As2	175.66 (4)	C11—As1—Pt	117.1 (3)
Cl—Pt—C11	173.7 (3)	C51—As2—Pt	110.5 (4)
As2—Pt—C11	88.66 (7)	C61—As2—Pt	118.5 (3)
As1—Pt—C11	88.84 (7)	C41—As2—Pt	118.7 (3)

The complex crystallizes in the non-centrosymmetric orthorhombic space group  $Pna2_1$ , which is a result of the substantial deviation from linearity of the As1—Pt—As2 moiety [bond angle  $175.66(4)^\circ$ ]. Attempted refinement in the centrosymmetric space group  $Pnma$  resulted in a disordered structure with non-relevant coordination and an  $R$  value (based on  $F$ )  $> 15\%$ . The phenyl H atoms and methyl H atoms were positioned assuming C—H distances of 0.96 and 0.93 Å, respectively, and were constrained riding on their parent C atoms. A single orientation parameter was refined for the methyl groups with an overall H-atom  $U_{\text{iso}}$  of  $0.062 \text{ \AA}^2$ . Scattering factors, dispersion corrections and absorption coefficients were taken from *International Tables for Crystallography* (1992, Vol. C, Tables 6.1.1.4, 4.2.6.8 and 4.2.4.2).

Data collection: CAD-4 diffractometer software. Cell refinement: CAD-4 diffractometer software. Data reduction: *Profit* (Streltsov & Zavodnik, 1989). Program(s) used to solve structure: *SHELXS86* (Sheldrick, 1990). Program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993). Molecular graphics: *ORTEPII* (Johnson, 1976). Software used to prepare material for publication: *SHELXL93*.

Financial assistance of the South African FRD and the Research Fund of the University of the Orange Free State is gratefully acknowledged.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: AB1292). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

## References

- Akselrud, L. G., Grin, Y. N., Zavalii, P. Y., Pecharsky, V. K. & Fundamensky, V. S. (1989). *XIIIth European Crystallographic Meeting, Moscow, Collected Abstracts*, Vol. 3, p. 155.
- Bardi, R. & Piazzesi, A. M. (1981). *Inorg. Chim. Acta*, **47**, 249–254.
- Cheesman, T. P., Odell, A. L. & Raethel, H. A. (1968). *J. Chem. Soc. Chem. Commun.* pp. 1496–1598.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Otto, S., Roodt, A. & Leipoldt, J. G. (1995). *S. Afr. J. Chem.* In the press.
- Roodt, A., Otto, S. & Leipoldt, J. G. (1995). *Acta Cryst.* **C51**, 1105–1106.
- Scott, J. D. & Puddephatt, R. J. (1983). *Organometallics*, **2**, 1643–1648.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Streltsov, V. A. & Zavodnik, V. E. (1989). *Kristallografiya*, **34**(6), 1369–1375.

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

$$U_{\text{eq}} = (1/3)\sum_i \sum_j U_{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j$$

	$x$	$y$	$z$	$U_{\text{eq}}$
Pt	0.40203 (1)	0.37767 (3)	0.49989 (3)	0.03683 (10)
As1	0.48665 (4)	0.49416 (9)	0.55231 (6)	0.0376 (2)
As2	0.32160 (4)	0.26029 (8)	0.43824 (5)	0.0348 (2)
Cl1	0.47212 (12)	0.2131 (3)	0.4639 (2)	0.0583 (7)
C1	0.3394 (4)	0.5127 (9)	0.5437 (7)	0.051 (3)
C11	0.4696 (4)	0.6631 (9)	0.5917 (6)	0.040 (2)
C12	0.4767 (4)	0.7659 (8)	0.5443 (6)	0.044 (2)
C13	0.4598 (5)	0.8854 (10)	0.5699 (7)	0.050 (2)
C14	0.4337 (4)	0.9031 (9)	0.6424 (7)	0.045 (2)
C141	0.4157 (6)	1.0304 (11)	0.6723 (8)	0.067 (3)
C15	0.4268 (5)	0.7959 (10)	0.6915 (7)	0.051 (3)
C16	0.4426 (5)	0.6821 (10)	0.6657 (6)	0.049 (3)
C21	0.5342 (4)	0.4230 (9)	0.6376 (5)	0.040 (2)
C22	0.5693 (5)	0.4937 (11)	0.6889 (6)	0.052 (3)
C23	0.6010 (5)	0.4416 (12)	0.7495 (7)	0.056 (3)
C24	0.5976 (5)	0.3125 (12)	0.7649 (7)	0.060 (3)
C241	0.6298 (8)	0.2549 (15)	0.8347 (8)	0.089 (4)
C25	0.5618 (6)	0.2421 (12)	0.7153 (7)	0.070 (3)
C26	0.5294 (6)	0.2937 (10)	0.6517 (7)	0.061 (3)
C31	0.5462 (4)	0.5296 (9)	0.4695 (6)	0.039 (2)
C32	0.5225 (5)	0.5418 (11)	0.3928 (6)	0.056 (3)
C33	0.5606 (5)	0.5763 (12)	0.3327 (7)	0.054 (3)
C34	0.6231 (5)	0.6008 (10)	0.3455 (7)	0.051 (3)
C341	0.6627 (7)	0.6400 (13)	0.2758 (9)	0.082 (4)
C35	0.6452 (5)	0.5870 (11)	0.4205 (7)	0.058 (3)
C36	0.6070 (4)	0.5518 (12)	0.4823 (6)	0.056 (3)
C41	0.2431 (4)	0.3438 (9)	0.4182 (5)	0.038 (2)
C42	0.2435 (5)	0.4667 (11)	0.3931 (6)	0.053 (3)
C43	0.1882 (6)	0.5266 (12)	0.3832 (7)	0.066 (3)
C44	0.1323 (5)	0.4695 (12)	0.3951 (6)	0.051 (3)
C441	0.0721 (6)	0.5413 (14)	0.3869 (8)	0.078 (4)
C45	0.1337 (5)	0.3460 (11)	0.4149 (7)	0.056 (3)
C46	0.1880 (4)	0.2825 (10)	0.4273 (7)	0.050 (3)
C51	0.2973 (3)	0.1203 (8)	0.5045 (9)	0.038 (2)
C52	0.2665 (6)	0.1502 (11)	0.5756 (7)	0.065 (3)
C53	0.2494 (6)	0.0517 (15)	0.6250 (8)	0.079 (4)
C54	0.2659 (6)	-0.0697 (13)	0.6108 (8)	0.068 (3)
C541	0.2496 (8)	-0.1733 (15)	0.6680 (11)	0.105 (6)
C55	0.2990 (5)	-0.0924 (10)	0.5443 (8)	0.060 (3)
C56	0.3157 (4)	0.0009 (9)	0.4908 (6)	0.045 (2)
C61	0.3381 (4)	0.1797 (9)	0.3377 (5)	0.038 (2)
C62	0.2947 (5)	0.1005 (9)	0.3035 (6)	0.047 (2)
C63	0.3048 (5)	0.0479 (10)	0.2305 (7)	0.055 (3)
C64	0.3566 (6)	0.0702 (11)	0.1880 (7)	0.055 (3)
C641	0.3652 (6)	0.0204 (14)	0.1047 (7)	0.075 (4)
C65	0.3992 (5)	0.1509 (11)	0.2211 (7)	0.059 (3)
C66	0.3903 (4)	0.2027 (10)	0.2947 (7)	0.048 (3)

Table 2. Selected geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Pt—C1	2.111 (9)	As1—C31	1.943 (9)
Pt—As1	2.3883 (10)	As1—C11	1.948 (9)
Pt—As2	2.3875 (10)	As2—C51	1.933 (11)
Pt—C11	2.397 (3)	As2—C61	1.934 (9)
As1—C21	1.928 (9)	As2—C41	1.954 (9)