

Refinement

Refinement on *F*
R = 0.049
wR = 0.040
S = 1.0
888 reflections
145 parameters
H atoms; see below
w = [1/*σ*²(*F*)]

(Δ/*σ*)_{max} = 0.00007
Δ*ρ*_{max} = 0.70 e Å⁻³
Δ*ρ*_{min} = -0.67 e Å⁻³
Extinction correction: none
Atomic scattering factors
from *International Tables for X-ray Crystallography*
(1974, Vol. IV)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (Å²)

$$B_{\text{eq}} = (8\pi^2/3) \sum_i \sum_j U_{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j$$

	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i> _{eq}
Ag1	1.1920 (1)	-0.0784 (1)	0.46845 (4)	3.29 (1)
O1	0.9120 (9)	-0.0906 (8)	0.3648 (3)	2.8 (1)
O2	0.599 (1)	0.0267 (7)	0.4121 (3)	2.8 (1)
O3	-0.082 (1)	0.1523 (7)	0.2097 (4)	3.1 (1)
O4	-0.066 (1)	0.0990 (9)	0.0788 (4)	4.7 (2)
O5	0.557 (1)	-0.2139 (7)	-0.0021 (4)	3.3 (2)
O6	0.864 (1)	-0.2670 (7)	0.0866 (4)	3.6 (2)
N1	0.014 (1)	0.0975 (9)	0.1526 (4)	2.6 (2)
N2	0.665 (1)	-0.2095 (9)	0.0680 (4)	2.8 (2)
C1	0.562 (1)	-0.0565 (9)	0.2728 (5)	2.1 (2)
C2	0.345 (2)	0.0171 (9)	0.2548 (5)	1.9 (2)
C3	0.235 (2)	0.014 (1)	0.1730 (5)	2.1 (2)
C4	0.338 (2)	-0.0597 (9)	0.1111 (5)	2.3 (2)
C5	0.549 (2)	-0.134 (1)	0.1330 (5)	2.1 (2)
C6	0.663 (2)	-0.1302 (9)	0.2118 (5)	2.0 (2)
C7	0.703 (1)	-0.0395 (9)	0.3582 (5)	1.8 (2)

Table 2. Selected geometric parameters (Å, °)

Ag1...Ag1 ⁱ	2.835 (2)	Ag1—O2 ⁱ	2.218 (6)
Ag1...Ag1 ⁱⁱ	3.746 (2)	Ag1—O2 ⁱⁱ	2.655 (7)
Ag1—O1	2.176 (5)		
Ag1—O1—C7	128.9 (5)	O1—Ag1—O2 ⁱⁱⁱ	108.0 (2)
Ag1 ⁱ —O2—C7	119.4 (5)	O2 ⁱ —Ag1—O2 ⁱⁱⁱ	79.9 (2)
Ag1 ⁱ —O2 ⁱⁱⁱ —C7 ⁱⁱ	118.07 (5)	Ag1—O2 ⁱ —Ag1 ⁱ	100.1 (3)
O1—Ag1—O2 ⁱ	164.0 (2)		

Symmetry codes: (i) 2-x, -y, 1-z; (ii) 3-x, -y, 1-z; (iii) 1+x, y, z.

All non-H atoms were refined with anisotropic displacement parameters. H atoms were placed geometrically 0.95 Å from their corresponding C atoms. A riding model was used for all H atoms, with *U*_{iso}(H) = 1.3 *B*_{eq}(C).

Data collection: *CAD-4 Express* (Enraf–Nonius, 1993). Data reduction: *MolEN* (Fair, 1990). Program(s) used to solve structure: *SIMPEL* in *MolEN*. Program(s) used to refine structure: *LSFM* in *MolEN*. Molecular graphics: *ORTEPII* (Johnson, 1976) in *MolEN*. Software used to prepare material for publication: *MolEN*.

The authors wish to acknowledge the purchase of the CAD-4 diffractometer under Grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: BM1093). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Enraf–Nonius (1993). *CAD-4 Express Software*. Version 1.1. Enraf–Nonius, Delft, The Netherlands.
Fair, C. K. (1990). *MolEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf–Nonius, Delft, The Netherlands.
Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
Mak, T. C. W., Smith, G. & Kennard, C. H. L. (1993). *Z. Kristallogr.* **207**, 33–38.
Smith, G., Kennard, C. H. L. & Mak, T. C. W. (1988). *Z. Kristallogr.* **184**, 275–280.
Tahir, M. N., Ülkü, D. & Movsumov, E. M. (1996). *Acta Cryst.* **C52**, 593–595.
Usubaliev, G. J., Movsumov, E. M., Amiraslanov, I. R., Akhmedov, A. I., Musaev, A. & Mamedov, Kh. S. (1981). *J. Struct. Chem.* **22**, 73–77.

Acta Cryst. (1996). **C52**, 2680–2682

Bis[N-(2,6-dimethylphenyl)amino-glyoximato-N,N']nickel(II)

DİNÇER ÜLKÜ,^{a*} FILİZ ERCAN,^a MUSTAFA MACIT^b AND AHMET GÜLCE^c

^a*Hacettepe University, Department of Engineering Physics, Beytepe 06532, Ankara, Turkey*, ^b*Ondokuz Mayıs University, Department of Chemistry, Samsun, Turkey*, and ^c*Süleyman Demirel University, Department of Chemistry, Isparta, Turkey*. E-mail: dulku@eti.cc.hun.edu.tr

(Received 30 May 1996; accepted 10 July 1996)


Abstract

The title compound consists of discrete centrosymmetric [Ni(C₁₀H₁₂N₂O₂)₂] molecules. The coordination around the Ni atom is square planar, involving four oxime N atoms of two chelating C₂H₂N₂O₂ groups. The oxime O atoms are linked by an intramolecular hydrogen bond [O...O 2.546 (2) Å]. The Ni—N distances are 1.891 (2) and 1.846 (2) Å, and the N—Ni—N angle within the NiC₂N₂ chelate ring is 82.75 (8)°. In the central portion of the molecule, the C=N distances are equal [1.297(3) and 1.298 (3) Å], while the two N—O distances are different [1.303 (3) and 1.386 (3) Å].

Comment

Aminoglyoximes and related ligands are of interest because of their free amino substituents, which gives them the ability to form new ligand systems or metal complexes. The structures of bis(oxamideoximato)-cobalt(II) oxamide oxime (Bekaroğlu, Sarışaban, Koray

& Ziegler, 1977), bis(oxamideoximato)nickel(II)-water-dimethylformamide (Enders, 1978) and tris(oxamideoxime)cobalt(III) trichloride (Bekaroğlu *et al.*, 1978) have been reported previously. The cobalt complexes have either tetragonal-bipyramidal or octahedral coordination, while the nickel complex exhibits a square-planar metal environment. The structure determination of the title complex, $[\text{Ni}(\text{C}_{10}\text{H}_{12}\text{N}_3\text{O}_2)_2]$, (I), has been undertaken in order to compare its structure with those of related compounds.



The molecule of (I) is a centrosymmetric monomer (Fig. 1) like the Ni complex mentioned above. The coordination around the Ni atom is square planar, involving four oxime N atoms of two chelating $\text{C}_2\text{H}_2\text{N}_2\text{O}_2$ moieties. The hydrogen bonds between the oxime O atoms are not symmetrical, with an $\text{O} \cdots \text{O}$ distance of 2.546(2) Å and an $\text{O}-\text{H}-\text{O}$ angle of 167(3)°. The $\text{N}-\text{Ni}-\text{N}$ angle within the NiC_2N_2 chelate ring is 82.75(8)°. Although the angles at the Ni atom and one $\text{Ni}-\text{N}$ distance have practically the same values as in bis(oxamideoximato)nickel(II)-water-dimethyl-formamide, the second $\text{Ni}-\text{N}$ distance is slightly longer [$\text{Ni}-\text{N}2$ 1.891(2) *versus* 1.860(4) Å]. The coordination plane around the Ni atom and the plane of the chelating moiety coincide within experimental error. The planes of the phenyl ring and the coordination plane are perpendicular to one another [dihedral angle 89.39(6)°]. A




Fig. 1. ORTEPII (Johnson, 1976) drawing showing the atom-numbering scheme. Displacement ellipsoids are shown at the 50% probability level.

comparison of the $\text{C}=\text{N}$ and $\text{N}-\text{O}$ distances with those of free oximes shows that at least one of the $\text{N}-\text{O}$ distances [$\text{N}2-\text{O}2$ 1.303(3) Å] is considerably shortened on complex formation, while the $\text{C}=\text{N}$ distance stays practically the same. On complex formation, the $\text{C}-\text{N}-\text{O}$ angle is also affected, opening by up to *ca* 10°. These observations are in agreement with those of Chakravorty (1974) for transition metal complexes of oximes and related ligands.

Experimental

A solution of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (1.20 g, 5 mmol) in ethanol–water (1:1) was added dropwise to a solution of *N*-(2,6-dimethylphenyl)aminoglyoxime (2.07 g, 10 mmol) in ethanol (40 ml) (pH 2.5–3.0). Then a 1% solution of KOH in water was dripped slowly into the mixture until the pH reached 5.5. The resulting precipitate was removed by filtration, washed and dried *in vacuo*. Recrystallization from a chloroform–ethanol mixture (2:1) gave reddish prismatic crystals.

Crystal data

$[\text{Ni}(\text{C}_{10}\text{H}_{12}\text{N}_3\text{O}_2)_2]$

$M_r = 471.15$

Triclinic

$P\bar{1}$

$a = 7.9572(7)$ Å

$b = 8.2383(4)$ Å

$c = 8.5846(6)$ Å

$\alpha = 110.416(5)$ °

$\beta = 92.067(6)$ °

$\gamma = 92.165(6)$ °

$V = 526.32(4)$ Å³

$Z = 1$

$D_x = 1.486$ Mg m⁻³

D_m not measured

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 25 reflections

$\theta = 11\text{--}18$ °

$\mu = 0.96$ mm⁻¹

$T = 296$ K

Prism

$0.48 \times 0.33 \times 0.13$ mm

Dark red

Data collection

Enraf–Nonius CAD-4 diffractometer

$\omega/2\theta$ scans

Absorption correction:

empirical *via* ψ scans

(North, Phillips &

Mathews, 1968)

$T_{\min} = 0.78$, $T_{\max} = 0.88$

2283 measured reflections

2196 independent reflections

2150 observed reflections

[$I > 3\sigma(I)$]

$R_{\text{int}} = 0.016$

$\theta_{\max} = 26.3$ °

$h = 0 \rightarrow 9$

$k = -10 \rightarrow 10$

$l = -10 \rightarrow 10$

3 standard reflections

frequency: 120 min

intensity decay: 1%

Refinement

Refinement on F

$R = 0.029$

$wR = 0.033$

$S = 1.25$

2150 reflections

145 parameters

If $F \geq 197.30$ then

$w = (197.30/F)^2$,

otherwise $w = 1.0$

$(\Delta/\sigma)_{\max} = 0.002$

$\Delta\rho_{\max} = 0.61$ e Å⁻³

$\Delta\rho_{\min} = -0.32$ e Å⁻³

Extinction correction: none

Atomic scattering factors

from International Tables

for X-ray Crystallography

(1974, Vol. IV)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (\AA^2)

$$B_{\text{eq}} = (8\pi^2/3) \sum_i \sum_j U_{ij} a_i^* a_j \cdot \mathbf{a}_i \cdot \mathbf{a}_j$$

	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i> _{eq}
Ni	0	0	0	2.635 (7)
N1	0.1230 (3)	-0.0968 (2)	-0.1853 (2)	3.18 (4)
O1	0.1776 (2)	-0.0196 (2)	-0.2957 (2)	4.29 (4)
C1	0.1783 (3)	-0.2488 (2)	-0.2064 (2)	2.98 (4)
C2	0.1188 (3)	-0.3209 (2)	-0.0846 (3)	3.32 (4)
N2	0.0314 (3)	-0.2148 (2)	-0.0277 (2)	3.22 (4)
O2	-0.0345 (3)	-0.2594 (2)	0.1447 (2)	4.57 (4)
N3	0.2754 (3)	-0.3352 (2)	-0.3301 (2)	3.64 (4)
C3	0.3832 (3)	-0.4663 (3)	-0.3165 (2)	3.13 (4)
C4	0.5340 (3)	-0.4165 (3)	-0.2233 (3)	3.96 (5)
C5	0.6395 (4)	-0.5440 (4)	-0.2187 (3)	5.00 (6)
C6	0.5949 (4)	-0.7160 (3)	-0.3014 (3)	5.02 (6)
C7	0.4427 (4)	-0.7626 (3)	-0.3893 (3)	4.51 (6)
C8	0.3323 (3)	-0.6407 (3)	-0.4010 (3)	3.47 (5)
C9	0.1677 (4)	-0.6917 (3)	-0.4951 (3)	5.11 (7)
C10	0.5811 (4)	-0.2276 (4)	-0.1289 (4)	6.28 (8)

Table 2. Selected geometric parameters (\AA , °)

Ni—N2	1.891 (2)	C1—N3	1.344 (3)
Ni—N1	1.846 (2)	N1—O1	1.386 (3)
N2—O2	1.303 (3)	N3—C3	1.438 (3)
N2—C2	1.298 (3)	C4—C10	1.508 (4)
C1—C2	1.455 (3)	C8—C9	1.479 (4)
C1—N1	1.297 (3)	O1—H1	0.95 (3)
N2—Ni—N1	82.75 (8)	Ni—N1—O1	127.6 (1)
Ni—N2—O2	123.9 (1)	C1—N1—O1	115.7 (2)
Ni—N2—C2	114.7 (2)	C1—N3—C3	121.9 (2)
O2—N2—C2	121.3 (2)	N3—C3—C4	119.2 (2)
C2—C1—N1	112.9 (2)	N3—C3—C8	118.4 (2)
C2—C1—N3	122.7 (2)	C3—C4—C10	120.7 (2)
N1—C1—N3	124.4 (2)	C5—C4—C10	121.0 (2)
N2—C2—C1	113.1 (2)	C3—C8—C9	121.6 (2)
Ni—N1—C1	116.3 (2)	C7—C8—C9	121.9 (2)
Ni—N2—C2—C1	1.6 (3)	C2—C1—N3—C3	-26.1 (3)
N3—C1—C2—N2	177.9 (2)	C1—N3—C3—C4	-77.5 (3)

H-atom positions were obtained from difference maps. For all H atoms, $U_{\text{iso}} = 1.3U_{\text{eq}}$ of the parent atom and a riding model was adopted except that the fractional coordinates of H1 (bonded to O1) were refined.

Data collection: CAD-4 Express Software (Enraf–Nonius, 1993). Data reduction: MolEN (Fair, 1990). Program(s) used to solve structure: SIMPEL in MolEN. Program(s) used to refine structure: LSFM in MolEN. Molecular graphics: ORTEPII (Johnson, 1976) in MolEN. Software used to prepare material for publication: MolEN.

The authors acknowledge the purchase of the CAD-4 diffractometer under Grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: MU1275). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Bekaroğlu, Ö., Sarsaban, S., Koray, A. R., Nuber, B., Weidenhammer, K., Weiss, J. & Ziegler, L. (1978). *Acta Cryst.* **B34**, 3591–3593.
 Bekaroğlu, Ö., Sarsaban, S., Koray, A. R. & Ziegler, M. L. (1977). *Z. Naturforsch. Teil. B*, **32**, 387–392.
 Chakravorty, A. (1974). *Coord. Chem. Rev.* **13**, 1–46.
- Enders, H. (1978). *Acta Cryst.* **B34**, 2306–2309.
 Enraf–Nonius (1993). CAD-4 Express Software. Version 1.1. Enraf–Nonius, Delft, The Netherlands.
 Fair, C. K. (1990). MolEN. An Interactive Intelligent System for Crystal Structure Analysis. Enraf–Nonius, Delft, The Netherlands.
 Johnson, C. K. (1976). ORTEPII. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Acta Cryst.* (1996). **C52**, 2682–2684
- (Diethylamine-N){4-methyl-2-[(2-oxido-benzylidene)amino]phenolato(2-)O,N,O' }-nickel(II)**

FILİZ ERCAN,^a MUSA SARI,^b RAHMI YAĞBASAN,^b MUSTAFA TAŞTEKİN^c AND LEVENT AKSU^c

^aHacettepe University, Faculty of Science, Department of Engineering Physics, 06532 Beytepe, Ankara, Turkey, ^bGazi University, Gazi Education Faculty, Department of Physics Education, 06500 Beşevler, Ankara, Turkey, and ^cAnkara University, Faculty of Science, Department of Chemistry, 06100 Tandoğan, Ankara, Turkey. E-mail: filiz@etit.cc.hun.edu.tr

(Received 24 November 1995; accepted 17 July 1996)

Abstract

In the title compound, [Ni(C₁₄H₁₁NO₂)(C₄H₁₁N)], the coordination of the Ni atom is distorted square planar, with Ni—O distances of 1.839 (7) and 1.812 (7) Å, Ni—N distances of 1.852 (8) and 1.951 (9) Å, and O—Ni—N angles between 86.7 (4) and 96.2 (4)°.

Comment

Salicylaldimine and its biologically interesting complexes have been extensively studied over the years (Stewart & Lingafelter, 1959; Calligaris, Nardin & Randaccio, 1972; Bhatia, Bindlish, Saini & Jain, 1981; Kessissoglu, Raptopoulou, Bakalbassis, Terzis & Mrozniski, 1992). In biological systems such as jack-bean urease and in some hydrogenases, nickel is in a related complexed form (Walsh & Orme-Johnson, 1987). Schiff base complexes have also been used in electrochemical research and catalytic reactions (Hamilton, Drago & Zombeck, 1987; Costamagna, Vargas, Latorre, Alvarado & Mena, 1992). The title compound, (I), is one of the square-planar complexes formed by the reaction of Ni^{II} with both a ligand having an ONO donor set and an aliphatic amine. The structure of this Ni^{II} complex was determined in order to compare the geometrical features with those of related compounds.