


References

- Enraf–Nonius (1989). *CAD-4 Software*. Version 5.0. Enraf–Nonius, Delft, The Netherlands.
- Evans, D. A., Bartoli, J. & Shih, T. L. (1981). *J. Am. Chem. Soc.* **103**, 2127–2129.
- Fernández, R., Rodríguez, J., Quiñoá, E., Riguera, R., Muñoz, L., Fernández-Suárez, M. & Debitus, C. (1996). *J. Am. Chem. Soc.* **118**, 11635–11643.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Frenz, B. A. (1983). *Enraf–Nonius Structure Determination Package: SDP User's Guide*. Version of 6 January 1983. Enraf–Nonius, Delft, The Netherlands.
- Gage, J. R. & Evans, D. A. (1989). *Org. Synth.* **68**, 77–91.
- Marsch, E. R., Schaefer, W. P., Kukkola, P. J. & Myers, A. G. (1992). *Acta Cryst.* **C48**, 1622–1624.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Rodríguez, J., Fernández, R., Quiñoá, E., Riguera, R., Debitus, C. & Bouchet, P. (1994). *Tetrahedron Lett.* **35**, 9239–9242.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Zsolnai, L. & Pritzkow, H. (1994). *ZORTEP. ORTEP Program for PC*. University of Heidelberg, Germany.

interactions. In contrast to the structure of (1) and other chlorins, the two pyrrole H atoms in (2) are located at rings *B* and *D*.

Comment

The structure of the phytychlorin (1) derived from chlorophyll *a* was determined in order to obtain information on the crystal packing of a pheophorbide *a* derivative containing no O functionalities. This was achieved in (1) by removal of the 13¹ oxo group and substitution of the 17-propionic acid group present in chlorophyll *a* with a propionyl group.



Acta Cryst. (1997). **C53**, 1314–1318

Structure and Conformation of Photosynthetic Pigments and Related Compounds. 10. Comparison of a Phytychlorin and Phytyporphyrin Derived from Chlorophyll *a*

MATHIAS O. SENGE^{a*} AND KEVIN M. SMITH^b

^a*Institut für Organische Chemie (WE02), Freie Universität Berlin, Takustrasse 3, D-14195 Berlin, Germany, and*

^b*Department of Chemistry, University of California, Davis, CA 95616, USA. E-mail: mosenge@chemie.fu-berlin.de*

(Received 6 January 1997; accepted 5 March 1997)

Abstract

The crystal and molecular structures of 17-decarboxyethyl-13¹-deoxo-17-propylphytychlorin, C₃₃H₄₀N₄, (1), and phytyporphyrin methyl ester, C₃₄H₃₆N₄O₃, (2), are compared. Compound (1) shows structural parameters similar to those of other naturally occurring phytychlorins. Owing to the absence of any heteroatom functionalities at the periphery, no close contacts are observed in the packing of (1), in contrast to those normally found in other chlorophyll derivatives. Compound (2) presents the first structure of a free-base pheophyrin and forms chains stabilized by C—H···O=C and π–π

Compound (1) crystallized with two crystallographically independent molecules in the asymmetric unit, mainly distinguished by the conformation of the C17 propionyl groups (Fig. 1). The structural parameters of the macrocycles agree well with those of other pheophorbides related to chlorophyll *a* (Hoppe, Will, Gassmann & Weichselgartner, 1969; Gassmann, Strell, Brandl, Sturm & Hoppe, 1971; Fischer, Templeton, Zalkin & Calvin, 1972; Kratky, Isenring & Dunitz, 1977; Smith, Goff, Fajer & Barkigia, 1982, 1983; Senge & Smith, 1992). The hydrophorphyrin character is clearly evidenced by the elongated C17—C18 and C37—C38 bond lengths [1.552 (5) and 1.551 (5) Å, respectively]. With the exception of the reduced ring *D*, both macrocycles are almost planar. The average deviation of the 26 atoms comprising the phytychlorin macrocycle system is 0.031 Å in (1) and 0.065 Å in (2). The largest deviations from the respective N₄ planes are observed for C18 [0.28 (5) Å] and C38 [0.42 (5) Å].

The chlorins form molecular stacks in the crystal, repeating the (1)–(1)–(2)–(2)–(1) motif. Whereas the interplanar separations of the N₄ planes are of the order of 3.5 Å, there is almost no π overlap of the ring systems as indicated by center-to-center separations of the order of 7.9 (1) Å and lateral shifts of the macrocycle centers against each other by about 7.03 (3) Å [geometrical parameters as defined by Scheidt & Lee (1987)]. Thus, in contrast to other phytychlorins, in the absence of any heteroatom functionalities at the phytychlorin periphery, no close contacts and only very weak π–π interactions are observed.




Fig. 1. Molecular structure and numbering scheme for the two crystallographically independent molecules of (1). H atoms have been omitted for clarity and ellipsoids are drawn for 50% occupancy.

Compound (2) presents the first example of a structure of a free-base pheophorphyrin. Compounds of this type, often called geo- or petroporphyrins, are found in geological deposits and derive from plant chlorophylls (Treibs, 1934; Callot, 1991). So far, only structures of some metallo-geoporphyrins have been published (Pettersen, 1969, 1971; Ekstrom *et al.*, 1983; Miller, Hambley & Taylor, 1984; Senge, Smith & Smith, 1993). These compounds are directly related to phytychlorins such as (1), the difference being the oxidation level of ring *D*. The molecular structure of (2) shows overall structural parameters that are similar to other porphyrins

(Scheidt & Lee, 1987) (Fig. 2). The oxidized character of ring *D* is shown by the shortened C17—C18 bond length [1.378 (3) Å] and the porphyrin character of the ring systems leads to an essentially planar macrocyclic system. The average deviation from planarity for the 26 atoms comprising the phytyporphyrin system is 0.041 Å. A significant difference from (1) or other chlorins is found in the location of the pyrrole H atoms. Whereas in phytychlorins they are located in rings *A* and *C*, (2) bears N—H groups in rings *B* and *D*. This is evidenced by differences in the C_{α} —N— C_{α} angles which are widened in rings *A* and *C* [average value 104.4 (2)°] compared with rings *B* and *D* [average angle 110.4 (2)°]. The respective values found in the corresponding phytychlorin methyl ester (Senge & Smith, 1992) are 110.5 (7) (rings *A* and *C*) and 107.1 (7)° (rings *B* and *D*).

The crystal packing of (2) is characterized by the formation of polymeric chains of dimeric porphyrin aggregates, which are held together by a combination of weak π and C—H...O interactions (Fig. 3). A dimer unit is stabilized by formation of a weak π — π aggregate [interplanar separation: 3.52 (5) Å; lateral shift of the porphyrin centers: 5.67 (5) Å] and by interactions between the C81 methylene group and the ester carbonyl O atom of the C17 substituent [H81B...O2 2.48 (1) Å]. The propionic acid methyl ester groups are folded towards the neighboring macrocycle in a clamp-like conformation. Neighboring units are stabilized *via* interaction of the C13¹ carbonyl O atom and the methyl




Fig. 2. Molecular structure and numbering scheme for (2). H atoms have been omitted for clarity and ellipsoids are drawn for 50% occupancy.

H atoms of the ester group [H17G···O1 2.42 (1) Å]. Owing to the folding back of the propionic acid ester groups towards each other, the outer faces of each polymer can form weak π - π aggregates with the next chain above and below. These interchain aggregates are slightly stronger (interplanar separation: 3.48 Å; lateral shift of the centers: 5.03 Å) than the intrachain aggregates mentioned above. Nevertheless, all aggregates described here are weak, especially when compared with the strong aggregates observed for metalloporphyrins (Scheidt & Lee, 1987).




Fig. 3. View of the chains formed by (2) in the crystal. Only H atoms involved in weak C—H···O interactions (dashed lines) are shown.

Experimental

Compound (1) was prepared synthetically as described by Abraham, Rowan, Smith & Smith (1993). Compound (2) was prepared synthetically as described by Fischer & Orth (1940) and Kenner, McCombie & Smith (1973). For both (1) and

(2), crystals were grown from CH₂Cl₂/*n*-hexane and mounted according to Hope (1994).

Compound (1)

Crystal data

C₃₃H₄₀N₄
M_r = 492.69
 Monoclinic
*P*2₁
a = 7.875 (3) Å
b = 24.756 (10) Å
c = 14.582 (6) Å
 β = 104.26 (3)°
V = 2755.2 (19) Å³
Z = 4
D_x = 1.188 Mg m⁻³
D_m not measured

Data collection

Siemens R3m/V diffractometer
 ω scans
 Absorption correction:
 ΔF (XABS2; Parkin, Moezzi & Hope, 1995)
T_{min} = 0.96, *T_{max}* = 0.98
 6946 measured reflections
 6474 independent reflections

Refinement

Refinement on *F*²
R[*F*² > 2 σ (*F*²)] = 0.058
 wR (*F*²) = 0.140
S = 1.022
 6474 reflections
 667 parameters
 H-atom parameters constrained

Mo *K* α radiation
 λ = 0.71073 Å
 Cell parameters from 35 reflections
 θ = 12–15°
 μ = 0.070 mm⁻¹
T = 126 (2) K
 Parallelepiped
 0.60 × 0.43 × 0.28 mm
 Green

4933 reflections with
 $I > 2\sigma(I)$
R_{int} = 0.051
 θ_{max} = 27.5°
h = -10 → 9
k = 0 → 32
l = 0 → 18
 2 standard reflections
 every 198 reflections
 intensity decay: none

$w = 1/[\sigma^2(F_o^2) + (0.0596P)^2 + 0.8366P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{max} = 0.003$
 $\Delta\rho_{max} = 0.34 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{min} = -0.28 \text{ e } \text{Å}^{-3}$
 Extinction correction: none
 Scattering factors from
International Tables for Crystallography (Vol. C)

Table 1. Selected geometric parameters (Å, °) for (1)

N21—C1	1.375 (5)	C13—C13A	1.496 (5)
N21—C4	1.377 (5)	C13A—C13B	1.566 (6)
N22—C6	1.365 (5)	C13B—C15	1.531 (5)
N22—C9	1.373 (5)	C14—C15	1.388 (6)
N23—C14	1.358 (5)	C15—C16	1.384 (5)
N23—C11	1.385 (5)	C16—C17	1.518 (5)
N24—C16	1.356 (5)	C17—C18	1.552 (5)
N24—C19	1.361 (5)	C18—C19	1.522 (5)
C12—C13	1.376 (6)	C37—C38	1.551 (5)
C13—C14	1.418 (5)		
C1—N21—C4	110.6 (3)	N23—C14—C13	109.0 (3)
C6—N22—C9	105.6 (3)	C15—C14—C13	114.2 (3)
C14—N23—C11	108.5 (3)	C16—C15—C14	127.1 (4)
C16—N24—C19	109.0 (3)	C16—C15—C13B	126.8 (4)
C10—C11—C12	130.9 (4)	C14—C15—C13B	106.0 (3)
N23—C11—C12	108.0 (3)	N24—C16—C15	121.6 (3)
C13—C12—C11	106.1 (3)	N24—C16—C17	113.0 (3)
C12—C13—C14	108.4 (3)	C15—C16—C17	125.4 (3)
C12—C13—C13A	142.1 (4)	C16—C17—C18	101.7 (3)
C14—C13—C13A	109.5 (3)	C19—C18—C17	102.4 (3)
C13—C13A—C13B	103.1 (3)	N24—C19—C20	124.1 (3)
C15—C13B—C13A	107.1 (3)	N24—C19—C18	112.1 (3)
N23—C14—C15	136.8 (4)		

Compound (2)*Crystal data*C₃₄H₃₆N₄O₃ $M_r = 548.67$

Triclinic

 $P\bar{1}$ $a = 9.703 (3) \text{ \AA}$ $b = 11.906 (2) \text{ \AA}$ $c = 13.685 (3) \text{ \AA}$ $\alpha = 99.11 (2)^\circ$ $\beta = 107.84 (2)^\circ$ $\gamma = 106.67 (2)^\circ$ $V = 1388.1 (6) \text{ \AA}^3$ $Z = 2$ $D_x = 1.313 \text{ Mg m}^{-3}$ D_m not measured*Data collection*

Siemens R3m/V diffractometer

 ω scans

Absorption correction:

 ΔF (XABS2; Parkin,

Moezzi & Hope, 1995)

 $T_{\min} = 0.95$, $T_{\max} = 0.98$

6753 measured reflections

6376 independent reflections

*Refinement*Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.065$ $wR(F^2) = 0.199$ $S = 1.037$

6376 reflections

370 parameters

H-atom parameters
constrainedMo $K\alpha$ radiation $\lambda = 0.71073 \text{ \AA}$ Cell parameters from 31
reflections $\theta = 11-14^\circ$ $\mu = 0.085 \text{ mm}^{-1}$ $T = 126 (2) \text{ K}$

Plate

 $0.6 \times 0.4 \times 0.2 \text{ mm}$

Red

4627 reflections with

 $I > 2\sigma(I)$ $R_{\text{int}} = 0.040$ $\theta_{\text{max}} = 27.5^\circ$ $h = -12 \rightarrow 11$ $k = -15 \rightarrow 15$ $l = 0 \rightarrow 17$

2 standard reflections

every 198 reflections

intensity decay: none

 $w = 1/[\sigma^2(F_o^2) + (0.091P)^2 + 1.4171P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} = -0.001$ $\Delta\rho_{\text{max}} = 0.59 \text{ e \AA}^{-3}$ $\Delta\rho_{\text{min}} = -0.62 \text{ e \AA}^{-3}$

Extinction correction: none

Scattering factors from

*International Tables for
Crystallography* (Vol. C)Table 2. Selected geometric parameters (\AA , $^\circ$) for (2)

N21—C4	1.368 (3)	C13—C14	1.434 (3)
N21—C1	1.374 (3)	C13—C131	1.458 (4)
N22—C6	1.362 (3)	C131—C132	1.538 (4)
N22—C9	1.374 (3)	C132—C15	1.517 (3)
N23—C14	1.351 (3)	C14—C15	1.403 (3)
N23—C11	1.383 (3)	C15—C16	1.391 (3)
N24—C19	1.356 (3)	C16—C17	1.437 (3)
N24—C16	1.380 (3)	C17—C18	1.378 (3)
C11—C12	1.455 (3)	C18—C19	1.440 (3)
C12—C13	1.368 (3)		
C4—N21—C1	105.0 (2)	N23—C14—C13	112.4 (2)
C6—N22—C9	110.2 (2)	C15—C14—C13	112.3 (2)
C14—N23—C11	103.8 (2)	C16—C15—C14	123.9 (2)
C19—N24—C16	110.6 (2)	C16—C15—C132	128.7 (2)
C10—C11—C12	124.5 (2)	C14—C15—C132	107.4 (2)
C13—C12—C11	104.1 (2)	N24—C16—C15	119.7 (2)
C12—C13—C14	107.3 (2)	N24—C16—C17	107.0 (2)
C12—C13—C131	144.3 (2)	C15—C16—C17	133.3 (2)
C14—C13—C131	108.4 (2)	C18—C17—C16	107.4 (2)
O1—C131—C132	124.0 (2)	C17—C18—C19	107.9 (2)
C13—C131—C132	106.7 (2)	N24—C19—C20	125.9 (2)
C15—C132—C131	105.2 (2)	N24—C19—C18	107.1 (2)
N23—C14—C15	135.3 (2)	C20—C19—C18	126.9 (2)

The pyrrole and *meso*-carbon H atoms were located in difference maps. For refinement, all H atoms were placed in calculated positions and refined using a riding model assuming planar geometry at the N atom. All calculations were performed using the *SHELXTL-Plus* program system (Sheldrick, 1994). The absolute structure of (1) could not be determined from the Flack (1983) parameter but was chosen with reference to the absolute structure of chlorophyll *a* from which this compound was derived chemically.

For both compounds, data collection: *P3* (Siemens, 1990); cell refinement: *P3*; data reduction: *XDISK* (Siemens, 1989) and *SHELXTL-Plus* (Sheldrick, 1994); program(s) used to solve structures: *SHELXS86* (Sheldrick, 1990); program(s) used to refine structures: *SHELXL93* (Sheldrick, 1993); molecular graphics: *XP* (Siemens, 1994) and *SHELXTL-Plus*; software used to prepare material for publication: *XCIF* and *SHELXTL-Plus*.

This work was supported by grants from the Deutsche Forschungsgemeinschaft (Se543/2-2 and Se543/2-3), the Fonds der Chemischen Industrie, and the National Science Foundation (CHE-96-23117).

Supplementary data for this paper are available from the IUCr electronic archives (Reference: JZ1193). Services for accessing these data are described at the back of the journal.

References

- Abraham, R. J., Rowan, A. E., Smith, N. W. & Smith, K. M. (1993). *J. Chem. Soc. Perkin Trans. 2*, pp. 1047–1059.
- Callot, H. J. (1991). *Chlorophylls*, edited by H. Scheer, pp. 339–379. Boca Raton, Florida: CRC Press.
- Ekstrom, A., Fookes, C. J. R., Hambley, T., Loeh, H. J., Miller, S. A. & Taylor, J. C. (1983). *Nature (London)*, **306**, 173–174.
- Fischer, H. & Orth, H. (1940). *Die Chemie des Pyrrols*, Vol. 2, Part 1, pp. 192–193. Leipzig: Akademische Verlagsgesellschaft.
- Fischer, M. S., Templeton, D. H., Zalkin, A. & Calvin, M. (1972). *J. Am. Chem. Soc.* **94**, 3613–3619.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Gassmann, J., Strell, I., Brandl, F., Sturm, M. & Hoppe, W. (1971). *Tetrahedron Lett.* **48**, 4609–4612.
- Hope, H. (1994). *Prog. Inorg. Chem.* **41**, 1–13.
- Hoppe, W., Will, G., Gassmann, J. & Weichselgartner, H. (1969). *Z. Kristallogr.* **128**, 18–35.
- Kenner, G. W., McCombie, S. W. & Smith, K. M. (1973). *J. Chem. Soc. Trans.* pp. 2517–2523.
- Kratky, C., Isenring, H. P. & Dunitz, J. D. (1977). *Acta Cryst.* **B33**, 547–549.
- Miller, S. A., Hambley, T. W. & Taylor, J. C. (1984). *Aust. J. Chem.* **47**, 761–766.
- Parkin, S. R., Moezzi, B. & Hope, H. (1995). *J. Appl. Cryst.* **28**, 53–56.
- Pettersen, R. C. (1969). *Acta Cryst.* **B25**, 2527–2539.
- Pettersen, R. C. (1971). *J. Am. Chem. Soc.* **93**, 5629–5634.
- Scheidt, W. R. & Lee, Y.-J. (1987). *Struct. Bonding (Berlin)*, **64**, 1–70.
- Senge, M. O. & Smith, K. M. (1992). *Z. Kristallogr.* **199**, 239–248.
- Senge, M. O., Smith, N. W. & Smith, K. M. (1993). *Inorg. Chem.* **32**, 1259–1265.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Sheldrick, G. M. (1994). *SHELXTL-Plus*. Release 5.03. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

- Siemens (1989). *XDISK. Data Reduction Program*. Version 3.11. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1990). *P3 Software*. Release 4.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1994). *XP. Molecular Graphics Program*. Version 5.03. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Smith, K. M., Goff, D. A., Fajer, J. & Barkigia, K. M. (1982). *J. Am. Chem. Soc.* **104**, 3747–3749.
- Smith, K. M., Goff, D. A., Fajer, J. & Barkigia, K. M. (1983). *J. Am. Chem. Soc.* **105**, 1674–1676.
- Treibs, A. (1934). *Liebigs Ann. Chem.* **509**, 103–114.

Acta Cryst. (1997). **C53**, 1318–1320

Diisopropyl Hydrazocarboxylate–Triphenylphosphine Oxide Adduct†

A. HÉROUX AND F. BRISSE

Département de Chimie, Université de Montréal, CP 6128, Succ. Centre-ville, Montréal, Québec, Canada H3C 3J7.
E-mail: brisse@ere.umontreal.ca

(Received 14 June 1996; accepted 1 April 1997)

Abstract

Crystals of the title complex, C₈H₁₆N₂O₄·C₁₈H₁₅OP, belong to the *P2₁/c* space group. The triphenylphosphine oxide and diisopropyl hydrazocarboxylate molecules form a stable centrosymmetric 2:2 complex through a series of hydrogen bonds.


Comment

Etter has undertaken a systematic study of complexes incorporating the triphenylphosphine oxide molecule (see, for example, Etter, Rasmussen, Gleason, Kress, Duerst & Gillard, 1986; Etter & Panunto, 1988; Etter & Reutzel, 1991, 1992; Etter, Urbanczyk-Lipkowska, Zia-Ebrahimi & Panunto, 1990). Since triphenylphosphine oxide is a very strong proton acceptor it often forms highly stable crystalline complexes with other molecules. These crystals are often reasonably large and well developed (Etter & Baures, 1988). Etter has also shown that triphenylphosphine oxide may be used to displace and modify the hydrogen-bonding pattern occurring in some of the organic molecules with which it forms complexes.

The bond distances and angles in the triphenylphosphine oxide molecule, *A*, are normal; the P=O bond is 1.492 (1) Å, while the P—C bonds average

† Alternative name: isopropyl 3-(isopropoxycarbonyl)carbazate–triphenylphosphine oxide (1/1).

1.797 Å. In the diisopropyl hydrazocarboxylate molecule, *B*, the central N41—N51 bond is 1.381 (2) Å and there is a *gauche* torsion angle, $\tau = 72^\circ$, around this bond. All the C—H distances in molecule *A* are in the range 0.88 (3)–1.04 (3) Å, with U_{iso} values between 0.056 (5) and 0.150 (11) Å². The two N—H distances in molecule *B* are 0.82 (2) and 0.90 (2) Å.



Four molecules (two *A* + two *B*) are connected to each other through hydrogen bonds of the N—H(*B*)···O(*A*) type and form a centrosymmetric ten-membered ring. There is a three-center hydrogen bond involving the phosphoryl O atom, H(*B*)···O(*A*)···H(*B*). This system is not symmetrical since the two N···O bonds are 2.896 (2) and 2.880 (2) Å, while the two N—H···O bond angles are 159 (2) and 172 (2)°. Each phosphoryl O atom is connected to NH groups of two centrosymmetrically related diisopropyl hydrazocarboxylate molecules. Although there is no report of such an arrangement with phosphines, a comparable eight-membered ring system has been observed in a diacetamide–benzamide host–guest assembly (Etter & Reutzel, 1991). The carbonyl O atoms of molecule *B* do not participate in hydrogen bonding. The C=O bond lengths are 1.202 (2) and 1.206 (2) Å.




Fig. 1. ORTEPII (Johnson, 1976) drawing of the two structural units, C₁₈H₁₅PO and C₈H₁₆N₂O₄, and the atomic numbering schemes. The ellipsoids are drawn at the 40% probability level, while the H atoms are represented as spheres of arbitrary size.