

Crystal data

[Cu(NCS)(C ₁₄ H ₃₂ N ₄)]ClO ₄	Mo K α radiation
$M_r = 477.50$	$\lambda = 0.7107 \text{ \AA}$
Monoclinic	Cell parameters from 22 reflections
$P2_1/c$	$\theta = 5.45\text{--}15.74^\circ$
$a = 8.9245 (10) \text{ \AA}$	$\mu = 1.30 \text{ mm}^{-1}$
$b = 14.9611 (10) \text{ \AA}$	$T = 293 \text{ K}$
$c = 15.7331 (13) \text{ \AA}$	Cubic
$\beta = 96.184 (10)^\circ$	$0.25 \times 0.13 \times 0.12 \text{ mm}$
$V = 2088.5 (3) \text{ \AA}^3$	Light blue
$Z = 4$	
$D_x = 1.519 \text{ Mg m}^{-3}$	
D_m not measured	

Data collection

Nonius CAD-4 diffractometer	2514 reflections with $I > 1.5\sigma(I)$
$\theta/2\theta$ scans	$R_{\text{int}} = 0.019$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$\theta_{\text{max}} = 24.93^\circ$
$T_{\text{min}} = 0.703$, $T_{\text{max}} = 0.867$	$h = -10 \rightarrow 10$
3926 measured reflections	$k = 0 \rightarrow 17$
3675 independent reflections	$l = 0 \rightarrow 18$
	3 standard reflections frequency: 60 min intensity decay: 1.0%

Refinement

Refinement on F^2	$\Delta\rho_{\text{max}} = 0.39 \text{ e \AA}^{-3}$
$R = 0.041$	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$
$wR = 0.052$	Extinction correction: Larson (1970)
$S = 1.13$	Extinction coefficient: $8 (2) \times 10^3$
2513 reflections	Scattering factors from <i>International Tables for X-ray Crystallography</i> (Vol. IV)
245 parameters	
H atoms not refined	
$w = 1/[\sigma^2(F) + 0.001F^2]$	
$(\Delta/\sigma)_{\text{max}} = 0.002$	

Table 1. Selected geometric parameters (\AA , $^\circ$)

Cu—N1	2.080 (4)	Cu—N4	2.097 (4)
Cu—N2	2.094 (4)	Cu—N5	2.116 (4)
Cu—N3	2.143 (4)	S—C15	1.628 (5)
N1—Cu—N2	85.45 (15)	Cu—N1—C3	106.4 (3)
N1—Cu—N3	145.46 (15)	Cu—N2—C4	104.4 (3)
N1—Cu—N4	92.40 (15)	Cu—N2—C5	115.3 (3)
N1—Cu—N5	114.57 (16)	Cu—N2—C6	112.1 (3)
N2—Cu—N3	92.80 (16)	Cu—N3—C8	112.1 (3)
N2—Cu—N4	174.37 (15)	Cu—N3—C9	114.7 (3)
N2—Cu—N5	93.53 (15)	Cu—N3—C10	103.8 (3)
N3—Cu—N4	86.02 (15)	Cu—N4—C11	103.0 (3)
N3—Cu—N5	99.98 (16)	Cu—N4—C12	115.7 (3)
N4—Cu—N5	92.09 (16)	Cu—N4—C13	112.8 (3)
Cu—N1—C1	114.0 (3)	Cu—N5—C15	144.6 (4)
Cu—N1—C2	111.2 (3)	S—C15—N5	178.9 (4)

Data reduction: *NRCVAX DATRD2* (Gabe *et al.*, 1989). Program(s) used to solve structure: *NRCVAX SOLVER*. Program(s) used to refine structure: *NRCVAX LSTSQ*. Molecular graphics: *ORTEPII* (Johnson, 1976) in *NRCVAX*. Software used to prepare material for publication: *NRCVAX TABLES*.

The authors thank the National Science Council, Republic of China, for support under grants NSC87-2112-M007-009 and NSC87-2113-M007-041.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: OA1047). Services for accessing these data are described at the back of the journal.

References

- Ahrland, S., Chatt, J. & Davics, N. R. (1958). *Q. Rev. Chem. Soc.* **12**, 265–269.
- Alcock, N. W., Herron, N. & Moore, P. (1978). *J. Chem. Soc. Dalton Trans.* pp. 1282–1288.
- Barefield, E. K. & Wagner, F. (1973). *Inorg. Chem.* **12**, 2435–2436.
- Burmeister, J. L. (1990). *Coord. Chem. Rev.* **105**, 77–105.
- D'Aniello, M. J., Mocella, M. T., Wagner, F., Barefield, E. K. & Paul, I. C. (1975). *J. Am. Chem. Soc.* **97**, 192–194.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Hodges, K. D., Wollmann, R. G., Kessel, S. L., Hendrickson, D. N., Van Deveer, D. G. & Barefield, E. K. (1979). *J. Am. Chem. Soc.* **101**, 906–917.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Kato, M. & Ito, T. (1985). *Inorg. Chem.* **24**, 509–514.
- Larson, A. C. (1970). *Crystallographic Computing*, p. 293. Copenhagen: Munksgaard.
- Lee, T.-J., Lee, T.-Y., Hong, C.-Y., Wu, D.-T. & Chung, C.-S. (1986). *Acta Cryst.* **C42**, 999–1001.
- Lu, T.-H., Tahirov, T.-H., Liu, Y.-L., Chung, C.-S., Huang, C.-C. & Hong, S.-H. (1996). *Acta Cryst.* **C52**, 1093–1095.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Pearson, R. G. (1963). *J. Am. Chem. Soc.* **85**, 3533–3539.
- Tahirov, T.-H., Lu, T.-H., Lan, W.-J. & Chung, C.-S. (1993). *Acta Cryst.* **C49**, 1908–1910.
- Tahirov, T.-H., Lu, T.-H., Luh, H., Lai, C.-Y. & Chung, C.-S. (1995). *Acta Cryst.* **C51**, 846–849.

Acta Cryst. (1998). **C54**, 1072–1074

Bis[*N*-(2-bromo-4-methylphenyl)-naphthaldiminato]copper(II)

YALÇIN ELERMAN,^a AYHAN ELMALI^a AND SÜHEYLA ÖZBEY^b

^aDepartment of Physics Engineering, Faculty of Sciences, University of Ankara, 06100 Besevler, Ankara, Turkey, and ^bDepartment of Physics Engineering, Hacettepe University, 06532 Beytepe, Ankara, Turkey. E-mail: elmali@science.ankara.edu.tr

(Received 6 November 1997; accepted 24 February 1998)


Abstract

The title compound, bis[1-(2-bromo-4-methylphenyl)iminomethyl]-2-naphtholato-*N,O*]copper(II), [Cu(C₁₈H₁₃-BrNO)₂], has crystallographic inversion symmetry. The Cu^{II} ion shows a pseudo-square-planar coordination. The Cu—N and Cu—O distances are 1.990 (4) and 1.880 (4) Å, respectively. The Cu^{II} ion is in an approximate octahedral environment if bromine is included in

the description, with a non-bonded $\text{Cu}\cdots\text{Br}$ distance of 3.296 (1) Å.

Comment

Because of their preparative accessibility and structural variability, a great number of Schiff base complexes of copper have been the subject of extensive studies (Garnovskii *et al.*, 1993). Despite this, there are few reported X-ray crystal structures of naphthalenic Schiff base-copper compounds (Acevedo-Arauz *et al.*, 1992). We report here the results of the reaction of copper(II) with the ligand *N*-(2-bromo-4-methylphenyl)naphthalimine, which produced the title compound, (I).



The title molecule (Fig. 1) has crystallographic inversion symmetry. The coordination of the Cu^{II} ion is necessarily planar by symmetry. A strictly planar or slightly

distorted *trans* configuration is characteristic for transition metal complexes of Cu^{II} with a CuN_2O_2 coordination sphere (Garnovskii *et al.*, 1993). In the title compound, the Cu^{II} ion is bonded to the O and N donor atoms of the two ligand molecules in a *trans* arrangement. The $\text{Cu}-\text{N}$ and $\text{Cu}-\text{O}$ distances are 1.990 (4) and 1.880 (4) Å, respectively. The $\text{O}-\text{Cu}-\text{N}$ angle is 90.2 (2)°. The $\text{Cu}-\text{N}$ distances range from 1.97 to 2.01 Å, while the $\text{Cu}-\text{O}$ distances have values between 1.87 and 1.92 Å in (1,2-naphthalaldiminato)copper complexes (Clark *et al.*, 1975, 1977; Martin & Waters, 1973; Shnulin *et al.*, 1978; Acevedo-Arauz *et al.*, 1992). The $\text{Cu}-\text{N}$ and $\text{Cu}-\text{O}$ values in (I) agree with those in other naphthalenic complexes. The Cu^{II} ion is in a pseudo-octahedral environment if bromine is included in the description. The non-bonded $\text{Cu}\cdots\text{Br}$ distance is 3.296 (1) Å. The title molecule is not planar; the Cu^{II} ion lies 0.238 (2) Å below the plane formed by the Cu1, O1, C1, C10, C11 and N1 atoms. This plane is inclined at an angle of 18.6 (1)° with respect to the coordination plane. The two Schiff base moieties, A (O1, C1-C11) and B (Br, N1, O2, C12-C18), are inclined at an angle of 43.0 (1)° with respect to one another, which is mainly the twist about N1-C12 [54.3 (7)°]. The N atom shows sp^2 hybridization, with a bond-angle sum of 359.5 (4)°.

Discrete monomeric molecules are held together in the crystal by van der Waals interactions. The minimum distances between two non-H atoms in neigh-




Fig. 1. The molecular structure of (I) showing 50% probability displacement ellipsoids (ORTEP II; Johnson, 1976).

bonding molecules are C8···O1, C8···C13, C9···C13 and O1···C7 with the values of 3.400 (7), 3.466 (9), 3.483 (9) and 3.518 (8) Å, respectively.

Experimental

For the preparation of (I), solutions of 2-hydroxynaphthalene-1-carbaldehyde (0.50 mmol) and 2-bromo-4-methylaniline (0.50 mmol) in 75 ml acetonitrile, and [Cu(O₂CCH₃)₂].4H₂O (0.25 mmol) in 40 ml methanol were heated to boiling, mixed and refluxed for 3 d. Crystals of the product were obtained on cooling.

Crystal data

[Cu(C₁₈H₁₃BrNO)₂]

$M_r = 741.95$

Monoclinic

$P2_1/c$

$a = 9.255 (1) \text{ \AA}$

$b = 12.524 (2) \text{ \AA}$

$c = 13.538 (2) \text{ \AA}$

$\beta = 104.32 (1)^\circ$

$V = 1520.4 (4) \text{ \AA}^3$

$Z = 2$

$D_x = 1.621 \text{ Mg m}^{-3}$

D_m not measured

Data collection

Enraf–Nonius CAD-4 diffractometer

$\theta/2\theta$ scans

Absorption correction: ψ scan (MolEN; Fair, 1990)

$T_{\min} = 0.496$, $T_{\max} = 0.713$

2762 measured reflections

2760 independent reflections

Refinement

Refinement on F^2

$R[F^2 > 2\sigma(F^2)] = 0.056$

$wR(F^2) = 0.097$

$S = 1.208$

2760 reflections

196 parameters

H-atom parameters were constrained to parent sites

Mo $K\alpha$ radiation

$\lambda = 0.71069 \text{ \AA}$

Cell parameters from 20 reflections

$\theta = 4.12\text{--}14.43^\circ$

$\mu = 3.383 \text{ mm}^{-1}$

$T = 293 (2) \text{ K}$

Prism

$0.28 \times 0.18 \times 0.10 \text{ mm}$

Purple–red

1866 reflections with

$I > 2\sigma(I)$

$R_{\text{int}} = 0.018$

$\theta_{\text{max}} = 26.14^\circ$

$h = 0 \rightarrow 11$

$k = 0 \rightarrow 15$

$l = -16 \rightarrow 16$

3 standard reflections

frequency: 120 min

intensity decay: 3.2%

$w = 1/[\sigma^2(F_o^2) + 3.7194P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 0.491 \text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.326 \text{ e \AA}^{-3}$

Extinction correction: none

Scattering factors from *International Tables for Crystallography* (Vol. C)

The title structure was solved by direct phase determination. The parameters of the complete structure could be refined by full-matrix anisotropic least squares. All phenyl rings were refined without constraints. Distance and angle values in the rings show no significant differences from those of an ideal benzene ring. All H-atom positions were calculated using a riding model with fixed U_{iso} values in all refinements.

Data collection: CAD-4 EXPRESS (Enraf–Nonius, 1994). Cell refinement: MolEN (Fair, 1990). Data reduction: MolEN. Program(s) used to solve structure: SHELXS86 (Sheldrick, 1990). Program(s) used to refine structure: SHELXL93 (Sheldrick, 1993). Molecular graphics: ORTEPII (Johnson, 1976). Software used to prepare material for publication: SHELXL93.

The authors wish to acknowledge the purchase of a CAD-4 diffractometer under grant DPT/TBAG1 of the Scientific and Technical Council of Turkey.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: AB1532). Services for accessing these data are described at the back of the journal.

References

- Acevedo-Arauz, E., Fernández-G., J. M., Rosales-Hoz, M. & Toscano, R. A. (1992). *Acta Cryst.* **C48**, 115–120.
- Clark, G. R., Waters, J. M. & Waters, T. N. (1975). *J. Inorg. Nucl. Chem.* **37**, 2455–2458.
- Clark, G. R., Waters, J. M. & Williams, G. J. (1977). *J. Inorg. Nucl. Chem.* **39**, 1971–1975.
- Enraf–Nonius (1994). CAD-4 EXPRESS. Version 5.1/1.2. Enraf–Nonius, Delft, The Netherlands.
- Fair, C. K. (1990). MolEN. *An Interactive Intelligent System for Crystal Structure Analysis*. Enraf–Nonius, Delft, The Netherlands.
- Garnovskii, A. D., Nivorozhkin, A. L. & Minkin, V. I. (1993). *Coord. Chem. Rev.* **126**, 1–69.
- Johnson, C. K. (1976). ORTEPII. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Martin, D. W. & Waters, T. N. (1973). *J. Chem. Soc. Dalton Trans.* pp. 2440–2443.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1993). SHELXL93. *Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Shnulin, A. N., Mamedov, Kh. S. & Stuchkov, Yu. Y. (1978). *Zh. Strukt. Khim.* **19**, 686–688.

Table 1. Selected geometric parameters (Å, °)

Cu1—O1 ¹	1.880 (4)	N1—C11	1.295 (6)
Cu1—N1 ¹	1.990 (4)	N1—C12	1.434 (6)
Br1—C18	1.900 (6)	O1—C1	1.293 (6)
O1—Cu1—N1 ¹	90.2 (2)	O1—C1—C2	116.0 (5)
C11—N1—C12	118.2 (4)	N1—C11—C10	126.2 (5)
C11—N1—Cu1	122.3 (4)	C13—C12—N1	122.1 (5)
C12—N1—Cu1	119.0 (3)	C18—C12—N1	120.9 (5)
C1—O1—Cu1	124.8 (3)	C17—C18—Br1	118.9 (5)
O1—C1—C10	124.8 (5)	C12—C18—Br1	119.3 (4)

Symmetry code: (i) $-x, -y, -z$.