

N11—C51 ¹	1.52 (2)	C11—O4	1.401 (3)
N2—C2	1.481 (4)	C11—O11	1.32 (1)
N2—C3	1.488 (4)	C11—O21	1.49 (1)
C1—C2	1.484 (6)	C11—O31	1.36 (2)
N1 ¹ —Cu1—N1	180.0	N2—C2—C1	108.0 (3)
N11 ¹ —Cu1—N11	180.0	N2—C3—C6	111.7 (3)
N1 ¹ —Cu1—N2	94.7 (1)	N2—C3—C4	111.0 (3)
N1—Cu1—N2	85.3 (1)	C6—C3—C4	111.8 (3)
N2 ¹ —Cu1—N11	91.7 (3)	C7—C4—C3	114.9 (4)
N2—Cu1—N11	88.3 (3)	C7—C4—C51	88.6 (6)
N2 ¹ —Cu1—N2	180.0	C3—C4—C51	117.7 (5)
N1—Cu1—O1	88.0 (2)	C7—C4—C5	117.7 (5)
N11—Cu1—O11	85.5 (5)	C3—C4—C5	112.4 (3)
N2—Cu1—O1	82.9 (2)	N1 ¹ —C5—C4	112.8 (4)
N2—Cu1—O11	81.8 (5)	N11 ¹ —C51—C4	106.3 (10)
O1 ¹ —Cu1—O1	180.0	O1—C11—O2	106.1 (3)
O11 ¹ —Cu1—O11	180.0	O1—C11—O3	109.0 (5)
C5 ¹ —N1—C1	110.0 (4)	O1—C11—O4	114.3 (3)
C5 ¹ —N1—Cu1	118.1 (3)	O2—C11—O3	107.9 (4)
C1—N1—Cu1	106.4 (3)	O2—C11—O4	110.4 (3)
C5 ¹ —N11—C1	103.8 (8)	O3—C11—O4	108.9 (3)
C5 ¹ —N11—Cu1	102.5 (7)	O11—C11—O21	106.2 (10)
C1—N11—Cu1	101.3 (5)	O11—C11—O31	111 (2)
C2—N2—C3	113.9 (3)	O11—C11—O4	108.7 (10)
C2—N2—Cu1	105.4 (2)	O21—C11—O31	114.9 (14)
C3—N2—Cu1	116.3 (2)	O21—C11—O4	104.7 (7)
C2—C1—N1	104.6 (3)	O31—C11—O4	111.4 (11)
C2—C1—N11	119.0 (4)		

Symmetry code: (i) $-x, -y, -z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989). Cell refinement: *CAD-4 Software*. Data reduction: *NRCVAX* (Gabe *et al.*, 1989). Program(s) used to solve structure: *SHELXS86* (Sheldrick, 1990). Program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993). Molecular graphics: *NRCVAX*. Software used to prepare material for publication: *SHELXL93*.

The authors thank the National Science Council of Taiwan for support under grants NSC86-2112-M007-028, NSC86-2811-M007-0041, NSC87-2112-M007-009 and NSC87-2113-M007-041.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: KA1262). Services for accessing these data are described at the back of the journal.

References

- Boeyens, J. C. A. & Dobson, S. M. (1987). *Stereochemical and Stereophysical Behavior of Macrocycles*, Vol. 2, pp. 48–70. New York: Elsevier.
- Bosnich, B., Poon, C. K. & Tobe, M. L. (1965). *Inorg. Chem.* **4**, 1102–1108.
- Cabbiness, D. K. & Margerum, D. W. (1969). *J. Am. Chem. Soc.* **91**, 6540–6541.
- Enraf–Nonius (1989). *CAD-4 Software*. Version 5.0. Enraf–Nonius, Delft, The Netherlands.
- Ferguson, G., Restivo, R. R. & Hay, R. W. (1979). *Acta Cryst.* **B35**, 159–162.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Hay, R. W., Piplani, D. P. & Jeragh, B. (1977). *J. Chem. Soc. Dalton Trans.* pp. 1951–1956.
- Hitchman, M. A. & Deeth, R. J. (1986). *Inorg. Chem.* **25**, 1225–1233.
- Krajewski, J. W., Urbanczyk-Lipkowska, Z. & Gluzinski, P. (1978). *Pol. J. Chem.* **52**, 1513–1523.
- Lee, T.-J., Lu, T.-H., Chung, C.-S. & Lee, T.-Y. (1984). *Acta Cryst.* **C40**, 70–72.

- Liang, B.-F. & Chung, C.-S. (1980). *Inorg. Chem.* **19**, 1867–1871.
- Lu, T.-H., Chung, C.-S. & Ashida, T. (1991). *J. Chin. Chem. Soc. (Taipei)*, **38**, 147–153.
- Murray-Rust, P. & Murray-Rust, J. (1979). *Acta Cryst.* **B35**, 1704–1706.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
- Sheldrick, G. M. (1993). *SHELXL 93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Tasker, P. A. & Sklar, L. (1975). *J. Cryst. Mol. Struct.* **5**, 329–344.
- Warner, L. G. & Busch, D. H. (1969). *J. Am. Chem. Soc.* **91**, 4092–4101.

Acta Cryst. (1998). **C54**, 1242–1244

trans-Diaquabis(nicotinamide-*N*¹)bis(4-nitrobenzoato-*O*)cobalt(II)

TUNCER HÖKELEK^{a*} AND HACALI NECEFOĞLU^b

^aHacettepe University, Department of Physics, 06532 Beytepe, Ankara, Turkey, and ^bKafkas University, Department of Chemistry, Kars, Turkey. E-mail: merzifon@eti.cc.hun.edu.tr

(Received 15 December 1997; accepted 24 March 1998)

Abstract

The title cobalt complex, [Co(C₇H₄NO₄)₂(C₆H₆N₂O)₂·(H₂O)₂], is monomeric and centrosymmetric, and contains two water molecules, two 4-nitrobenzoate and two nicotinamide (NA) ligands, all acting as monodentate ligands. The four nearest O atoms around the Co atom form a slightly distorted square-planar arrangement, with a distorted octahedral coordination completed by the pyridine N atoms of the NA ligand, at a distance of 2.134 (2) Å. The water molecules are hydrogen bonded to the carboxyl O atoms [O...O 2.634 (5) Å].

Comment

There are only a few examples of structure determinations of Co^{II} complexes with nicotinic and/or benzoic acid derivatives as ligands, *e.g.* [Co(C₇H₅O₃)₂(DENA)₂(H₂O)₂] (where DENA is diethylnicotinamide, C₁₀H₁₄N₂O; Hökelek & Necefoğlu, 1997), [Co(CH₃CO₂)₂(DENA)₂(H₂O)₂] (Mikelashvili, 1982), [Co(H₂O)₄(*p*-O₂NC₆H₄COO)₂] (Nadzhafov *et al.*, 1981) and [Co(H₂O)₄(*p*-H₂NC₆H₄COO)₂] (Amiraslanov *et al.*, 1979). The structure–function–coordination relationships of the arylcarboxylate ion in Co^{II} complexes of benzoic acid derivatives change depending on the





Fig. 1. An ORTEP (Johnson, 1976) drawing of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

nature and position of the substituted groups in the phenyl ring, the nature of the additional ligand molecule or solvent, and the pH and temperature of synthesis (Shnulin *et al.*, 1981; Antsyshkina *et al.*, 1980; Adiwidjaja *et al.*, 1978). When pyridine and its derivatives are used instead of water molecules, the structure is completely different (Catterick *et al.*, 1974). In the $[\text{Co}(p\text{-O}_2\text{NC}_6\text{H}_4\text{COO})_2(\text{H}_2\text{O})_4]$ complex, Nadzhafov *et al.* (1981) found that the Co atom is situated on a centre of symmetry, and is surrounded by six O atoms forming a slightly distorted octahedron. Four positions are occupied by O atoms from the water molecules, and the other two by the O atoms of the carboxyl groups of the *p*-nitrobenzoate anions. The structure determination of the title molecule, (I), was undertaken to determine the ligand properties of the nicotinamide (NA) and benzoate ligands, and to compare the coordination geometries when the NA ligands are substituted for water in the Nadzhafov *et al.* (1981) complex. The

the NA ligand complete a Jahn–Teller-distorted octahedron at distances of 2.134 (2) Å (Fig. 1). There are hydrogen bonds between the carboxyl O4 atoms and water O1 atoms [O4...O1 2.634 (5) Å]. Similar hydrogen bonds are observed in $[\text{Co}(\text{C}_7\text{H}_5\text{O}_3)_2(\text{DENA})_2(\text{H}_2\text{O})_2]$ [O...O 2.687 (5) Å; Hökelek & Necefoğlu, 1997], $[\text{Co}(\text{H}_2\text{O})_4(p\text{-O}_2\text{NC}_6\text{H}_4\text{COO})_2]$ [O...O 2.59 Å; Nadzhafov *et al.*, 1981] and $[\text{Co}(\text{H}_2\text{O})_4(p\text{-H}_2\text{NC}_6\text{H}_4\text{COO})_2]$ [O...O 2.592 (3) Å; Amiraslanov *et al.*, 1979]. The Co atom lies 0.414 (3) Å out of the C7,O3,O4 least-squares plane. In the carboxylate group, the C7—O3 and C7—O4 bond lengths (Table 1) are the same as in the $[\text{Co}(\text{C}_7\text{H}_5\text{O}_3)_2(\text{DENA})_2(\text{H}_2\text{O})_2]$ complex [1.251 (6) and 1.254 (7) Å; Hökelek & Necefoğlu, 1997], but they are different from the corresponding ones in the $[\text{Co}(\text{H}_2\text{O})_4(p\text{-O}_2\text{NC}_6\text{H}_4\text{COO})_2]$ complex [1.292 (6) and 1.246 (8) Å; Nadzhafov *et al.*, 1981]. The dihedral angle between the planar carboxyl group (C7,O3,O4) and the phenyl ring (C8—C13) is 23.7 (3)°. The corresponding angle is 2.2 (6)° in $[\text{Co}(\text{C}_7\text{H}_5\text{O}_3)_2(\text{DENA})_2(\text{H}_2\text{O})_2]$ (Hökelek & Necefoğlu, 1997).



title compound is a monomeric complex, with Co on a centre of symmetry. All the ligands are monodentate, and the O atoms of each water molecule and benzoate group form a slightly distorted square-planar coordination around the Co atom. The pyridine N atoms of

Experimental

The title compound was prepared from the reaction of $\text{CoSO}_4 \cdot 7\text{H}_2\text{O}$ (0.01 mol) and NA (0.02 mol) in sodium *p*-nitrobenzoate solution (0.02 mol). The mixture was filtered and set aside to crystallize at ambient temperature for a few days. Pink crystals suitable for X-ray diffraction were obtained.

Crystal data

$[\text{Co}(\text{C}_7\text{H}_4\text{NO}_4)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2 \cdot (\text{H}_2\text{O})_2]$
 $M_r = 671.442$
 Triclinic
 $P\bar{1}$

Mo $K\alpha$ radiation
 $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 25 reflections
 $\theta = 10\text{--}18^\circ$

$a = 7.5815(8) \text{ \AA}$
 $b = 9.9861(9) \text{ \AA}$
 $c = 10.2413(12) \text{ \AA}$
 $\alpha = 78.417(9)^\circ$
 $\beta = 88.559(9)^\circ$
 $\gamma = 71.491(8)^\circ$
 $V = 719.66(14) \text{ \AA}^3$
 $Z = 1$
 $D_x = 1.5493 \text{ Mg m}^{-3}$
 D_m not measured

$\mu = 0.671 \text{ mm}^{-1}$
 $T = 298 \text{ K}$
 Rod
 $0.30 \times 0.25 \times 0.25 \text{ mm}$
 Pink

son, 1976). Software used to prepare material for publication: *MolEN*.

The authors wish to acknowledge the purchase of a CAD-4 diffractometer under Grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey, and are grateful to the Kafkas University Research Fund for the financial support of this work.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: OS1013). Services for accessing these data are described at the back of the journal.

Data collection

Enraf–Nonius CAD-4 diffractometer
 $\omega/2\theta$ scans
 Absorption correction: empirical via ψ scans (*MolEN*; Fair, 1990)
 $T_{\min} = 0.828$, $T_{\max} = 0.846$
 3086 measured reflections
 2913 independent reflections
 2842 reflections with $F > 0$

$R_{\text{int}} = 0.064$
 $\theta_{\text{max}} = 26.3^\circ$
 $h = 0 \rightarrow 9$
 $k = -11 \rightarrow 12$
 $l = -12 \rightarrow 12$
 3 standard reflections every 250 reflections
 frequency: 120 min
 intensity decay: 1%

Refinement

Refinement on F
 $R = 0.037$
 $wR = 0.046$
 $S = 1.03$
 2842 reflections
 205 parameters
 H atoms riding
 $w = 1/[\sigma(F^2) + (0.02F)^2 + 1]$

$(\Delta/\sigma)_{\text{max}} = 0.01$
 $\Delta\rho_{\text{max}} = 0.50 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$
 Extinction correction: none
 Scattering factors from *International Tables for X-ray Crystallography* (Vol. IV)

Table 1. Selected geometric parameters (\AA , $^\circ$)

Co—O1	2.136 (1)	O6—N3	1.199 (3)
Co—O3	2.091 (2)	N1—C1	1.323 (3)
Co—N2	2.134 (2)	N2—C3	1.337 (3)
O2—C1	1.234 (2)	N2—C4	1.335 (2)
O3—C7	1.254 (2)	N3—C11	1.475 (4)
O4—C7	1.251 (2)	C7—C8	1.512 (3)
O5—N3	1.195 (3)		
O1—Co—O3	92.67 (6)	O2—C1—N1	122.2 (2)
O1—Co—N2	92.88 (6)	O2—C1—C2	119.8 (2)
O3—Co—N2	90.19 (6)	N1—C1—C2	118.0 (2)
Co—O3—C7	126.8 (1)	N3—C11—C10	118.5 (2)
Co—N2—C3	119.1 (1)	N3—C11—C12	118.5 (2)
Co—N2—C4	123.0 (4)	N2—C3—C2	123.4 (2)
C3—N2—C4	118.0 (2)	N2—C4—C5	122.4 (2)
O5—N3—O6	122.9 (3)	O3—C7—O4	126.0 (2)
O5—N3—C11	118.8 (2)	O3—C7—C8	117.7 (2)
O6—N3—C11	118.2 (2)	O4—C7—C8	116.3 (2)
O1—Co—O3—C7	-18.3 (2)	O3—Co—N2—C3	34.7 (2)
N2—Co—O3—C7	74.6 (2)	O3—Co—N2—C4	-144.7 (2)
O1—Co—N2—C3	127.4 (2)	Co—O3—C7—O4	14.3 (3)
O1—Co—N2—C4	-52.0 (2)	Co—O3—C7—C8	-162.2 (1)

The title structure was solved by the Patterson method. The H-atom positions were calculated geometrically, with $U_{\text{eq}}(\text{H}) = 1.3U_{\text{eq}}$ of the parent non-H atom. A riding model was used in the refinement.

Data collection: *MolEN* (Fair, 1990). Cell refinement: *MolEN*. Data reduction: *MolEN*. Program(s) used to solve structure: *SHELXS86* (Sheldrick, 1990). Program(s) used to refine structure: *MolEN*. Molecular graphics: *ORTEPII* (John-

References

- Adiwidjaja, G., Rossmanith, E. & Kupperts, H. (1978). *Acta Cryst.* **B34**, 3079–3081.
 Amiraslano, I. R., Mamedov, Kh. S., Movsumov, E. M., Musaev, F. N. & Nadzhafov, G. N. (1979). *Zh. Strukt. Khim.* **20**, 1075–1080.
 Antsyshkina, A. S., Chiragov, F. M. & Poray-Koshits, M. A. (1980). *Koord. Khim.* **15**, 1098–1103.
 Catterick, J., Hursthouse, M. B., New, D. B. & Thornton, P. (1974). *J. Chem. Soc. Chem. Commun.* pp. 843–844.
 Fair, C. K. (1990). *MolEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf–Nonius, Delft, The Netherlands.
 Hökelek, T. & Necefoglu, H. (1997). *Acta Cryst.* **C53**, 187–189.
 Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
 Mikelashvili, Z. V. (1982). Dissertation, Tbilisi State University, Georgia.
 Nadzhafov, G. N., Shnulin, A. N. & Mamedov, Kh. S. (1981). *Zh. Strukt. Khim.* **22**, 124–128.
 Sheldrick, G. M. (1990). *Acta Cryst.* **A46**, 467–473.
 Shnulin, A. N., Nadzhafov, G. N., Amiraslano, I. R., Usabaliev, B. T. & Mamedov, Kh. S. (1981). *Koord. Khim.* **7**, 1409–1416.

Acta Cryst. (1998). **C54**, 1244–1247

Bis[μ -methylenebis(dicyclohexylphosphine)-*P:P'*]bis[chloropalladium(I)](*Pd—Pd*) Diacetonitrile Solvate

MIHO HASHIOKA, KEN SAKAI AND TARO TSUBOMURA*

Department of Industrial Chemistry, Seikei University, Kichijoji-Kitamachi, Musashino, Tokyo 180, Japan. E-mail: tsubo@ch.seikei.ac.jp

(Received 28 November 1997; accepted 24 March 1998)

Abstract

A dinuclear palladium(I) complex bridged by two diphosphine ligands, bis[μ -methylenebis(dicyclohexylphosphine)-*P:P'*]bis[chloropalladium(I)](*Pd—Pd*) diacetonitrile solvate, $[\text{Pd}_2\text{Cl}_2(\text{C}_{25}\text{H}_{46}\text{P}_2)_2]_2 \cdot 2\text{C}_2\text{H}_3\text{N}$, has been prepared, and the crystal structure determined.